REVIEW

The role of sclerostin and dickkopf-1 in oral tissues – A review from the perspective of the dental disciplines [version 1; peer review: 2 approved]

Mohammad Samiei¹⁻³, Klara Janjić^{id}^{2,3}, Barbara Cviki^{2,3}, Andreas Moritz^{2,3}, Hermann Agis^{id}^{2,3}

¹Department of Endodontics, Faculty of Dentistry, Tabriz University of Medical Sciences, Tabriz, Iran

²Department of Conservative Dentistry and Periodontology, University Clinic of Dentistry, Medical University of Vienna, Vienna, 1090, Austria

³Austrian Cluster for Tissue Regeneration, Vienna, 1200, Austria

V1 First published: 30 Jan 2019, 8:128
<https://doi.org/10.12688/f1000research.17801.1>
 Latest published: 30 Jan 2019, 8:128
<https://doi.org/10.12688/f1000research.17801.1>

Abstract

Wnt signaling is of high relevance in the development, homeostasis, and regeneration of oral tissues. Therefore, Wnt signaling is considered to be a potential target for therapeutic strategies. The action of Wnt is tightly controlled by the inhibitors sclerostin (SOST) and Dickkopf (DKK)-1. Given the impact of SOST and DKK-1 in hard tissue formation, related diseases and healing, it is of high relevance to understand their role in oral tissues. The clinical relevance of this knowledge is further underlined by systemic and local approaches which are currently in development for treating a variety of diseases such as osteoporosis and inflammatory hard tissue resorption. In this narrative review, we summarize the current knowledge and understanding on the Wnt signaling inhibitors SOST and DKK-1, and their role in physiology, pathology, and regeneration in oral tissues. We present this role from the perspective of the different specialties in dentistry, including endodontics, orthodontics, periodontics, and oral surgery.

Keywords

Sclerostin, Dickkopf-1, Wnt Pathway, Regeneration, Periodontology, Endodontology

Open Peer Review

Approval Status

	1	2
version 1
30 Jan 2019	view	view

1. **Francisco Javier Rodríguez-Lozano** ^{id},

University of Murcia, Murcia, Spain

2. **Qiming Jin**, University of Michigan, Ann Arbor, USA

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Hermann Agis (hermann.agis@meduniwien.ac.at)

Author roles: **Samiei M:** Conceptualization, Investigation, Methodology, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Janjić K:** Conceptualization, Investigation, Methodology, Project Administration, Supervision, Validation, Writing – Review & Editing; **Cvikl B:** Conceptualization, Funding Acquisition, Investigation, Methodology, Resources, Validation, Writing – Review & Editing; **Moritz A:** Conceptualization, Investigation, Methodology, Resources, Supervision, Validation, Writing – Review & Editing; **Agis H:** Conceptualization, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: We thank the European Society of Endodontology (ESE) for financial support of our research [ESE research grant 2015]. The authors deny any conflict of interest.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 Samiei M *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Samiei M, Janjić K, Cvikl B *et al.* **The role of sclerostin and dickkopf-1 in oral tissues – A review from the perspective of the dental disciplines [version 1; peer review: 2 approved]** F1000Research 2019, **8**:128 <https://doi.org/10.12688/f1000research.17801.1>

First published: 30 Jan 2019, **8**:128 <https://doi.org/10.12688/f1000research.17801.1>

Introduction

Research and development of novel treatment strategies for regenerative dentistry has become crucial to improve oral health due to the growing numbers of dental and maxillofacial problems which require treatment. Therefore, cell-therapy modalities, biologicals, and gene therapy approaches are in development with the aim to provide novel tools for the clinical demands (Dissanayaka *et al.*, 2014; Fretwurst *et al.*, 2018; Itoh *et al.*, 2018; Kaigler *et al.*, 2015; Nevins *et al.*, 2005; Plonka *et al.*, 2017; Taut *et al.*, 2013). This research needs to be guided by an appropriate understanding of the cell biological mechanisms underlying pathological changes and regeneration in the oral tissue, including the periodontium and the dental tissue. Due to the major role of Wnt signaling and the respective regulators in development and healing, research in regenerative medicine and dentistry has evaluated the feasibility of targeting the pathway for therapeutic approaches (Florio *et al.*, 2016; Heiland *et al.*, 2010; Taut *et al.*, 2013; Yu *et al.*, 2018).

The Wnts comprise a family of at least 19 lipid-modified glycoproteins and short-range ligands. Wnts can activate canonical and non-canonical pathways of signaling in the cells (Mohammed *et al.*, 2016; Tang *et al.*, 2009; Yang *et al.*, 2016). The canonical Wnt path involves the interaction of Wnt with frizzled (Frz) and low density lipoprotein receptor-related protein 5/6 (LRP5/6). Thereby β -catenin accumulation is induced leading to the translocation of β -catenin into the nucleus where target genes are activated (Mohammed *et al.*, 2016; Tang *et al.*, 2009; Yang *et al.*, 2016) (Figure 1).

Wnt signaling regulates various cellular functions which include cell migration, proliferation, differentiation, apoptosis, and morphogenesis. Given this broad involvement of Wnt signaling it is not surprising that it has a key role in organ development, regeneration and homeostasis of tissues (Logan & Nusse, 2004; Sarkar & Sharpe, 2000; Seo *et al.*, 2012). In the maxillofacial region, Wnt signaling modulates morphological patterns in tooth development including teeth number, shape, size, and positioning (Han *et al.*, 2011; Kim *et al.*, 2013; Kratochwil *et al.*, 2002; Liu *et al.*, 2008; Sarkar *et al.*, 2000; Wang *et al.*, 2009;

Zhang *et al.*, 2013). Maturation of dental mesenchyme into odontoblasts and cementoblasts is also modulated by Wnt signaling which highlights the relevance to dentistry (Janjić *et al.*, 2018; Zhang *et al.*, 2013). Given this dominant role of Wnts, Wnt signaling needs to be tightly controlled involving inhibitors.

Sclerostin (SOST) and dickkopf-1 (DKK-1) are the primary inhibitors controlling the Wnt signaling pathways (Figure 1). These inhibitors can directly bind to LRP5/6 and inhibit the activation of LRP5- and LRP6- related signaling. Dkk-1 binds to a larger region on LRP5 and LRP6 extracellular surface and thereby can inhibit binding with other Wnts than SOST. (Ahn *et al.*, 2011; Bourhis *et al.*, 2010; Bourhis *et al.*, 2011; Cheng *et al.*, 2011; Florio *et al.*, 2016; Li *et al.*, 2005; van Bezooijen *et al.*, 2004) Thereby SOST and DKK-1 play critical roles in the formation of hard tissue and associated diseases. SOST and DKK-1 are consequently considered potential therapeutic targets for regenerative approaches (Janjić *et al.*, 2018; Li *et al.*, 2005; Semenov *et al.*, 2005; Taut *et al.*, 2013).

The glycoprotein SOST is mainly secreted by osteocytes, the mechanosensor cells of the bone, and has a major effect on bone and dental tissue (Kuchler *et al.*, 2014; Pflanz *et al.*, 2017; Saag *et al.*, 2017; Taut *et al.*, 2013; Winkler *et al.*, 2003; Yu *et al.*, 2018). In recent years, SOST production in oral tissues has gained more attention and led to *in vitro* and *in vivo* studies on the regulation of SOST (Pflanz *et al.*, 2017; Taut *et al.*, 2013; Yu *et al.*, 2018). SOST production in mineralizing periodontal ligament cells is increased; this observation suggests that targeting of SOST might support periodontal regeneration. SOST has also been demonstrated to have a role in the dimension of the periodontal ligament (Kuchler *et al.*, 2014). This further supports the hypothesis of a key role of SOST in oral tissues. DKK-1, another antagonist of Wnt signaling, modulates the dimension of oral tissues, including the periodontium and teeth (Bao *et al.*, 2013; Jäger *et al.*, 2010; MacDonald *et al.*, 2007).

According to the findings of 10 years analysis of publications regarding SOST and DKK-1 in medical fields (Figure 2), it becomes evident that there is a growing set of literature on SOST

Figure 1. Inhibition of Wnt signaling by sclerostin and dickkopf-1. Scheme of the Wnt signaling pathway (A) which is inhibited by sclerostin (SOST, B) and dickkopf-1 (DKK-1, C). Adopted from (Yorgan & Schinke, 2014) and modified.

Figure 2. The numbers of publication on sclerostin and dickkopf-1 increased rapidly over the past 10 years. The number of publications on sclerostin (SOST, **A**) and dickkopf-1 (DKK-1, **C**) per year as found in Pubmed.org (Search terms SOST or DKK-1 with and without Medicine or Dentistry). The total number of publications on SOST (**B**) and DKK-1 (**D**) in Dentistry and the respective specialties Endodontics, Periodontics, Orthodontics, and Oral Surgery as found in Pubmed.org (Search terms SOST with and without Dentistry, Endodontics, Periodontics, Orthodontics, or Oral Surgery).

and DKK-1. Within dentistry the majority of publications on SOST and DKK-1 is from the field of oral surgery. This review aims to provide a synopsis of the existing knowledge on the Wnt signaling inhibitors SOST and DKK-1 from endodontics, orthodontics, periodontics, and oral surgery perspectives.

Endodontics perspective

Research and developments in endodontics lead to the continuous improvement of the field, as the introduction of new techniques and biomaterials drive profound progress in endodontic clinical practice. In particular regenerative endodontics is an exciting dental specialty. Treatment protocols of infected root canals in immature teeth were established which allow continued apical closure and root maturation (Chueh & Huang, 2006; Huang, 2008; Huang *et al.*, 2008). In these regenerative endodontic procedures, after treatment with ethylenediaminetetraacetic acid (EDTA) signaling factors embedded in the dentin matrix such

as TGF- β are exposed and released into the canal space (Galler *et al.*, 2015). These factors can guide the migration of dental pulp stem cells and support differentiation into odontoblast-like cells, leading to the production of mineralized matrix also termed reparative dentin (Nakashima *et al.*, 1994; Rutherford *et al.*, 1993; Sloan & Smith, 1999; Tziafas *et al.*, 1998). Interestingly TGF- β has the capacity to induce SOST expression (Gruber *et al.*, 2017; Manokawinchoke *et al.*, 2015); therefore, it is possible that dentin conditioning can interfere with the regulation system of Wnt signaling (Gruber *et al.*, 2017; Manokawinchoke *et al.*, 2015).

Clinically applied regenerative protocols, such as revascularization, utilize the regenerative capacity of endogenous stem cells. During treatment, over instrumentation into the periapical tissues causes stem cells to migrate and enter into the canal system via the blood clot. This process leads to the treatment of immature teeth with

pulp necrosis by replacing dentin, root structures and pulp-dentin complex cells (Diogenes *et al.*, 2016). However, revascularization can be accompanied by intracanal calcification (Song *et al.*, 2017). It is unclear how Wnts and their inhibitors SOST and DKK-1 are involved in this process. Therefore, studies which investigate the regulation of SOST and DKK-1 in pulp cells and tissue under conditions which are present in the early phase of healing are of clear relevance (Janjić *et al.*, 2018).

Results of Zhang *et al.* indicate that Wnt/ β -catenin signaling is required for odontoblastic differentiation and also promotes proliferation of pre-odontoblasts and odontogenesis during root development (Zhang *et al.*, 2013). Based on these findings, the integrity of Wnt/ β -catenin signaling in odontoblasts is vital for proliferation and differentiation through root formation. Targeted deletion of β -catenin in odontoblasts leads to incomplete incisors and rootless molars. Furthermore, β -catenin deficiency disrupts the differentiation of odontoblasts and cementoblasts. (Zhang *et al.*, 2013) Epithelial expression of DKK-1 or epithelium-specific inactivation of β -catenin causes abnormal tooth development at the early bud stage (Andl *et al.*, 2002; Chen *et al.*, 2009; Han *et al.*, 2011). Mesenchyme-specific inhibition of β -catenin indicates the critical role of Wnt/ β -catenin signaling in the potential mesenchymal odontogenic activation throughout early tooth growth (Chen *et al.*, 2009).

After tooth development odontoblasts secrete dentin and the pulp chamber system narrows with age (Arana-Chavez & Massa, 2004; Foster *et al.*, 2013; Lee *et al.*, 2013; Sakai *et al.*, 2010). While dentin gradually thickens, the pulp chamber space is reduced and a massive bone loss can be observed during aging. (Boskey & Coleman, 2010; Carvalho & Lussi, 2017; Gabet & Bab, 2011) In response to injuries and stress the formation of mineralized tissue by odontoblasts, also termed reparative dentinogenesis, is stimulated. The odontoblasts which are involved in this repair process are clearly responsive to Wnt (Zhao *et al.*, 2018). SOST knockout mice demonstrate dramatically enhanced formation of reparative mineralized bridges and increased mineralization in dental pulp cells compared with wild-type mice (Collignon *et al.*, 2017). These findings are related to an increased SOST expression in wild type cells. Further more these results show that SOST deficiency accelerates reparative dentinogenesis after pulp damage and therefore inhibition of SOST may provide a promising therapeutic strategy to improve the healing of injured pulps (Collignon *et al.*, 2017).

Expression of DKK-1 is up-regulated in a rodent model of induced periapical lesions, suggesting that DKK-1 is involved in the inflammatory processes and bone resorption in periapical lesions (Zhang *et al.*, 2014). Interestingly several potential therapeutic approaches rely on the modulation of Wnt signaling (Ishimoto *et al.*, 2015; Lee *et al.*, 2016; Zhao *et al.*, 2018), including several plant-derived molecules. Baicalein promotes odontoblastic differentiation and angiogenesis of human dental pulp cells. It was suggested that via the inhibition of DKK-1 Baicalein can contribute to regenerative endodontics and dental pulp repair. (Lee *et al.*, 2016).

Given the importance of hypoxia-induced signaling in the early phase of pulp healing we investigated the production of SOST and DKK-1 in dental pulp cells upon treatment with hypoxia or the hypoxia mimetic agent L-mimosine in monolayer, spheroid, and tooth slice cultures. Our results show that the response with regard to SOST and DKK-1 production depends on the culture model.

Taken together, the literature highlights a major role of SOST and DKK-1 in tooth development and as a potential target for regenerative strategies.

Orthodontics perspective

Orthodontic tooth movement is a result of external force applications to the teeth, an intervention which involves remodeling in dental and surrounding oral tissues, such as periodontal ligament, alveolar bone, and gingiva (Antoun *et al.*, 2017; Krishnan & Davidovitch, 2006). The applied forces induce displacement of the teeth in the periodontal ligament space, thereby establishing sites where the tissue is compressed and sites where traction establishes (Antoun *et al.*, 2017; Krishnan & Davidovitch, 2006). Thereby, modeling of the alveolar bone through the processes of bone resorption and bone formation is induced leading to clinical changes in the position of the tooth. (Krishnan & Davidovitch, 2006; Krishnan & Davidovitch, 2009; Reitan, 1967)

Mechanical forces can regulate the expression of SOST by osteocytes; for instance the expression of SOST is reduced when loading is increased, and increased when loading is decreased leading to an increase in bone formation or bone loss, respectively (Moustafa *et al.*, 2012; Robling *et al.*, 2008). This feature of SOST indicates that it is a crucial protein in bone formation under mechanical stimulation (Agholme *et al.*, 2011; Mantila Roosa *et al.*, 2011; Robling *et al.*, 2008;). Given that SOST is produced by osteocytes and cementocytes in alveolar bone and in cellular cementum, respectively, an involvement of SOST in tooth movement is likely (Jäger *et al.*, 2010).

During orthodontic tooth movement, one of the essential processes is remodeling of the periodontal ligament. Periostin is one of the factors which increased in the periodontal ligament during initial stages of orthodontic tooth movement (Wilde *et al.*, 2003), a matricellular protein and collagen-rich tissues which are highly expressed in periosteum under persistent mechanical stress (Horiuchi *et al.*, 1999). Periodontal cells have shown to respond to these mechanical forces with an increase of TGF- β which can stimulate SOST (Gruber *et al.*, 2017; Manokawinchoke *et al.*, 2015). The increase in the levels of SOST in the compression side and the decrease in the tension side during tooth movement can be suggestive of an interplay between periostin from Sharpey's fibres and SOST in alveolar bone during orthodontic therapy (Nishiyama *et al.*, 2015).

Cementoblasts are highly differentiated cells from the mesenchymal lineage which generate cementum (Bosshardt, 2005). Diseases that influence bone characteristics often interfere with

the properties of the cementum, which highlights the similarities of these tissues. Although SOST is expressed in both osteocytes and cementocytes in dental tissues (Jäger *et al.*, 2010), it is still not clear if cementocytes can have similar mechanical stress sensing capacity as osteocytes. However, SOST which is released by osteocytes in the alveolar bone may affect the function of cementoblasts. This process might be modulated upon orthodontic treatment. Research on the role of SOST can also help to understand the impact of therapeutics on the periodontium. Strontium supports differentiation of cementoblasts (Bao *et al.*, 2014) with one possible underlying mechanism being that strontium decreases the production of SOST. Thus, strontium was proposed for regenerative approaches to support cementum production in cases of root resorption. (Bao *et al.*, 2014) However, if SOST or DKK-1 can serve effectively as targets for orthodontic approaches and cementum repair requires further studies.

Periodontics perspective

Periodontitis is an inflammatory disease of the teeth supporting tissues, has a multifactorial etiology. The inflammation created by specific microorganisms extends deep into the tissues and causes the destruction of the tooth supporting connective tissue and alveolar bone. This progressive process leads to the pathological impairment of collagen fibres, loss of periodontal ligament and alveolar bone recession (Armitage, 2004; Highfield, 2009; Pihlstrom *et al.*, 2005). Evaluation of SOST and DKK-1 in chronic periodontitis patients showed that both of these inhibitors were up-regulated in the periodontal tissues of these subjects (Napimoga *et al.*, 2014).

There are hints from animal studies that inflammation can not only trigger bone resorption, but also decrease bone formation. This anti-anabolic effects seem to be mediated by increased levels of DKK-1 (Heiland *et al.*, 2010). *In vitro* models of oral soft tissue augmentation suggest that DKK-1 is also increased in oral soft tissue wound healing (Agis *et al.*, 2014). Thus, Wnt inhibitors can play a major role in pathological processes and regeneration in the periodontal tissue. Although periodontitis can be treated in early stages, mostly because of the chronic entity of problem, it is diagnosed in the advanced phase of destruction of the periodontal ligament and the prognosis of maintaining the teeth are poor. Conventional regenerative approaches use biomaterials of natural or synthetic origin as filler for the defect thereby aiding the host to replace lost periodontal tissue and bone. While these interventions can stimulate tissue repair and stop the destruction of the periodontium, methods to archive full regeneration are still the focus of research (Hernández-Monjaraz *et al.*, 2018).

The blocking of Wnt signaling impairs the periodontal ligament and alveolar bone (Lim *et al.*, 2014), while enhancing Wnt signaling by SOST knock out stimulates alveolar bone formation and reduces the width of periodontal ligament (Kuchler *et al.*, 2014). Expression of SOST by cementocytes suggests that these cells may regulate cell activity on the cementum surface (Bao *et al.*, 2013; Lehen *et al.*, 2012). TGF- β can increase the production of SOST in fibroblasts from periodontal ligament and gingiva (Gruber *et al.*, 2017). This mechanism seems to be involved in the impact of mechanical loading on mineralized tissue formation in

the periodontal ligament (Manokawinchoke *et al.*, 2015). Deletion of SOST leads to more cellular cementum, in parallel to more dramatically increased alveolar bone deposition (Kuchler *et al.*, 2014). Blocking SOST by application of a SOST-specific antibody enhances healing of alveolar bone in experimental periodontitis (Chen *et al.*, 2015; Liu *et al.*, 2018; Taut *et al.*, 2013). In addition, it was reported that reduced SOST in periostin knock-out mice can re-establish periodontal ligament and alveolar bone (Rangiani *et al.*, 2016; Ren *et al.*, 2015). This evidence supports that targeting of SOST is a feasible approach for periodontal therapy.

Dental cementum is a mineralized hard tissue on the surface of root dentin and present either in acellular or cellular form. Defective cementum results in periodontal breakdown, tooth dysfunction, and finally leads to tooth loss. Cementogenesis is a key element in the process of periodontal tissue regeneration (Bosshardt, 2005; Kao & Fiorellini, 2012). SOST was detected only in cementocytes of cellular cementum in the late stages of cementum development (Lehen *et al.*, 2012). SOST levels in cementocytes increased in periodontal ligament cultures, following mineralization treatment (Jäger *et al.*, 2010). Interestingly, in periodontal ligament cells Baicalein can promote osteoblastic differentiation involving Wnt/ β -catenin signaling (Chen *et al.*, 2017). DKK-1 significantly reversed the effects of Baicalein on human periodontal ligament cells (Chen *et al.*, 2017). It is possible that this mechanism can be exploited in regenerative approaches.

The here presented literature supports the significant effects of SOST and DKK-1 in the periodontium system and periodontal diseases. As a result, they could be the main targets in future periodontics regenerative therapies.

Oral surgery perspective

The alveolar bone supports the tooth in the maxilla and mandible and is characterized by continuous and rapid remodeling in response to mechanical forces (Javed *et al.*, 2010; Pagni *et al.*, 2012). Thereby alveolar bone continuously adapts to functional load. If this mechanical stimuli is lacking the alveolar bone undergoes a resorptive process (Einhorn & Gerstenfeld, 2015; Pagni *et al.*, 2012; Sodek & McKee, 2000). Following trauma due to overloading or surgery bone has the capacity to regenerate. While long bone healing occurs by endo-chondral ossification, alveolar bone healing typically occurs without histological cartilage formation (Devlin *et al.*, 1997). The success of oral surgery procedures, such as implants, depends on the proper healing of alveolar bone and strategies which stimulate bone regeneration (Lin *et al.*, 2011). Thus understanding the cell and molecular biological background of bone healing is clearly of clinical relevance.

In bone, SOST is mainly secreted by osteocytes and represents a key modulator of bone homeostasis (Brunkow *et al.*, 2001; van Bezooijen *et al.*, 2004). The importance of SOST in bone formation is illustrated by sclerosteosis, a rare autosomal recessive disorder with a loss-of-function mutation in SOST (Sebastian & Loots, 2018; Yavropoulou *et al.*, 2014). Further evidence comes from Van Buchem Disease, which is characterized by a noncoding deletion which removes a SOST-specific regulator

(Sebastian & Loots, 2018; Yavropoulou *et al.*, 2014). These diseases show bone overgrowth, particularly in the craniofacial bones and the jaw bone (Balemans *et al.*, 2002; Brunkow *et al.*, 2001). There is also a phenotype in oral tissue; partial anodontia, malocclusion, and delayed tooth eruption is seen in subjects with Van Buchem Disease or SOST (Stephen *et al.*, 2001; van Bezooijen *et al.*, 2009). Animal studies on the role of SOST in loss of function or gain function models indicate SOST decreases bone formation and can stimulate bone resorption (Canalis, 2013; Li *et al.*, 2008; O'Brien *et al.*, 2013). SOST knockout mice show a dramatically increased basal mandibular bone and less effect on the coronal and apical part of the alveolar bone (Kuchler *et al.*, 2014).

The response of bone to mechanical loading is highly important in implantology. Since SOST is a key player in the regulation of bone formation in the response to mechanical loading it is important to understand its role in the alveolar bone. Upon loading, osteocytes reduce the expression of SOST permitting Wnts to bind their receptors. (Burgers & Williams, 2013; Galli *et al.*, 2010; Zhao *et al.*, 2013). Interestingly, SOST expression is up-regulated around implants without primary stability (Shu *et al.*, 2017). Thus, it is very likely that SOST regulates the adaptation of bone around dental implants to the mechanical forces of loading. Application of SOST specific antibodies has been shown to stimulate bone formation around dental implants (Yu *et al.*, 2018).

DKK-1 can decrease Wnt/ β catenin signalling and reduced the production of type II collagen in chondrocytes under mechanical loading (Niu *et al.*, 2016). Interestingly, DKK-1 has been found to play a distinct role in inflammation-induced bone loss (Heiland *et al.*, 2010). However, when evaluating genetic markers for peri-implantitis clinically, no direct relation of DKK-1 and peri-implantitis is seen (Hall *et al.*, 2011). Thus, further research is required to understand the role of DKK-1 in the alveolar bone.

From a clinical standpoint, osteoporosis is a risk factor in implantology. Inspired by the fact that inhibition of SOST can effectively improve bone formation in osteoporotic patients strategies which harness this capacity to improve bone formation around implants have been evaluated. (Canalis, 2013; Costa & Bilezikian, 2012; Recker *et al.*, 2015; Shu *et al.*, 2017). Implant

osseointegration is superior in the SOST knockout mice suggesting that SOST is a promising target to enhance implant osseointegration in osteoporosis. (Shu *et al.*, 2017) Also application of antibodies specific for SOST can improve bone formation around implants (Yu *et al.*, 2018).

Taken together these data show the relevance of the Wnt signaling inhibitors of SOST and DKK-1 for implantology and bone augmentation, as well as for tumor development and cancer therapy. Future studies will show how effective targeting of SOST and DKK-1 can be applied to stimulate regeneration and for tumor therapy.

Conclusion

The here presented review highlights the relevance of Wnt signaling, the inhibitors SOST and DKK-1, and its key role in oral tissue homeostasis throughout life (Florio *et al.*, 2016; Heiland *et al.*, 2010; Witcher *et al.*, 2018). Major sources of Wnt inhibitors are the mechanosensors of the bone, the osteocytes (Moustafa *et al.*, 2012; Odagaki *et al.*, 2018; Robling *et al.*, 2008). Cells from oral tissue including cementocytes can modulate Wnt signaling by these inhibitors and mice deficient of SOST or DKK-1 have highlighted the important role in the oral tissue (Jäger *et al.*, 2010; Lehnen *et al.*, 2012). Based on this knowledge concepts to antagonize inhibitors Wnt signaling have been developed to support oral tissue regeneration (Taut *et al.*, 2013; Yu *et al.*, 2018). The future will reveal the capacity of these strategies which target these Wnt signaling inhibitors for regenerative therapy in dentistry. Given the interplay of SOST and DKK-1, a bidirectional approach which targets both SOST and DKK-1 locally has high potential (Florio *et al.*, 2016).

Data availability

No data is associated with this article.

Grant information

We thank the European Society of Endodontology (ESE) for financial support of our research [ESE research grant 2015]. The authors deny any conflict of interest.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Agholme F, Isaksson H, Li X, *et al.*: **Anti-sclerostin antibody and mechanical loading appear to influence metaphyseal bone independently in rats.** *Acta Orthop.* 2011; **82**(5): 628–632.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Agis H, Collins A, Taut AD, *et al.*: **Cell population kinetics of collagen scaffolds in ex vivo oral wound repair.** *PLoS One.* 2014; **9**(11): e112680.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ahn VE, Chu ML, Choi HJ, *et al.*: **Structural basis of Wnt signaling inhibition by Dickkopf binding to LRP5/6.** *Dev Cell.* 2011; **21**(5): 862–873.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Andl T, Reddy ST, Gaddapara T, *et al.*: **WNT signals are required for the initiation of hair follicle development.** *Dev Cell.* 2002; **2**(5): 643–653.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Antoun JS, Mei L, Gibbs K, *et al.*: **Effect of orthodontic treatment on the periodontal tissues.** *Periodontol 2000.* 2017; **74**(1): 140–157.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Arana-Chavez VE, Massa LF: **Odontoblasts: the cells forming and maintaining dentine.** *Int J Biochem Cell Biol.* 2004; **36**(8): 1367–1373.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Armitage GC: **Periodontal diagnoses and classification of periodontal**

diseases. *Periodontol* 2000. 2004; **34**(1): 9–21.

[PubMed Abstract](#) | [Publisher Full Text](#)

Balemans W, Patel N, Ebeling M, *et al.*: Identification of a 52 kb deletion downstream of the *SOST* gene in patients with van Buchem disease. *J Med Genet*. 2002; **39**(2): 91–97.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Bao X, Liu Y, Han G, *et al.*: The effect on proliferation and differentiation of cementoblast by using sclerostin as inhibitor. *Int J Mol Sci*. 2013; **14**(10): 21140–21152.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Bao X, Liu X, Zhang Y, *et al.*: Strontium promotes cementoblasts differentiation through inhibiting sclerostin expression *in vitro*. *Biomed Res Int*. 2014; **2014**: 487535.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Boskey AL, Coleman R: Aging and bone. *J Dent Res*. 2010; **89**(12): 1333–1348.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Bosshardt DD: Are cementoblasts a subpopulation of osteoblasts or a unique phenotype? *J Dent Res*. 2005; **84**(5): 390–406.

[PubMed Abstract](#) | [Publisher Full Text](#)

Bourhis E, Tam C, Franke Y, *et al.*: Reconstitution of a frizzled8.Wnt3a.LRP6 signaling complex reveals multiple Wnt and Dkk1 binding sites on LRP6. *J Biol Chem*. 2010; **285**(12): 9172–9179.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Bourhis E, Wang W, Tam C, *et al.*: Wnt antagonists bind through a short peptide to the first β -propeller domain of LRP6. *Structure*. 2011; **19**(10): 1433–1442.

[PubMed Abstract](#) | [Publisher Full Text](#)

Brunkow ME, Gardner JC, Van Ness J, *et al.*: Bone dysplasia sclerosteosis results from loss of the *SOST* gene product, a novel cysteine knot-containing protein. *Am J Hum Genet*. 2001; **68**(3): 577–589.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Burgers TA, Williams BO: Regulation of Wnt/ β -catenin signaling within and from osteocytes. *Bone*. 2013; **54**(2): 244–249.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Canalis E: Wnt signalling in osteoporosis: mechanisms and novel therapeutic approaches. *Nat Rev Endocrinol*. 2013; **9**(10): 575–583.

[PubMed Abstract](#) | [Publisher Full Text](#)

Carvalho TS, Lussi A: Age-related morphological, histological and functional changes in teeth. *J Oral Rehabil*. 2017; **44**(4): 291–298.

[PubMed Abstract](#) | [Publisher Full Text](#)

Chen H, Xu X, Liu M, *et al.*: Sclerostin antibody treatment causes greater alveolar crest height and bone mass in an ovariectomized rat model of localized periodontitis. *Bone*. 2015; **76**: 141–148.

[PubMed Abstract](#) | [Publisher Full Text](#)

Chen J, Lan Y, Baek JA, *et al.*: Wnt/ β -catenin signaling plays an essential role in activation of odontogenic mesenchyme during early tooth development. *Dev Biol*. 2009; **334**(1): 174–185.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Chen LJ, Hu BB, Shi XL, *et al.*: Baicalein enhances the osteogenic differentiation of human periodontal ligament cells by activating the Wnt/ β -catenin signaling pathway. *Arch Oral Biol*. 2017; **78**: 100–108.

[PubMed Abstract](#) | [Publisher Full Text](#)

Cheng Z, Biechele T, Wei Z, *et al.*: Crystal structures of the extracellular domain of LRP6 and its complex with DKK1. *Nat Struct Mol Biol*. 2011; **18**(11): 1204–1210.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Chueh LH, Huang GT: Immature teeth with periradicular periodontitis or abscess undergoing apexogenesis: a paradigm shift. *J Endod*. 2006; **32**(12): 1205–1213.

[PubMed Abstract](#) | [Publisher Full Text](#)

Collignon AM, Amri N, Lesieur J, *et al.*: Sclerostin Deficiency Promotes Reparative Dentinogenesis. *J Dent Res*. 2017; **96**(7): 815–821.

[PubMed Abstract](#) | [Publisher Full Text](#)

Costa AG, Bilezikian JP: Sclerostin: therapeutic horizons based upon its actions. *Curr Osteoporos Rep*. 2012; **10**(1): 64–72.

[PubMed Abstract](#) | [Publisher Full Text](#)

Devlin H, Hoyland J, Newall JF, *et al.*: Trabecular bone formation in the healing of the rodent molar tooth extraction socket. *J Bone Miner Res*. 1997; **12**(12): 2061–2067.

[PubMed Abstract](#) | [Publisher Full Text](#)

Diogenes A, Ruparel NB, Shiloah Y, *et al.*: Regenerative endodontics: A way forward. *J Am Dent Assoc*. 2016; **147**(5): 372–380.

[PubMed Abstract](#) | [Publisher Full Text](#)

Dissanayaka WL, Zhu L, Hargreaves KM, *et al.*: Scaffold-free Prevascularized Microtissue Spheroids for Pulp Regeneration. *J Dent Res*. 2014; **93**(12): 1296–1303.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Einhorn TA, Gerstenfeld LC: Fracture healing: mechanisms and interventions. *Nat Rev Rheumatol*. 2015; **11**(1): 45–54.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Florio M, Gunasekaran K, Stolina M, *et al.*: A bispecific antibody targeting sclerostin and DKK-1 promotes bone mass accrual and fracture repair. *Nat Commun*. 2016; **7**: 11505.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Foster BL, Nagatomo KJ, Tso HW, *et al.*: Tooth root dentin mineralization

defects in a mouse model of hypophosphatasia. *J Bone Miner Res*. 2013; **28**(2): 271–282.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Fretwurst T, Larsson L, Yu SH, *et al.*: Periodontal Tissue Bioengineering: Is the Future Now? *Compend Contin Educ Dent*. 2018; **39**(4): 218–223; quiz 224.

[PubMed Abstract](#)

Gabet Y, Bab I: Microarchitectural changes in the aging skeleton. *Curr Osteoporos Rep*. 2011; **9**(4): 177–183.

[PubMed Abstract](#) | [Publisher Full Text](#)

Galler KM, Buchalla W, Hiller KA, *et al.*: Influence of root canal disinfectants on growth factor release from dentin. *J Endod*. 2015; **41**(3): 363–368.

[PubMed Abstract](#) | [Publisher Full Text](#)

Galli C, Passeri G, Macaluso GM: Osteocytes and WNT: the mechanical control of bone formation. *J Dent Res*. 2010; **89**(4): 331–343.

[PubMed Abstract](#) | [Publisher Full Text](#)

Gruber M, Gruber R, Agis H: Transforming growth factor- β 1 increases sclerostin in fibroblasts of the periodontal ligament and the gingiva – Matters. *Matters Archive*. 2017.

[Reference Source](#)

Hall J, Britse AO, Jemt T, *et al.*: A controlled clinical exploratory study on genetic markers for peri-implantitis. *Eur J Oral Implantol*. 2011; **4**(4): 371–382.

[PubMed Abstract](#)

Han XL, Liu M, Voisey A, *et al.*: Post-natal effect of overexpressed DKK1 on mandibular molar formation. *J Dent Res*. 2011; **90**(11): 1312–1317.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Heiland GR, Zwerina K, Baum W, *et al.*: Neutralisation of Dkk-1 protects from systemic bone loss during inflammation and reduces sclerostin expression. *Ann Rheum Dis*. 2010; **69**(12): 2152–2159.

[PubMed Abstract](#) | [Publisher Full Text](#)

Hernández-Monjaraz B, Santiago-Osorio E, Monroy-García A, *et al.*: Mesenchymal Stem Cells of Dental Origin for Inducing Tissue Regeneration in Periodontitis: A Mini-Review. *Int J Mol Sci*. 2018; **19**(4): pii: E944.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Highfield J: Diagnosis and classification of periodontal disease. *Aust Dent J*. 2009; **54** Suppl 1: S11–26.

[PubMed Abstract](#) | [Publisher Full Text](#)

Horiuchi K, Amizuka N, Takeshita S, *et al.*: Identification and characterization of a novel protein, periostin, with restricted expression to periosteum and periodontal ligament and increased expression by transforming growth factor beta. *J Bone Miner Res*. 1999; **14**(7): 1239–1249.

[PubMed Abstract](#) | [Publisher Full Text](#)

Huang GT: A paradigm shift in endodontic management of immature teeth: conservation of stem cells for regeneration. *J Dent*. 2008; **36**(6): 379–386.

[PubMed Abstract](#) | [Publisher Full Text](#)

Huang GT, Sonoyama W, Liu Y, *et al.*: The hidden treasure in apical papilla: the potential role in pulp/dentin regeneration and bioroot engineering. *J Endod*. 2008; **34**(6): 645–651.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Ishimoto K, Hayano S, Yanagita T, *et al.*: Topical application of lithium chloride on the pulp induces dentin regeneration. *PLoS One*. 2015; **10**(3): e0121938.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Itoh Y, Sasaki JI, Hashimoto M, *et al.*: Pulp Regeneration by 3-dimensional Dental Pulp Stem Cell Constructs. *J Dent Res*. 2018; **97**(10): 1137–1143.

[PubMed Abstract](#) | [Publisher Full Text](#)

Jäger A, Götz W, Lossdörfer S, *et al.*: Localization of *SOST*/sclerostin in cementocytes *in vivo* and in mineralizing periodontal ligament cells *in vitro*. *J Periodontol Res*. 2010; **45**(2): 246–254.

[PubMed Abstract](#) | [Publisher Full Text](#)

Janjić K, Cvikić B, Kurzmann C, *et al.*: Do hypoxia and L-mimosine modulate sclerostin and dickkopf-1 production in human dental pulp-derived cells? Insights from monolayer, spheroid and tooth slice cultures. *BMC Oral Health*. 2018; **18**(1): 36.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Javed A, Chen H, Ghori FY: Genetic and transcriptional control of bone formation. *Oral Maxillofac Surg Clin North Am*. 2010; **22**(3): 283–93, v.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Kaigler D, Avila-Ortiz G, Travan S, *et al.*: Bone Engineering of Maxillary Sinus Bone Deficiencies Using Enriched CD90+ Stem Cell Therapy: A Randomized Clinical Trial. *J Bone Miner Res*. 2015; **30**(7): 1206–1216.

[PubMed Abstract](#) | [Publisher Full Text](#)

Kao DW, Fiorellini JP: Regenerative periodontal therapy. *Front Oral Biol*. 2012; **15**: 149–159.

[PubMed Abstract](#) | [Publisher Full Text](#)

Kim TH, Bae CH, Lee JC, *et al.*: β -catenin is required in odontoblasts for tooth root formation. *J Dent Res*. 2013; **92**(3): 215–221.

[PubMed Abstract](#) | [Publisher Full Text](#)

Kratochwil K, Galceran J, Tontsch S, *et al.*: FGF4, a direct target of LEF1 and Wnt signaling, can rescue the arrest of tooth organogenesis in *Lef1*⁺ mice. *Genes Dev*. 2002; **16**(24): 3173–3185.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Krishnan V, Davidovitch Z: Cellular, molecular, and tissue-level reactions to orthodontic force. *Am J Orthod Dentofacial Orthop*. 2006; **129**(4): 469.e1–32.

[PubMed Abstract](#) | [Publisher Full Text](#)

- Krishnan V, Davidovitch Z: **On a path to unfolding the biological mechanisms of orthodontic tooth movement.** *J Dent Res.* 2009; **88**(7): 597–608.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kuchler U, Schwarze UY, Dobsak T, *et al.*: **Dental and periodontal phenotype in sclerostin knockout mice.** *Int J Oral Sci.* 2014; **6**(2): 70–76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lee SI, Kim SY, Park KR, *et al.*: **Baicalein Promotes Angiogenesis and Odontoblastic Differentiation via the BMP and Wnt Pathways in Human Dental Pulp Cells.** *Am J Chin Med.* 2016; **44**(7): 1457–1472.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lee YH, Kim GE, Cho HJ, *et al.*: **Aging of *in vitro* pulp illustrates change of inflammation and dentinogenesis.** *J Endod.* 2013; **39**(3): 340–345.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lehnen SD, Götz W, Baxmann M, *et al.*: **Immunohistochemical evidence for sclerostin during cementogenesis in mice.** *Ann Anat.* 2012; **194**(5): 415–421.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Li X, Ominsky MS, Niu QT, *et al.*: **Targeted deletion of the sclerostin gene in mice results in increased bone formation and bone strength.** *J Bone Miner Res.* 2008; **23**(6): 860–869.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Li X, Zhang Y, Kang H, *et al.*: **Sclerostin binds to LRP5/6 and antagonizes canonical Wnt signaling.** *J Biol Chem.* 2005; **280**(20): 19883–19887.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lim WH, Liu B, Cheng D, *et al.*: **Wnt signaling regulates homeostasis of the periodontal ligament.** *J Periodontol Res.* 2014; **49**(6): 751–759.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lin Z, Rios HF, Volk SL, *et al.*: **Gene expression dynamics during bone healing and osseointegration.** *J Periodontol.* 2011; **82**(7): 1007–1017.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Liu F, Chu EY, Watt B, *et al.*: **Wnt/ β -catenin signaling directs multiple stages of tooth morphogenesis.** *Dev Biol.* 2008; **313**(1): 210–224.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Liu M, Kurimoto P, Zhang J, *et al.*: **Sclerostin and DKK1 Inhibition Preserves and Augments Alveolar Bone Volume and Architecture in Rats with Alveolar Bone Loss.** *J Dent Res.* 2018; **97**(9): 1031–1038.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Logan CY, Nusse R: **The Wnt signaling pathway in development and disease.** *Annu Rev Cell Dev Biol.* 2004; **20**: 781–810.
[PubMed Abstract](#) | [Publisher Full Text](#)
- MacDonald BT, Joiner DM, Oyserman SM, *et al.*: **Bone mass is inversely proportional to *Dkk1* levels in mice.** *Bone.* 2007; **41**(3): 331–339.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Manokawinchoke J, Limjeerajarus N, Limjeerajarus C, *et al.*: **Mechanical Force-induced *TGF β 1* Increases Expression of *SOST/POSTN* by hPDL Cells.** *J Dent Res.* 2015; **94**(7): 983–989.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mantila Roosa SM, Liu Y, Turner CH: **Gene expression patterns in bone following mechanical loading.** *J Bone Miner Res.* 2011; **26**(1): 100–112.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mohammed MK, Shao C, Wang J, *et al.*: **Wnt/ β -catenin signaling plays an ever-expanding role in stem cell self-renewal, tumorigenesis and cancer chemoresistance.** *Genes Dis.* 2016; **3**(1): 11–40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Moustafa A, Sugiyama T, Prasad J, *et al.*: **Mechanical loading-related changes in osteocyte sclerostin expression in mice are more closely associated with the subsequent osteogenic response than the peak strains engendered.** *Osteoporos Int.* 2012; **23**(4): 1225–1234.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nakashima M, Nagasawa H, Yamada Y, *et al.*: **Regulatory role of transforming growth factor- β , bone morphogenetic protein-2, and protein-4 on gene expression of extracellular matrix proteins and differentiation of dental pulp cells.** *Dev Biol.* 1994; **162**(1): 18–28.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Napimoga MH, Nametala C, da Silva FL, *et al.*: **Involvement of the Wnt/ β -catenin signalling antagonists, sclerostin and dickkopf-related protein 1, in chronic periodontitis.** *J Clin Periodontol.* 2014; **41**(6): 550–557.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Nevins M, Giannobile WV, McGuire MK, *et al.*: **Platelet-derived growth factor stimulates bone fill and rate of attachment level gain: results of a large multicenter randomized controlled trial.** *J Periodontol.* 2005; **76**(12): 2205–2215.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Nishiyama Y, Matsumoto T, Lee JW, *et al.*: **Changes in the spatial distribution of sclerostin in the osteocytic lacuno-canalicular system in alveolar bone due to orthodontic forces, as detected on multimodal confocal fluorescence imaging analyses.** *Arch Oral Biol.* 2015; **60**(1): 45–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Niu Q, Li F, Zhang L, *et al.*: **Role of the Wnt/ β -catenin signaling pathway in the response of chondrocytes to mechanical loading.** *Int J Mol Med.* 2016; **37**(3): 755–762.
[PubMed Abstract](#) | [Publisher Full Text](#)
- O'Brien CA, Nakashima T, Takayanagi H: **Osteocyte control of osteoclastogenesis.** *Bone.* 2013; **54**(2): 258–263.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Odagaki N, Ishihara Y, Wang Z, *et al.*: **Role of Osteocyte-PDL Crosstalk in Tooth Movement via *SOST/Sclerostin*.** *J Dent Res.* 2018; **97**(12): 1374–1382.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Pagni G, Pellegrini G, Giannobile WV, *et al.*: **Postextraction alveolar ridge preservation: biological basis and treatments.** *Int J Dent.* 2012; **2012**: 151030.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Pflanz D, Birkhold AI, Albiol L, *et al.*: ***Sost* deficiency led to a greater cortical bone formation response to mechanical loading and altered gene expression.** *Sci Rep.* 2017; **7**(1): 9435.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Pihlstrom BL, Michalowicz BS, Johnson NW: **Periodontal diseases.** *Lancet.* 2005; **366**(9499): 1809–1820.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Plonka AB, Khorsand B, Yu N, *et al.*: **Effect of sustained PDGF nonviral gene delivery on repair of tooth-supporting bone defects.** *Gene Ther.* 2017; **24**(1): 31–39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rangiani A, Jing Y, Ren Y, *et al.*: **Critical roles of periostin in the process of orthodontic tooth movement.** *Eur J Orthod.* 2016; **38**(4): 373–378.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Recker RR, Benson CT, Matsumoto T, *et al.*: **A randomized, double-blind phase 2 clinical trial of blosozumab, a sclerostin antibody, in postmenopausal women with low bone mineral density.** *J Bone Miner Res.* 2015; **30**(2): 216–224.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Reitan K: **Clinical and histologic observations on tooth movement during and after orthodontic treatment.** *Am J Orthod.* 1967; **53**(10): 721–745.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ren Y, Han X, Ho SP, *et al.*: **Removal of *SOST* or blocking its product sclerostin rescues defects in the periodontitis mouse model.** *FASEB J.* 2015; **29**(7): 2702–2711.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Robling AG, Niziolek PJ, Baldridge LA, *et al.*: **Mechanical stimulation of bone *in vivo* reduces osteocyte expression of *Sost/sclerostin*.** *J Biol Chem.* 2008; **283**(9): 5866–5875.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rutherford RB, Wahle J, Tucker M, *et al.*: **Induction of reparative dentine formation in monkeys by recombinant human osteogenic protein-1.** *Arch Oral Biol.* 1993; **38**(7): 571–576.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Saag KG, Petersen J, Brandi ML, *et al.*: **Romosozumab or Alendronate for Fracture Prevention in Women with Osteoporosis.** *N Engl J Med.* 2017; **377**(15): 1417–1427.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Sakai VT, Zhang Z, Dong Z, *et al.*: **SHED differentiate into functional odontoblasts and endothelium.** *J Dent Res.* 2010; **89**(8): 791–796.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Sarkar L, Cobourne M, Naylor S, *et al.*: **Wnt/Shh interactions regulate ectodermal boundary formation during mammalian tooth development.** *Proc Natl Acad Sci U S A.* 2000; **97**(9): 4520–4524.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sarkar L, Sharpe PT: **Inhibition of Wnt signaling by exogenous *Mfrzb1* protein affects molar tooth size.** *J Dent Res.* 2000; **79**(4): 920–925.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Sebastian A, Loots GG: **Genetics of *Sost/SOST* in sclerosteosis and van Buchem disease animal models.** *Metabolism.* 2018; **80**: 38–47.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Semēnov M, Tamai K, He X: ***SOST* is a ligand for LRP5/LRP6 and a Wnt signaling inhibitor.** *J Biol Chem.* 2005; **280**(29): 26770–26775.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Seo MS, Hwang KG, Kim H, *et al.*: **Analysis of gene expression during odontogenic differentiation of cultured human dental pulp cells.** *Restor Dent Endod.* 2012; **37**(3): 142–148.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Shu R, Ai D, Bai D, *et al.*: **The effects of *SOST* on implant osseointegration in ovariectomy osteoporotic mice.** *Arch Oral Biol.* 2017; **74**: 82–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Sloan AJ, Smith AJ: **Stimulation of the dentine-pulp complex of rat incisor teeth by transforming growth factor- β isoforms 1-3 *in vitro*.** *Arch Oral Biol.* 1999; **44**(2): 149–156.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Sodek J, McKee MD: **Molecular and cellular biology of alveolar bone.** *Periodontol 2000.* 2000; **24**(1): 99–126.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Song M, Cao Y, Shin SJ, *et al.*: **Revascularization-associated Intracanal Calcification: Assessment of Prevalence and Contributing Factors.** *J Endod.* 2017; **43**(12): 2025–2033.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Stephen LX, Hamersma H, Gardner J, *et al.*: **Dental and oral manifestations of sclerosteosis.** *Int Dent J.* 2001; **51**(4): 287–290.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tang N, Song WX, Luo J, *et al.*: **BMP-9-induced osteogenic differentiation of**

mesenchymal progenitors requires functional canonical Wnt/beta-catenin signalling. *J Cell Mol Med.* 2009; **13**(8B): 2448–2464.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Taut AD, Jin Q, Chung JH, *et al.*: **Sclerostin antibody stimulates bone regeneration after experimental periodontitis.** *J Bone Miner Res.* 2013; **28**(11): 2347–2356.

[PubMed Abstract](#) | [Publisher Full Text](#)

Tziatas D, Alvanou A, Papadimitriou S, *et al.*: **Effects of recombinant basic fibroblast growth factor, insulin-like growth factor-II and transforming growth factor-beta 1 on dog dental pulp cells in vivo.** *Arch Oral Biol.* 1998; **43**(6): 431–444.

[PubMed Abstract](#) | [Publisher Full Text](#)

van Bezooijen RL, Bronckers AL, Gortzak RA, *et al.*: **Sclerostin in mineralized matrices and van Buchem disease.** *J Dent Res.* 2009; **88**(6): 569–574.

[PubMed Abstract](#) | [Publisher Full Text](#)

van Bezooijen RL, Roelen BA, Visser A, *et al.*: **Sclerostin is an osteocyte-expressed negative regulator of bone formation, but not a classical BMP antagonist.** *J Exp Med.* 2004; **199**(6): 805–814.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Wang XP, O'Connell DJ, Lund JJ, *et al.*: **Apc inhibition of Wnt signaling regulates supernumerary tooth formation during embryogenesis and throughout adulthood.** *Development.* 2009; **136**(11): 1939–1949.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Wilde J, Yokozeki M, Terai K, *et al.*: **The divergent expression of periostin mRNA in the periodontal ligament during experimental tooth movement.** *Cell Tissue Res.* 2003; **312**(3): 345–351.

[PubMed Abstract](#) | [Publisher Full Text](#)

Winkler DG, Sutherland MK, Geoghegan JC, *et al.*: **Osteocyte control of bone formation via sclerostin, a novel BMP antagonist.** *EMBO J.* 2003; **22**(23): 6267–6276.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Witcher PC, Miner SE, Horan DJ, *et al.*: **Sclerostin neutralization unleashes the**

osteoblastic effects of Dkk1 inhibition. *JCI Insight.* 2018; **3**(11): pii: 98673.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Yang K, Wang X, Zhang H, *et al.*: **The evolving roles of canonical WNT signaling in stem cells and tumorigenesis: implications in targeted cancer therapies.** *Lab Invest.* 2016; **96**(2): 116–136.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Yavropoulou MP, Xygonakis C, Lolou M, *et al.*: **The sclerostin story: from human genetics to the development of novel anabolic treatment for osteoporosis.** *Hormones (Athens).* 2014; **13**(4): 323–337.

[PubMed Abstract](#) | [Publisher Full Text](#)

Yorgan TA, Schinke T: **Relevance of Wnt signaling for osteoblastic therapy.** *Mol Cell Ther.* 2014; **2**: 22.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Yu SH, Hao J, Fretwurst T, *et al.*: **Sclerostin-Neutralizing Antibody Enhances Bone Regeneration Around Oral Implants.** *Tissue Eng Part A.* 2018; **24**(21–22): 1672–1679.

[PubMed Abstract](#) | [Publisher Full Text](#)

Zhang R, Huang S, Wang L, *et al.*: **Histochemical localization of Dickkopf-1 in induced rat periapical lesions.** *J Endod.* 2014; **40**(9): 1394–1399.

[PubMed Abstract](#) | [Publisher Full Text](#)

Zhang R, Yang G, Wu X, *et al.*: **Disruption of Wnt/β-catenin signaling in odontoblasts and cementoblasts arrests tooth root development in postnatal mouse teeth.** *Int J Biol Sci.* 2013; **9**(3): 228–236.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Zhao L, Shim JW, Dodge TR, *et al.*: **Inactivation of Lrp5 in osteocytes reduces young's modulus and responsiveness to the mechanical loading.** *Bone.* 2013; **54**(1): 35–43.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Zhao Y, Yuan X, Liu B, *et al.*: **Wnt-Responsive Odontoblasts Secrete New Dentin after Superficial Tooth Injury.** *J Dent Res.* 2018; **97**(9): 1047–1054.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 15 April 2019

<https://doi.org/10.5256/f1000research.19462.r46090>

© 2019 Jin Q. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Qiming Jin

Department of Cariology, Restorative Sciences, and Endodontics, School of Dentistry, University of Michigan, Ann Arbor, MI, USA

This manuscript is a review paper on sclerostin and dickkopf-1 impacts on dentistry. The authors made good publication search work, and reviewed the expressions and biological functions of sclerostin and DKK-1 in each parts of tooth and surrounding tissues. Furthermore, the authors explored the correlation between SOST and DKK-1 and various pathologies. More importantly, the treatment pathways to target SOST and DKK-1 were discussed profoundly. The only flaw is the lack of SOST and DKK-1 in enamel formation, which does not affect the merits of this paper.

Is the topic of the review discussed comprehensively in the context of the current literature?

Yes

Are all factual statements correct and adequately supported by citations?

Yes

Is the review written in accessible language?

Yes

Are the conclusions drawn appropriate in the context of the current research literature?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: periodontal regeneration and dental implant researches

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 04 February 2019

<https://doi.org/10.5256/f1000research.19462.r43827>

© 2019 Rodríguez-Lozano F. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Francisco Javier Rodríguez-Lozano

Cellular Therapy and Hematopoietic Transplant Unit, Hematology Department, Virgen de la Arrixaca Clinical University Hospital, IMIB-Arrixaca, University of Murcia, Murcia, Spain

The manuscript fits into the aim of the journal. The manuscript is well written and the results presented interesting.

I would like to know the search strategy. Inclusion/exclusion criteria.

Minor aspects:

Endodontic perspective: "Given the importance of hypoxia-induced signaling in the early phase of pulp healing we investigated the production of SOST and DKK-1 in dental pulp cells upon treatment with hypoxia or the hypoxia mimetic agent L-mimosine in monolayer, spheroid, and tooth slice cultures". Citation?

Periodontic Perspective: "Dental cementum is a mineralized hard tissue on the surface of root dentin and present either in acellular or cellular form. Defective cementum results in periodontal breakdown, tooth dysfunction, and finally leads to tooth loss." I recommend adding a reference.

Finally, I recommend re-writing the conclusion without citations

Is the topic of the review discussed comprehensively in the context of the current literature?

Yes

Are all factual statements correct and adequately supported by citations?

Yes

Is the review written in accessible language?

Yes

Are the conclusions drawn appropriate in the context of the current research literature?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: endodontics; stem cells; dental stem cells

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research