REVIEW

REVISED Recent advances in managing fecal incontinence

[version 2; peer review: 3 approved]

Giovanna Da Silva , Anne Sirany

Department of Colon and Rectal Surgery, Cleveland Clinic Florida, Weston, FL, 33331, USA

v2 First published: 31 Jul 2019, 8(F1000 Faculty Rev):1291
<https://doi.org/10.12688/f1000research.15270.1>Latest published: 23 Aug 2019, 8(F1000 Faculty Rev):1291
<https://doi.org/10.12688/f1000research.15270.2>**Abstract**

Fecal incontinence (FI) is the uncontrolled passage of feces or gas in an individual who previously had control. The prevalence of the problem varies but can be as high as 50% of institutionalized individuals. The severity varies among individuals, but the negative impact on self-esteem and quality of life can have devastating effects. The goals of treatment are to decrease the frequency and severity of episodes as well as to improve quality of life. At present, several therapies, ranging from medical management to more invasive surgical interventions, are offered for the management of FI. In this article, we review the most recent advances in the management of FI.

Keywords

Fecal incontinence, Cleveland Clinic Incontinence, Magnetic anal sphincter augmentation, Percutaneous nerve evaluation, Percutaneous tibial nerve stimulation, Sacral nerve neuromodulation.

Open Peer Review**Approval Status**

	1	2	3
version 2 (revision) 23 Aug 2019			
version 1 31 Jul 2019			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Charles H. Knowles**, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, Queen Mary University of London, UK
2. **Stacy B. Menees**, University of Michigan, Ann Arbor, USA
3. **Arnold Wald**, University of Wisconsin School of Medicine and Public Health, Madison, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Giovanna Da Silva (dasilvg@ccf.org)

Author roles: **Da Silva G:** Conceptualization, Methodology, Project Administration, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Sirany A:** Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2019 Da Silva G and Sirany A. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Da Silva G and Sirany A. **Recent advances in managing fecal incontinence [version 2; peer review: 3 approved]** F1000Research 2019, 8(F1000 Faculty Rev):1291 <https://doi.org/10.12688/f1000research.15270.2>

First published: 31 Jul 2019, 8(F1000 Faculty Rev):1291 <https://doi.org/10.12688/f1000research.15270.1>

REVISED Amendments from Version 1

We have replaced [Figure 3](#) in this new version as the figure in version 1 was incorrect.

See referee reports

Introduction

Fecal incontinence (FI) is defined as the uncontrolled passage of feces or gas over at least one month's duration in an individual who had previously achieved control¹. The severity of FI varies widely but can have significant negative impacts, including reduced quality of life, negative psychological effects, and associated social stigma². The etiology of FI may be multifactorial, and risk factors include old age, female sex, sphincter injury, obstetrical trauma, post-surgical complications, diarrhea, and constipation^{3,4}. Prevalence rates vary but have been reported to range from 1.4 to 18% and up to 50% in institutionalized patients⁵.

The goals of treatment are to decrease the frequency and severity of episodes and improve quality of life. The decision of which treatment to employ is based on the severity of symptoms and integrity of the anal sphincter. A continence scoring system such as the Cleveland Clinic Florida-Fecal Incontinence

Score (CCF-FIS) is helpful in the evaluation of these patients ([Table 1](#)). The American Society of Colon and Rectal Surgeons (ASCRS) Clinical Practice Guidelines recommend that first-line therapy for FI include dietary modifications, medical management with anti-diarrheal or fiber supplement (or both) to bulk the stool, and biofeedback exercises, which can lead to improvement in a significant portion of patients with mild FI. Adjuncts to these conservative measures have recently been reported and include the vaginal bowel control system (VBS) and anal plugs. Patients with more severe disease or sphincter defects will require more invasive procedures which can be categorized into methods that repair (sphincteroplasty), augment (injectables, radiofrequency remodeling, stem cell therapy, and posterior anal sling), or replace (adynamic muscle transfers, dynamic graciloplasty, artificial bowel sphincter, and magnetic anal sphincter) the anal sphincter or neuromodulate bowel function (sacral nerve neuromodulation [SNM] and posterior tibial nerve stimulation) or divert fecal transit (stoma and antegrade stoma procedure)⁶. This article will focus on the most recent advances in the management of FI. [Table 2](#) shows a list of the advancements in the area and whether or not there are available published randomized trials to support them.

Anal insertion device

The use of anal plugs was described in the early 2000s. The concept is simple: the goal is to block the involuntary passage of

Table 1. Cleveland Clinic Florida-Fecal Incontinence Score.

Type of incontinence	Frequency				
	Never	Rarely	Sometimes	Usually	Always
Solid	0	1	2	3	4
Liquid	0	1	2	3	4
Gas	0	1	2	3	4
Wears pad	0	1	2	3	4
Lifestyle alteration	0	1	2	3	4

Never, 0; Rarely, <1/month; Sometimes, <1/week, ≥1/month; Usually, <1/day, ≥1/week; Always, ≥1/day. Minimum score 0 (perfect continence), maximum score 20 (complete incontinence).

Table 2. Advancements in the treatment of fecal incontinence and the availability of published randomized trials.

Recent advance	Randomized trial results available	Comments
Anal insert	Yes	Total of 136 patients. Two studies compared plug versus no plug, and two studies compared different types of plugs. Data limited and incomplete ^{7,8} .
Vaginal bowel control system	No	
Stem cell therapy	Yes	The largest trial included 24 patients (12 in the treatment arm) and showed superior results with stem cell therapy ¹⁷⁻²² .
Anal sling	No	
Sacral nerve neuromodulation (SNM)	Yes	See below.
Percutaneous tibial nerve stimulation	Yes	Not better than placebo or SNM
Magnetic anal sphincter augmentation	Pending	

stool from the anus. The device is self-inserted by patients and self-expelled or manually removed with a bowel movement. The data regarding the use of anal insertion devices for the treatment of FI are limited. A Cochrane Review in 2015⁷ found only four studies that met eligibility criteria and noted that plugs can be difficult to tolerate. In patients who are able to tolerate the device, there is limited evidence that plugs may be helpful in alleviating problems due to FI. The most recent device on the market, not included in the Cochrane Review, is a soft silicone anal insert⁸. In a multicenter prospective study, 77% of the 73 patients who completed the study and 62% of the 91 intention-to-treat patients achieved at least a 50% reduction in incontinence frequency over a 12-week period. The device, however, was used for patients with moderate to severe FI. It is the authors' opinion that patients with mild FI or leakage are most likely to benefit from this type of device (Figure 1).

Vaginal bowel control system

The VBS consists of a vaginal insert and a pressure-regulated pump introduced into the vagina (Figure 2). Patients are able to deflate the vaginal insert for defecation versus having to remove the device altogether, as is the case with the anal plug. In 2015, Richter *et al.*⁹ reported the results of 61 women successfully fitted and treated with the VBS. At 1 month, 79% of patients demonstrated treatment success and 41% reported complete continence. At 3 months, 86% reported success and

Figure 1. Renew insert device.

Figure 2. Vaginal bowel control system. The vaginal bowel control insert, folded and deflated for insertion (right) and inflated as it would sit in the vagina; the insert is inflated with a handheld pressure-regulated pump. Reprinted with permission from Pelvalon Inc. (Sunnyvale, CA, USA).

45% reported complete continence. The VBS device was found to be easy to use and to have a high degree of efficacy and relatively few adverse effects. It is limited by the fact that not all women can be fitted with the device. In the study by Richter *et al.*, only 54.5% of participants could be fitted with the device; however, once fitted, 96% found the device comfortable. More recently, the results of a multicenter, open-label trial including 73 patients with a mean CCF-FIS of 14.1 ± 12.15 showed a success rate of 72.6% at 3 months in the intention-to-treat analysis and of 84.1% in the per-protocol (patients able to keep the device) analysis. At 6 and 12 months, success remained high, improving to 90% and 94%, respectively, in the per-protocol population. There was significant improvement in the Fecal Incontinence Quality of Life score and the satisfaction rate reached over 94% at 12 months. This study showed that the VBS seemed to have a durable efficacy in patients who were successfully fitted and who had initial treatment response¹⁶.

Bulking agents

Injection of a bulking agent to augment the function of the anal sphincter for the treatment of FI was first described in 1993¹¹. The ideal agent should be biocompatible, non-immunogenic, and small enough to inject and have minimal migration of the agent¹. Multiple agents have been proposed over the years. A Cochrane Review in 2010¹² concluded that there was little evidence to support the use of bulking agents in the treatment of FI. In 2011, the US Food and Drug Administration (FDA) approved a non-animal-stabilized hyaluronic acid dextranomer gel for submucosal injection (NASHA Dx) for the treatment of FI. The largest series to date compared injection of NASHA Dx versus sham injection in a randomized, sham-controlled trial¹³. Again, the primary endpoint was a more than 50% reduction in the number of incontinence episodes and an increase in incontinence-free days. Eighty percent of patients in the NASHA Dx group actually had a second injection within 1 month after no improvement with the first injection. Those in the NASHA Dx group had a 52% reduction compared with 31% in the sham group ($P = 0.0089$). Despite this reduction, there was no significant difference in the FI scores between the groups at 2 months. In the published long-term follow-up¹⁴, the reduction in incontinence episodes was sustained at 12 (57%) and 36 (52%) months and the mean CCF-FIS was lower at 36 months. Despite improvement in FI, patients still had what is considered important FI (CCF-FIS of 11). NASHA Dx may be more beneficial for patients with minor, passive FI such as those with limited internal sphincter injury rather than for patients with more severe FI as in the case of most of the studies in the literature. Its final role in the treatment of FI, however, remains uncertain as a randomized trial comparing NASHA Dx with biofeedback did not show difference between the treatments¹⁵. A new bulking agent consisting of copolymer particle of polyacrylate-polyalcohol immersed in a carrier of glycerol and saline was described in 2015¹⁶. In total, 58 patients were enrolled in the study, which followed patients up to 36 months post-injection. Sixty percent of patients were successfully treated (with 50% more improvement in the CCF-FIS), although only 41.5% of patients achieved long-term follow-up to 36 months. Further research is needed to confirm these results and demonstrate the efficacy and safety of this new substance in the treatment

of FI. Injectable agents are contraindicated in patients with inflammatory bowel disease, rectocele, prolapse, previous pelvic irradiation, or anorectal malformations.

Stem cell therapy

Stem cell therapy has been evaluated for use in a variety of clinical settings such as cardiovascular, hematological, neurological, digestive, and trauma-associated conditions. The most commonly used stem cells are hematopoietic stem cells, mesenchymal stem cells, and adipose-derived stem cells¹⁷. The use of stem cells for the treatment of FI has been postulated as a means of providing recovery of sphincter function through regeneration of damaged striated sphincter muscle and allowing reinnervation of newly formed myofibers¹⁸. The early literature on stem cells in FI was based on animal models demonstrating the safety and efficacy of stem cells for FI^{18–20}. The first report of stem cell therapy and FI was by Lorenzi *et al.* in 2008²⁰. In this animal model of FI, the authors demonstrated that bone marrow–derived mesenchymal stem cell injection improved muscle regeneration and increased contractile function of anal sphincters after injury and repair.

The first study in humans was an observational study by Frudinger *et al.*²¹. They injected autologous myoblasts into the external anal sphincter in 10 women with lesions that had not been operated on and were refractory to conservative management. The authors demonstrated improved Wexner Incontinence Scores and quality of life at 12 months but did not demonstrate any significant physiological change to account for these improvements. These results were sustained at 5-year follow-up²². There were no significant side effects or adverse events noted in any patient. In 2015, Bisson *et al.*¹⁸ published a proof-of-concept study evaluating the injection of myoblasts into the anal sphincter of rats. The authors demonstrated sustained increase in sphincter pressures after injection. The follow-up, phase II, randomized, double-blind, placebo-controlled study in humans was published in 2018²³. In total, 24 patients were enrolled: 12 in the treatment arm and 12 in the control arm. At 6 months, both groups demonstrated reduction in the Cleveland Clinic Incontinence (CCI) score, but at 12 months, the treatment group continued to show improvement in the CCI score whereas the control arm did not. No severe adverse events were reported. Injection of stem cells appears to be safe, does not have major adverse side effects, and provides clinical benefit in the treatment of FI. To date, there have been no phase III trials comparing stem cell therapy with the SNM and this option of treatment is not currently approved by the FDA in the US.

Anal sling

Parks *et al.* emphasized the importance of the anorectal angle in the maintenance of continence²⁴. Based on this theory, the use of an artificial sling to support the posterior rectal wall has been developed for the treatment of FI^{25–27}. Most recently, Mellgren *et al.*²⁷ reported on the 1-year data of a prospective, multicenter study involving 152 patients at 14 centers in the US using the Transobturator Posterior Anal Sling (TOPAS) (American Medical Systems, Minnetonka, MN, USA). The mesh is placed behind the anorectum via two small incisions in the buttocks, and each arm of the mesh exits through the obturator foramen²⁸. The mechanism of action is not exactly clear but is

thought to support the puborectalis muscle and reinforce the anorectal angle. At 12 months, 69% of patients met the requirement for treatment success (at least 50% reduction in the FI episodes from baseline to 12 months post-operatively) and 19% reported complete continence. The posterior anal sling has not been compared with other treatment options for FI, such as SNM, and is not available in the US.

Sacral nerve neuromodulation

SNM was approved for use in the management of FI by the FDA in 2011 (Figure 3). SNM works by electrical stimulation of the sacral nerve roots, producing anal sphincter augmentation and modulation of spinal/supraspinal pathways²⁹. Wexner *et al.*³⁰ demonstrated the efficacy of this device in the treatment of FI in a prospective, multicenter study in the US. Success was defined as at least 50% reduction of incontinent episodes per week over 12 weeks in at least 50% of patients. The authors demonstrated an 83% therapeutic success at 12 months, and 41% of patients achieved 100% continence. In long-term follow-up at 3 years³¹, 83 patients were evaluated, success was demonstrated in 86% of patients, and 40% achieved complete continence. The most common reported adverse events were implant site pain (28%), paresthesias (15%), change in sensation of stimulation (12%), and infection (10%). The majority of adverse events (67%) occurred within the first year of implant. Of patients with infections, only six required surgical intervention: five device explants and one device replacement. A pooled analysis of all studies to date found that 69 to 83% of patients achieved success³².

For many years, the only option for patients with a sphincter defect of up to 120 degrees was sphincter repair, but results are known to deteriorate over time. Soon after its advent, SNM was also demonstrated to be effective in patients with FI in the setting of sphincter injury. Brouwer and Duthie³³ found that SNM provided significant improvement of FI even in the presence of a sphincter defect or pudendal neuropathy. In the largest study that evaluated this, 91 patients with no sphincter defect were compared with 54 patients with imaging-documented external defect. There was no significant difference in the comparison of baseline and 12-month CCF-FIS between the two groups. The ASCRS strongly recommends SNM as a first-line surgical option of FI in patients with or without sphincter defects.

Figure 3. Sacral nerve neuromodulation. Courtesy of Lucia Oliveira, Rio de Janeiro, Brazil.

SNM is a two-stage procedure involving a test phase followed by permanent implantation. The test phase is predictive of success with the permanent device. The test phase can be completed with placement of an external stimulator in the operating room (OR) and a subsequent trip to the OR for permanent implantation or by percutaneous nerve evaluation (PNE) in the office setting. PNE is commonly used among surgeons in Europe, whereas a two-stage procedure is used in the US. Rice *et al.*³⁴ compared PNE with the traditional staged approach used in the US. They demonstrated that success was similar between the two approaches (90.2% for staged and 82.2% for PNE, $P = 0.36$), and the 3-month CCF-FIS was not significantly different (4.4 versus 4.1, $P = 0.74$). However, they did find that the infection rate was higher in the staged approach (10.5% versus 0%, $P \leq 0.005$). On the other hand, a prospective randomized trial comparing one-stage to two-stage SNM for urinary incontinence found no difference in infection rates and significantly higher failure rates in patients undergoing the one-stage procedure³⁵. A systematic review by Maeda *et al.*³⁶ reported that the most common complication after PNE was lead displacement in 5.3% of patients and an overall complication rate of 6.4%, although 60% of studies did not report any complications related to PNE.

Percutaneous tibial nerve stimulation

Although SNM is the first-line surgical intervention for FI, it does currently require two trips to the OR and has associated high equipment costs. Alternatives to SNM involving neuromodulation have recently been investigated. Percutaneous tibial nerve stimulation (PTNS) is thought to cause similar changes in anorectal neuromuscular function as SNM because of the shared sacral segmental innervation³⁷. This treatment is not currently available in the US. PTNS is a minimally invasive technique that can be performed in the outpatient setting. The initial data come from several case series, but more recently, two randomized controlled trials have evaluated this therapy^{37,38}, one with placebo and one with SNM³⁹. In the CONFIDeNT trial³⁷, 227 patients were randomly assigned to PTNS versus sham PTNS over a 12-week period. There was no significant clinical benefit of PTNS over sham stimulation in the treatment of FI as measured by a more than 50% reduction in the number of FI episodes, although the absolute number of FI episodes per week was significantly reduced in the PTNS group.

An additional randomized control trial by Van der Wilt *et al.*³⁸ compared PTNS with sham stimulation. In total, 59 patients were included in the trial. The authors demonstrated a statistically significant reduction in the median and mean number of FI episodes of severity over 9 weeks. However, this was not clinically significant when response to treatment was based on a more than 50% reduction in FI episodes per week. A third trial³⁹ compared 17 patients undergoing SNM with 16 patients receiving PTNS and showed slightly superior results with SNM in the short term. All trials demonstrate that PTNS may be beneficial for some patients who have failed conservative management, but the trials have not established PTNS as a clinically meaningful treatment option, especially compared with SNM.

Magnetic anal sphincter augmentation

A few methods and devices have been developed for patients who have severe FI or major sphincter disruption and who have failed previous treatments. These have included dynamic graciloplasty and artificial bowel sphincter. Patients who have retained the devices experienced significant improvement, but owing to high rates of complications and explantations, these procedures are no longer available.

The magnetic anal sphincter augmentation (MAS) is a newer therapeutic option for patients with FI and was approved by the FDA in 2015 as a humanitarian use device. The MAS device is a ring of magnetic beads which is surgically implanted around the anal sphincter to reinforce weakened sphincter muscles. Several studies have reported promising short-term outcomes^{40,41}. A 2017 publication by Sugrue *et al.*⁴² reported on the long-term overall outcomes at a median follow-up of 5 years. A total of 35 patients underwent implantation of the device. Therapeutic success rates were reported as 63% at 1 year, 66% at 3 years, and 53% at 5 years. The device was explanted in seven patients because of major adverse events. In total, 30 adverse events ranging from pain to device erosion were reported in 20 patients. The authors state that the majority (73%) of these were minor events and require little to no intervention and occurred during the first year after implantation. Two ongoing randomized controlled trials are comparing MAS with SNM and hopefully will shed light as to where MAS fits into the treatment algorithm of FI^{43,44}.

Conclusions

At present, several therapies, ranging from medical management to more invasive surgical intervention, are offered for the management of FI. Not all therapies are appropriate for every patient. Therefore, it is important to select the correct therapy for the specific patient. For patients with mild FI, we recommend conservative and non-invasive adjuncts. Patients with minor leakage secondary to an internal sphincter defect may benefit from injectables. Those with moderate to severe FI should be evaluated for SNM. Lastly, those with severe FI or those who have failed several prior therapies may be good candidates for a trial of MAS prior to committing to a permanent stoma.

Abbreviations

ASCRS, American Society of Colon and Rectal Surgeons; CCF-FIS, Cleveland Clinic Florida-Fecal Incontinence Score;; FDA, CCI, Cleveland Clinic Incontinence; US Food and Drug Administration; FI, fecal incontinence; MAS, magnetic anal sphincter augmentation; NASHA Dx, non-animal-stabilized hyaluronic acid/dextranomer; OR, operating room; PNE, percutaneous nerve evaluation; PTNS, percutaneous tibial nerve stimulation; SNM, sacral nerve neuromodulation; VBS, vaginal bowel control system

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

1. **F** Paquette IM, Varma MG, Kaiser AM, *et al.*: **The American Society of Colon and Rectal Surgeons' Clinical Practice Guideline for the Treatment of Fecal Incontinence.** *Dis Colon Rectum.* 2015; **58**(7): 623–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
2. **NIH state-of-the-science conference statement on prevention of fecal and urinary incontinence in adults.** *NIH Consens State Sci Statements.* 2007; **24**(1): 1–37.
[PubMed Abstract](#)
3. Whitehead WE, Borrud L, Goode PS, *et al.*: **Fecal incontinence in US adults: epidemiology and risk factors.** *Gastroenterology.* 2009; **137**(2): 512–7, 517.e1–2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Shamlan T, Wyman J, Bliss DZ, *et al.*: **Prevention of urinary and fecal incontinence in adults.** *Evid Rep Technol Assess (Full Rep).* 2007; (161): 1–379.
[PubMed Abstract](#) | [Free Full Text](#)
5. Ditah I, Devaki P, Luma HN, *et al.*: **Prevalence, trends, and risk factors for fecal incontinence in United States adults, 2005–2010.** *Clin Gastroenterol Hepatol.* 2014; **12**(4): 636–43.e1–2.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Wexner SD: **Percutaneous tibial nerve stimulation in faecal incontinence.** *Lancet.* 2015; **386**(10004): 1605–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Deutekom M, Dobben AC: **Plugs for containing faecal incontinence.** *Cochrane Database Syst Rev.* 2012; (4): CD005086.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Lukacz ES, Segall MM, Wexner SD: **Evaluation of an Anal Insert Device for the Conservative Management of Fecal Incontinence.** *Dis Colon Rectum.* 2015; **58**(9): 892–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Richter HE, Matthews CA, Muir T, *et al.*: **A vaginal bowel-control system for the treatment of fecal incontinence.** *Obstet Gynecol.* 2015; **125**(3): 540–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. **F** Richter HE, Dunivan G, Brown HW, *et al.*: **A 12-Month Clinical Durability of Effectiveness and Safety Evaluation of a Vaginal Bowel Control System for the Nonsurgical Treatment of Fecal Incontinence.** *Female Pelvic Med Reconstr Surg.* 2019; **25**(2): 113–119.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
11. Shafik A: **Polytetrafluoroethylene injection for the treatment of partial fecal incontinence.** *Int Surg.* 1993; **78**(2): 159–61.
[PubMed Abstract](#)
12. Maeda Y, Laurberg S, Norton C: **Perianal injectable bulking agents as treatment for faecal incontinence in adults.** *Cochrane Database Syst Rev.* 2010; **12**(5): CD007959.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Graf W, Mellgren A, Matzel KE, *et al.*: **Efficacy of dextranomer in stabilised hyaluronic acid for treatment of faecal incontinence: a randomised, sham-controlled trial.** *Lancet.* 2011; **377**(9770): 997–1003.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. **F** Mellgren A, Matzel KE, Pollack J, *et al.*: **Long-term efficacy of NASHA Dx injection therapy for treatment of fecal incontinence.** *Neurogastroenterol Motil.* 2014; **26**(8): 1087–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
15. Dehli T, Stordahl A, Vatten LJ, *et al.*: **Sphincter training or anal injections of dextranomer for treatment of anal incontinence: a randomized trial.** *Scand J Gastroenterol.* 2013; **48**(3): 302–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. **F** Rosato G, Piccinini P, Oliveira L, *et al.*: **Initial results of a new bulking agent for fecal incontinence: a multicenter study.** *Dis Colon Rectum.* 2015; **58**(2): 241–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
17. **F** Trébol J, Carabias-Ortiz A, García-Arranz M, *et al.*: **Stem cell therapy for faecal incontinence: Current state and future perspectives.** *World J Stem Cells.* 2018; **10**(7): 82–105.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
18. Bisson A, Fréret M, Drouot L, *et al.*: **Restoration of anal sphincter function after myoblast cell therapy in incontinent rats.** *Cell Transplant.* 2015; **24**(2): 277–86.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Kang SB, Lee HN, Lee JY, *et al.*: **Sphincter contractility after muscle-derived stem cells autograft into the cryoinjured anal sphincters of rats.** *Dis Colon Rectum.* 2008; **51**(9): 1367–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Lorenzi B, Pessina F, Lorenzoni P, *et al.*: **Treatment of experimental injury of anal sphincters with primary surgical repair and injection of bone marrow-derived mesenchymal stem cells.** *Dis Colon Rectum.* 2008; **51**(4): 411–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Frudinger A, Kölle D, Schwaiger W, *et al.*: **Muscle-derived cell injection to treat anal incontinence due to obstetric trauma: pilot study with 1 year follow-up.** *Gut.* 2010; **59**(1): 55–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. **F** Frudinger A, Pfeifer J, Paede J, *et al.*: **Autologous skeletal-muscle-derived cell injection for anal incontinence due to obstetric trauma: a 5-year follow-up of an initial study of 10 patients.** *Colorectal Dis.* 2015; **17**(9): 794–801.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
23. **F** Boyer O, Bridoux V, Giverne C, *et al.*: **Autologous Myoblasts for the Treatment of Fecal Incontinence: Results of a Phase 2 Randomized Placebo-controlled Study (MIAS).** *Ann Surg.* 2018; **267**(3): 443–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
24. Parks AG, Porter NH, Hardcastle J: **The syndrome of the descending perineum.** *Proc R Soc Med.* 1966; **59**(6): 477–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Yamana T, Takahashi T, Iwadare J: **Perineal puborectalis sling operation for fecal incontinence: preliminary report.** *Dis Colon Rectum.* 2004; **47**(11): 1982–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. **F** Brochard C, Queralto M, Cabarro P, *et al.*: **Technique of the transobturator puborectal sling in fecal incontinence.** *Tech Coloproctol.* 2017; **21**(4): 315–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
27. **F** Mellgren A, Zutshi M, Lucente VR, *et al.*: **A posterior anal sling for fecal incontinence: results of a 152-patient prospective multicenter study.** *Am J Obstet Gynecol.* 2016; **214**(3): 349.e1–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
28. Rosenblatt P: **New developments in therapies for fecal incontinence.** *Curr Opin Obstet Gynecol.* 2015; **27**(5): 353–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. **F** Weledji EP: **Electrophysiological Basis of Fecal Incontinence and Its Implications for Treatment.** *Ann Coloproctol.* 2017; **33**(5): 161–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
30. Wexner SD, Collier JA, Devroede G, *et al.*: **Sacral nerve stimulation for fecal incontinence: results of a 120-patient prospective multicenter study.** *Ann Surg.* 2010; **251**(3): 441–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. **F** Mellgren A, Wexner SD, Collier JA, *et al.*: **Long-term Efficacy and Safety of Sacral Nerve Stimulation for Fecal Incontinence.** *Dis Colon Rectum.* 2011; **54**(9): 1065–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
32. **F** Thin NN, Horrocks EJ, Hotouras A, *et al.*: **Systematic review of the clinical effectiveness of neuromodulation in the treatment of faecal incontinence.** *Br J Surg.* 2013; **100**(11): 1430–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
33. **F** Brouwer R, Duthie G: **Sacral nerve neuromodulation is effective treatment for fecal incontinence in the presence of a sphincter defect, pudendal neuropathy, or previous sphincter repair.** *Dis Colon Rectum.* 2010; **53**(3): 273–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
34. **F** Rice TC, Quezada Y, Rafferty JF, *et al.*: **Dynamic Article: Percutaneous Nerve Evaluation Versus Staged Sacral Nerve Stimulation for Fecal Incontinence.** *Dis Colon Rectum.* 2016; **59**(10): 962–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
35. Everaert K, Kerckhaert W, Caluwaerts H, *et al.*: **A prospective randomized trial comparing the 1-stage with the 2-stage implantation of a pulse generator in patients with pelvic floor dysfunction selected for sacral nerve stimulation.** *Eur Urol.* 2004; **45**(5): 649–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Maeda Y, Matzel K, Lundby L, *et al.*: **Postoperative issues of sacral nerve stimulation for fecal incontinence and constipation: A systematic literature review and treatment guideline.** *Dis Colon Rectum.* 2011; **54**(11): 1443–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. **F** Knowles CH, Horrocks EJ, Bremner SA, *et al.*: **Percutaneous tibial nerve stimulation versus sham electrical stimulation for the treatment of faecal incontinence in adults (CONFIDeNT): A double-blind, multicentre, pragmatic, parallel-group, randomised controlled trial.** *Lancet.* 2015; **386**(1004): 1640–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
38. van der Wilt AA, Giuliani G, Kubis C, *et al.*: **Randomized clinical trial of percutaneous tibial nerve stimulation versus sham electrical stimulation in patients with faecal incontinence.** *Br J Surg.* 2017; **104**(9): 1167–76.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. **F** Thin NN, Taylor SJ, Bremner SA, *et al.*: **Randomized clinical trial of sacral versus percutaneous tibial nerve stimulation in patients with faecal incontinence.** *Br J Surg.* 2015; **102**(4): 349–58.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
40. **F** Wong MT, Meurette G, Stangherlin P, *et al.*: **The magnetic anal sphincter versus the artificial bowel sphincter: A comparison of 2 treatments for fecal incontinence.** *Dis Colon Rectum.* 2011; **54**(7): 773–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
41. Lehur PA, McNevin S, Buntzen S, *et al.*: **Magnetic anal sphincter augmentation for the treatment of fecal incontinence: A preliminary report from a feasibility**

study. *Dis Colon Rectum*. 2010; **53**(12): 1604–10.

[PubMed Abstract](#) | [Publisher Full Text](#)

42. **F** Sugrue J, Lehur PA, Madoff RD, *et al.*: Long-term Experience of Magnetic Anal Sphincter Augmentation in Patients With Fecal Incontinence. *Dis Colon Rectum*. 2017; **60**(1): 87–95.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
43. Williams AE, Croft J, Napp V, *et al.*: SaFaRI: Sacral nerve stimulation versus

the FENIX magnetic sphincter augmentation for adult faecal incontinence: a randomised investigation. *Int J Colorectal Dis*. 2016; **31**(2): 465–72.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

44. Lehur PA, Wyart V, Riche VP: SaFaRI: Sacral nerve stimulation versus the Fenix® magnetic sphincter augmentation for adult faecal incontinence: a randomised investigation. *Int J Colorectal Dis*. 2016; **31**(8): 1505.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Arnold Wald

Division of Gastroenterology and Hepatology, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Competing Interests: No competing interests were disclosed.

2. Stacy B. Menees

Division of Gastroenterology, University of Michigan, Ann Arbor, MI, USA

Competing Interests: No competing interests were disclosed.

3. Charles H. Knowles

National Bowel Research Centre, Blizard Institute, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, Queen Mary University of London, UK

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research