REVIEW

Interactions between the hippocampus, prefrontal cortex, and amygdala support complex learning and memory [version 1; peer review: 3 approved]

Ersin Yavas¹, Sarah Gonzalez¹, Michael S. Fanselow^{1,2}

¹Psychology, University of California, Los Angeles, Los Angeles, CA, 90095, USA

²Psychiatry and Biobehavioral Sciences, University of California, Los Angeles, Los Angeles, CA, 90095, USA

V1 First published: 31 Jul 2019, 8(F1000 Faculty Rev):1292
<https://doi.org/10.12688/f1000research.19317.1>

Latest published: 31 Jul 2019, 8(F1000 Faculty Rev):1292
<https://doi.org/10.12688/f1000research.19317.1>

Abstract

One of the guiding principles of memory research in the preceding decades is multiple memory systems theory, which links specific task demands to specific anatomical structures and circuits that are thought to act orthogonally with respect to each other. We argue that this view does not capture the nature of learning and memory when any degree of complexity is introduced. In most situations, memory requires interactions between these circuits and they can act in a facilitative manner to generate adaptive behavior.

Keywords

fear, learning, amygdala, hippocampus, prefrontal cortex

Open Peer Review

Approval Status

	1	2	3
version 1			
31 Jul 2019			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Shantanu P Jadhav**, Brandeis University, Massachusetts, USA
2. **Stephen Maren**, Texas A&M University, Texas, USA
3. **Mouna Maroun**, University of Haifa, Haifa, Israel

Any comments on the article can be found at the end of the article.

Corresponding author: Michael S. Fanselow (fanselow@ucla.edu)

Author roles: **Yavas E:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Gonzalez S:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Fanselow MS:** Conceptualization, Funding Acquisition, Resources, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: MSF is Director of Research at Neurovation Labs. EY and SG declare that they have no competing interests.

Grant information: This work was supported by National Institute of Mental Health grant RO1MH62122, National Science Foundation Division of Graduate Education grant 1650604, and the Staglin Center for Brain and Behavioral Health.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 Yavas E *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Yavas E, Gonzalez S and Fanselow MS. **Interactions between the hippocampus, prefrontal cortex, and amygdala support complex learning and memory [version 1; peer review: 3 approved]** F1000Research 2019, 8(F1000 Faculty Rev):1292 <https://doi.org/10.12688/f1000research.19317.1>

First published: 31 Jul 2019, 8(F1000 Faculty Rev):1292 <https://doi.org/10.12688/f1000research.19317.1>

Introduction

There is no doubt that most animal species depend on the ability to learn for survival. Learning and the behaviors through which learning is expressed require interactions between large-scale brain networks. In mammals, simple forms of learning, such as when X perfectly predicts Y, often can be handled by subcortical networks. Take, for example, basic Pavlovian fear conditioning in which a tone signals aversive shock 100% of the time and shock never occurs in the absence of tone. In this case, projections from auditory thalamus to amygdala to midbrain periaqueductal gray are sufficient for learning¹. However, as learning becomes more complex, memory and recall depend on interactions between the cortex and hippocampus and through those interactions these structures modulate the subcortical circuits that support simple learning. Healthy behavior depends on these interactions and abnormal structure and function in these regions are associated with a variety of neuropathology and psychopathology, including anxiety disorders, schizophrenia, and Alzheimer's dementia²⁻⁴. Although the roles of both the hippocampus and frontal cortex in regulating complex behavior have been studied extensively, much has been learned recently about how the interactions between these regions impact learning and memory-related functions.

When are hippocampal–cortical interactions necessary for memory?

Traditionally, the relationship between brain structures and forms of memory has been conceptualized under a multiple memory systems taxonomy, in which specific structures are linked to specific classes of learned behavior⁵. In this taxonomy, the hippocampus is associated with episodic memory, the striatum with skill learning, the neocortex with perceptual learning, and the amygdala with emotional memory. Although this taxonomy has been a major impetus to memory research and spurred considerable advancement in our understanding of learning, it fosters the idea that there is a simple one-to-one correspondence between a single structure and a type of memory and that these separate systems act in an orthogonal manner⁶. Such a view fails to capture the fact that even modestly complex memory emerges from a network of interactions between these structures.

Complex versus simple stimuli

Above we described fear conditioning, in which a tone perfectly predicts shock, as an example of simple learning that can be carried out by a subcortical circuit. However, when experiencing fear conditioning, the subject simultaneously learns that the context or place where this learning occurs is also associated with shock, enabling that context to trigger fear and avoidance⁷. Contexts are complex stimuli made of many features, all of which may lack the salience of the sudden onset of a simple auditory stimulus such as a tone⁸. The context is also less precisely paired with the stimulus that reinforces learning (in this case, the shock). This modest increase in complexity cannot be handled by the simple subcortical circuit and instead requires an orchestrated dance between hippocampus, prefrontal cortex, and amygdala⁹.

Consistent with their role in spatial memory, dorsal hippocampal neurons appear to encode the details of the context supporting recognition of the trained environment. Two groups of rats were placed in a novel chamber for 5 min where they received shock. One group received the shock in a manner that promoted fear learning (at the end of the session) and the other in a manner that did not promote fear learning (at the beginning of the session). When these rats were returned to the same chamber neurons that were active during the initial exposure tended to reactivate during the second exposure indicating that the hippocampus recognized the chamber as familiar. Interestingly, the same number of neurons reactivated in both groups even though they showed large differences in fear expression, suggesting that hippocampus is not essential for the coding of affective features of the context⁹. On the other hand, basolateral amygdala neurons reactivate only if the animal has learned fear of the context, indicating that the amygdala is involved in the affective component of memory.

These two sources of information appear to come together in the prelimbic portion of the prefrontal cortex as activity in that region reflects a sum of that found in hippocampus and amygdala⁹. In support of this, coordinated 4-Hz rhythmic activity between the prelimbic and amygdala regions accurately predicts freezing, a behavioral manifestation of fear¹⁰. Coordinated oscillatory activity between hippocampus and prelimbic regions may be the method by which the hippocampus provides spatial information to the prelimbic cortex¹¹.

The dynamics between the hippocampus, cortex, and amygdala depend on the length of memory retention^{7,12}. Shortly after learning, the memory is dependent on the hippocampus and there is more hippocampal activity during memory retrieval¹³. However, at later time points (weeks to months), the prefrontal cortex takes on the larger role. A recent study indicated that the development of this prefrontal representation of contextual memory is dependent on inputs from the entorhinal cortex, hippocampus, and basolateral amygdala during learning¹⁴. The changes that occur over these long time periods are referred to as systems consolidation.

This coordination of regions during systems consolidation may occur off-line during the replay of events that occurs during sleep¹⁵. The integration of spatial information with emotional memory may also occur during non-REM sleep when there is a coordinated replay of the amygdala and hippocampal activity that occurred during learning¹⁶.

Interestingly, the prefrontal cortex can compensate for loss of hippocampal function to a certain degree, but the resulting memories differ from those formed using an intact hippocampus. Following damage to the dorsal hippocampus, contextual fear memories can be formed after extensive training but these memories lack the permanence of those formed using an intact hippocampus^{17,18}. This compensation depends on communication between the infralimbic and prelimbic regions of the prefrontal cortex. Damage to and disconnection between these two

adjacent prefrontal regions abolish contextual fear learning entirely when combined with dorsal hippocampal damage¹⁸.

Certain versus uncertain predictors

While learning that an auditory stimulus consistently signals shock can be accomplished by a subcortical circuit, this changes when the relationship between the tone and shock becomes ambiguous. The simplest and most translationally relevant example of this is fear extinction, in which a stimulus that previously received consistent reinforcement is now presented without reinforcement. Because of this inconsistent reinforcement, the animal treats the tone as ambiguous and uses context to resolve the decision of whether or not to respond¹⁹. This decision also relies on interactions between hippocampus, prefrontal cortex, and amygdala. When communication between the ventral hippocampus and prelimbic cortex or between the ventral hippocampus and amygdala is prevented, the subject treats the tone as safe regardless of context^{20,21}. If these connections are intact but communication between the hippocampus and infralimbic cortex or between the amygdala and infralimbic cortex is prevented, the animal treats the tone as consistently dangerous^{22,23}. Additionally, infralimbic lesions lead to greater generalization between two contexts and simultaneously reduce the context's ability to modulate responses to an extinguished stimulus¹⁸. Recent evidence suggests that the hippocampus may help solve the problem by encoding two distinct contextual memories of the same context: one where the tone is dangerous and one where it is safe²⁴. This may allow the rapid switching that rats can accomplish between reacting to the tone as dangerous or safe on the basis of the current context²⁵.

Such a role for the prefrontal cortex in resolving ambiguity has also been shown in contextual biconditional discrimination procedures, which require subjects to use contextual information to resolve the meaning of ambiguous stimuli. In these procedures,

two different stimuli are presented in two different contexts; one stimulus is predictive of an outcome in one context whereas the second stimulus is predictive of that outcome in the second context^{26–28}. Lesions of the prelimbic cortex impair the ability of subjects to use contextual stimuli to determine which of two auditory stimuli predicts footshock, instead producing an intermediate level of fear to both stimuli²⁸.

Conclusions

Learning is often conceptualized as occurring in distinct and independently acting brain regions. However, to navigate a complex and ever-changing environment, animals must be able to learn about complex stimuli and complex relationships between stimuli. In these instances, mammals use environmental cues to resolve the meaning of ambiguous stimuli. Rather than being dependent on any one brain region, this ability arises from a network of cortical and subcortical structures and their interactions. Here, we focused on Pavlovian conditioning because it is often thought of as a simple form of learning. However, introducing even slight complexities in this learning, such as those that occur when contexts are used as stimuli or following the change in meaning that occurs during extinction, recruits and requires long-range interactions between regions that are often thought to serve distinct domains of memory.

Grant information

This work was supported by National Institute of Mental Health grant RO1MH62122, National Science Foundation Division of Graduate Education grant 1650604, and the Staglin Center for Brain and Behavioral Health.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Romanski LM, LeDoux JE: **Equipotentiality of thalamo-amygdala and thalamo-cortico-amygdala circuits in auditory fear conditioning.** *J Neurosci.* 1992; **12**(11): 4501–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mennesson M, Rydgren E, Lipina T, *et al.*: **Kainate receptor auxiliary subunit NETO2 is required for normal fear expression and extinction.** *Neuropsychopharmacol.* 2019.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Tellez-Merlo G, Morales-Medina JC, Camacho-Ábreo I, *et al.*: **Prenatal immune challenge induces behavioral deficits, neuronal remodeling, and increases brain nitric oxide and zinc levels in the male rat offspring.** *Neuroscience.* 2019; **406**: 594–605.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Nuzzo T, Feligioni M, Cristino L, *et al.*: **Free d-aspartate triggers NMDA receptor-dependent cell death in primary cortical neurons and perturbs JNK activation, Tau phosphorylation, and protein SUMOylation in the cerebral cortex of mice lacking d-aspartate oxidase activity.** *Exp Neurol.* 2019; **317**: 51–65.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Squire LR, Zola SM: **Structure and function of declarative and nondeclarative memory systems.** *Proc Natl Acad Sci U S A.* 1996; **93**(24): 13515–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- McDonald RJ, White NM: **A triple dissociation of memory systems: hippocampus, amygdala, and dorsal striatum.** *Behav Neurosci.* 1993; **107**: 3–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kim JJ, Fanselow MS: **Modality-specific retrograde amnesia of fear.** *Science.* 1992; **256**(5057): 675–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fanselow MS: **Contextual fear, gestalt memories, and the hippocampus.** *Behav Brain Res.* 2000; **110**(1–2): 73–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zelikowsky M, Hersman S, Chawla MK, *et al.*: **Neuronal ensembles in amygdala, hippocampus, and prefrontal cortex track differential components of contextual fear.** *J Neurosci.* 2014; **34**(25): 8462–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Karalis N, Dejean C, Chaudun F, *et al.*: **4-Hz oscillations synchronize prefrontal-amygdala circuits during fear behavior.** *Nat Neurosci.* 2016; **19**(4): 605–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Shin JD, Jadhav SP: **Multiple modes of hippocampal-prefrontal interactions**

- in memory-guided behavior. *Curr Opin Neurobiol.* 2016; **40**: 161–169.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
12. Wiltgen BJ, Brown RA, Talton LE, *et al.*: **New circuits for old memories: the role of the neocortex in consolidation.** *Neuron.* 2004; **44**(1): 101–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 13. Lopez J, Herbeaux K, Cosquer B, *et al.*: **Context-dependent modulation of hippocampal and cortical recruitment during remote spatial memory retrieval.** *Hippocampus.* 2012; **22**(4): 827–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
 14. Kitamura T, Ogawa SK, Roy DS, *et al.*: **Engrams and circuits crucial for systems consolidation of a memory.** *Science.* 2017; **356**(6333): 73–78.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 15. Penagos H, Varela C, Wilson MA: **Oscillations, neural computations and learning during wake and sleep.** *Curr Opin Neurobiol.* 2017; **44**: 193–201.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 16. Girardeau G, Inema I, Buzsáki G: **Reactivations of emotional memory in the hippocampus-amygdala system during sleep.** *Nat Neurosci.* 2017; **20**(11): 1634–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 17. Zelikowsky M, Bissiere S, Fanselow MS: **Contextual fear memories formed in the absence of the dorsal hippocampus decay across time.** *J Neurosci.* 2012; **32**(10): 3393–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 18. Zelikowsky M, Bissiere S, Hast TA, *et al.*: **Prefrontal microcircuit underlies contextual learning after hippocampal loss.** *Proc Natl Acad Sci U S A.* 2013; **110**(24): 9938–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 19. Bouton ME: **Context, ambiguity, and unlearning: sources of relapse after behavioral extinction.** *Biol Psychiatry.* 2002; **52**(10): 976–86.
[PubMed Abstract](#) | [Publisher Full Text](#)
 20. Jin J, Maren S: **Prefrontal-Hippocampal Interactions in Memory and Emotion.** *Front Syst Neurosci.* 2015; **9**: 170.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 21. Orsini CA, Kim JH, Knapska E, *et al.*: **Hippocampal and prefrontal projections to the basal amygdala mediate contextual regulation of fear after extinction.** *J Neurosci.* 2011; **31**(47): 17269–77.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 22. Adhikari A, Lerner TN, Finkelstein J, *et al.*: **Basomedial amygdala mediates top-down control of anxiety and fear.** *Nature.* 2015; **527**(7577): 179–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 23. Marek R, Jin J, Goode TD, *et al.*: **Hippocampus-driven feed-forward inhibition of the prefrontal cortex mediates relapse of extinguished fear.** *Nat Neurosci.* 2018; **21**(3): 384–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 24. Lacagnina AF, Brockway ET, Crovetti CR, *et al.*: **Distinct hippocampal engrams control extinction and relapse of fear memory.** *Nat Neurosci.* 2019; **22**(5): 753–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 25. Bouton ME, Swartzentruber D: **Analysis of the associative and occasion-setting properties of contexts participating in a Pavlovian discrimination.** *Journal of Experimental Psychology: Animal Behavior Processes.* 1986; **12**(4): 333–50.
[Publisher Full Text](#)
 26. Haddon JE, Killcross S: **Prefrontal cortex lesions disrupt the contextual control of response conflict.** *J Neurosci.* 2006; **26**(11): 2933–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
 27. Gonzalez ST, Welch ES, Colwill RM: **Pavlovian contextual and instrumental biconditional discrimination learning in mice.** *Behav Brain Res.* 2013; **256**: 398–404.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 28. Sharpe MJ, Killcross S: **The prelimbic cortex uses higher-order cues to modulate both the acquisition and expression of conditioned fear.** *Front Syst Neurosci.* 2015; **8**: 235.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Mouna Maroun**

Sagol Department of Neurobiology, Faculty of Natural Sciences, University of Haifa, Haifa, Israel

Competing Interests: No competing interests were disclosed.

2. **Stephen Maren**

Department of Psychological and Brain Sciences, Texas A&M University, Texas, USA

Competing Interests: No competing interests were disclosed.

3. **Shantanu P Jadhav**

Neuroscience Program, Department of Psychology and Volen National Center for Complex Systems, Brandeis University, Massachusetts, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research