REVIEW

Vasopressin-aquaporin-2 pathway: recent advances in understanding water balance disorders [version 1; peer review: 3 approved]

Marianna Ranieri ¹, Annarita Di Mise¹, Grazia Tamma^{1,2}, Giovanna Valenti ¹⁻³

¹Department of Biosciences, Biotechnologies and Biopharmaceutics, University of Bari, Bari, Italy, 70125, Italy

²Istituto Nazionale di Biostrutture e Biosistemi, Rome, Roma, Italy, 00136, Italy

³Center of Excellence in Comparative Genomics (CEGBA), University of Bari, Bari, Italy, 70125, Italy

V1 First published: 04 Feb 2019, 8(F1000 Faculty Rev):149
<https://doi.org/10.12688/f1000research.16654.1>

Latest published: 04 Feb 2019, 8(F1000 Faculty Rev):149
<https://doi.org/10.12688/f1000research.16654.1>

Abstract

The alteration of water balance and related disorders has emerged as being strictly linked to the state of activation of the vasopressin-aquaporin-2 (vasopressin-AQP2) pathway. The lack of responsiveness of the kidney to the vasopressin action impairs its ability to concentrate the urine, resulting in polyuria, polydipsia, and risk of severe dehydration for patients. Conversely, non-osmotic release of vasopressin is associated with an increase in water permeability in the renal collecting duct, producing water retention and increasing the circulatory blood volume. This review highlights some of the new insights and recent advances in therapeutic intervention targeting the dysfunctions in the vasopressin-AQP2 pathway causing diseases characterized by water balance disorders such as congenital nephrogenic diabetes insipidus, syndrome of inappropriate antidiuretic hormone secretion, nephrogenic syndrome of inappropriate antidiuresis, and autosomal dominant polycystic kidney disease. The recent clinical data suggest that targeting the vasopressin-AQP2 axis can provide therapeutic benefits in patients with water balance disorders.

Keywords

Vasopressin, AQP2, NDI, SIADH, NSIAD, ADPKD

Open Peer Review

Approval Status

	1	2	3
version 1 04 Feb 2019			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Enno Klusmann**, Max Delbrück Center for Molecular Medicine Berlin (MDC), Berlin, Germany
2. **Kenichi Ishibashi**, Meiji Pharmaceutical University, Tokyo, Japan
3. **Tae-Hwan Kwon**, Kyungpook National University, Taegu, South Korea

Any comments on the article can be found at the end of the article.

Corresponding author: Giovanna Valenti (giovanna.valenti@uniba.it)

Author roles: **Ranieri M:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Di Mise A:** Writing – Original Draft Preparation; **Tamma G:** Writing – Original Draft Preparation; **Valenti G:** Conceptualization, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This study was supported in part by Telethon funding (grant number GGP13227) and by ASI (Italian Space Agency, grant number 2013-091-R.0). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 Ranieri M *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Ranieri M, Di Mise A, Tamma G and Valenti G. **Vasopressin-aquaporin-2 pathway: recent advances in understanding water balance disorders [version 1; peer review: 3 approved]** F1000Research 2019, 8(F1000 Faculty Rev):149 <https://doi.org/10.12688/f1000research.16654.1>

First published: 04 Feb 2019, 8(F1000 Faculty Rev):149 <https://doi.org/10.12688/f1000research.16654.1>

Introduction

The maintenance of water balance is essential for all physiological processes and is critically dependent on water intake via thirst and water output in the kidney under the control of the antidiuretic hormone vasopressin. An increase in plasma osmolality, sensed by osmoreceptors situated in the brain, represents the most important input to cause thirst and stimulation of vasopressin release. Vasopressin is secreted into the circulation by the posterior pituitary gland in response to an increase in serum osmolality or a decrease in blood volume. In the kidney, vasopressin binds to the type 2 vasopressin receptor (V2R) and increases osmotic water transport through the regulation of the aquaporin-2 (AQP2) water channel localized in the kidney connecting tubules and collecting ducts^{1,2}. V2R is a G-protein-coupled receptor (GPCR) localized at the basolateral plasma membrane of the principal cells of the kidney collecting duct. A recent transcriptome study of 14 microdissected nephron segments of rat kidney demonstrated the V2R mRNA expression from the medullary thick ascending limb to the inner medullary collecting duct³. Upon binding of vasopressin to V2R, a G_s protein is activated, leading to stimulation of adenylyl cyclases, increase in intracellular cAMP, and activation of protein kinase A (PKA). The cAMP/PKA signal transduction cascade results in multiple phosphorylating events in the C-terminus of the water channel AQP2 regulating its trafficking and the water luminal permeability^{2,4-6}. Vasopressin also triggers increases in intracellular calcium required for AQP2 trafficking^{7,8}.

Most of the effect of vasopressin is thought to be related to PKA-mediated phosphorylation currently explored by large-scale phosphoproteomics to identify regulated proteins' downstream PKA activation⁹. On the other hand, several proteins participating in the control of cAMP-dependent AQP2 trafficking, including SNAREs, annexin-2, hsc70, A-kinase anchoring proteins (AKAPs), and small GTPases of the Rho family proteins controlling cytoskeletal dynamics, have been identified¹⁰⁻¹⁸. In addition to phosphorylation, AQP2 undergoes different regulated post-translational modifications, such as ubiquitination and glutathionylation, which are likely to be fundamental for controlling AQP2 cellular localization, stability, and function¹⁹⁻²³.

Besides short-term regulation of AQP2 trafficking, vasopressin regulates the total amount of the water channel within the cell and alters the protein half-life of AQP2²⁴⁻²⁹. Alterations in AQP2 abundance as well as defects in vasopressin signaling in the renal collecting duct can seriously compromise renal function and the maintenance of water balance in the body.

This review highlights some of the new insights and recent advances in targeting the vasopressin–AQP2 pathway in some relevant diseases associated with water balance disorders, such as congenital nephrogenic diabetes insipidus (NDI), idiopathic syndrome of inappropriate antidiuretic hormone secretion (SIADH), nephrogenic syndrome of inappropriate antidiuresis (NSIAD), and autosomal dominant polycystic kidney disease (ADPKD).

Vasopressin–AQP2 pathway in water balance disorders Plasma copeptin as a surrogate marker of vasopressin secretion in renal disorders

Significant progress in studying the role of vasopressin in renal disorders came from the identification of copeptin, a stable surrogate marker of vasopressin secretion that is relatively easily measured³⁰⁻³³. Copeptin is a peptide corresponding to the COOH-terminal portion of pro-vasopressin and is co-secreted in equimolar amounts with vasopressin, representing a good biomarker for vasopressin³⁴. Indeed, it has been reported that plasma concentrations of copeptin correlate strongly in several clinical conditions^{32,33,35,36}. In a large cross-sectional study, plasma copeptin and microalbuminuria positively correlated³⁷, and this correlation was also found to persist with a 16-year follow-up³⁸. The increase in urinary albumin excretion is likely to reflect not only glomerular damage but also systemic endothelial dysfunction and is consistent with the hypothesis that vasopressin induces urinary albumin excretion as previously reported in rats and humans^{39,40}. Conversely, suppressing vasopressin by administering a V2R antagonist or by simply increasing water intake might be beneficial for renal function and diabetes^{39,41}.

As shown previously, vasopressin levels are also increased in diabetic nephropathy characterized by dysregulation of water balance displaying water depletion as a consequence of osmotic diuresis due to glycosuria⁴², probably to limit water loss. Of interest, in patients with type 2 diabetes, plasma copeptin was found to be associated with a faster decline in glomerular filtration rate (GFR) in two distinct studies^{43,44}.

The role of vasopressin is particularly central in the pathogenesis of another severe disease, ADPKD, characterized by the expansion of renal cysts eventually leading to loss of renal function. Association of urinary copeptin with the severity of ADPKD was also recently demonstrated⁴⁴, suggesting that copeptin can represent a novel marker to predict renal prognosis in ADPKD. Copeptin levels are negatively associated with GFR, kidney size, and number of renal cysts^{33,45,46}. While ADPKD is the most advanced disease for the therapeutic use of vasopressin receptor blockade, this strategy is also currently being explored in chronic kidney diseases (CKDs).

In summary, the vasopressin–AQP2 system has a critical role in various stages of CKD⁴⁷⁻⁴⁹ and in several kidney diseases⁴⁶, making this pathway very promising from a therapeutic perspective. Indeed, high water intake (particularly plain water) has been proven to be beneficial in preventing CKD progression in several studies (reviewed in 46), based on the expected suppression of vasopressin release that can be easily monitored by plasma copeptin measurements. On the other hand, the association between total fluid intake (excluding water intake) and change in kidney function requires additional prospective studies, since other data do not support the hypothesis that total fluid consumption (excluding water intake) protects against CKD⁵⁰; therefore, additional study with prospective evaluation of kidney function is required.

Targeting defective vasopressin–AQP2 pathway in nephrogenic diabetes insipidus

NDI is a disease characterized by the kidney's inability to concentrate urine despite elevated concentrations of vasopressin. The congenital form of NDI can be due to mutation of either the vasopressin receptor or AQP2 and is associated with marked polyuria, polydipsia, and electrolyte imbalance with a constant risk of severe dehydration for patients^{51,52}. Most of the forms of NDI are due to non-functional V2R (X-linked NDI)⁵³. X-linked NDI accounts for 90% of cases of congenital NDI and occurs with a frequency of 4 to 8 per 1 million male live births. Autosomal NDI accounts for about 10% of the remaining cases⁵⁴.

NDI can also be acquired and is a frequent side effect of lithium therapy⁵⁵ or other drugs^{56–60}. Hypokalemia or hypercalcemia with associated hypercalciuria also cause acquired NDI^{28,61–68}.

The year 2013 was the 100th anniversary of vasopressin treatment for central diabetes insipidus due to vasopressin hormone deficiency⁶⁹. Despite extensive research in the field, a real cure for NDI is still missing and treatment is rather symptomatic with a continuous supply of water and drugs such as ibuprofen and indomethacin in combination with hydrochlorothiazide, which however only partially (by 25% to 50%) reduce the abundant polyuria^{51,68,70}.

The therapeutic approaches under investigation regard the rescue of mutated V2R or AQP2, in case of defective trafficking of these proteins, or bypassing the V2R signaling, in case of inactive V2Rs, finalized to increasing the cell surface expression of AQP2^{51,71–73}.

In patients with NDI due to V2R mutations or as a consequence of lithium therapy, AQP2 is in fact theoretically functioning, suggesting the possibility of increasing its apical membrane abundance independently from vasopressin or cAMP. Some studies suggest that this may be possible. Sildenafil, a phosphodiesterase inhibitor causing an increase in cyclic guanosine monophosphate, reduced polyuria in rats with lithium-induced NDI⁷⁴. Moreover, a child affected by X-linked NDI treated with sildenafil had a beneficial effect in decreasing urine volume and increased urine osmolality⁷⁵.

Another approach to bypass defective V2R was altering the cholesterol content of the apical membrane to increase AQP2 accumulation. To this end, different statins have been used. Statins are a family of 3-hydroxy-3-methylglutaryl-CoA reductase inhibitors commonly used to inhibit cholesterol biosynthesis. Besides this, *in vitro* studies showed that statins reduced endocytosis possibly by inhibiting isoprenylation of Rho GTPase, which leads to actin cytoskeletal reorganization during protein trafficking⁷⁶. Of note, in renal cells and *in vivo*, statins have been shown to promote AQP2 localization to the apical membrane^{77,78}, a process known to require apical depolymerization of the actin network and Rho inhibition^{11,12}.

Interestingly, a combination of two statins—fluvastatin and secretin—drastically reduced urine output by nearly 90% and doubled

urine osmolality in mouse models of NDI in a mouse model⁷⁹. Statin treatment was also found to be beneficial in NDI patients with inactivating mutation of the V2R⁷⁸ and in patients under lithium therapy⁸⁰. Besides the effect in reducing urinary output in patients with NDI, the efficacy of the treatment with statins in increasing apical expression of AQP2 has been monitored by measuring the urinary excretion of AQP2 (u-AQP2), which was found to be increased in a time- and dose-dependent manner⁸¹.

The most recent approach for treating congenital X-linked NDI is based on treatment with metformin, a drug used for the treatment of diabetes which is able to activate the adenosine monophosphate kinase (AMPK)^{82,83}. In the renal inner medullary collecting duct, AMPK phosphorylates AQP2 and urea transporter UT-A1, resulting in water and urea reabsorption⁸³, and improves renal concentrating ability in X-linked NDI mice⁸². Therefore, AMPK activation with metformin might represent an alternative therapeutic approach to treat X-linked NDI. More recently, it was shown that inhibition of AKAPs binding to PKA increases PKA activity and activates AQP2 channels in cortical collecting duct cells⁸⁴. In line with this finding, *in vivo* experiments revealed that the compound 3,3'-diamino-4,4'-dihydroxydiphenylmethane (FMP-API-1) induced AQP2 phosphorylation, trafficking, and water reabsorption to the same extent as vasopressin in the context of V2R inhibition. This study points to AKAP-PKA disruptors as a potential novel category of therapeutic drugs for congenital NDI⁸⁴.

Targeting selected GPCRs with agonists to increase cAMP has also been considered an option to treat NDI. Among the GPCRs expressed in the inner medulla collecting duct, activation of the calcitonin receptor caused an increase in cAMP and accumulation of AQP2 in the plasma membrane in LLC-PK1 cells⁸⁵ and reduced urinary output in vasopressin-deficient Brattleboro rats⁷⁹. Other GPCRs whose activation was found to be associated with increased AQP2 expression are the angiotensin II AT1 receptor⁸⁶ and secretin receptor^{79,87}. However, it has to be underlined that *in vivo* testing of this approach targeting GPCRs revealed that the effect is transient and this is probably due to desensitization of the receptors^{79,85}. In parallel, four other GPCRs, the protein-coupled E-prostanoid receptors EP1–EP4, have been considered for their ability to increase cAMP and AQP2 in MDCK cells⁸⁸. Selective silencing of EP4 in mice resulted in reduced renal concentrating ability⁸⁹. Interestingly, in renal collecting duct cell models, EP4 activation increased AQP2 trafficking independently from cAMP elevation^{73,90}, although the mechanisms responsible for EP4-induced stimulation of AQP2 trafficking are not yet clarified.

The calcium signaling pathway in the vasopressin response has also been considered a major target in the treatment of NDI. Specifically, activators of calcium signaling in collecting duct principal cells may represent a therapeutic strategy in NDI. In a recent seminal study, Uchida *et al.* demonstrated that activation of the calcium signal transducer Wnt5a, through a calcium/calmodulin/calcineurin signaling pathway, induced phosphorylation, trafficking, and mRNA expression of AQP2 in mpkCCDC14 cells⁹¹. Of note, in isolated cortical collecting ducts from a V2R-inhibited NDI mouse model, Wnt5a increased osmotic water permeability and urine osmolality, activating a different pathway of the

vasopressin-induced cAMP elevation, involving a functional role of calcineurin/arachidonic acid known to induce vasopressin-like effects in mpkCCD cells. These new data point to calcineurin activators as possible drugs for the treatment of congenital NDI.

Vasopressin–AQP2 pathway in syndrome of inappropriate antidiuretic hormone secretion and nephrogenic syndrome of inappropriate antidiuresis

SIADH may be considered an opposite disease with respect to NDI. Patients with SIADH have high levels of vasopressin even in the presence of low serum osmolality, leading to dilutional hyponatremia secondary to elevated renal water reabsorption^{92,93}.

Hyponatremia is defined as a serum Na level below 135 mmol/L and is associated with an increase in mortality in hospitalized patients⁹⁴. In SIADH, high levels of vasopressin are present even when plasma osmolality reaches values that normally suppress vasopressin release and the resulting hyponatremia can be ascribed to a non-osmotic release of vasopressin⁹⁵. Nevertheless, a phenomenon called “vasopressin escape” allows the kidney to excrete free water through reduced expression of the gene encoding AQP2, thus increasing water excretion^{96,97}. Interestingly, a recent study uncovered some signaling mechanisms that defend against hyponatremia in SIADH²¹. A SIADH-like condition is frequently related to aging⁹³. SIADH has several major etiologies, including central nervous system and pulmonary disorders, tumors, and drugs, the last of which is related mainly to psychopharmacological treatment. An interesting observation is that treatment with antipsychotics is associated with elevated activity of calcineurin and enhanced vasopressin release, which may contribute to activation of AQP2 trafficking causing drug-induced SIADH. Moreover, cyclophosphamide, an anti-cancer drug, activates V2R and induces AQP2 expression in rat kidney despite the absence of vasopressin stimulation⁹⁸. This finding suggested the possibility of drug-induced NSIAD. The syndrome of inappropriate antidiuresis has been suggested as an alternative to SIADH, since a subgroup of patients with features of SIADH does not have high plasma vasopressin levels⁹⁹. Therefore, dilutional hyponatremia may result from either excessive AVP release or constitutive activation of V2R.

The use of antagonists of the vasopressin receptors (vaptans) represents the most direct treatment for hyponatremia in SIADH. In this respect, over the last 10 years, significant advances in the treatment of SIADH have been made¹⁰⁰.

Among vaptans, conivaptan represents the first vasopressin receptor blocker approved by the US Food and Drug Administration⁹³. Conivaptan is given intravenously and acts on both V1 and V2 vasopressin receptors, resulting in solute-free water diuresis (aquaresis). Tolvaptan instead is orally active and is a selective antagonist of the V2R. It has been shown to be effective and safe for up to 3 years¹⁰¹. Both conivaptan and tolvaptan are approved for the treatment of hypervolemic hyponatremia in patients with heart failure or cirrhosis and of euvolemic hyponatremia in patients with SIADH. We recently demonstrated that tolvaptan prevents AQP2 trafficking and function in collecting duct principal cells and reduces AQP2 excretion in two patients with SIADH paralleled by normalization

of plasma sodium concentration, clearly demonstrating the central role of AQP2 blockade in the aquaretic effect of tolvaptan¹⁰². An alternative approach with broad potential clinical application is the identification of small-molecule inhibitors of aquaporins¹⁰³; however, so far, progress in the field has been disappointing.

As discussed, SIADH is characterized by hypersecretion of vasopressin. However, 10% to 20% of patients with inappropriate antidiuresis display low or undetectable vasopressin circulating levels. This condition was found to be associated with gain-of-function mutations of the V2R, first discovered in 2005¹⁰⁴, and the associated disease was defined as NSIAD to distinguish it from SIADH. NSIAD is a very rare disease representing the mirror image of NDI; 21 cases have been reported to date and the majority of them come from five families. In the first two cases of NSIAD, the mutations of the V2R were found at residue R137.

In the first two cases of NSIAD, the mutations of the V2R gene (*AVPR2*) result in changes in codon 137 from arginine to cysteine or leucine (R137C and R137L). Of note, whereas substitution of R137 to cysteine or leucine results in NSIAD, conversion of R137 to histidine (R137H) is a well-known loss-of-function mutation associated with NDI, leading to water loss and inability to concentrate urine¹⁰⁵. Thus, mutations of the same amino acid of the V2R, R137, can have opposite clinical outcomes. Other constitutively activating mutations causing NSIAD—namely F229V¹⁰⁶, I130N¹⁰⁷, and L312S¹⁰⁸—have been subsequently identified. Theoretically, NSIAD should be treated with a V2R inverse agonist; however, tolvaptan and satavaptan had no efficacy in patients carrying mutations of R137¹⁰⁹. In contrast, both drugs have an effect on F229V, I130N mutants, and L312S *in vitro*, suggesting that they could be effective in patients carrying these mutations^{106–108}.

Targeting the vasopressin–AQP2 pathway in autosomal dominant polycystic kidney disease

ADPKD is an inherited disorder with an estimated frequency of 1 in 400 to 1,000 live births¹¹⁰ and is characterized by the progressive growth of renal cysts causing disruption of renal architectures and eventually leading to end-stage renal disease. Despite high baseline vasopressin levels, patients frequently present defective urinary concentration, and this is probably due to a peripheral resistance to vasopressin^{111,112}. Interestingly, the blockade of V2R using tolvaptan in patients with rapidly progressing ADPKD has been proven to slow cyst growth, supporting the involvement of the V2R pathway in ADPKD^{113,114}.

The hereditary form of ADPKD is due to mutations in *PKD1* or *PKD2* genes, encoding for polycystin 1 (PC1) and polycystin 2 (PC2), respectively. PC1 is localized to the primary cilium and in cell–cell contacts, suggesting a role as an adhesion protein¹¹⁵. PC2, instead, is a non-selective cation channel permeable to calcium, expressed in the endoplasmic reticulum and in primary cilium, where it forms a complex with PC1 and has a role in intracellular Ca²⁺ homeostasis. At the cellular level, mutations in *PKD1* or *PKD2* are associated with a reduction in intracellular calcium, increase in cAMP, and constitutive activation of PKA,

making the collecting duct principal cells under constant tonic effect of vasopressin. The disruption in calcium and cAMP signaling cascades activates pathways causing cell proliferation, increased fluid secretion, and interstitial inflammation^{116,117}. Our recent *in vitro* study demonstrated that in human conditionally immortalized proximal tubular epithelial cells silenced for PKD1 (ciPTEC-PC1KD) or generated from a patient with ADPKD1 (ciPTEC-PC1Pt), selective activation of the calcium-sensing receptor increases cytosolic calcium, reduces intracellular cAMP and mTOR activity¹¹⁸, and rescues defective ATP mitochondrial production¹¹⁹, reversing the principal ADPKD dysregulations.

Currently, there is no cure for the disease; it is clinically manageable only through the control of its many complications, and the existing therapeutic approaches are rather supportive. Nevertheless, as mentioned, some drugs targeting the vasopressin–AQP2 pathway have been found to slow the progression of ADPKD in animal models and clinical trials¹²⁰. Tolvaptan has been tested on ADPKD patients with higher total kidney volume and was found to delay the progression of ADPKD, supporting the link between V2R signaling and ADPKD development^{113,121}. Tolvaptan has been approved to delay the progression of ADPKD in Japan, Canada, and the European Union, and very recently the US Food and Drug Administration also approved tolvaptan as the first drug treatment to slow kidney function decline in adults at risk of rapidly progressing ADPKD.

Conclusions and perspectives

Over the past decade, interest in the vasopressin–AQP2 pathway has been renewed, mainly because of the availability of vaptans,

the orally active vasopressin receptor antagonists, and of the use of copeptin as a surrogate marker of vasopressin. The dysregulation of the vasopressin–AQP2 system is clearly and tightly associated with specific diseases such as NDI, SIADH, NSIAD, and ADPKD, briefly discussed in this review (Figure 1).

With regard to NDI, many target molecules for the treatment of congenital NDI have been proposed; however, no specific pharmacological drugs have yet reached clinical application. In the development of drugs for the treatment of congenital NDI, it is important to identify/design drugs that directly activate AQP2 without toxic effects and to pay attention to preserving the medullary osmotic gradient, representing the driving force for water reabsorption. In this scenario, the screening for calcineurin activators like Wnt5a is a potentially promising therapeutic strategy to develop novel drugs for the treatment of heritable NDI.

Conversely, for the management of opposite diseases such as SIADH, characterized by hyperactivation of the vasopressin–AQP2 pathway, as discussed in this review, the use of vaptans to block vasopressin receptors represents the main avenue for the direct treatment of hyponatremia. Although among vaptans tolvaptan is well tolerated, the design of new V2R blockers reducing the reported side effects should be encouraged. Moreover, co-targeting V2R and other GPCRs known to increase intracellular calcium might be a successful approach for ADPKD treatments.

Novel AQP-targeting therapies through modulation of micro-RNA (miRNA) function have recently been suggested (reviewed in 122). The possibility of using miRNA alone or in combination

Figure 1. Targeting the vasopressin–AQP2 pathway in water balance disorders. Alterations of the vasopressin–AQP2 axis, causes of the disease, and principal effects observed in selected water balance disorders. AQP2, aquaporin-2; CNS, central nervous system; V2R, type 2 vasopressin receptor.

with other drugs to modulate the vasopressin–AQP2 pathway specifically provides new hints for AQP-based therapeutics^{27,123}. In this respect, the recent identification of miR-132 as a first miRNA target which blocks vasopressin gene expression has opened new avenues for drug development¹²⁴.

Abbreviations

ADPKD, autosomal dominant polycystic kidney disease; AKAP, A-kinase anchoring protein; AMPK, adenosine monophosphate kinase; AQP2, aquaporin-2; CKD, chronic kidney disease; GFR, glomerular filtration rate; GPCR, G-protein-coupled receptor; miRNA, microRNA; NDI, nephrogenic diabetes insipidus;

NSIAD, nephrogenic syndrome of inappropriate antidiuresis; PKA, protein kinase A; SIADH, syndrome of inappropriate antidiuretic hormone secretion; V2R, type 2 vasopressin receptor

Grant information

This study was supported in part by Telethon funding (grant number GGP13227) and by ASI (Italian Space Agency, grant number 2013-091-R.0).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Knepper MA, Kwon TH, Nielsen S: **Molecular physiology of water balance.** *N Engl J Med.* 2015; **372**(14): 1349–58.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jung HJ, Kwon TH: **Molecular mechanisms regulating aquaporin-2 in kidney collecting duct.** *Am J Physiol Renal Physiol.* 2016; **311**(6): F1318–F1328.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Lee JW, Chou CL, Knepper MA: **Deep Sequencing in Microdissected Renal Tubules Identifies Nephron Segment-Specific Transcriptomes.** *J Am Soc Nephrol.* 2015; **26**(11): 2669–77.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nedvetsky PI, Tamma G, Beulshausen S, et al.: **Regulation of aquaporin-2 trafficking.** *Handb Exp Pharmacol.* 2009; (190): 133–57.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rojek A, Fuchtbauer EM, Kwon TH, et al.: **Severe urinary concentrating defect in renal collecting duct-selective AQP2 conditional-knockout mice.** *Proc Natl Acad Sci U S A.* 2006; **103**(15): 6037–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Valenti G, Procino G, Tamma G, et al.: **Minireview: aquaporin 2 trafficking.** *Endocrinology.* 2005; **146**(12): 5063–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chou CL, Yip KP, Michea L, et al.: **Regulation of aquaporin-2 trafficking by vasopressin in the renal collecting duct. Roles of ryanodine-sensitive Ca²⁺ stores and calmodulin.** *J Biol Chem.* 2000; **275**(47): 36839–46.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Balasubramanian L, Sham JS, Yip KP: **Calcium signaling in vasopressin-induced aquaporin-2 trafficking.** *Pflügers Arch.* 2008; **456**(4): 747–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hoffert JD, Pisitkun T, Knepper MA: **Phosphoproteomics of vasopressin signaling in the kidney.** *Expert Rev Proteomics.* 2011; **8**(2): 157–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Isobe K, Jung HJ, Yang CR, et al.: **Systems-level identification of PKA-dependent signaling in epithelial cells.** *Proc Natl Acad Sci U S A.* 2017; **114**(42): E8875–E8884.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Klussmann E, Tamma G, Lorenz D, et al.: **An inhibitory role of Rho in the vasopressin-mediated translocation of aquaporin-2 into cell membranes of renal principal cells.** *J Biol Chem.* 2001; **276**(23): 20451–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tamma G, Klussmann E, Maric K, et al.: **Rho inhibits cAMP-induced translocation of aquaporin-2 into the apical membrane of renal cells.** *Am J Physiol Renal Physiol.* 2001; **281**(6): F1092–101.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Procino G, Barbieri C, Tamma G, et al.: **AQP2 exocytosis in the renal collecting duct – involvement of SNARE isoforms and the regulatory role of Munc18b.** *J Cell Sci.* 2008; **121**(Pt 12): 2097–106.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Tamma G, Procino G, Mola MG, et al.: **Functional involvement of Annexin-2 in cAMP induced AQP2 trafficking.** *Pflügers Arch.* 2008; **456**(4): 729–36.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Park EJ, Lim JS, Jung HJ, et al.: **The role of 70-kDa heat shock protein in dDAVP-induced AQP2 trafficking in kidney collecting duct cells.** *Am J Physiol Renal Physiol.* 2013; **304**(7): F958–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lu HA, Sun TX, Matsuzaki T, et al.: **Heat shock protein 70 interacts with aquaporin-2 and regulates its trafficking.** *J Biol Chem.* 2007; **282**(39): 28721–32.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Schrade K, Tröger J, Eldahshan A, et al.: **An AKAP-Lbc-RhoA interaction inhibitor promotes the translocation of aquaporin-2 to the plasma membrane of renal collecting duct principal cells.** *PLoS One.* 2018; **13**(1): e0191423.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Henn V, Edemir B, Stefan E, et al.: **Identification of a novel A-kinase anchoring protein 18 isoform and evidence for its role in the vasopressin-induced aquaporin-2 shuttle in renal principal cells.** *J Biol Chem.* 2004; **279**(25): 26654–65.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Trimpert C, Wesche D, de Groot T, et al.: **NDFIP allows NEDD4/NEDD4L-induced AQP2 ubiquitination and degradation.** *PLoS One.* 2017; **12**(9): e0183774.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Moeller HB, Olesen ET, Fenton RA: **Regulation of the water channel aquaporin-2 by posttranslational modification.** *Am J Physiol Renal Physiol.* 2011; **300**(5): F1062–73.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lee JW, Alsady M, Chou CL, et al.: **Single-tubule RNA-Seq uncovers signaling mechanisms that defend against hyponatremia in SIADH.** *Kidney Int.* 2018; **93**(1): 128–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Medvar B, Raghuram V, Pisitkun T, et al.: **Comprehensive database of human E3 ubiquitin ligases: application to aquaporin-2 regulation.** *Physiol Genomics.* 2016; **48**(7): 502–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Tamma G, Ranieri M, Di Mise A, et al.: **Glutathionylation of the aquaporin-2 water channel: a novel post-translational modification modulated by the oxidative stress.** *J Biol Chem.* 2014; **289**(40): 27807–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sandoval PC, Claxton JS, Lee JW, et al.: **Systems-level analysis reveals selective regulation of Aqp2 gene expression by vasopressin.** *Sci Rep.* 2016; **6**: 34863.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Sandoval PC, Slentz DH, Pisitkun T, et al.: **Proteome-wide measurement of protein half-lives and translation rates in vasopressin-sensitive collecting duct cells.** *J Am Soc Nephrol.* 2013; **24**(11): 1793–805.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nedvetsky PI, Tabor V, Tamma G, et al.: **Reciprocal regulation of aquaporin-2 abundance and degradation by protein kinase A and p38-MAP kinase.** *J Am Soc Nephrol.* 2010; **21**(10): 1645–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Ranieri M, Zahedi K, Tamma G, et al.: **CaSR signaling down-regulates AQP2 expression via a novel microRNA pathway in pendrin and NaCl cotransporter knockout mice.** *FASEB J.* 2018; **32**(4): 2148–59.
[PubMed Abstract](#) | [Publisher Full Text](#)

28. Centrone M, Ranieri M, Di Mise A, *et al.*: **AQP2 Abundance is Regulated by the E3-Ligase CHIP Via HSP70.** *Cell Physiol Biochem.* 2017; **44**(2): 515–31.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. **F** Wu Q, Moeller HB, Stevens DA, *et al.*: **CHIP Regulates Aquaporin-2 Quality Control and Body Water Homeostasis.** *J Am Soc Nephrol.* 2018; **29**(3): 936–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
30. Morgenthaler NG, Struck J, Alonso C, *et al.*: **Assay for the measurement of copeptin, a stable peptide derived from the precursor of vasopressin.** *Clin Chem.* 2006; **52**(1): 112–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Morgenthaler NG, Struck J, Jochberger S, *et al.*: **Copeptin: clinical use of a new biomarker.** *Trends Endocrinol Metab.* 2008; **19**(2): 43–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Szinnai G, Morgenthaler NG, Berneis K, *et al.*: **Changes in plasma copeptin, the c-terminal portion of arginine vasopressin during water deprivation and excess in healthy subjects.** *J Clin Endocrinol Metab.* 2007; **92**(10): 3973–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
33. Roussel R, Fezeu L, Marre M, *et al.*: **Comparison between copeptin and vasopressin in a population from the community and in people with chronic kidney disease.** *J Clin Endocrinol Metab.* 2014; **99**(12): 4656–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. de Bree FM, Burbach JP: **Structure-function relationships of the vasopressin prohormone domains.** *Cell Mol Neurobiol.* 1998; **18**(2): 173–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Balanescu S, Kopp P, Gaskill MB, *et al.*: **Correlation of plasma copeptin and vasopressin concentrations in hypo-, iso-, and hyperosmolar States.** *J Clin Endocrinol Metab.* 2011; **96**(4): 1046–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. **F** Nakajima A, Lu Y, Kawano H, *et al.*: **Association of arginine vasopressin surrogate marker urinary copeptin with severity of autosomal dominant polycystic kidney disease (ADPKD).** *Clin Exp Nephrol.* 2015; **19**(6): 1199–205.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
37. **F** Meijer E, Bakker SJ, Halbesma N, *et al.*: **Copeptin, a surrogate marker of vasopressin, is associated with microalbuminuria in a large population cohort.** *Kidney Int.* 2010; **77**(1): 29–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
38. Enhörning S, Bankir L, Bouby N, *et al.*: **Copeptin, a marker of vasopressin, in abdominal obesity, diabetes and microalbuminuria: The prospective Malmö Diet and Cancer Study cardiovascular cohort.** *Int J Obes (Lond).* 2013; **37**(4): 598–603.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Bardoux P, Bichet DG, Martin H, *et al.*: **Vasopressin increases urinary albumin excretion in rats and humans: Involvement of V2 receptors and the renin-angiotensin system.** *Nephrol Dial Transplant.* 2003; **18**(3): 497–506.
[PubMed Abstract](#) | [Publisher Full Text](#)
40. Fernandes S, Bruneval P, Hagege A, *et al.*: **Chronic V2 vasopressin receptor stimulation increases basal blood pressure and exacerbates deoxycorticosterone acetate-salt hypertension.** *Endocrinology.* 2002; **143**(7): 2759–66.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Sugiyama T, Yamauchi A, Kitamura H, *et al.*: **High water intake ameliorates tubulointerstitial injury in rats with subtotal nephrectomy: Possible role of TGF-beta.** *Kidney Int.* 1999; **55**(5): 1800–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
42. Bankir L, Bardoux P, Ahloulay M: **Vasopressin and diabetes mellitus.** *Nephron.* 2001; **87**(1): 8–18.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Boertien WE, Riphagen IJ, Drion I, *et al.*: **Copeptin, a surrogate marker for arginine vasopressin, is associated with declining glomerular filtration in patients with diabetes mellitus (ZODIAC-33).** *Diabetologia.* 2013; **56**(8): 1680–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Pikkemaat M, Melander O, Bengtsson Boström K: **Association between copeptin and declining glomerular filtration rate in people with newly diagnosed diabetes. The Skaraborg Diabetes Register.** *J Diabetes Complicat.* 2015; **29**(8): 1062–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
45. Ponte B, Pruijm M, Ackermann D, *et al.*: **Copeptin is associated with kidney length, renal function, and prevalence of simple cysts in a population-based study.** *J Am Soc Nephrol.* 2015; **26**(6): 1415–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. **F** Clark WF, Devuyst O, Roussel R: **The vasopressin system: New insights for patients with kidney diseases: Epidemiological evidence and therapeutic perspectives.** *J Intern Med.* 2017; **282**(4): 310–21.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
47. **F** Clark WF, Sontrop JM, Huang SH, *et al.*: **Hydration and Chronic Kidney Disease Progression: A Critical Review of the Evidence.** *Am J Nephrol.* 2016; **43**(4): 281–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
48. Kwon TH, Frøkier J, Knepper MA, *et al.*: **Reduced AQP1, -2, and -3 levels in kidneys of rats with CRF induced by surgical reduction in renal mass.** *Am J Physiol.* 1998; **275**(5 Pt 2): F724–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Teitelbaum I, McGuinness S: **Vasopressin resistance in chronic renal failure. Evidence for the role of decreased V2 receptor mRNA.** *J Clin Invest.* 1995; **96**(1): 378–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Palmer SC, Wong G, Iff S, *et al.*: **Fluid intake and all-cause mortality, cardiovascular mortality and kidney function: A population-based longitudinal cohort study.** *Nephrol Dial Transplant.* 2014; **29**(7): 1377–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
51. **F** Sands JM, Klein JD: **Physiological insights into novel therapies for nephrogenic diabetes insipidus.** *Am J Physiol Renal Physiol.* 2016; **311**(6): F1149–F1152.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
52. Sands JM, Bichet DG, American College of Physicians, *et al.*: **Nephrogenic diabetes insipidus.** *Ann Intern Med.* 2006; **144**(3): 186–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
53. **F** Bichet DG, Bockenhauer D: **Genetic forms of nephrogenic diabetes insipidus (NDI): Vasopressin receptor defect (X-linked) and aquaporin defect (autosomal recessive and dominant).** *Best Pract Res Clin Endocrinol Metab.* 2016; **30**(2): 263–76.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
54. Di Iorgi N, Napoli F, Allegri AE, *et al.*: **Diabetes insipidus—diagnosis and management.** *Horm Res Paediatr.* 2012; **77**(2): 69–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
55. Bedford JJ, Weggerly S, Ellis G, *et al.*: **Lithium-induced nephrogenic diabetes insipidus: Renal effects of amiloride.** *Clin J Am Soc Nephrol.* 2008; **3**(5): 1324–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. CASTELL DO, SPARKS HA: **NEPHROGENIC DIABETES INSIPIDUS DUE TO DEMETHYLCHLORTETRACYCLINE HYDROCHLORIDE.** *JAMA.* 1965; **193**: 237–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
57. Metzger NL, Varney Gill KL: **Nephrogenic diabetes insipidus induced by two amphotericin B liposomal formulations.** *Pharmacotherapy.* 2009; **29**(5): 613–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
58. Roth H, Becker KL, Shalhoub RJ, *et al.*: **Nephrotoxicity of demethylchlortetracycline hydrochloride. A prospective study.** *Arch Intern Med.* 1967; **120**(4): 433–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
59. Singer I, Rotenberg D: **Demeclocycline-induced nephrogenic diabetes insipidus. In-vivo and in-vitro studies.** *Ann Intern Med.* 1973; **79**(5): 679–83.
[PubMed Abstract](#) | [Publisher Full Text](#)
60. Torin DE: **Nephrogenic diabetes insipidus induced by demethylchlortetracycline (Declomycin).** *Calif Med.* 1967; **107**(5): 420–2.
[PubMed Abstract](#) | [Free Full Text](#)
61. Marples D, Frøkier J, Dørup J, *et al.*: **Hypokalemia-induced downregulation of aquaporin-2 water channel expression in rat kidney medulla and cortex.** *J Clin Invest.* 1996; **97**(8): 1960–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
62. Wang W, Li C, Kwon TH, *et al.*: **AQP3, p-AQP2, and AQP2 expression is reduced in polyuric rats with hypercalcemia: prevention by cAMP-PDE inhibitors.** *Am J Physiol Renal Physiol.* 2002; **283**(6): F1313–25.
[PubMed Abstract](#) | [Publisher Full Text](#)
63. Sands JM, Flores FX, Kato A, *et al.*: **Vasopressin-elicited water and urea permeabilities are altered in IMCD in hypercalcemic rats.** *Am J Physiol.* 1998; **274**(5 Pt 2): F978–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
64. Bustamante M, Hasler U, Leroy V, *et al.*: **Calcium-sensing receptor attenuates AVP-induced aquaporin-2 expression via a calmodulin-dependent mechanism.** *J Am Soc Nephrol.* 2008; **19**(1): 109–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
65. Procino G, Mastrofrancesco L, Tamma G, *et al.*: **Calcium-sensing receptor and aquaporin 2 interplay in hypercalcemia-associated renal concentrating defect in humans. An in vivo and in vitro study.** *PLoS One.* 2012; **7**(3): e33145.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
66. Ranieri M, Tamma G, Di Mise A, *et al.*: **Negative feedback from CaSR signaling to aquaporin-2 sensitizes vasopressin to extracellular Ca²⁺.** *J Cell Sci.* 2015; **128**(13): 2350–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
67. Tamma G, Di Mise A, Ranieri M, *et al.*: **A decrease in aquaporin 2 excretion is associated with bed rest induced high calciuria.** *J Transl Med.* 2014; **12**: 133.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
68. Procino G, Carosino M, Tamma G, *et al.*: **Extracellular calcium antagonizes forskolin-induced aquaporin 2 trafficking in collecting duct cells.** *Kidney Int.* 2004; **66**(6): 2245–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
69. Qureshi S, Galiveti S, Bichet DG, *et al.*: **Diabetes insipidus: celebrating a century of vasopressin therapy.** *Endocrinology.* 2014; **155**(12): 4605–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
70. Monnens L, Jonkman A, Thomas C: **Response to indomethacin and hydrochlorothiazide in nephrogenic diabetes insipidus.** *Clin Sci (Lond).* 1984; **66**(6): 709–15.
[PubMed Abstract](#)

71. **F** Milano S, Carmosino M, Gerbino A, *et al.*: **Hereditary Nephrogenic Diabetes Insipidus: Pathophysiology and Possible Treatment. An Update.** *Int J Mol Sci.* 2017; **18** (11): pii: E2385.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
72. **F** Bockenhauer D, Bichet DG: **Nephrogenic diabetes insipidus.** *Curr Opin Pediatr.* 2017; **29**(2): 199–205.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
73. **F** Ando F, Uchida S: **Activation of AQP2 water channels without vasopressin: therapeutic strategies for congenital nephrogenic diabetes insipidus.** *Clin Exp Nephrol.* 2018; **22**(3): 501–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
74. Sanches TR, Volpini RA, Massola Shimizu MH, *et al.*: **Sildenafil reduces polyuria in rats with lithium-induced NDI.** *Am J Physiol Renal Physiol.* 2012; **302**(1): F216–25.
[PubMed Abstract](#) | [Publisher Full Text](#)
75. Assadi F, Sharbat FG: **Sildenafil for the Treatment of Congenital Nephrogenic Diabetes Insipidus.** *Am J Nephrol.* 2015; **42**(1): 65–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
76. Sidaway JE, Davidson RG, McTaggart F, *et al.*: **Inhibitors of 3-hydroxy-3-methylglutaryl-CoA reductase reduce receptor-mediated endocytosis in opossum kidney cells.** *J Am Soc Nephrol.* 2004; **15**(9): 2258–65.
[PubMed Abstract](#) | [Publisher Full Text](#)
77. Procino G, Barbieri C, Carmosino M, *et al.*: **Fluvastatin modulates renal water reabsorption in vivo through increased AQP2 availability at the apical plasma membrane of collecting duct cells.** *PLoS Arch.* 2011; **462**(5): 753–66.
[PubMed Abstract](#) | [Publisher Full Text](#)
78. Li W, Zhang Y, Bouley R, *et al.*: **Simvastatin enhances aquaporin-2 surface expression and urinary concentration in vasopressin-deficient Brattleboro rats through modulation of Rho GTPase.** *Am J Physiol Renal Physiol.* 2011; **301**(2): F309–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
79. Procino G, Milano S, Carmosino M, *et al.*: **Combination of secretin and fluvastatin ameliorates the polyuria associated with X-linked nephrogenic diabetes insipidus in mice.** *Kidney Int.* 2014; **86**(1): 127–38.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
80. Elie D, Segal M, Low NC, *et al.*: **Statins in the prevention of lithium-associated diabetes insipidus: preliminary findings.** *Kidney Int.* 2015; **87**(4): 862.
[PubMed Abstract](#) | [Publisher Full Text](#)
81. Procino G: **Fluvastatin Increases AQP2 Urine Excretion in a Dyslipidemic Patient with Nephrogenic Diabetes Insipidus: An In Vivo and In Vitro Study.** *J Diabetes Metab.* 2014; **5**: 408.
[Publisher Full Text](#)
82. **F** Efe O, Klein JD, LaRocque LM, *et al.*: **Metformin improves urine concentration in rodents with nephrogenic diabetes insipidus.** *JCI Insight.* 2016; **1**(11): pii: e88409.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
83. **F** Klein JD, Wang Y, Blount MA, *et al.*: **Metformin, an AMPK activator, stimulates the phosphorylation of aquaporin 2 and urea transporter A1 in inner medullary collecting ducts.** *Am J Physiol Renal Physiol.* 2016; **310**(10): F1008–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
84. **F** Ando F, Mori S, Yui N, *et al.*: **AKAPs-PKA disruptors increase AQP2 activity independently of vasopressin in a model of nephrogenic diabetes insipidus.** *Nat Commun.* 2018; **9**(1): 1411.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
85. Bouley R, Lu HA, Nunes P, *et al.*: **Calcitonin has a vasopressin-like effect on aquaporin-2 trafficking and urinary concentration.** *J Am Soc Nephrol.* 2011; **22**(1): 59–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
86. Kwon TH, Nielsen J, Knepper MA, *et al.*: **Angiotensin II AT₁ receptor blockade decreases vasopressin-induced water reabsorption and AQP2 levels in NaCl-restricted rats.** *Am J Physiol Renal Physiol.* 2005; **288**(4): F673–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
87. Li C, Wang W, Rivard CJ, *et al.*: **Molecular mechanisms of angiotensin II stimulation on aquaporin-2 expression and trafficking.** *Am J Physiol Renal Physiol.* 2011; **300**(5): F1255–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
88. **F** Olesen ET, Rützler MR, Moeller HB, *et al.*: **Vasopressin-independent targeting of aquaporin-2 by selective E-prostanoid receptor agonists alleviates nephrogenic diabetes insipidus.** *Proc Natl Acad Sci U S A.* 2011; **108**(31): 12949–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
89. **F** Gao M, Cao R, Du S, *et al.*: **Disruption of prostaglandin E2 receptor EP4 impairs urinary concentration via decreasing aquaporin 2 in renal collecting ducts.** *Proc Natl Acad Sci U S A.* 2015; **112**(27): 8397–402.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
90. **F** Olesen ET, Moeller HB, Assentoft M, *et al.*: **The vasopressin type 2 receptor and prostaglandin receptors EP2 and EP4 can increase aquaporin-2 plasma membrane targeting through a cAMP-independent pathway.** *Am J Physiol Renal Physiol.* 2016; **311**(5): F935–F944.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
91. **F** Ando F, Sohara E, Morimoto T, *et al.*: **Wnt5a induces renal AQP2 expression by activating calcineurin signalling pathway.** *Nat Commun.* 2016; **7**: 13636.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
92. Hannon MJ, Thompson CJ: **The syndrome of inappropriate antidiuretic hormone: prevalence, causes and consequences.** *Eur J Endocrinol.* 2010; **162** Suppl 1: S5–12.
[PubMed Abstract](#) | [Publisher Full Text](#)
93. Schrier RW, Sharma S, Shchekochikhin D: **Hyponatraemia: more than just a marker of disease severity?** *Nat Rev Nephrol.* 2013; **9**(1): 37–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
94. Gheorghiadu M, Abraham WT, Albert NM, *et al.*: **Relationship between admission serum sodium concentration and clinical outcomes in patients hospitalized for heart failure: an analysis from the OPTIMIZE-HF registry.** *Eur Heart J.* 2007; **28**(8): 980–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
95. Anderson RJ, Chung HM, Kluge R, *et al.*: **Hyponatremia: a prospective analysis of its epidemiology and the pathogenetic role of vasopressin.** *Ann Intern Med.* 1985; **102**(2): 164–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
96. Ecelbarger CA, Nielsen S, Olson BR, *et al.*: **Role of renal aquaporins in escape from vasopressin-induced antidiuresis in rat.** *J Clin Invest.* 1997; **99**(8): 1852–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
97. Ecelbarger CA, Chou CL, Lee AJ, *et al.*: **Escape from vasopressin-induced antidiuresis: role of vasopressin resistance of the collecting duct.** *Am J Physiol.* 1998; **274**(6 Pt 2): F1161–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
98. **F** Kim S, Choi HJ, Jo CH, *et al.*: **Cyclophosphamide-induced vasopressin-independent activation of aquaporin-2 in the rat kidney.** *Am J Physiol Renal Physiol.* 2015; **309**(5): F474–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
99. Robertson GL: **Regulation of arginine vasopressin in the syndrome of inappropriate antidiuresis.** *Am J Med.* 2006; **119**(7 Suppl 1): S36–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
100. **F** Cuesta M, Thompson CJ: **The syndrome of inappropriate antidiuresis (SIAD).** *Best Pract Res Clin Endocrinol Metab.* 2016; **30**(2): 175–87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
101. **F** Berl T, Quinatat-Pelletier F, Verbalis JG, *et al.*: **Oral tolvaptan is safe and effective in chronic hyponatremia.** *J Am Soc Nephrol.* 2010; **21**(4): 705–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
102. Tamma G, Di Mise A, Ranieri M, *et al.*: **The V2 receptor antagonist tolvaptan raises cytosolic calcium and prevents AQP2 trafficking and function: an in vitro and in vivo assessment.** *J Cell Mol Med.* 2017; **21**(9): 1767–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
103. **F** Tradtrantip L, Jin BJ, Yao X, *et al.*: **Aquaporin-Targeted Therapeutics: State-of-the-Field.** *Adv Exp Med Biol.* 2017; **969**: 239–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
104. Feldman BJ, Rosenthal SM, Vargas GA, *et al.*: **Nephrogenic syndrome of inappropriate antidiuresis.** *N Engl J Med.* 2005; **352**(18): 1884–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
105. Rosenthal W, Seibold A, Antaramian A, *et al.*: **Mutations in the vasopressin V2 receptor gene in families with nephrogenic diabetes insipidus and functional expression of the Q-2 mutant.** *Cell Mol Biol (Noisy-le-grand).* 1994; **40**(3): 429–36.
[PubMed Abstract](#)
106. **F** Carpentier E, Greenbaum LA, Rochdi D, *et al.*: **Identification and characterization of an activating F229V substitution in the V2 vasopressin receptor in an infant with NSIAD.** *J Am Soc Nephrol.* 2012; **23**(10): 1635–40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
107. Erdélyi LS, Mann WA, Morris-Rosendahl DJ, *et al.*: **Mutation in the V2 vasopressin receptor gene, AVPR2, causes nephrogenic syndrome of inappropriate diuresis.** *Kidney Int.* 2015; **88**(5): 1070–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
108. **F** Tiulpakov A, White CW, Abhayawardana RS, *et al.*: **Mutations of Vasopressin Receptor 2 Including Novel L312S Have Differential Effects on Trafficking.** *Mol Endocrinol.* 2016; **30**(8): 889–904.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
109. **F** Decaux G, Vandergheynst F, Bouko Y, *et al.*: **Nephrogenic syndrome of inappropriate antidiuresis in adults: high phenotypic variability in men and women from a large pedigree.** *J Am Soc Nephrol.* 2007; **18**(2): 606–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
110. Harris PC, Torres VE: **Genetic mechanisms and signaling pathways in autosomal dominant polycystic kidney disease.** *J Clin Invest.* 2014; **124**(6): 2315–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
111. Zitterma D, Boertien WE, van Beek AP, *et al.*: **Vasopressin, copeptin, and renal concentrating capacity in patients with autosomal dominant polycystic kidney disease without renal impairment.** *Clin J Am Soc Nephrol.* 2012; **7**(6): 906–13.
[PubMed Abstract](#) | [Publisher Full Text](#)
112. Gabow PA, Kaehny WD, Johnson AM, *et al.*: **The clinical utility of renal concentrating capacity in polycystic kidney disease.** *Kidney Int.* 1989; **35**(2): 675–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
113. **F** Chebib FT, Perrone RD, Chapman AB, *et al.*: **A Practical Guide for Treatment**

- of Rapidly Progressive ADPKD with Tolvaptan. *J Am Soc Nephrol.* 2018; 29(10): 2458–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
114. **F** van Gastel MDA, Torres VE: **Polycystic Kidney Disease and the Vasopressin Pathway.** *Ann Nutr Metab.* 2017; 70 Suppl 1: 43–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
115. Ong AC, Harris PC: **A polycystin-centric view of cyst formation and disease: the polycystins revisited.** *Kidney Int.* 2015; 88(4): 699–710.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
116. Chebib FT, Sussman CR, Wang X, *et al.*: **Vasopressin and disruption of calcium signalling in polycystic kidney disease.** *Nat Rev Nephrol.* 2015; 11(8): 451–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
117. **F** Torres VE, Chapman AB, Devuyst O, *et al.*: **Tolvaptan in patients with autosomal dominant polycystic kidney disease.** *N Engl J Med.* 2012; 367(25): 2407–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
118. Di Mise A, Tamma G, Ranieri M, *et al.*: **Activation of Calcium-Sensing Receptor increases intracellular calcium and decreases cAMP and mTOR in PKD1 deficient cells.** *Sci Rep.* 2018; 8(1): 5704.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
119. Di Mise A, Ranieri M, Centrone M, *et al.*: **Activation of the Calcium-Sensing Receptor Corrects the Impaired Mitochondrial Energy Status Observed in Renal Polycystin-1 Knockdown Cells Modeling Autosomal Dominant Polycystic Kidney Disease.** *Front Mol Biosci.* 2018; 5: 77.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
120. Happé H, Peters DJ: **Translational research in ADPKD: lessons from animal models.** *Nat Rev Nephrol.* 2014; 10(10): 587–601.
[PubMed Abstract](#) | [Publisher Full Text](#)
121. **F** Devuyst O, Chapman AB, Gansevoort RT, *et al.*: **Urine Osmolality, Response to Tolvaptan, and Outcome in Autosomal Dominant Polycystic Kidney Disease: Results from the TEMPO 3:4 Trial.** *J Am Soc Nephrol.* 2017; 28(5): 1592–602.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
122. **F** Gomes A, da Silva IV, Rodrigues CMP, *et al.*: **The Emerging Role of microRNAs in Aquaporin Regulation.** *Front Chem.* 2018; 6: 238.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
123. Kim JE, Jung HJ, Lee YJ, *et al.*: **Vasopressin-regulated miRNAs and AQP2-targeting miRNAs in kidney collecting duct cells.** *Am J Physiol Renal Physiol.* 2015; 308(7): F749–64.
[PubMed Abstract](#) | [Publisher Full Text](#)
124. **F** Bijkerk R, Trimpert C, van Solingen C, *et al.*: **MicroRNA-132 controls water homeostasis through regulating MECP2-mediated vasopressin synthesis.** *Am J Physiol Renal Physiol.* 2018; 315(4): F1129–F1138.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Tae-Hwan Kwon**

Department of Biochemistry and Cell Biology, Kyungpook National University, Taegu, South Korea

Competing Interests: No competing interests were disclosed.

2. **Kenichi Ishibashi**

Department of Pathophysiology, Meiji Pharmaceutical University, Tokyo, Japan

Competing Interests: No competing interests were disclosed.

3. **Enno Klussmann**

Max Delbrück Center for Molecular Medicine Berlin (MDC), Berlin, Germany

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research