Check for updates

CASE REPORT

Case Report: Photo-activated oral disinfection in regenerative endodontics [version 1; peer review: 1 approved with reservations, 1 not approved]

Al-Shaimaa Abdel Hafiz Abdel Rahim ¹, Fatma Abdelgawad ²,
Ahmed M. Abd Alsamed ³, Dalia Mohamed Moheb²,
Norhan Abdel Wahab El-Dokky²

¹Pediatric Dentistry and Dental Public Health Department, Faculty of Dentistry, Fayoum University, Fayoum, 63514, Egypt

²Pediatric Dentistry and Dental Public Health Department, Faculty of Dentistry, Cairo University, Cairo, 11553, Egypt

³Oral and Maxillofacial Radiology Department, Faculty of Dentistry, Cairo University, Cairo, 11553, Egypt

V1 First published: 28 Aug 2019, 8:1519
<https://doi.org/10.12688/f1000research.20118.1>
Latest published: 03 Jun 2020, 8:1519
<https://doi.org/10.12688/f1000research.20118.2>

Abstract

Background: Root canal disinfection is considered critical for achieving successful regenerative endodontic procedures. Photo-activated oral disinfection is a novel disinfection method that can help to achieve the goal of regenerative endodontics. This article reports the clinical and radiographic results after single visit regenerative endodontics using photo-activated oral disinfection.

Methods: An 8.5-year-old girl complained of fractured upper right central incisor. Pulp necrosis was diagnosed on the basis of clinical findings. The root canal was irrigated with sodium hypochlorite solution (1.5%) followed by saline. Then, the canal was dried with paper points. A combination of a photosensitizer solution and low power laser light were applied. EDTA solution was used as a final irrigant. Bleeding was induced, followed by placement of collagen resorbable matrix and white mineral trioxide aggregate. Two days later, the tooth was sealed and restored with permanent filling.

Results: Clinical findings revealed no pain on percussion or palpation tests. Radiographic examination revealed an increase in root length, an increase of apical root thickness and apical closure at the 12-month follow-up period.

Conclusion: Regenerative endodontics using photo-activated oral disinfection achieved successful outcomes in the necrotic immature permanent tooth.

Keywords

regenerative endodontics, necrotic tooth, open apex, root canal disinfection

Open Peer Review

Approval Status

	1	2
version 2		
(revision)		
03 Jun 2020		
version 1		
28 Aug 2019		

1. **Gianluca Gambarini**, Sapienza University of Rome, Rome, Italy

Dario Di Nardo, Sapienza University of Rome, Rome, Italy

2. **Mohamed Mokhtar Nagy** , Ain Shams University, Cairo, Egypt

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Al-Shaimaa Abdel Hafiz Abdel Rahim (aaa50@fayoum.edu.eg)

Author roles: **Abdel Hafiz Abdel Rahim AS:** Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation; **Abdelgawad F:** Methodology, Supervision, Validation, Visualization, Writing – Review & Editing; **Abd Alsamed AM:** Investigation, Software, Supervision, Visualization; **Moheb DM:** Conceptualization, Supervision, Writing – Review & Editing; **Wahab El-Dokky NA:** Supervision, Validation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2019 Abdel Hafiz Abdel Rahim AS *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Abdel Hafiz Abdel Rahim AS, Abdelgawad F, Abd Alsamed AM *et al.* **Case Report: Photo-activated oral disinfection in regenerative endodontics [version 1; peer review: 1 approved with reservations, 1 not approved]** F1000Research 2019, 8:1519 <https://doi.org/10.12688/f1000research.20118.1>

First published: 28 Aug 2019, 8:1519 <https://doi.org/10.12688/f1000research.20118.1>

Introduction

The term regenerative endodontic procedures (REPs) has been widely endorsed. This term describes all procedures that aim to achieve organized repair of the dental pulp¹. The clinical considerations for REPs include: disinfecting the root canal system; providing a scaffold with periapical tissue laceration to get a blood clot and introduce stem cell activity within the root canal; and sealing the coronal access properly to prevent reinfection^{2,3}.

Antibiotics appear to be suitable intra-canal medication. Triple antibiotic paste (TAP) consisting of metronidazole, ciprofloxacin and minocycline is considered to be a successful regimen in managing the root canal pathogen of necrotic immature permanent teeth⁴. Recently, new methods of disinfection have been described to control the limitations of conventional disinfecting methods by neither diminishing the bacterial number to an adequate level or eliminating the toxicity to periapical stem/progenitors⁵. Other concerns regarding the use of TAP include tooth discoloration and bacterial resistance⁶.

Photodynamic therapy (PDT)/photo-activated oral disinfection (PAD) involves a photosensitizer (photoactive dye) that is activated by exposure to light of a specific wavelength in the presence of oxygen. The energy transferred from the activated photosensitizer to available oxygen leads to toxic oxygen species formation, such as singlet oxygen and free radicals. These very reactive chemical species can destroy proteins, nucleic acids, lipids and other cellular components⁷.

Several *in vitro* studies have shown that PDT is effective in root canal disinfection^{8–10}. A clinical study by Johns *et al.*¹¹ reported successful outcomes for using PAD in root canal treatment of completely formed root. The case report by Johns *et al.*¹² was the first to document pulp revascularization using PDT and platelet-rich fibrin. Root lengthening, continued thickening of the canal walls and apical closure were demonstrated at 10-month follow-up¹².

This article reports the successful use of PAD in regenerative endodontics as a novel and effective disinfection method, which might present a solution to the problems associated with triple antibiotic paste¹².

Case report

An 8.5-year-old Egyptian girl came to the outpatient clinic of Pediatric Dentistry and Dental Public Health Department, Faculty of Dentistry, Cairo University with the chief complaint of fractured upper right central incisor due to trauma one and half months previously (Figure 1A and B). The patient's medical history was non-contributory. On clinical examination (which

included a visual examination for any abnormalities, palpation of labial vestibule, percussion test and sensibility test), the tooth was sensitive to percussion, which was determined by tapping the tooth with the back of the mirror. The surrounding soft tissue had no tenderness to finger palpation and the tooth had no response to the hot test in comparison to the contralateral tooth. Preoperative radiographic examination revealed a wide root canal with an open apex (Figure 2A) using conventional periapical radiograph.

The case was managed by single visit regenerative endodontic procedure using PAD after obtaining written informed consent from the parents (including usage of data and publication) and assent from the child. The procedure was performed according to the American Association of Endodontics (AAE) guidelines¹³, except PAD instead of TAP was used. A 1.8ml carpule of mepivacaine 3% (Mepivacaine hydrochloride, Alexandria Co. for Pharmaceuticals, Egypt) was administered by infiltration. A rubber dam was applied, followed by access cavity preparation and tooth length detection. Gentle, copious irrigation was done using 20ml NaOCl (1.5% NaOCl 20ml/canal for five minutes). NaOCl of lower concentration was advised and then flushed with saline (20ml/canal for minutes), using irrigating needle located about 1mm from root end then the canal was dried using paper points.

Aseptim solution (SciCan Ltd, Denfotex Light Systems Ltd, Inverkeithing, Scotland) was applied to the canal, followed by a low power laser diode red light system (Aseptim system, SciCan Ltd, Denfotex Light Systems Ltd, Inverkeithing, Scotland) with a specific wavelength (635nm) to activate the aseptic solution for 150 seconds. Canals was washed with saline to remove the aseptic solution. Gentle, copious, irrigation with 20ml of 17% EDTA was applied. Dryness with paper points to remove excess EDTA was done. Bleeding was initiated into the canal system by rotating a K-file at 2mm beyond the apical foramen. A resorbable matrix (Collacote dressing Zimmer Biomet, USA) was placed over the blood clot. Then, white mineral trioxide aggregate (MTA) (Angelus, Brazil) was placed over the matrix with placement of a moistened pellet of cotton and glass ionomer (Figure 2B). After two days, the tooth was double sealed using glass ionomer cement (Kromoglass 2, LASCOD- Italy) and composite restoration 3M composite (3M, America Inc).

Clinical examination of the patient revealed no adverse signs and symptoms at three, six-, nine- and 12-month follow-up periods (Figure 1C and 1D). Radiographic examination revealed an increase in root length and root thickness at six months (Figure 2C) and complete root closure at 12 months (Figure 2D) using digital radiography (Digora™ Optime UV) and digital software (Soredex, Finland) (Table 1).

Figure 1. **A)** front view of preoperative intra-oral photo of traumatized upper right central incisor; **B)** occlusal preoperative photo; **C)** three month follow up; and **D)** 12-month follow-up.

Figure 2. **A)** preoperative periapical radiograph; **B)** immediate postoperative radiograph after mineral trioxide aggregate coronal plug application; **C)** follow up radiograph at six months; and **D)** follow-up radiograph at 12 months.

Table 1. The patient's timeline of symptoms, treatment and follow-up periods.

Timeline	Event	Findings
0	Patient comes to the clinic. Patient's medical history obtained. Clinical and radiographic examination performed.	Fractured necrotic traumatized upper right central incisor, sensitive to percussion and no response to hot test with a wide root canal and an open apex
0	Regenerative endodontic procedure using photo-activated oral disinfection	
+ 2 days	Composite restoration	
+ 3 months	1 st Follow up (clinical assessment)	No pain, no swelling (symptom free)
+ 6 months	2 nd Follow up (clinical and radiographic assessment)	No pain, no swelling (symptom free) Increase in root length and dentin root thickness, and no adverse radiographic evidence.
+ 9 months	3 rd follow up (clinical assessment)	No pain, no swelling (symptom free).
+ 12 months	4 th follow up (clinical and radiographic assessment)	No pain, no swelling (symptom free). Complete apical root closure

Discussion

This case report documents the successful application of a novel and innovative disinfection technique in regenerative endodontics for management of a necrotic immature permanent tooth in a single visit, although a longer follow up period is recommended.

Root canal system disinfection is an integral step in the success of REPs. Chemical disinfection of the root canal is dependent on bacteriostatic/bactericidal properties of the agents as well as avoid harming the patient's stem cells³. Copious and gentle irrigation was done using 20ml NaOCl. The use of NaOCl at lower concentrations was advised (1.5% NaOCl) with the irrigating needle adjusted to be 1mm from the root end to reduce the cytotoxicity to stem cells and to minimize the possibility of irrigant extrusion into the periapical tissues¹³. Furthermore, 1.5% NaOCl was effective in bacterial reduction of root canals as reported by Trevino *et al.*¹⁴.

The PAD technique was effective in removing high bacterial concentrations from infected root canals¹⁵. Aseptim solution was applied to the canal and agitated in the canal for 60 seconds using an endodontic file to ensure maximum penetration of the dye, since it is essential that the aseptim solution comes in close contact with the bacteria, otherwise the photosensitivity process does not occur¹⁶.

Blood clot formation inside the canal is considered a scaffold and a source of growth factors. Inducing bleeding to promote blood clot formation is a commonly used method in many of the reported cases^{17–20}. A resorbable matrix (collagen wound dressing) should be carefully placed on the top of the blood clot to serve as an internal matrix and ease the placement of MTA¹³. White MTA was applied to act as a coronal plug and

achieve effective coronal seal. It was the most frequently chosen material in published regenerative endodontic studies^{21–24}.

No adverse clinical signs and symptoms were noted during follow up periods, which is considered as a primary goal for regenerative endodontics as reported by AAE¹³. Apical closure and increasing apical dentin thickness were observed at the 12-month follow-up period. This case report demonstrates that PAD is a promising method for controlling infection in a single visit regenerative endodontic procedure. Further randomized clinical studies are needed to assess this disinfection technique.

Data availability

All data underlying the results are available as part of the article and no additional source data are required.

Consent

Written informed consent for publication of their clinical details and clinical images was obtained from the parents of the patient.

Grant information

The author(s) declared that no grants were involved in supporting this work.

Acknowledgements

I dedicate this work to the soul of Prof. Mervat Abdelmonem Rashed, Professor of Pediatric Dentistry, Pediatric Dentistry and Dental Public Health Department, Faculty of Dentistry, Cairo University that her death prevents her to see the result of this report.

References

- Diogenes A, Henry MA, Teixeira FB, *et al.*: **An update on clinical regenerative endodontics.** *Endod Top.* 2013; **28**(1): 2–23.
[Publisher Full Text](#)
- Bezzin T, Sönmez H: **Review of current concepts of revascularization/revitalization.** *Dent Traumatol.* 2015; **31**(4): 267–73.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Galler KM: **Clinical procedures for revitalization: current knowledge and considerations.** *Int Endod J.* 2016; **49**(10): 926–36.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Vijayaraghavan R, Mathian VM, Sundaram AM, *et al.*: **Triple antibiotic paste in root canal therapy.** *J Pharm Bioallied Sci.* 2012; **4**(Suppl 2): S230–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kim SG: **Dental pulp regeneration.** *Curr Ther Endod.* 2016; 271–80.
[Publisher Full Text](#)
- Huang GT: **A paradigm shift in endodontic management of immature teeth: conservation of stem cells for regeneration.** *J Dent.* 2008; **36**(6): 379–86.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Konopka K, Goslinski T: **Photodynamic therapy in dentistry.** *J Dent Res.* 2007; **86**(8): 694–707.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bergmans L, Moisiadis P, Huybrechts B, *et al.*: **Effect of photo-activated disinfection on endodontic pathogens ex vivo.** *Int Endod J.* 2008; **41**(3): 227–39.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Garcez AS, Ribeiro MS, Tegos GP, *et al.*: **Antimicrobial photodynamic therapy combined with conventional endodontic treatment to eliminate root canal biofilm infection.** *Lasers Surg Med.* 2007; **39**(1): 59–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Williams JA, Pearson GJ, John CM: **Antibacterial action of photoactivated disinfection {PAD} used on endodontic bacteria in planktonic suspension and in artificial and human root canals.** *J Dent.* 2006; **34**(6): 363–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Johns DA, Varughese JM, Thomas K, *et al.*: **Clinical and radiographical evaluation of the healing of large periapical lesions using triple antibiotic paste, photo activated disinfection and calcium hydroxide when used as root canal disinfectant.** *J Clin Exp Dent.* 2014; **6**(3): e230–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Johns DA, Shivashankar VY, Krishnamma S, *et al.*: **Use of photoactivated disinfection and platelet-rich fibrin in regenerative Endodontics.** *J Conserv Dent.* 2014; **17**(5): 487–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Consent Informed, Appointment First: **AAE Clinical Considerations for a Regenerative Procedure Revised 4-12-15.** 2015: 1–6.
- Trevino EG, Patwardhan AN, Henry MA, *et al.*: **Effect of irrigants on the survival of human stem cells of the apical papilla in a platelet-rich plasma scaffold in human root tips.** *J Endod.* 2011; **37**(8): 1109–15.
[PubMed Abstract](#) | [Publisher Full Text](#)

15. Plotino G, Cortese T, Grande NM, *et al.*: **New Technologies to Improve Root Canal Disinfection.** *Braz Dent J.* 2016; **27**(1): 3–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Bonsor SJ, Nichol R, Reid TM, *et al.*: **Microbiological evaluation of photo-activated disinfection in endodontics (an *in vivo* study).** *Br Dent J.* 2006; **200**(6): 337–41, discussion 329.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Reynolds K, Johnson JD, Cohenca N: **Pulp revascularization of necrotic bilateral bicuspid using a modified novel technique to eliminate potential coronal discolouration: a case report.** *Int Endod J.* 2009; **42**(1): 84–92.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Shin SY, Albert JS, Mortman RE: **One step pulp revascularization treatment of an immature permanent tooth with chronic apical abscess: a case report.** *Int Endod J.* 2009; **42**(12): 1118–26.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Cehreli ZC, Sara S, Aksoy B: **Revascularization of immature permanent incisors after severe extrusive luxation injury.** *J Can Dent Assoc.* 2012; **78**(1): c4.
[PubMed Abstract](#)
20. Mao JJ, Kim SG, Zhou J, *et al.*: **Regenerative endodontics: barriers and strategies for clinical translation.** *Dent Clin North Am.* 2012; **56**(3): 639–49.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Banchs F, Trope M: **Revascularization of immature permanent teeth with apical periodontitis: new treatment protocol?** *J Endod.* 2004; **30**(4): 196–200.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Thibodeau B, Trope M: **Pulp revascularization of a necrotic infected immature permanent tooth: case report and review of the literature.** *Pediatr Dent.* 2007; **29**(1): 47–50.
[PubMed Abstract](#)
23. Ding RY, Cheung GS, Chen J, *et al.*: **Pulp revascularization of immature teeth with apical periodontitis: a clinical study.** *J Endod.* 2009; **35**(5): 745–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Lenzi R, Trope M: **Revitalization procedures in two traumatized incisors with different biological outcomes.** *J Endod.* 2012; **38**(3): 411–4.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 30 March 2020

<https://doi.org/10.5256/f1000research.22088.r61455>

© 2020 Nagy M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Mohamed Mokhtar Nagy

Endodontic Department, Faculty of Dentistry, Ain Shams University, Cairo, Egypt

The manuscript focuses upon two main issues which are single visit regeneration and photo activated disinfection.

1. The title lacks the "the single visit concept".
2. The title states "oral" disinfection, this would be better as "canal" disinfection.
3. The introduction and discussion did not include the justification of the idea of single visit regeneration.
4. The introduction and discussion did not include the possible effects of laser as heat generation and effects on stem cells.
5. The manuscript did not state radiographic standardization during follow up.
6. The discussion did not justify the supposed mechanism of root canal disinfection by laser (PAD).

Is the background of the case's history and progression described in sufficient detail?

Partly

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Partly

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

No

Is the case presented with sufficient detail to be useful for other practitioners?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Endodontic regeneration

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 04 Apr 2020

Shaimaa Abdel Hafiz, Fayoum University, Fayoum, Egypt

We are pleased to submit the new version of **Case Report: Photo-activated oral disinfection in regenerative endodontics**. We appreciate the constructive criticisms of the reviewer that helped us to improve our manuscript. Herein we address their concerns. All changes appear by track changes in the revised version. Hope you find our replies satisfactory.

Response to comments of Reviewer:**1. The title lacks the "the single visit concept".**

Single visit photo-activated disinfection in regenerative endodontics. The title was modified.

2. The title states "oral" disinfection, this would be better as "canal" disinfection

I omit the word oral because the most common terminology used in literature is Photo-activated disinfection and according to the remaining title it represents the site of disinfection applied on. So its preferable to state Photoactivated disinfection as mentioned in published literature of endodontics field **(1-5)**.

3.

The introduction and discussion did not include the justification of the idea of singlevisit regeneration

For introduction:

Photodynamic therapy (PDT)/photo-activated disinfection (PAD) is considered one of the optimized single visit approaches **(6)**. This part was added to introduction section. Several *in vitro* studies have shown that PDT is effective in root canal disinfection **(1,2,7)**. A clinical study by Johns *et al.* **(3)** reported successful outcomes for using PAD in root canal treatment of completely formed root. The case report by Johns *et al.* **(4)** was the first to document pulp revascularization using PDT and platelet-rich fibrin. Root lengthening, continued thickening of the canal walls and apical closure were demonstrated at 10-month follow-up **(4)**. All of these mentioned studies applied the PAD system in single visit. As mentioned in introduction section.

For Discussion:

Regenerative endodontic procedures were completed during a single visit because PAD is

considered one of the optimized single visit disinfection approaches (6). A successful single-visit regenerative endodontic therapy of an immature permanent tooth with a chronic apical abscess was published by Shin et al. (8). They stated that a single-visit revascularization procedure has some advantages. It eliminates subsequent appointments to access the root canal environment, thus reducing the possibility of further bacterial contamination of the root canal. It also diminishes the detrimental consequences of poor patient compliance with regular follow-up evaluation (8). Topçuoğlu and Topçuoğlu (9) reported that a single-visit regenerative endodontic procedure may be a favorable treatment option for an asymptomatic immature tooth with a necrotic pulp and no periapical lesion (9). This part was added to discussion section

4. The introduction and discussion did not include the possible effects of laser as heat generation and effects on stem cells.

For introduction:

Moreover, PAD acts in a selective way. Both photosensitizer and oxygen released during bacterial cell death do not exhibit toxicity to the viable tissues (10). This part was added to introduction section.

For Discussion:

Photosensitizer and oxygen released during bacterial cell death are not reported to produce any toxicity to normal cells(10).

Dickers et al. (11) stated that the average temperature rise was lower than the 7 degrees C safety level for periodontal injury. Moura-Netto et al.(12)reported that studies on tissue engineering using stem cells from human exfoliated deciduous teeth have yielded promising results. Laser phototherapy is able to influence the proliferation and differentiation of these cells. This part was added to discussion section.

5. The manuscript did not state radiographic standardization during follow up

An individual acrylic XCP (Extension Cone Parallelizing) index was prepared by registering the bite and placed around the XCP plastic tip for radiographic standardization during follow up. This part was added to the case report section.

6. The discussion did not justify the supposed mechanism of root canal disinfection by laser (PAD).

Response:

The photosensitizer stains the bacterial cells in soft and hard tissues and the photo-activated cells release molecular oxygen causing disruption of the bacterial cell wall (10). Moreover, PAD is a newer antimicrobial strategy that involves the combination of a non-toxic PS or dyes and a non-harmful visible light source to disinfect the root canal. Low power laser in itself is not particularly lethal to bacteria, but is useful for photochemical activation of oxygen-releasing dyes (13). This part was added to discussion section.

References

- 1 Bergmans L., Moisiadis P., Huybrechts B., Van Meerbeek B., Quirynen M., Lambrechts P. Effect of photo-activated disinfection on endodontic pathogens ex vivo. *Int Endod J* 2008;41(3):227-39. Doi: 10.1111/j.1365-2591.2007.01344.x.

- 2 Williams Jill A., Pearson Gavin J., John Colles M. Antibacterial action of photoactivated disinfection {PAD} used on endodontic bacteria in planktonic suspension and in artificial and human root canals. *J Dent* 2006;34(6):363–71. Doi: 10.1016/j.jdent.2005.08.002.
- 3 Johns Dexton Antony, Varughese Jolly Mary, Thomas Kunjamma, Abraham Aby, James Elizabeth Prabha, Maroli Ramesh Kumar. Clinical and radiographical evaluation of the healing of large periapical lesions using triple antibiotic paste, photo activated disinfection and calcium hydroxide when used as root canal disinfectant. *J Clin Exp Dent* 2014;6(3). Doi: 10.4317/jced.51324.
- 4 Johns Dexton Antony, Shivashankar VasundaraYayathi, Krishnamma Shoba, Johns Manu. Use of photoactivated disinfection and platelet-rich fibrin in regenerative Endodontics. *J Conserv Dent* 2014;17(5):487. Doi: 10.4103/0972-0707.139850.
- 5 Bonsor S. J., Nichol R., Reid T. M S, Pearson G. J. Microbiological evaluation of photo-activated disinfection in endodontics (An in vivo study). *Br Dent J* 2006;200(6):337–41. Doi: 10.1038/sj.bdj.4813371.
- 6 Siqueira José F., Rôças Isabela N. Optimising single-visit disinfection with supplementary approaches: A quest for predictability. *Aust Endod J* 2011. Doi: 10.1111/j.1747-4477.2011.00334.x.
- 7 Garcez Aguinaldo S., Ribeiro Martha S., Tegos George P., Núñez Silvia C., Jorge Antonio O C, Hamblin Michael R. Antimicrobial photodynamic therapy combined with conventional endodontic treatment to eliminate root canal biofilm infection. *Lasers Surg Med* 2007;39(1):59–66. Doi: 10.1002/lsm.20415.
- 8 Shin S. Y., Albert J. S., Mortman R. E. One step pulp revascularization treatment of an immature permanent tooth with chronic apical abscess: A case report. *Int Endod J* 2009;1118–26. Doi: 10.1111/j.1365-2591.2009.01633.x.
- 9 Topçuoğlu Gamze, Topçuoğlu Hüseyin Sinan. Regenerative Endodontic Therapy in a Single Visit Using Platelet-rich Plasma and Biodentine in Necrotic and Asymptomatic Immature Molar Teeth: A Report of 3 Cases. *J Endod* 2016;42(9):1344–6. Doi: 10.1016/j.joen.2016.06.005.
- 10 López Fernanda Ullmann, Kopper Patrícia Maria Poli, Bona Alvaro Della, Steier Liviu, de Figueiredo José Antônio Poli, Vier-Pelisser Fabiana Vieira. Effect of different irrigating solutions and photo-activated therapy for in vivo root canal treatment. *Braz Dent J* 2015. Doi: 10.1590/0103-6440201300154.
- 11 Dickers B., Lamard L., Peremans A., et al. Temperature rise during photo-activated disinfection of root canals. *Lasers Med Sci* 2009;24(1):81–5. Doi: 10.1007/s10103-007-0526-y.
- 12 MOURA-NETTO Cacio, FERREIRA Leila Soares, MARANDUBA Carlos Magno, MELLO-MOURA Anna Carolina Volpi, MARQUES Marcia Martins. Low-intensity laser phototherapy enhances the proliferation of dental pulp stem cells under nutritional deficiency. *Braz Oral Res* 2016;30(1). Doi: 10.1590/1807-3107BOR-2016.vol30.0080.
- 13 Mohan Dennis, Maruthingal Sunith, Indira Rajamani, et al. Photoactivated disinfection (PAD) of dental root canal system - An ex-vivo study. *Saudi J Biol Sci* 2016. Doi: 10.1016/j.sjbs.2015.01.013.

Competing Interests: No competing interests were disclosed.

<https://doi.org/10.5256/f1000research.22088.r58468>

© 2020 Gambarini G et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Gianluca Gambarini

Department of Oral and Maxillo-Facial Sciences, Sapienza University of Rome, Rome, Italy

Dario Di Nardo

Department of Oral and Maxillo-Facial Sciences, Sapienza University of Rome, Rome, Italy

The article presents a case of apicogenesis performed by gold standard methodologies. The novelty is the use of a photoactivated disinfectant instead of a triple antibiotic paste. Since NaOCl and EDTA used are sufficient to remove all the tissues that could avoid a proper regeneration of the apex, the effectiveness of the light activated disinfectant is doubtful.

The title of the article is on regenerative endodontics, that is a procedure that aims to regenerate dental pulp, but no evidences of regeneration are presented in the article instead of the production of new dentine, which is obtainable with a simple apicogenesis procedure and also without the use of triple antibiotic paste.

The overall novelty of the article is doubtful and the evidences of the regeneration of pulp are not explained at all. Dentine formation and the closure of the apex are evidences that could be obtained with gold standard techniques that do not involve the use of other expensive appliances like diode lasers.

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Partly

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Partly

Is the case presented with sufficient detail to be useful for other practitioners?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Endodontics

We confirm that we have read this submission and believe that we have an appropriate level of expertise to state that we do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Author Response () 12 Mar 2020

Shaimaa Abdel Hafiz, Fayoum University, Fayoum, Egypt

We appreciate the comments of the reviewer. Herein we address the concerns of the reviewer.

1- The article presents a case of apicogenesis performed by gold standard methodologies. The novelty is the use of a photoactivated disinfectant instead of a triple antibiotic paste. Since NaOCl and EDTA used are sufficient to remove all the tissues that could avoid a proper regeneration of the apex, the effectiveness of the light activated disinfectant is doubtful.

Response:

- Apicogenesis is a root complementation therapy performed in teeth with pulp vitality and consists in the removal of infected coronal pulp, maintenance of vital root pulp and its protection with biocompatible material (1).

Pulpal necrosis was diagnosed on the basis of clinical findings in our case report as mentioned in the methods section of abstract. So regenerative endodontic procedure was applied for necrotic tooth with open apex as recommended by American Association of Endodontists (AAE)(2).

- Naocl and EDTA may be sufficient to remove all necrotic tissues but in this case report, it was mentioned that the procedure was performed according to the American Association of Endodontics (AAE) guidelines (2), except PAD instead of TAP was used. The AAE guidelines recommend using intracanal medication in adjunctive to the irrigants.

- There are several in vitro and in vivo studies proved the effectiveness of PAD system in root canal disinfection. And this article mentioned some of them (4–9).

2-

The title of the article is on regenerative endodontics, that is a procedure that aims to regenerate dental pulp, but no evidences of regeneration are presented in the article instead of the production of new dentine, which is obtainable with a simple apicogenesis procedure and also without the use of triple antibiotic paste.

Response:

- Regenerative Endodontics is defined as a biologically-based procedure designed to predictably replace damaged, diseased or missing structures, including dentin and root structures as well as cells of the pulp-dentin complex, with living viable tissues, preferably of the same origin, that restore the normal physiologic functions of the pulp-dentin complex (10)

- The term regenerative endodontic procedures (REPs) has been widely adopted and refers to all procedures that aim to attain organized repair of the dental pulp and include future therapies yet to regenerative endodontic treatment of necrotic immature permanent teeth resulting in continued root development, increased thickness in the dentinal walls and apical closure. evolve in the field of regenerative endodontics (3).

3-

The overall novelty of the article is doubtful and the evidences of the regeneration of pulp are not

explained at all.

Dentine formation and the closure of the apex are evidences that could be obtained with gold standard techniques that do not involve the use of other expensive appliances like diode lasers.

Response:

- Regenerative endodontic treatment of necrotic immature permanent teeth resulting in continued root development, increased thickness in the dentinal walls and apical closure as mentioned by AAE.

- AAE Clinical Considerations for a Regenerative Procedure (2) stated that the degree of success of Regenerative Endodontic Procedures is largely measured by the extent to which it is possible to attain primary, secondary, and tertiary goals:

1. Primary goal: The elimination of symptoms and the evidence of bony healing.
2. Secondary goal: Increased root wall thickness and/or increased root length (desirable, but perhaps not essential)
3. Tertiary goal: Positive response to vitality testing (which if achieved, could indicate a more organized vital pulp tissue).

- Root canal system disinfection is an integral step in the success of REPs (2). Antimicrobial approaches to disinfect root canals have been proposed that include Photo-activated disinfection (PAD) as it offers a means of destroying bacteria remaining after using conventional irrigants in endodontic therapy (4).

References

- 1 Mente Johannes, Leo Meltem, Panagidis Dimos, et al. Treatment outcome of mineral trioxide aggregate in open apex teeth. *J Endod* 2013. Doi: 10.1016/j.joen.2012.10.007.
- 2 AAE. Clinical Considerations for a Regenerative Procedure (New, Revised 6-18-16). AAE 2016.
- 3 Diogenes Anibal, Henry Michael A., Teixeira Fabricio B., Hargreaves Kenneth M. An update on clinical regenerative endodontics. *Endod Top* 2013;28(1):2-23. Doi: 10.1111/etp.12040.
- 4 Bonsor S. J., Nichol R., Reid T. M S, Pearson G. J. Microbiological evaluation of photo-activated disinfection in endodontics (An in vivo study). *Br Dent J* 2006;200(6):337-41. Doi: 10.1038/sj.bdj.4813371.
- 5 Williams Jill A., Pearson Gavin J., John Colles M. Antibacterial action of photoactivated disinfection {PAD} used on endodontic bacteria in planktonic suspension and in artificial and human root canals. *J Dent* 2006;34(6):363-71. Doi: 10.1016/j.jdent.2005.08.002.
- 6 Garcez Aguinaldo S., Ribeiro Martha S., Tegos George P., Núñez Silvia C., Jorge Antonio O C, Hamblin Michael R. Antimicrobial photodynamic therapy combined with conventional endodontic treatment to eliminate root canal biofilm infection. *Lasers Surg Med* 2007;39(1):59-66. Doi: 10.1002/lsm.20415.
- 7 Bergmans L., Moisiadis P., Huybrechts B., Van Meerbeek B., Quirynen M., Lambrechts P. Effect of photo-activated disinfection on endodontic pathogens ex vivo. *Int Endod J* 2008;41(3):227-39. Doi: 10.1111/j.1365-2591.2007.01344.x.
- 8 Johns Dexton Antony, Varughese Jolly Mary, Thomas Kunjamma, Abraham Aby, James Elizabeth Prabha, Maroli Ramesh Kumar. Clinical and radiographical evaluation of the healing of large periapical lesions using triple antibiotic paste, photo activated disinfection and calcium hydroxide when used as root canal disinfectant. *J Clin Exp Dent* 2014;6(3). Doi:

10.4317/jced.51324.

9 Johns DextonAntony, Shivashankar VasundaraYayathi, Krishnamma Shoba, Johns Manu. Use of photoactivated disinfection and platelet-rich fibrin in regenerative Endodontics. J Conserv Dent 2014;17(5):487. Doi: 10.4103/0972-0707.139850.

10 Murray Peter E., Garcia-Godoy Franklin, Hargreaves Kenneth M. Regenerative Endodontics: A Review of Current Status and a Call for Action. J Endod 2007;33(4):377-90. Doi: 10.1016/j.joen.2006.09.013.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research