Check for updates

RESEARCH ARTICLE

REVISED Prevalence of malaria and quantification of cytokine levels during infection in East Nile locality, Khartoum State: a cross-sectional study [version 2; peer review: 2 approved]

Hwida Barkat¹, Ahmed Bakheet Abd Alla ², Ahmed Galander², Tagwa Salah², Tayseer Elfaki², Ali Nasir²

¹Laboratory Administration, Ministry of Health, Sudan, East Nile Locality, Khartoum, Sudan

²Parasitology and Medical Entomology, College of Medical Laboratory Science, Sudan University of Science and Technology, Khartoum, Sudan

v2 First published: 29 Aug 2019, 8:1529
<https://doi.org/10.12688/f1000research.19217.1>
 Latest published: 16 Aug 2021, 8:1529
<https://doi.org/10.12688/f1000research.19217.2>

Abstract

Background: The cytokines interferon gamma (IFN- γ), interleukin-10 (IL-10) and tumor necrosis factor- α (TNF- α) play an important role in malaria infection. The aim of this study was to determine the prevalence of malaria and to evaluate cytokine responses to malaria infection in patients from the East Nile locality of Khartoum State.

Methods: This study was carried out from May to July 2018 in the East Nile Locality, Khartoum State. Blood samples were collected from 384 randomly selected patients for blood film analysis. Of these, 39 were selected for cytokine level analysis (10 control and 29 patient samples), determined using enzyme-linked immunosorbent assays.

Results: The malaria prevalence rate among 384 patients was 18.5%. *Plasmodium falciparum* was the most prevalent (13%), while the prevalence of *Plasmodium vivax* was 4.6%. The rate of mixed infection was 0.8%. There was a higher prevalence rate (22.7%) in males than females (15.6%). However, we found no significant correlation between cytokine levels and parasitemia in the study group. Nevertheless, our study demonstrated a significant correlation between cytokine levels and recurrent infections.

Conclusions: Together, our data show that malaria remains a public health problem in East Nile locality with a high prevalence. Additionally, serum levels of IFN- γ and TNF- α were significantly higher in malaria-infected individuals compared to non-infecting individuals and cytokine levels were found to be correlated with recurrent malaria infection.

Keywords

cytokines, IL - 10, TNF- α , IFN- γ , recurrent, malaria, Plasmodium

Open Peer Review

Approval Status

	1	2
version 2		
(revision)
16 Aug 2021
version 1
29 Aug 2019

1. **Yaowapa Maneerat**, Mahidol University, Bangkok, Thailand

2. **Rusliza Basir** , University Putra Malaysia (UPM), Serdang, Malaysia

Any reports and responses or comments on the article can be found at the end of the article.

This article is included in the **Pathogens** gateway.

This article is included in the **Emerging Diseases and Outbreaks** gateway.

Corresponding author: Ahmed Bakheet Abd Alla (ahmed.hassanab@gmail.com)

Author roles: **Barkat H:** Formal Analysis; **Abd Alla AB:** Supervision, Writing – Original Draft Preparation; **Galander A:** Methodology; **Salah T:** Data Curation; **Elfaki T:** Software; **Nasir A:** Visualization

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2021 Barkat H *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Barkat H, Abd Alla AB, Galander A *et al.* **Prevalence of malaria and quantification of cytokine levels during infection in East Nile locality, Khartoum State: a cross-sectional study [version 2; peer review: 2 approved]** F1000Research 2021, 8:1529 <https://doi.org/10.12688/f1000research.19217.2>

First published: 29 Aug 2019, 8:1529 <https://doi.org/10.12688/f1000research.19217.1>

REVISED Amendments from Version 1

In this new version we do some editing, we edit the write name of IFN- γ in all manuscript, as we add in conclusion in abstract section the result of cytokines which was not found in version 1.

In introduction part we add the rational and objective of our study, also we correlate between the level of cytokines between *P.falciparum* and *P.vivax*, also we merge the Table 1, Table 2 and Table 3.

Any further responses from the reviewers can be found at the end of the article

Introduction

Malaria is a mosquito-borne disease that affects humans and animals. This condition is caused by a protozoan parasite of the genus *Plasmodium*. Malaria symptoms include fever and headache, which result from parasite invasion of red blood cells. In severe cases, malaria infection may progress to coma or death (Elmardi *et al.*, 2011). There are five *Plasmodium* species that are known to cause disease in humans (*P. falciparum*, *P. vivax*, *P. ovale*, *P. malariae*, and the recently described *P. knowlesi*). The species that causes the most severe cases is *P. falciparum*. Malaria control efforts in Sudan began in 1904, when Dr. A. Balfour succeeded in eradicating malaria from Khartoum (Malik *et al.*, 2006). Determination of the correct prevalence of malaria is essential in implementing effective control strategies to curb its dissemination (Malik *et al.*, 2006).

Immunity to *Plasmodium* develops slowly and protection against *Plasmodium* occurs later than protection against malaria symptoms. The immune responses will not be same for the liver and blood stages because *Plasmodium* expresses various antigens at the liver and blood stages (Langhorne, 2005). Cytokines may assume a significant role in protection and pathology in malaria. The early effective inflammatory reaction is regulated by IFN- γ , IL-12. TNF- α appears to be crucial for parasitemia control in malaria infection (Artavanis-Tsakonas *et al.*, 2003). However, pro-inflammatory cytokines (TNF- α , IFN- γ , IL-1 and IL-6) were associated with severe malaria (Malaguarnera & Musumeci, 2002). The high production of proinflammatory cytokines may increase cytoadherence of parasitized red blood cells to the endothelium through upregulation of adhesion molecules in *P. falciparum* infections (Day *et al.*, 1999).

The expression of cytokines (pro- and anti-inflammatory) are said to be involved in malaria pathogenesis. Severe malaria has been associated with low serum levels of IL-12 and low IL-10 to TNF- α serum concentration ratios in a few studies of childhood malaria in holoendemic areas.

In order to explore the effect of the immune response to malaria and the development of clinical immunity, this study aimed to measure and determine the prevalence of malaria and quantify cytokine levels in patients with malaria infection in the East Nile locality.

Methods

This cross-sectional study was carried out in the East Nile locality, which is located in the eastern part of Khartoum State, Sudan. This study was conducted in different pre-urban areas during the period from 1st May to 23rd of July 2018.

Study population

The study population included participants of all ages and genders from a population admitted to Elbanjaded Hospital, Aldebaba Medical Health Center, Eid Babekir Medical Health Center, Helat Koko Medical Health Center and Omdom Medical Health Center. In total, 384 participants were asked to participate in this study, all of whom were admitted for malaria diagnosis. Participants were selected using quota sampling and all patients admitted for malaria diagnosis were eligible to be included in the study. The age groups were categorized as follows: less than 10 years, 11–49 years and over 50 years old. After participants signed an informed consent form for participation in the study, a questionnaire was used by expert laboratory technician to collect demographic data (information about age, sex, residence and occupation) and medical history of chronic disease (renal disease, heart disease and diabetes mellitus or other chronic disease) from patients enrolled in the study.

Sample size

In total, 384 samples were collected from patients admitted for malaria infection at hospitals and health centers. The sample size was calculated on the following formula (Daniel, 1999):

$$N = Z^2 p(1-p)/d^2$$

where N = sample size, Z = statistic for a level of confidence (1.96), P = prevalence in study area (50%) and d = precision (5%).

Sample collection

From each patient, thick and thin blood films were prepared using finger prick blood samples taken as part of the routine diagnosis of infection with *Plasmodium* species. Thin films were fixed with methanol and slides were placed face down on a drying rack for five minutes to allow the methanol to fix. Thick and thin blood films were stained with Giemsa stain at a concentration of 10% for 10 minutes. The stain was flushed from the slides by adding drops of buffered water until all the stain has been washed away. Where the blood film analysis was positive for malaria, 5 ml of venous blood was collected from each patient into a sterile container. For cytokine analysis, 5ml of venous blood was also collected from 10 participants who tested negative for malaria and agreed to participate further in the study. These participants were selected using stratified random sampling, with two participants being randomly selected from each of the five health centers. Following collection, blood samples were centrifuged at 3000 rpm for 10 minutes. After centrifugation, the serum was separated and transferred to another labeled sterile container and stored in refrigerator at 4°C until use.

Microscopic examination

After the films dried, they were examined microscopically by experienced personal to determine the parasite stages (ring,

trophozoite, gametocyte and schizont), using the thin blood film to identify the species of *Plasmodium* and the thick film to classify parasitemia as follows:

- + : 1–10 parasites per 100 thick film fields
- + + : 11–100 parasites per 100 thick film fields
- + + + : 1–10 parasites per one thick film field
- + + + + : more than 10 parasites per one thick film field

Measurement of cytokines

For cytokine analysis, 29 of the patient serum samples were selected using simple stratified sampling, with five positive samples randomly selected from each health center and nine randomly selected from the hospital. Cytokine analysis was not performed for all samples due to financial restrictions. The stored serum was brought to the laboratory and was allowed to thaw. Serum concentrations of IFN- γ , IL-10, TNF- α were determined using an enzyme-linked immunosorbent assay (ELISA) according to manufacturer's instructions (BioLegend's ELISA MAXTM Deluxe Sets, catalog numbers 430104, 430604, and 340204 for IFN- γ , IL-10, and TNF- α , respectively) for the patient samples and 10 control samples.

Briefly, 100 μ L of diluted capture antibody solution was added to each well and the sealed plate was incubated overnight between 2–8°C. The plates were washed four times and then blocked by adding 200 μ L assay diluents to each well, then were sealed and incubated for one hour with shaking on a palate shaker at 500 rpm with a 0.3cm circular orbit. The plates were washed four times and then 100 μ L diluted standards and samples were added to each well. The plate was sealed and incubated at room temperature for two hours with shaking. The plate was washed four times, then 100 μ L diluted detection antibody solution was added to each well. Plates were sealed and incubated at room temperature for one hour with shaking. The plate was washed four times, then 100 μ L diluted avidin–HRP solution was added to each well. The plate sealed and was incubated at room temperature for 30 minutes with shaking. The plate was washed five times and then soaked for 30 seconds to one minute per wash. Then, 100 μ L fresh TMB substrate solution was added to each well and incubated in the dark for 20 minutes. Finally, 100 μ L of stop solution was added to each well and the absorbance was read with the SPECTROstar Nano Microplate Reader at 540 nm and 570 nm within 15 minutes.

Statistical analysis

Data were analyzed using SPSS version-20. The Chi-squared test was performed to determine statistical significance and a *P*-value of less than 0.05 was considered statistically significant.

Ethical statement

Ethical clearance for this study was obtained from Committee of Scientific Research Deanship, Sudan University of Science and Technology, ethical approval number (DSR – IEC – 12 – 07). Written informed consent for participation and publication

of the data was obtained from all participants included in this study or for children, from their guardian.

Results

A total of 384 patients were enrolled in this study, of which 154 were male and 230 were female. Their ages were grouped into three categories: less than 10 years old (134), 11 – 49 years old (198) and more than 50 years old (52).

Prevalence of malaria in patients from the study area

First, we sought to determine the prevalence of malaria in our study population. To this end, we collected blood samples from our patients and examined them using blood films. Out of 384 blood samples collected from different pre-urban areas (medical centers and hospitals) in the East Nile locality during the period from May to July 2018 (pre- malaria season), 71 (18.5%) were found to be positive and 313 (81.5 %) were negative for malaria (Table 1) (Abd Alla & Brakat, 2019). Moreover, we observed a higher prevalence of malaria among males (22.7%) compared to females (15.6%) in our study population (Table 1). In addition, analysis of the prevalence among different age groups revealed that highest prevalence rate was in the under 10 age group (20.1%), followed by the 11–49 age group with a prevalence of 19.7%, while the lowest prevalence rate was reported among the over 50 age group (2%) (Table 1).

Distribution of *Plasmodium* species in the study area

Next, we determined the species distribution of *Plasmodium* species in our study population. We observed that *P. falciparum* had the highest prevalence rate (13%), followed by *P. vivax* (4.6%). However, mixed infection by *P. falciparum* and *P. vivax* had the lowest prevalence rate (0.8%). We failed to

Table 1. Overall prevalence of malaria in study area, assessed using blood films.

Blood film	Frequency	Percentage
Positive	71	18.5
Negative	313	81.5
Total	384	100.0
Gender	Number examined	Number positive (%)
Male	154	35 (22.7)
Female	230	36 (15.6)
Total	384	71 (18.5)
Age group		
≤ 10	134	27 (20.1)
11 – 49	198	39 (19.7)
≥ 50	52	5 (2)
Total	384	71 (18.5)

detect any positive results for *P. malariae* or *P. ovale* infection in our study population (Table 2).

Our data reveal that there was no significant statistical association between *P. falciparum* and *P. vivax* in cytokines level, P-value were (0.436, 0.789 and 0.530) for $INF-\gamma$, $TNF-\alpha$ and IL-10 respectively.

Severity of malaria infection among the study population

Next, we determined the distribution of severe malaria in the studied population. Our data showed that most (58%) of the study population had a low parasite count (mild parasitemia), while 13% had a moderate parasitemia (++). We detected no cases that exhibited severe parasitemia (+++ and +++) (Table 3) and no statistically significant association between age group and parasitemia (Table 4).

Cytokine profiles in malaria-infected patients

Cytokines may play a role in protection and pathology in malaria. Thus, we investigated the cytokine profile in 29 patients and 10 controls from our study population. In particular, we investigated serum levels of $INF-\gamma$, $TNF-\alpha$, and IL-10 with malaria infection in the study population. Interestingly, mean serum levels of $INF-\gamma$ were significantly higher in malaria-infected individuals compared to non-infected individuals (P value = 0.026). However, $TNF-\alpha$ serum levels were comparable between patients and non-infected individuals (P value = 0.646). Mean serum levels of IL-10 were higher in patients compared to non-infected individuals, although this

Table 2. Distribution of *Plasmodium* species in the study area.

Species	Frequency (%)
<i>P. falciparum</i>	51 (13)
<i>P. vivax</i>	17 (4.6)
<i>P. ovale</i>	0 (0)
<i>P. malariae</i>	0 (0)
<i>P. falciparum</i> and <i>P. vivax</i> co-infection	3 (0.8)
Total	71 (18.5)

Table 3. Prevalence of malaria according to parasite count.

Parasite count	Frequency (%)
+	58 (81.7)
++	13 (18.3)
+++	0 (0)
++++	0 (0)
Total	71 (18.5)

Table 4. Correlation between parasitemia and age.

Density	Age group (%)			Total	P value
	≤ 10	11–49	≥ 50		
+	20(34.5)	31(53.4)	7(12.1)	58	0.108
++	7(53.8)	6(46.2)	0	13	
+++	0	0	0	0	
++++	0	0	0	0	

difference was not statistically significant (P value = 0.071) (Table 5).

Next, we sought to determine whether there is a correlation between the cytokine profiles of individuals enrolled in our study and severity of malaria and/or recurrent infection. However, we found no correlation between levels of $INF-\gamma$, $TNF-\alpha$, or IL-10 and level of parasitemia (Table 6). Intriguingly, levels of $TNF-\alpha$ and IL-10 were significantly higher in patients who suffered from recurrent malaria infection compared to those who did not (Table 7). However, we failed to detect a significant correlation between levels of $INF-\gamma$ and recurrent malaria infection (Table 7).

Discussion

The findings of our study revealed a prevalence rate of malaria of 18.5%. This rate was greater than the rate reported in Khartoum by El Mekki *et al.* (2012), who reported the prevalence of malaria in 5% and 11% in Dar Al Salam Camp and Jabal Awlia Camp, respectively. El Sayed *et al.* (2000) reported that Khartoum, which was formerly malaria free, can be considered as a hypoendemic or mesoendemic area in which malaria is unstable and epidemic outbreaks are common; our results agree with their findings.

Our study show that Falciparum malaria is the most prevalent and constitutes about 13% of all infections, benign tertian Vivax malaria has prevalence of 4.6%, and the lowest prevalence rate of 0.8% is observed for mixed infection (*P. falciparum* and *P. vivax*). However, we observed no cases of *P. malariae* and *P. ovale* infection. Moreover, males had higher prevalence rate (22.7%) of malaria infection than females (15.6%). Our study findings agree with a study in Khartoum by Abdalla *et al.* (2007), who reported that the overall prevalence of malaria was 28.2 % and was higher in males than in females.

The highest prevalence rate (53.8%) of moderate parasitemia was in the under 10 years age group. Although children are more susceptible to malaria infection due to a slow developing immune system, a high prevalence rate (34.5%) of mild parasitemia was reported among the 11–49 years age group. A lower prevalence rate of 2% and mild parasitemia 12.1% was reported among the over 50 age group. This finding was closer to the finding of Igwe *et al.* (2014) in Nigeria, who reported that the highest prevalence of asymptomatic malaria parasitemia

Table 5. Cytokine profiles of the study population.

Sample	Number	Cytokine profiles (mean $\pm$ SD)		
		IFN- γ (ng/ml)	TNF- α (ng/ml)	IL-10 (ng/ml)
Patients	29	61.98 $\pm$ 71.93	5.91 $\pm$ 5.67	48.87 $\pm$ 52.99
Controls	10	11.05 $\pm$ 11.05	5.36 $\pm$ 3.78	12.48 $\pm$ 8.08
P value		0.026	0.646	0.071

Table 6. Correlation between parasitemia and cytokine profiles.

Density	Number	Mean		
		IFN- γ (ng/ml)	TNF- α (Ng/ml)	IL-10 (Ng/ml)
+	25	79.49	5.80	60.50
++	4	79.94	7.98	67.17
P value		0.922	0.525	0.380

Table 7. Correlation between recurrent malaria infection and cytokine profiles.

Recurrent malaria infection (yes/no)	Number	Mean (ng/ml)	SD (ng/ml)	P value
IFN- γ				
Yes	15	83.06	70.61	0.151
No	24	48.82	71.02	
TNF- α				
Yes	15	8.76	8.10	0.011
No	24	4.13	2.03	
IL-10				
Yes	15	77.21	65.55	0.007
No	24	31.16	34.26	

(87.5%) was found in parturient women who were ≤ 19 years, while the lowest prevalence (68.2%) occurred in those who were 40–49 years old. In the present study we observed that age group was not significantly associated with parasitemia. This is in agreement with a study done by [El Khalifa et al. \(2008\)](#), who found no significant difference in parasitemia among those aged five years and above.

In this study, serum levels IFN- γ , TNF- α and IL-10 were measured in healthy controls and in patients with *P. falciparum* and *P. vivax* infection and IFN- γ was found to be significantly higher in patients than in non-infected individuals. This finding is in line with a study in Poland by [Wroczynska et al. \(2005\)](#), who reported that the mean serum level of IFN- γ was found to be significantly higher in severe and uncomplicated malaria groups compared to the controls. Also, another study done by

[Favre et al. \(1997\)](#) reported that these findings are consistent with a requirement for an early production of IFN- γ to mount resistance against infection. Interestingly, in this study, a significant correlation between IL-10 with gender and age was found. This association between initial IL-10 levels and parasite densities agreed in part with the findings of [Hugosson et al. \(2004\)](#), who reported similar findings during patient treatment, indicating that IL-10 levels may play a role in clearance of parasites during treatment. In addition, they also suggest that there are age-related differences in immunity and the development of partial clinical tolerance.

The present study reported no association between parasite density and levels of IFN- γ , TNF- α and IL-10. These findings are in line with study done by [Jason et al. \(2001\)](#) in which serum IL-10 levels had statistically significant association with level of parasitemia. Furthermore, our data agrees with a study by [Nnaemaka et al. \(2009\)](#), who found no significant correlation between IL-10, IL-12 and IFN- γ in asymptomatic individuals with parasitemia; however, [Wroczynska et al. \(2005\)](#) found that IL-10 and IL-12 were associated with malaria. In this study, there was significantly increased production of IL-10 and TNF- α in patients with recurrent malaria. This finding was in agreement with a study done by [Edward et al. \(2008\)](#), who reported that the high levels of IL-10 observed during malarial episodes may be beneficial, acting to reduce the inflammatory response. However, they may also be detrimental and decrease antiparasitic cellular immune responses. This is suggested by our data, with significant levels of TNF- α found in patients with recurrent malaria infection. This is in line with [Medzhitov et al. \(2012\)](#), who reported that these observations are predictable, with the possibility that recurrent malaria may drive the host towards a disease tolerant state, so as to diminish the negative effects of disease-related pathology. In subjects who are routinely infected by malaria, the pro-inflammatory response may be immediately controlled by regulatory mechanisms. This impact might be particularly exaggerated in this study area, where transmission is particularly extreme.

Data availability

Underlying data

Figshare: hoda datta.sav. <https://doi.org/10.6084/m9.figshare.8986178.v2> (Abd Alla & Brakat, 2019)

This project contains the following underlying data:

- hoda datta.sav (demographic, behavioral and medical data for each participant and results of the microscopic examination)
- samle and control.sav (cytokine levels for 39 patient and control samples, determined using ELISA)

- Data dictionary_FL.docx

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC0 1.0 Public domain dedication).

Acknowledgements

We would like to express our immense gratitude and appreciation to the Deanship of Scientific Research-Sudan University of Science and Technology, who supported this project by providing space in which to carry out the cytokine analysis performed in this study.

References

- Abd Alla A, Brakat HE: **hoda datta.sav**. 2019. <http://www.doi.org/10.6084/m9.figshare.8986178.v1>
- Abdalla SI, Malik EM, Ali KM: **The burden of malaria in Sudan: incidence, mortality and disability-adjusted life-years**. *Malar J.* 2007; **6**(1): 97. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Artavanis-Tsakonas K, Tongren JE, Riley EM: **The war between the malaria parasite and the immune system: immunity, immunoregulation and immunopathology**. *Clin Exp Immunol.* 2003; **133**(2): 145–52. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Daniel WW: **Biostatistics: A Foundation for analysis in the health sciences**. 7th ed, R Wiley. New York. 1999. [Reference Source](#)
- Day NP, Hien TT, Schollaardt T, et al.: **The prognostic and pathophysiologic role of pro- and antiinflammatory cytokines in severe malaria**. *J Infect Dis.* 1999; **180**(4): 1288–97. [PubMed Abstract](#) | [Publisher Full Text](#)
- Edward RK, Atis M, Michal F, et al.: **Maternal peripheral blood level of IL-10 as a marker for inflammatory placental malaria**. *Malar J.* 2008; **7**(1): 26. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- El Khalifa SM, Mustafa IO, Wais M, et al.: **Malaria control in an urban area: a success story from Khartoum, 1995-2004**. *East Mediterr Health J.* 2008; **14**(1): 206–215. [PubMed Abstract](#)
- Elmardi KA, Noor AM, Githinji S, et al.: **Self-reported fever, treatment actions and malaria infection prevalence in the northern states of Sudan**. *Malar J.* 2011; **10**(1): 128. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- El Mekki MA, Aburas NA, Alghaithy AA, et al.: **Prevalence and Molecular Identification of Malaria Parasite in Displaced Camps in Khartoum State, Sudan**. 2012; **4**(1): 7–12. [Publisher Full Text](#)
- El Sayed BB, Arnot DE, Mukhtar MM, et al.: **A study of the urban malaria transmission problem in Khartoum**. *Acta Trop.* 2000; **75**(2): 163–71. [PubMed Abstract](#) | [Publisher Full Text](#)
- Favre N, Ryffel B, Bordmann G, et al.: **The course of *Plasmodium chabaudi* infections in interferon-gamma receptor deficient mice**. *Parasite Immunol.* 1997; **19**(8): 375–83. [PubMed Abstract](#) | [Publisher Full Text](#)
- Hugosson E, Montgomery SM, Premji Z, et al.: **Higher IL-10 levels are associated with less effective clearance of *Plasmodium falciparum* parasites**. *Parasite Immunol.* 2004; **26**(3): 111–7. [PubMed Abstract](#) | [Publisher Full Text](#)
- Igwe NM, Joannes UO, Chukwuma OB, et al.: **Prevalence and parasite density of asymptomatic malaria parasitemia among unbooked patients at Abakaliki, Nigeria**. *Journal of Basic and Clinical Reproductive Sciences.* 2014; **3**(1): 44–8. [Reference Source](#)
- Jason J, Archibald LK, Nwanyanwu OC, et al.: **Cytokines and malaria parasitemia**. *Clin Immunol.* 2001; **100**(2): 208–18. [PubMed Abstract](#) | [Publisher Full Text](#)
- Langhorne J: **Immunology and immune pathogenesis of malaria**. National institute for medical research, the Ridgeway, Mill Hill, London, NW7 1AA, UK. 2005. [Publisher Full Text](#)
- Malaguarnera L, Musumeci S: **The immune response to *Plasmodium falciparum* malaria**. *Lancet Infect Dis.* 2002; **2**(8): 472–8. [PubMed Abstract](#) | [Publisher Full Text](#)
- Malik EM, Hanafi K, Ali SH, et al.: **Treatment-seeking behaviour for malaria in children under five years of age: implication for home management in rural areas with high seasonal transmission in Sudan**. *Malar J.* 2006; **5**(1): 60. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Medzhitov R, Schneider DS, Soares MP: **Disease tolerance as a defense strategy**. *Science.* 2012; **335**(6071): 936–41. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nnaemeka C, Okafor CM, Idowu Ayede Y, et al.: **Cytokine profiles and antibody responses to *Plasmodium falciparum* malaria infection in individuals living in Ibadan, southwest Nigeria**. *Afr Health Sci.* 2009; **9**(2): 66–74. [PubMed Abstract](#) | [Free Full Text](#)
- Wroczyńska A, Nahorski W, Bąkowska A, et al.: **Cytokines and clinical manifestations of malaria in adults with severe and uncomplicated disease**. *Int Marit Health.* 2005; **56**(1–4): 103–14. [PubMed Abstract](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 14 September 2021

<https://doi.org/10.5256/f1000research.76573.r92038>

© 2021 Basir R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Rusliza Basir

Department of Human Anatomy, Faculty of Medicine and Health Sciences, University Putra Malaysia (UPM), Serdang, Malaysia

The authors have attended to all the comments satisfactorily.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Immunotherapeutic target for malaria infection.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 20 August 2021

<https://doi.org/10.5256/f1000research.76573.r92037>

© 2021 Maneerat Y. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Yaowapa Maneerat

Department of Tropical Pathology, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand

I confirm that I have read the revised article carefully. The authors have already corrected some points according to my suggestion. I approve the revised article. However, I feel that Table 2 and 3 (with only 2 columns) are not necessary. If possible, the authors should describe completely using 2-3 sentences.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Pathology of malaria

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 03 July 2020

<https://doi.org/10.5256/f1000research.21057.r66205>

© 2020 Basir R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Rusliza Basir

Department of Human Anatomy, Faculty of Medicine and Health Sciences, University Putra Malaysia (UPM), Serdang, Malaysia

1. It was mentioned that patients serum samples were selected randomly for cytokines analysis. How does this random selection carry out? Does it take into consideration the age of patients since they are categorized according to age at the beginning? There would be differences in cytokine release profile between child, adult and elderly as the level of immune response between the three groups towards invading organisms would be different. So how does the selection carry out among the three groups of patients?
2. The symbol for gamma for IFN must be presented properly with the right symbol... not "y".
3. What is the age profile or category of the 29 patients in whom the cytokine profile was carried out? It would be interesting to see if the cytokines release is influenced by age.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Immunotherapeutic target for malaria infection.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 16 January 2020

<https://doi.org/10.5256/f1000research.21057.r58145>

© 2020 Maneerat Y. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Yaowapa Maneerat

Department of Tropical Pathology, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand

As per my knowledge, there have been several studies in roles of cytokine profiles in human and animal malaria. This study aimed to determine important cytokines in malaria. These included IFN- γ , IL-12, TNF- α . They used rather appropriate sample size. However, as follows, it was found some points are unclear and should be clarified.

1. The conclusion in the abstract is not clear. The authors did not indicate any cytokines.
2. The rationale and the benefit of study is not clear.
3. The author did not show the difference in any cytokine levels between *P. falciparum* and *P. vivax* infected patients.
4. It seems that there are too many tables. Is it possible to merge Table 1 and 2; 3 and 6?
5. Correlation between parasitemia and cytokines in Table 8 is not reliable. Sample size (n=4) is too small for determining correlation.
6. The authors did not mention about detail of patients in ELISA results. Why sample size was 10 control and 29 patients?

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

No

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Immunopathology in malaria.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research