RESEARCH ARTICLE

Health and saliva microbiomes of a semi-urbanized indigenous tribe in Peninsular Malaysia [version 1; peer review: 2 approved with reservations]

Li-Fang Yeo ^{1,2}, Farhang F. Aghakhanian^{1,2}, James S. Y. Tan³, Han Ming Gan ⁴, Maude E. Phipps¹

¹Jeffrey Cheah School of Medicine and Health Sciences, Monash University, Bandar Sunway, Selangor, 46150, Malaysia

²Tropical Medicine and Biology Platform, Monash University, Bandar Sunway, Selangor, 46150, Malaysia

³Imperial College London, South Kensington Campus, London, SW7 2AZ, UK

⁴School of Life & Env Sciences, Deakin University, Geelong Waurn Ponds Campus, Australia

V1 First published: 11 Feb 2019, 8:175
<https://doi.org/10.12688/f1000research.17706.1>
 Second version: 09 May 2019, 8:175
<https://doi.org/10.12688/f1000research.17706.2>
 Latest published: 28 May 2019, 8:175
<https://doi.org/10.12688/f1000research.17706.3>

Abstract

Background: The indigenous people of Peninsular Malaysia, also known as Orang Asli, have gradually been urbanized. A shift towards non-communicable diseases commonly associated with sedentary lifestyles have been reported in many tribes. This study engaged with a semi-urbanized Temiar tribe from Kampong Pos Piah, Perak, who are experiencing an epidemiological transition.

Methods: Weight, height, waist circumference, blood pressure, HbA1C and lipid levels were measured as indicators of cardio-metabolic health. DNA was extracted from saliva using salting-out method followed by PCR amplification of the V3-V4 region of the 16S rRNA gene and sequencing on Illumina MiSeq. Microbiome analysis was conducted on Qiime v1.9. Statistical analysis was conducted using Qiime v1.9 and R.

Results: The study revealed that 60.4% of the Temiar community were overweight/obese, with a higher prevalence among women. HbA1C levels showed that 45% of Temiar had pre-diabetes. Insulin resistance was identified in 21% of Temiar by using a surrogate marker, TG/HDL. In total, 56.5% of Temiar were pre-hypertensive, and the condition was prevalent across all age-groups. The saliva microbiome profiles of Temiar revealed significant differences by gender, BMI, abdominal obesity as well as smoking status. The relative abundance of *Bifidobacterium*, bacteria commonly found in dairy products, was increased in men. *Prevotella*, associated with consumption of plant-rich diets, was increased in women. *Mogibacteriaceae* and *Mycoplasma* levels were significantly elevated in overweight individuals. Proteobacteria was significantly depleted in smokers.

Open Peer Review

Approval Status

	1	2
version 3		
(revision)	view	
28 May 2019		
version 2		
(revision)	view	view
09 May 2019		
version 1		
11 Feb 2019	view	view

1. **Andres Gomez** , University of Minnesota, Saint Paul, USA

2. **Siti Nursheena Mohd Zain** , University of Malaya, Kuala Lumpur, Malaysia

Any reports and responses or comments on the article can be found at the end of the article.

Conclusions: Temiar from Pos Piah had a high prevalence of cardio-metabolic risks, including general and abdominal obesity, pre-diabetes, prehypertension and hypertension. This phenomenon has not been previously reported in this tribe. The saliva microbiome profiles were significantly different for individuals of different gender, BMI scores and abdominal obesity and smoking status.

Keywords

Orang Asli, saliva microbiome, anthropometrics, cardio-metabolic health, indigenous people

Corresponding author: Maude E. Phipps (maude.phipps@monash.edu)

Author roles: **Yeo LF:** Formal Analysis, Investigation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Aghakhanian FF:** Formal Analysis, Investigation, Resources, Validation, Writing – Review & Editing; **Tan JSY:** Data Curation, Investigation, Validation, Writing – Review & Editing; **Gan HM:** Funding Acquisition, Investigation, Methodology, Resources, Supervision, Writing – Review & Editing; **Phipps ME:** Conceptualization, Funding Acquisition, Methodology, Project Administration, Resources, Supervision, Validation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This study was funded by Tropical Medicine and Biology Platform (TMB), Monash University Malaysia seed grant TMB -2016-HG3185140920-YLF/MP awarded to LFY and MEP. Ms Li-Fang Yeo is supported by a scholarship awarded by Jeffrey Cheah School of Medicine and Health Sciences, Monash University Malaysia.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 Yeo LF *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Yeo LF, Aghakhanian FF, Tan JSY *et al.* **Health and saliva microbiomes of a semi-urbanized indigenous tribe in Peninsular Malaysia [version 1; peer review: 2 approved with reservations]** F1000Research 2019, 8:175 <https://doi.org/10.12688/f1000research.17706.1>

First published: 11 Feb 2019, 8:175 <https://doi.org/10.12688/f1000research.17706.1>

Introduction

The Orang Asli (OA), which means “original people” in the Malay language, comprise approximately 0.5% (150,000) of the total Malaysian population¹. They are categorized into three main groups, namely Negrito, Senoi and Proto Malay. OAs are widely spread across the Peninsular and range from semi-nomadic deep forest hunter-gatherers such as the Jahai to resettled communities such as Mah Meri to urbanized city-fringe dwellers such as Orang Seletar¹. This study focused on the Temiar who are a subtribe of Senoi and are believed to be descendants of the first Neolithic farmers who migrated to the Malay Peninsula².

In recent years, many OA communities were resettled by the government in effort to improve their lives. As the OA became more urbanized and by large left their ancestral habitats and practices, they led more sedentary lifestyles. These factors, coupled with loss of access to forest resources and increasing pressures to turn to store-bought food, may largely explain the rise in cardio-metabolic diseases such as hypertension, diabetes and obesity shown in recent studies^{1,3,4}.

The launch of the Human Microbiome Project heralded the unprecedented investigations of various microbiomes. Of these, oral microbiomes had been widely studied in human health and diseases^{5,6}. Studies implied an oral origin to systemic diseases such as cardiovascular diseases as the oral cavity is a major gateway into the body^{7,8}. Studies also investigated associations between oral microbiome and diabetes^{9,10} and obesity¹⁰, with mixed results. There were suggestions that obese people may

have a different salivary bacterial composition perhaps akin to inflammation, which contributed to periodontal diseases and caries¹¹.

Little is known about the microbiomes of indigenous communities in Asia. To our knowledge, this was a pioneering investigation of their saliva microbiomes. Furthermore, biomedical studies of Temiar were sparse and outdated, despite them being a very large community. With this impetus, our study aimed to address the gap in knowledge by reporting on the anthropometrics and cardio-metabolic health of a resettled Temiar community and investigated their saliva microbiome in association with their health.

Results

Anthropometrics and cardio-metabolic health

A total of 72 Temiars, 33 men and 39 women, participated in the study. The median age was 34 years old. General and abdominal obesity had higher prevalence among Temiar women (Table 1). Notably, 71.4% (n=25) of women and 28.1% (n=9) of men displayed abdominal obesity.

HbA1C levels indicated 44.9% of Temiar to be pre-diabetic, with a higher prevalence in men. The high prevalence of prediabetes is worrying because it indicates a rise in non-communicable diseases that was previously of low prevalence in rural communities. Using a TG/HDL as a surrogate marker for insulin resistance, 22% of Temiar were at risk of IR, mostly affecting men.

Table 1. Anthropometrics and cardio-metabolic risk factors among Temiar.

Variable	Whole, n (%)	Men, n (%)	Women, (%)
General obesity (BMI)			
Underweight (<18.5)	7 (10.3%)	4 (12.5%)	3 (8.3%)
Normal (18.5-22.9)	20 (29.4%)	12 (37.5%)	8 (22.2%)
Overweight (23-24.9)	15 (22.1%)	6 (18.8%)	9 (25%)
Pre-obese (25-29.9)	22 (32.4%)	10 (31.3%)	12 (33.3%)
Obese (≥30)	4 (5.9%)	0 (0%)	4 (11.1%)
Abdominal obesity (waist circumference)			
Healthy (M: <90cm; W: <80cm)	-	23 (71.9%)	10 (28.6%)
Risk (M: ≥90cm; W: ≥80cm)	-	9 (28.1%)	25 (71.4%)
HbA1C level			
Normal (4.0-5.6%)	36 (52.2%)	15 (46.9%)	21 (56.8%)
Pre-diabetes (5.7-6.4%)	31 (44.9%)	17 (53.1%)	14 (37.8%)
Diabetes	2 (2.9%)	0 (0%)	2 (5.4%)
Insulin resistance (TG/HDL 0.9-1.7)			
Normal	51 (78%)	20 (75%)	32 (94%)
Risk	14 (22%)	10 (15%)	4 (6%)
Blood pressure			
Normotensive	14 (20.3%)	7 (21.9%)	7 (18.9%)
Pre-hypertensive	39 (56.5%)	14 (43.8%)	24 (64.9%)
Stage 1	12 (17.4%)	9 (28.1%)	3 (8.1%)
Stage 2	4 (5.8%)	2 (6.3%)	3 (8.1%)

Blood pressure measurements showed that 56.5% (n=39) had pre-hypertension, which was more prevalent among women and was prevalent across all age groups. Stage 1 hypertension prevalence rate was 17.4% (n=12) and was found more prevalent among men. Raw data for these measurements are available in figshare¹².

Saliva microbiome analysis

To analyse the saliva microbiota diversity, the V3-V4 hypervariable region on the 16S rRNA gene was amplified and sequenced on Illumina MiSeq. After data quality control (QC), a total of 991,006 reads with mean $14,362 \pm 78$ reads per individual remained.

To investigate whether the samples were sequenced to a sufficient depth, a rarefaction curve was plotted using the alpha diversity metric, Phylogenetic Distance. The plateau showed in the rarefaction curve indicated that all 69 samples were sequenced to a sufficient depth (Figure 1). Reads were aligned to Greengenes database V13. The major phyla observed include Actinobacteria, Bacteroidetes, Firmicutes, Fusobacteria and Proteobacteria (Figure 2). These are the features of common oral microbiomes¹³.

PERMANOVA was used to investigate the saliva microbiome compositions and were found to be associated with gender, waist circumference and smoking habits. To determine whether the relative abundance of individual bacterial species was differently represented in association with the factors investigated, we used the Kruskal-Wallis test on both weighted and unweighted

UniFrac distance metrics. UniFrac¹⁴ is a phylogeny-based distance matrix that takes into account the relatedness between each bacterial species. Unweighted UniFrac is a qualitative metric that considers only the presence or absence of the bacteria whereas weighted UniFrac is a quantitative metric that considers the abundance of the bacteria.

Weighted UniFrac revealed that the dominant bacterial species present in the saliva were not significantly different among

Figure 1. Rarefaction curve plotted using alpha diversity metric, Phylogenetic Distance against number of sequences per sample.

Figure 2. Relative abundance of bacteria found in the oral microbiome of Temiar at the phylum level.

gender (p-value = 0.165). However, we found that the saliva microbiomes differed significantly between men and women for unweighted UniFrac (p-value = 0.028, [Figure 3a](#)). Kruskal-Wallis test revealed that the relative abundance of *Prevotella*, *Capnocytophaga*, *Leptotrichia*, *Neisseria* and *Streptococcus* were significantly increased in women's saliva microbiomes ([Table 2](#)). These commensal oral bacteria may become opportunistic pathogens in immune-compromised states. The relative abundance of *Bifidobacterium*, the 'good bacteria' commonly found in dairy products, was found to be highly elevated in men.

The saliva microbiome profiles also differed significantly with BMI (weighted UniFrac, p-value = 0.01; unweighted UniFrac, p-value = 0.029, [Figure 3b](#)). A post-hoc Dunn's test that conducted pairwise comparisons between underweight, normal and overweight groups revealed that the oral microbiome profiles of underweight individuals differed significantly from both overweight and normal individuals (underweight vs normal group, p-value = 0.0179; underweight vs overweight group, p-value = 0.0007). There was no significant difference between the saliva microbiome profiles of normal and overweight individuals

Figure 3. PCoA plots with the larger shapes representing the mean of distance matrix for (a) gender; (b) body mass index (BMI); (c) smoking status.

Table 2. Relative abundance of bacteria (genus level) differently represented in Temiar.

Bacteria (genus level)	FDR (q-value)	Abundance
<i>Prevotella</i>	0.006	More abundant in women
<i>Capnocytophaga</i>	0.021	
<i>Leptotrichia</i>	0.026	
<i>Neisseria</i>	0.026	
<i>Streptococcus</i>	0.036	
<i>Bifidobacterium</i>	0.043	More abundant in men

(p-value = 0.0819). Use of the Kruskal-Wallis test using R revealed higher relative abundance of *Mogibacteriaceae* and *Mycoplasma* in the microbiomes of overweight individuals (FDR q-value = 0.035, FDR q-value = 0.035). The Kruskal-Wallis test generated by QIIME revealed no significant difference.

There was a significant difference between waist circumference and saliva microbiome composition (weighted UniFrac, p-value = 0.022). However, there was no significant difference in unweighted UniFrac (p-value = 0.286) as well as in individual bacterial taxa (p-value > 0.05) among healthy individuals and those with abdominal obesity. The saliva microbiome composition of non-diabetic, pre-diabetic and diabetic individuals suggested some differences, but they were not significant (unweighted p-value = 0.069, weighted p-value = 0.122). The saliva microbiome composition and relative abundance of specific bacterial species did not statistically differ by age group, lipid levels nor blood pressure levels.

There was a perceptible difference in the saliva microbiomes and smoking habits using weighted UniFrac (p-value = 0.016, Figure 3c) but no difference detected when using unweighted UniFrac on QIIME (p-value = 0.059). Further testing showed the relative abundance of Proteobacteria and Firmicutes (phylum level) were significantly different among smokers and non-smokers. Within the phylum Proteobacteria, the relative abundance of *Neisseria* and *Aggregatibacter* was decreased in smokers compared to never-smokers and former smokers. Current smokers had a lower abundance of *Neisseria* and *Aggregatibacter* than former smokers, but the difference was not statistically significant (p-value > 0.05). The relative abundance of *Campylobacter* and bacteria of the class Clostridia, under the phylum Firmicutes were higher in both current and former smokers compared to never-smokers (Table 3).

Overall, the relative abundance of *Neisseria* and *Aggregatibacter* was decreased in current and former smokers, whereas the relative abundance of *Campylobacter* and Clostridia was greater in smokers. The saliva microbiome showed no significant difference between former smokers and never-smokers, suggesting perhaps the saliva microbiome may recover partially to an environment prior to smoking. Relative abundance of bacteria such as *Neisseria*, *Aggregatibacter* and *Campylobacter* was most affected in smokers, followed by former smokers.

Discussion

We reported a high prevalence of cardio-metabolic diseases such as obesity, pre-diabetes, insulin resistance and pre-hypertension among Temiar. These non-communicable diseases were previously not reported in OA, but recent studies have indicated their high prevalence, especially in OA tribes living near cities^{1,3,15}. Increased cardio-metabolic risks were also reported in aboriginal Torres Straits Islanders from Australia¹⁶, the Jaguapiru indigenous community in Brazil¹⁷ and the Rang Bothiya tribe in India¹⁸.

Many OA tribes lead relatively more sedentary lifestyles compared to their hunter-gatherer ancestors. They can no longer and perhaps have no need to rely entirely on the forest and its resources for survival. Rapid development had given them easier access to high-calorie processed foods, which may have contributed to obesity and cardio-metabolic diseases.

National Health and Morbidity Survey 2015 reported obesity prevalence among the major ethnic groups in Malaysia to be 30.6%, which is comparable to the findings in this study (Temiar obesity = 32.4%); the national prevalence rate of diabetes was reported to be 22.9%¹⁹, which is very much higher than that of the Temiar (2.9%) in this study. Generally, reported prevalence rates of obesity among OA are still low. Nonetheless, the high prevalence rate of pre-diabetes reported among the Temiar (44.9%) indicates that rural OA communities are in dire need of awareness education and medical intervention.

Saliva microbiome analysis revealed significant difference in microbial composition among men and women, where *Prevotella* was significantly higher in women. Hitherto, no studies have reported increased prevalence in women's saliva microbiome. *Prevotella* in the gut has been associated with plant-rich diets and possibly with improved glucose tolerance²⁰. Temiar women had a lower prevalence of pre-diabetes and insulin resistance, despite the majority of them presenting with either general or abdominal obesity. Perhaps the women consumed a traditional, indigenous diet, which is richer in plant-fibre and less meat compared to men. However, although *Prevotella* is a naturally occurring member of the oral microbiota²¹, it is also associated with inflammatory conditions such as rheumatoid arthritis and periodontal infections²².

Table 3. Relative abundance of bacteria found among smokers vs former and never-smokers generated by QIIME.

Bacteria species		FDR(q-value)	Relative abundance of bacteria
Genus	<i>Neisseria</i>	0.025	Depleted in smokers and former smokers
	<i>Aggregatibacter</i>	0.035	Depleted in smokers and former smokers
	<i>Campylobacter</i>	0.037	Increased in smokers and former smokers
Class	Clostridia	0.037	Increased in smokers and former smokers

The relative abundance of *Bifidobacterium*, healthy bacteria found in dairy products and used in probiotics, was shown to be elevated in Temiar men. While it is uncertain whether Temiar men were exposed to more dairy products than women, food taboos practiced among Temiar may contribute to the differences observed among gender²³. Several studies that investigated oral microbiomes of urbanized cohorts in association with gender have reported no differences in oral microbiome profiles^{24–26}. This may be explained by the relatively homogenous environment that urbanized cohorts were exposed to. Studies have shown that salivary microbiomes are most affected by environmental factors, as the oral microbiome of twins which were similar became highly dissimilar when they lived apart²⁴.

Temiar, on the other hand, lived in a traditional environment where men and women had different social standings. Men went out to the forest to hunt or forage while women stayed in the village with the children. They also observed certain food taboos, where the bush meat consumption of animals such as river terrapin, gibbons and porcupine were reserved only for men²³.

This preliminary investigation suggested links between saliva microbiomes and gender where differences may be attributed to cultural, dietary and environmental factors. Even though the bacteria driving the differences in obesity and gender were of different species, it should be noted that most of the women were overweight/obese, which could be a confounding factor in gender-driven disparities in the saliva microbiome.

Studies have suggested an association between obesity and altered oral microbiome^{27–29}, concurring with the findings our study. Similar to our findings, a prior study found that *Mogibacterium* was reported to be significantly more abundant in an obese group³⁰. *Mycoplasma* is a commensal, opportunistic oral pathogen reported to be elevated in overweight Temiar, perhaps due to dysbiosis of the oral microbiome. However, a significant difference was noted only when comparing overweight and underweight individuals. Both states are considered to be ‘unhealthy’ and thus assumed to be at dysbiosis.

The oral hygiene practices and oral health among Temiar were unknown, although due to their geographical isolation, it was highly unlikely they have regular access to dental health care. Our study revealed that the relative abundance of Proteobacteria, including *Neisseria* and *Aggregatibacter*, were decreased in smokers, compared to non-smokers. This was in line with the findings of Wu *et al.* (2016)³¹, although our study did not detect differences in the several other genera reported by them. This was probably due to the differences in sample size of both studies.

Wu *et al.* (2016) reported that Proteobacteria were associated with the breakdown of toxic hydrocarbons found in cigarette smoke, hence a depletion of the bacteria genus in smokers may prove detrimental to oral health³¹. An interesting difference noted was an increase in pathogenic anaerobe *Streptococcus* observed in their study³¹, whereas Hernandez *et al.* (2017)³² reported depletion of *Streptococcus* among betel-nut chewers, even after

controlling for smoking. Our investigations revealed no significant difference, even though many of the Temiar smokers were also self-reported betel-nut chewers, a practice frequently associated with oral cancer³³. Further investigations may be required to distinguish the effects of betel-nut chewing and smoking on the saliva microbiome.

Conclusion

Our study revealed a high prevalence of cardio-metabolic diseases among the Temiar, including general and abdominal obesity, pre-diabetes and insulin resistance. Pre-hypertension was found highly prevalent across all age groups.

Interestingly, the saliva microbiome profiles were significantly different for gender where the relative abundance of *Prevotella*, *Capnocytophaga*, *Leptotrichia*, *Neisseria*, *Streptococcus* and *Bifidobacterium* were concerned. Our study also noted a significant difference between the saliva microbiome compositions of underweight vs overweight and normal individuals. *Mogibacteriaceae* and *Mycoplasma* were found to be elevated in overweight individuals. The oral microbiome was not significantly different among non-diabetic, pre-diabetic and diabetic individuals. The microbiome profiles differed significantly among smokers and non-smokers where further investigation showed that Proteobacteria were significantly decreased in smokers. Investigation towards other health parameters such as pre-diabetes were inconclusive.

The sample size was limited, but findings from this pilot study warrants further and larger studies into other Malaysian indigenous tribes which may present unexpected findings perhaps attributed to differences in their cultural practices, lifestyle and diet.

Methods

Data and sample collection

The study was approved by Ministry of Health Malaysia under National Medical Research Registry, MNDR ID #09—23-3913, Department of Orang Asli Development, Malaysia (JAKOA) and Monash University Human Research Ethics Committee (MUHREC). Before sampling, a courtesy visit to the Temiar elders in Kampong Pos Piah, Perak was conducted to explain the rationale of the study. Upon agreement to participate in our study, a medical team returned to the village on an agreed date and conducted health screening and sampling. Participants who were over 18 years old with no visible health ailments and able to provide informed consent were recruited for the study through convenience sampling, that is whoever who turned up and was eligible. Participants who were pregnant, with a history of alcohol/drug abuse, or with chronic illness (e.g. kidney failure, cancer, heart disease) were excluded from the study.

The consent form was read aloud by interviewers and queries were addressed before either a signature or thumbprint was provided as a sign of consent. A total of 72 Temiar provided informed consent to participate. Interviews were conducted in Bahasa Malaysia using a questionnaire¹² to collect information about their socio-demography, medical history and diet. Height, weight,

waist circumference and blood pressure were measured¹. Participants were also examined by clinicians. Acanthosis nigricans, which is darkening of the skin around the neck and creases of elbows indicative of insulin resistance, was noted.

Saliva samples were collected in sterile 50ml polypropylene Falcon tube. Participants were requested to rinse their mouths with water thoroughly 30 minutes prior to collecting saliva. Venous blood samples were taken for biochemical analyses.

Anthropometrics and Biochemical analysis

BMI, waist circumference and blood pressure cut-off values were in accordance to WHO recommendations³⁴. We measured their HbA1C and blood lipid levels (cholesterol, HDL, LDL, Triglyceride). We used TG/HDL ratio as a surrogate marker for insulin resistance with a cut-off value of 0.9-1.7³⁵.

DNA extraction and PCR

DNA was extracted from saliva using a modified high salt-method³⁶. The V3-V4 region of the 16S rRNA gene were targeted, resulting in a PCR product of approximately 550 bp³⁷.

Sequencing on Illumina MiSeq

DNA sequencing was done by Genomics Facility in Monash University Malaysia on Illumina MiSeq to produce paired end reads of approximately 230 bp each.

Data analysis

Microbiome analysis was conducted on QIIME 1.9³⁸. Chimeras were filtered using UCHIME 1.39.3³⁹ before being aligned to Greengenes database V13.8⁴⁰. The reads were then clustered into operational taxonomic units (OTUs) with open-reference method at 97% similarity level using UCLUST⁴¹ in the QIIME pipeline. OTU clusters were assigned taxonomy with RDP classifier⁴². The reads were normalized and OTUs that were present at less than 0.05% were filtered off.

Alpha diversity and beta diversity of the samples were reported using phylogenetic distance (PD) and UniFrac¹⁴, respectively. PCoA plots were generated to visualize beta diversity of the samples.

Statistical analysis

Statistical analyses were completed on QIIME and R 3.4.4. Information taken from the mapping file included gender, BMI and smoking status. PERMANOVA, a non-parametric test was used to test for differences in median among the groups using weighted and unweighted UniFrac distance matrix Usi/ng the R packages *vegan* (v2.4-2), *readr* (v1.1.0) and *dplyr* (v0.5.0).

Kruskal-Wallis test was used to test for differences in the relative abundance of OTUs among the different groups. A post-hoc test, Dunn's test was done for pairwise comparison when testing factors like BMI and smoking, as they had more than two groups. False discovery rate, reported as q-value, was used to control for multiple hypothesis testing and was statistically significant at 5%.

Data availability

Underlying data

Saliva microbiomes of the individuals in this study are available from the Sequence Read Archive, BioProject accession number PRJNA515166; <https://identifiers.org/bioproject/PRJNA515166>.

Anthropometric data, along with the other variables measured, are available on figshare. DOI: <https://doi.org/10.26180/5c453ff43588312>.

Extended data

The questionnaire used in this study is available on figshare. DOI: <https://doi.org/10.26180/5c453ff43588312>.

Data are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

Grant information

This study was funded by Tropical Medicine and Biology Platform (TMB), Monash University Malaysia seed grant TMB-2016-HG3185140920-YLF/MP awarded to LFY and MEP. Ms Li-Fang Yeo is supported by a scholarship awarded by Jeffrey Cheah School of Medicine and Health Sciences, Monash University Malaysia.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We thank the Temiar community for their participation and the Department of Orang Asli Development (JAKOA) of Malaysia. We are especially grateful to Prof. Sunil K. Lal for helpful suggestions, Ms Yin Peng Lee for technical advice, and Dr Christina Yap for conducting biochemical analysis, Prof Daniel Reidpath for his assistance in statistical analysis on R, Dr Amreeta Dhanoa for editing the manuscript, Prof Yvonne Lim for advice and Dr Eustacia Lee and the University of Malaya team for participating in sample collection. Furthermore, our special thanks to Dr Badariah Ahmad and Dr Siti Noraida for participating in fieldwork.

References

1. Phipps ME, Chan KK, Naidu R, *et al.*: **Cardio-metabolic health risks in indigenous populations of Southeast Asia and the influence of urbanization.** *BMC Public Health*. 2015; **15**: 47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
2. McColl H, Racimo F, Vinner L, *et al.*: **The prehistoric peopling of Southeast Asia.** *Science*. 2018; **361**(6397): 88–92.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Wong CY, Zailiah MS, Chua EY, *et al.*: **Double-burden of malnutrition among the**

- indigenous peoples (Orang Asli) of Peninsular Malaysia. *BMC Public Health*. 2015; 15: 680.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Tuan Abdul Aziz TA, Teh LK, Md Idris MH, *et al.*: Increased risks of cardiovascular diseases and insulin resistance among the Orang Asli in Peninsular Malaysia. *BMC Public Health*. 2016; 16: 284.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 5. Olsen I: The oral microbiome in health and disease. *Infections And General Health*. 2016; 97–114.
[Publisher Full Text](#)
 6. Yamashita Y, Takeshita T: The oral microbiome and human health. *J Oral Sci*. 2017; 59(2): 201–206.
[PubMed Abstract](#) | [Publisher Full Text](#)
 7. Sudhakara P, Gupta A, Bhardwaj A, *et al.*: Oral Dysbiotic Communities and Their Implications in Systemic Diseases. *Dent J (Basel)*. 2018; 6(2): pii: E10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 8. Idris A, Hasnain SZ, Huat LZ, *et al.*: Human diseases, immunity and the oral microbiota—Insights gained from metagenomic studies. *Oral Science International*. 2017; 14(2): 27–32.
[Publisher Full Text](#)
 9. Long J, Cai Q, Steinwandel M, *et al.*: Association of oral microbiome with type 2 diabetes risk. *J Periodontal Res*. 2017; 52(3): 636–643.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 10. Zeigler CC, Persson GR, Wondimu B, *et al.*: Microbiota in the oral subgingival biofilm is associated with obesity in adolescence. *Obesity (Silver Spring)*. 2012; 20(1): 157–164.
[PubMed Abstract](#) | [Publisher Full Text](#)
 11. Choromańska K, Choromańska B, Dąbrowska E, *et al.*: Saliva of obese patients - Is it different? *Postępy Hig Med Dosw (Online)*. 2015; 69: 1190–1195.
[PubMed Abstract](#)
 12. Yeo LF, Phipps M, Aghakhanian FF, *et al.*: Health and Saliva Microbiomes of a Semi-urbanized Indigenous Tribe in Peninsular Malaysia. *figshare*. Fileset. 2019. <http://www.doi.org/10.26180/5c453ff435883>
 13. Dewhirst FE, Chen T, Izard J, *et al.*: The human oral microbiome. *J Bacteriol*. 2010; 192(19): 5002–5017.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 14. Lozupone C, Knight R: UniFrac: a new phylogenetic method for comparing microbial communities. *Appl Environ Microbiol*. 2005; 71(12): 8228–8235.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 15. Pell C, Allotey P, Evans N, *et al.*: Coming of age, becoming obese: a cross-sectional analysis of obesity among adolescents and young adults in Malaysia. *BMC Public Health*. 2016; 16(1): 1082.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 16. Burns J, Thomson N: Summary of overweight and obesity among Indigenous peoples. Retrieved [03 November 2015]; 2006.
 17. Oliveira GF, Oliveira TR, Ikejiri AT, *et al.*: Prevalence of Obesity and Overweight in an Indigenous Population in Central Brazil: A Population-Based Cross-Sectional Study. *Obes Facts*. 2015; 8(5): 302–310.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 18. Kandpal V, Sachdeva MP, Saraswathy KN: An assessment study of CVD related risk factors in a tribal population of India. *BMC Public Health*. 2016; 16(1): 434.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 19. Institute of Public Health: National Health and Morbidity Survey. Institute of Public Health; 2015.
[Reference Source](#)
 20. Kovatcheva-Datchary P, Nilsson A, Akrami R, *et al.*: Dietary Fiber-Induced Improvement in Glucose Metabolism Is Associated with Increased Abundance of *Prevotella*. *Cell Metab*. 2015; 22(6): 971–982.
[PubMed Abstract](#) | [Publisher Full Text](#)
 21. Palmer RJ Jr: Composition and development of oral bacterial communities. *Periodontol 2000*. 2014; 64(1): 20–39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 22. Larsen JM: The immune response to *Prevotella* bacteria in chronic inflammatory disease. *Immunology*. 2017; 151(4): 363–374.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 23. Benjamin G: Temiar Religion, 1964-2012: Enchantment, Disenchantment and Re-enchantment in Malaysia's Uplands. 2014.
[Reference Source](#)
 24. Stahnger SS, Clemente JC, Corley RP, *et al.*: Nurture trumps nature in a longitudinal survey of salivary bacterial communities in twins from early adolescence to early adulthood. *Genome Res*. 2012; 22(11): 2146–2152.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 25. Chen H, Jiang W: Application of high-throughput sequencing in understanding human oral microbiome related with health and disease. *Front Microbiol*. 2014; 5: 508.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 26. Li J, Quinke D, Horz HP, *et al.*: Comparative analysis of the human saliva microbiome from different climate zones: Alaska, Germany, and Africa. *BMC Microbiol*. 2014; 14(1): 316.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 27. Takeshita T, Kageyama S, Furuta M, *et al.*: Bacterial diversity in saliva and oral health-related conditions: the Hisayama Study. *Sci Rep*. 2016; 6(1): 22164.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 28. Goodson JM, Hartman ML, Shi P, *et al.*: The salivary microbiome is altered in the presence of a high salivary glucose concentration. *PLoS One*. 2017; 12(3): e0170437.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 29. Janem WF, Scannapieco FA, Sabharwal A, *et al.*: Salivary inflammatory markers and microbiome in normoglycemic lean and obese children compared to obese children with type 2 diabetes. *PLoS One*. 2017; 12(3): e0172647.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 30. Wu Y, Chi X, Zhang Q, *et al.*: Characterization of the salivary microbiome in people with obesity. *PeerJ*. 2018; 6: e4458.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 31. Wu J, Peters BA, Dominianni C, *et al.*: Cigarette smoking and the oral microbiome in a large study of American adults. *ISME J*. 2016; 10(10): 2435–2446.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 32. Hernandez BY, Zhu X, Goodman MT, *et al.*: Betel nut chewing, oral premalignant lesions, and the oral microbiome. *PLoS One*. 2017; 12(2): e0172196.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 33. Guha N, Warnakulasuriya S, Vlaanderen J, *et al.*: Betel quid chewing and the risk of oral and oropharyngeal cancers: a meta-analysis with implications for cancer control. *Int J Cancer*. 2014; 135(6): 1433–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
 34. World Health Organization: Use of glycated haemoglobin (HbA1c) in the diagnosis of diabetes mellitus. Abbreviated report of a WHO consultation 2011. Geneva: World Health Organization. 2013.
[Reference Source](#)
 35. Mostafa SA, Davies MJ, Morris DH, *et al.*: The association of the triglyceride-to-HDL cholesterol ratio with insulin resistance in White European and South Asian men and women. *PLoS One*. 2012; 7(12): e50931.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 36. Quinke D, Kittler R, Kayser M, *et al.*: Evaluation of saliva as a source of human DNA for population and association studies. *Anal Biochem*. 2006; 353(2): 272–277.
[PubMed Abstract](#) | [Publisher Full Text](#)
 37. PCR I: 16S Metagenomics Sequencing Library Preparation. *Illumina*. 2013.
[Reference Source](#)
 38. Caporaso JG, Kuczynski J, Stombaugh J, *et al.*: QIIME allows analysis of high-throughput community sequencing data. *Nat Methods*. 2010; 7(5): 335–336.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 39. Edgar RC, Haas BJ, Clemente JC, *et al.*: UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics*. 2011; 27(16): 2194–2200.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 40. DeSantis TZ, Hugenholtz P, Larsen N, *et al.*: Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl Environ Microbiol*. 2006; 72(7): 5069–5072.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 41. Edgar RC: Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*. 2010; 26(19): 2460–2461.
[PubMed Abstract](#) | [Publisher Full Text](#)
 42. Wang Q, Garrity GM, Tiedje JM, *et al.*: Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol*. 2007; 73(16): 5261–5267.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status: ? ?

Version 1

Reviewer Report 10 April 2019

<https://doi.org/10.5256/f1000research.19359.r46035>

© 2019 Mohd Zain S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Siti Nursheena Mohd Zain

Institute of Biological Science, Faculty of Science, University of Malaya, Kuala Lumpur, Malaysia

An interesting work looking at non-communicable disease risks link to microbiome in saliva as an indicator. The discussion could be written better to pull the results together. What is missing is a comparative study with another sedentary population. I would like if the conclusion is added with recommendations to improve the issue facing the Temiar tribe. Noted that sample size is small to make small but still warrants some finding to improve their health status.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Parasitology and molecular parasitology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 13 March 2019

<https://doi.org/10.5256/f1000research.19359.r44294>

© 2019 Gomez A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Andres Gomez

Department of Animal Science, University of Minnesota, Saint Paul, MN, USA

Results

The QIIME microbiome analyses should be briefly described as far as the bioinformatics pipeline used and its details: clustering of OTUS? OTU picking? amplicon sequence variants (DADA2-ASV)? These should be specifically detailed.

It does not seem that the rarefaction curve shows compete plateau, also what do the colors mean?

PERMANOVA is used to mine for distinctions in microbiome composition.

It is unclear how kruskal-wallis test was used in conjunction with Unifrac - KW is used for pairwise comparisons in multivariate space, not for multivariate data.

PERMANOVA results (P values) should be accompanied by R2 and, Pseudo F values also. PERMANOVA models used should also be reported.

"Kruskal-Wallis test revealed that the relative abundance of *Prevotella*, *Capnocytophaga*, *Leptotrichia*, *Neisseria* and *Streptococcus*"

Were these discriminant analyses made at genus or OTU/ASV levels? Also, it is recommended to show these results in box plots showing medians and variation of data, not in tables

Authors make several assertions as to the character and typically reported nature of the markers discovered, in the context of "good/bad bacteria" - but many of the claims made are in the context of gut microbiomes not oral communities.

For instance:

"*Prevotella*, associated with consumption of plant-rich diets"

This claim may be true only for the gut microbiome not for oral communities. Also: "The relative abundance of *Bifidobacterium*, the 'good bacteria' commonly found in dairy products, was found to be highly elevated in men" - this may be associated with effect of dairy products in the gut not in the oral cavity.

"These commensal oral bacteria may become opportunistic pathogens in immune-compromised states." please cite

Fig 3- legends should indicate that a-b were based on Unw-UniFrac, and that C on W-UniFrac - Please depict % variation on axes

It is not clear how the post-hoc Dun's test were conducted-0 is this still based on PERMANOVA?

"The Kruskal-Wallis test generated by QIIME revealed no significant difference." This is not clear - compared to what?

"There was a significant difference between waist circumference and saliva microbiome composition (weighted UniFrac, p-value = 0.022). " This is not clear - you mean differences between the microbiomes of subjects with different waist circumference?

QIIME is just a software with plugins to perform statistical analyses - it is not clear when authors are using qiime or R and how these platforms would yield different results. Please avoid jargon as "generated by QIIME"

"The saliva microbiome showed no significant difference between former smokers and never-smokers, suggesting perhaps the saliva microbiome may recover partially to an environment prior to smoking". This is unclear but also speculative.

The discussion is highly speculative and should be toned down, mainly as far as the hypotheses behind the differences reported.

Authors also seem to interchangeably extrapolate hypothesis on oral microbiome changes with gut microbiome changes in the context of lifestyle and diet (e.g. Prevotella) although diet may also be linked to oral microbiome modulation, this relationship is less clear, and taxa such as Prevotella, abundant in both gut and oral cavity, cannot be assumed to be modulated by diet in both sites

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Microbial ecology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 18 Mar 2019

Li-Fang Yeo, Monash University, Bandar Sunway, Malaysia

Details on the bioinformatics pipeline will be addressed in version 2. The different colours on the rarefaction curve represent different individuals. Other metrics such as Shannon and Chao1 shows an obvious plateau. We felt that PD was more informative because it included phylogenetic distances.

Our statistics were done mostly with reference to this paper:

Wu, J., Peters, B.A., Dominianni, C., Zhang, Y., Pei, Z., Yang, L., Ma, Y., Purdue, M.P., Jacobs, E.J., Gapstur, S.M. and Li, H., 2016. "Cigarette smoking and the oral microbiome in a large study of American adults". *The ISME journal*. <https://www.nature.com/articles/ismej201637>

However, more recent methods such as using balances and log ratio shall be considered in a larger, extended study. Test scores for PERMANOVA results shall be addressed in version 2 as will several questions that were unclear in version 1.

To date, we have none found literature that was comparable (ie. Having Bifidobacterium elevated in men, Prevotella elevated in women). Hence we attempted to suggest possible explanations to this observation. The discussion will be re-worded to take into account reviewer's comments.

Competing Interests: No competing interest.

Reviewer Response 20 May 2019

Andres Gomez, University of Minnesota, Saint Paul, USA

Dear authors,

Could you please provide point by point responses to the comments raised in the previous version, along with where changes were made in the manuscript?
That would facilitate reviewing of the new manuscript.

Competing Interests: No competing interests were disclosed.

Author Response 20 May 2019

Li-Fang Yeo, Monash University, Bandar Sunway, Malaysia

Dear reviewer,

As per request:

1. The QIIME microbiome analyses should be briefly described as far as the bioinformatics pipeline used and its details: clustering of OTUS? OTU picking? amplicon sequence variants (DADA2-ASV)? These should be specifically detailed.

Microbiome analysis pipeline edited and detailed in Methods section.

2. It does not seem that the rarefaction curve shows compete plateau, also what do the colors mean?

Rarefaction curve now shows Shannon index instead of phylogenetic diversity for better indicator of plateau. Each colour is a biological sample.

3. It is unclear how kruskal-wallis test was used in conjunction with Unifrac - KW is used for pairwise comparisons in multivariate space, not for multivariate data.

KW test was conducted on an OTU table, not UniFrac. Refer to Ying & Sun (2017)

<https://www.sciencedirect.com/science/article/pii/S2352304217300351>

Edits made in Results section.

4. PERMANOVA results (P values) should be accompanied by R² and, Pseudo F values also. PERMANOVA models used should also be reported.

Pseudo F values are reported. R² value not available for PERMANOVA but available for adonis. All available values are reported in supplementary material.

5. "Kruskal-Wallis test revealed that the relative abundance of *Prevotella*, *Capnocytophaga*, *Leptotrichia*, *Neisseria* and *Streptococcus*"

Were these discriminant analyses made at genus or OTU/ASV levels? Also, it is recommended to show these results in box plots showing medians and variation of data, not in tables

These analyses were made at OTU level 6, equivalent to genus level. Addressed in Results section. Also, due to time constraint and admittedly technical limitations on our part, the results will be shown in tables for this paper. Reviewer's advice are definitely well-received and will be considered for our next paper.

6. Authors make several assertions as to the character and typically reported nature of the markers discovered, in the context of "good/bad bacteria" - but many of the claims made are in the context of gut microbiomes not oral communities.

We have acknowledged the reviewer's comments and rephrased the discussion to be more cautious and less speculative.

7. "These commensal oral bacteria may become opportunistic pathogens in immune-compromised states." please cite

Idris A, Hasnain SZ, Huat LZ, Koh D. Human diseases, immunity and the oral microbiota—Insights gained from metagenomic studies. Oral Science International. 2017 Jul;14(2):27–32.

Cited in-text as well.

8. Fig 3- legends should indicate that a-b were based on Unw-UniFrac, and that C on W-UniFrac - Please depict % variation on axes

Done.

9. It is not clear how the post-hoc Dun's test were conducted-0 is this still based on PERMANOVA?

Post-hoc Dunn's test was conducted using an OTU table. Permanova does not conduct pairwise comparison hence Dunn's test was chosen to test for differences between people with different BMI status (underweight, normal, overweight). And the test revealed that underweight individuals had microbiomes that was significantly different from both normal and overweight individuals.

10. "The Kruskal-Wallis test generated by QIIME revealed no significant difference." This is not clear - compared to what?

12. QIIME is just a software with plugins to perform statistical analyses - it is not clear when authors are using qiime or R and how these platforms would yield different results. Please avoid jargon as "generated by QIIME"

For 10 & 12, previously qiime and R yielded different results (one not significant, one significantly different) for oral microbiomes and BMI. Upon closer inspection we realized this happened because of mismatching samples and OTU table upon input to R. We checked the other sets of data and the qiime results matches with R. Changes addressed in Results section.

11. "There was a significant difference between waist circumference and saliva microbiome composition (weighted UniFrac, p-value = 0.022). " This is not clear - you mean differences between the microbiomes of subjects with different waist circumference?

The above sentence has been rephrased to "There was a significant difference in the saliva microbiome of Temiar who had a healthy waist circumference compared to those with abdominal obesity."

13. "The saliva microbiome showed no significant difference between former smokers and never-smokers, suggesting perhaps the saliva microbiome may recover partially to an environment prior to smoking". This is unclear but also speculative.

The discussion is highly speculative and should be toned down, mainly as far as the hypotheses behind the differences reported.

Authors also seem to interchangeably extrapolate hypothesis on oral microbiome changes with gut microbiome changes in the context of lifestyle and diet (e.g. Prevotella) although diet may also be linked to oral microbiome modulation, this relationship is less clear, and taxa such as Prevotella, abundant in both gut and oral cavity, cannot be assumed to be modulated by diet in both sites

This comment has been well-received and edits have been done to Discussion section.

TQ.

Competing Interests: No competing interests.

Comments on this article

Version 1

Author Response 30 Apr 2019

Li-Fang Yeo, Monash University, Bandar Sunway, Malaysia

Thank you Andres and Siti Nursheena for your comments. We have addressed all your concerns, hopefully to your satisfaction, with one exception regarding Andres's suggestion to use boxplots to show median and variation of data. For this publication, we have decided to maintain the results in table-form showing q-values due to time constraint and technical limitations on our side. We will definitely take this into consideration to improve our future work.

Competing Interests: No competing interests.

Author Response 24 Apr 2019

Maude Phipps, Monash University, Bandar Sunway, Malaysia

Thanks Andres and Siti Nursheena for your constructive comments. We endeavor to improve this paper and hope it will be confirmed and citable soon.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research