REVIEW

Recent advances in understanding neocortical development

[version 1; peer review: 2 approved]

Victor Borrell

Institute of Neuroscience, Consejo Superior de Investigaciones Científicas (CSIC) - Universidad Miguel Hernández, Ramon y Cajal s/n, 03550 San Juan de Alicante, Spain

V1 First published: 23 Oct 2019, 8(F1000 Faculty Rev):1791
<https://doi.org/10.12688/f1000research.20332.1>

Latest published: 23 Oct 2019, 8(F1000 Faculty Rev):1791
<https://doi.org/10.12688/f1000research.20332.1>

Abstract

The neocortex is the largest part of the mammalian brain and is the seat of our higher cognitive functions. This outstanding neural structure increased massively in size and complexity during evolution in a process recapitulated today during the development of extant mammals. Accordingly, defects in neocortical development commonly result in severe intellectual and social deficits. Thus, understanding the development of the neocortex benefits from understanding its evolution and disease and also informs about their underlying mechanisms. Here, I briefly summarize the most recent and outstanding advances in our understanding of neocortical development and focus particularly on dorsal progenitors and excitatory neurons. I place special emphasis on the specification of neural stem cells in distinct classes and their proliferation and production of neurons and then discuss recent findings on neuronal migration. Recent discoveries on the genetic evolution of neocortical development are presented with a particular focus on primates. Progress on all these fronts is being accelerated by high-throughput gene expression analyses and particularly single-cell transcriptomics. I end with novel insights into the involvement of microglia in embryonic brain development and how improvements in cultured cerebral organoids are gradually consolidating them as faithful models of neocortex development in humans.

Keywords

Radial Glia, cell lineage, OSVZ, cortex folding, basal progenitors, neurogenesis, human-specific neocortex, neocortical development, cerebral cortex evolution, progenitor proliferation

Open Peer Review

Approval Status

	1	2
version 1 23 Oct 2019		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Simon Hippenmeyer**, Institute of Science and Technology Austria (IST Austria), Klosterneuburg, Austria
2. **Laurent Nguyen**, University of Liège, Liège, Belgium

Any comments on the article can be found at the end of the article.

Corresponding author: Victor Borrell (vborell@umh.es)

Author roles: Borrell V: Conceptualization, Data Curation, Project Administration, Writing – Original Draft Preparation

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2019 Borrell V. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Borrell V. **Recent advances in understanding neocortical development [version 1; peer review: 2 approved]** F1000Research 2019, 8(F1000 Faculty Rev):1791 <https://doi.org/10.12688/f1000research.20332.1>

First published: 23 Oct 2019, 8(F1000 Faculty Rev):1791 <https://doi.org/10.12688/f1000research.20332.1>

Introduction

The cerebral cortex, or neocortex, is the part of our brains primarily responsible for abstract thinking and our unique cognitive abilities as human beings. It is by far the most complex biological structure, and it forms during embryonic development following a sequence of genetically predefined molecular and cellular events. This process of embryonic development recapitulates the emergence of the human cerebral cortex during evolution. Thus, understanding neocortical development in humans and other species tells us about the magnitude of its complexity, how it differs from neocortical development in non-humans, how it emerged during evolution from simpler brains, and what goes wrong in developmental brain disease.

Through the use of informative experimental animal species, much has been learned about the basic processes of neocortical development: neural stem and progenitor cell (NSPC) proliferation, generation, and migration of neurons from their place of birth to their final location in neocortical layers and the growth of neural processes and establishment of nerve connections. Recent advances in genome editing, tissue culture, and DNA sequencing have accelerated our understanding of neocortical development at many levels. Here, I briefly summarize some of the most significant advances in our understanding of neocortical development generated recently by the international community, and I give special attention to progenitor proliferation and neurogenesis of dorsal progenitors and projection neurons. Although much remains to be learned, the emerging picture shows that the evolutionary and developmental emergence of the human neocortex involved a plethora of genetic and epigenetic changes, including novel genes and variants, functional gene networks, and novel cell types and cellular specializations.

Neurogenesis

Development of the neocortex begins with the proliferation of NSPCs, which amplify their own pool prior to generating neurons. Recent work shows that the proliferation of NSPCs and their mode of neuron production vary across species, and previously unsuspected mechanisms of molecular regulation have been identified. Novel insights about the diversity of progenitor cells and how their neuronal output contributes to build the cerebral cortex are also emerging.

Progenitor cell diversity

Work in the last decade has identified a diversity of NSPC types and their lineage relationships. Following the seminal discovery of apical radial glial cells (aRGCs) as the primary type of cortical progenitor cell^{1,2}, intermediate progenitor cells (IPCs) were identified³⁻⁵. IPCs are transit-amplifying progenitors expressing the transcription factor *Tbr2* and producing the majority of excitatory neurons in the mouse and rat neocortex^{3,4,6,7}. Most IPCs cluster and undergo mitosis in a layer basal from the ventricular zone (VZ), the subventricular zone (SVZ), although a subset reside in the VZ and undergo apical mitosis, named short neural precursors⁸. Further studies in primates led to the discovery of an expanded and specialized SVZ, subdivided in inner and outer domains (ISVZ and OSVZ, respectively)⁹. These layers were later identified in other species with an expanded neocortex, like ferret, cat, and

sheep¹⁰ and even in New World monkeys and the Amazonian rodent agouti^{11,12}. Progenitor cell lineage tracings in human, macaque, ferret, and mouse led to the discovery of additional types of basal progenitor cells in the developing neocortex, which are bound by complex lineage relationships. These include various types of basal radial glial cells (bRGCs)^{10,11,13-15} and subapical progenitor cells¹⁶. The relative abundance and proliferative capacity of these progenitor cell types are much greater in species with a neocortex that is large and folded (carnivores and primates) than in those where it is small and smooth (mouse and rat). This has been proposed to contribute to the evolutionary increase in complexity, or complexification, of the neocortex¹⁶⁻¹⁸.

Pioneer transcriptomic studies of the developing cerebral cortex using bulk tissue, identified differences in gene expression between germinal layers, cortical areas, developmental stages, and mammalian species^{15,19-23}. This illustrated the profound diversity of transcriptional landscapes in cortical development at all levels, including folding and across phylogeny, and set the conceptual foundations for the next technological leap: transcriptomics of single cells. The advent of single-cell transcriptomic analyses has revolutionized our approach to studying cortical development by providing a global and unbiased picture of cell diversity with unprecedented resolution²⁴⁻²⁶. This technology has enabled identification of multiple sets of transcriptionally distinct progenitor cell classes in the cortical primordium, generating excitatory neurons²⁷⁻²⁹, and in the basal ganglia, generating inhibitory interneurons^{30,31}.

Single-cell analyses are also beginning to shed light on long-standing hypotheses about the heterogeneity of cortical progenitor cells and the dynamics of their lineage and fate potential during development³²⁻³⁶. Early experimental studies of these fundamental questions indicated that the fate potential of cortical progenitors is temporally restricted, such that early progenitors can produce neurons for all cortical layers but late progenitors can produce neurons for superficial layers only^{34,37}. Such late fate restriction would be cell-autonomous as late cortical progenitors continued producing superficial-layer neurons even when transplanted into the new cellular environment of a young host cortex^{34,37}. The identification of a subset of RGCs that expressed *Cux2* and that were fate-restricted to produce upper-layer neurons further supported this model³⁵. However, such fate-restricted progenitors have not been identified in single-cell transcriptomic studies^{28,33,38}. Rather, some of these studies support the existence of epigenetically regulated temporal molecular birthmarks in RGCs, which act in their daughter neurons as seeds for neuronal diversity. It is proposed that these conserved differentiation programs may then be integrated with environmental (non-cell-autonomous) cues to ultimately define the identity of the neuronal progeny^{28,39,40}. Nevertheless, controversial points of view on these and related issues remain because of differences in single-cell data processing, analysis, and interpretation^{30,31,41}.

Modes of neurogenesis and influence on cortex size

Cortical excitatory neurons may be generated directly from the primary progenitor cells (aRGCs) or indirectly via secondary basal progenitors such as IPCs and bRGCs. Indirect neurogenesis

is considered a milestone of mammalian cortical evolution, enabling a phenomenal increase in the numbers of neurons produced, particularly those destined to superficial layers, and leading to cortical expansion^{42,43}. Recent studies reveal that indirect neurogenesis is the major mode of producing deep as well as superficial cortical layers in mouse^{40,44} and that it also exists in non-mammals, albeit at lower levels and mostly in birds^{44,45}. At the molecular level, regulation of the balance between direct and indirect neurogenesis is phylogenetically conserved across amniotes, from snakes to birds and mammals, including humans, where high Robo1/2 signaling promotes expression of the Notch ligands Jag1/2 while repressing the canonical ligand Dll1⁴⁴. Blockade of this signaling system drives indirect neurogenesis, increasing neuron numbers and cortex size. Intriguingly, indirect neurogenesis also correlates with hyperpolarization of apical progenitor cells, which represses Wnt signaling⁴⁰. Changes in the mode of neurogenesis are responsible for the reduced cortex size in developmental brain disease, like microcephaly induced by Zika virus infection. In this case, activation of the unfolded protein response drives direct neurogenesis, leading to premature and limited neuron production and to a small cortex^{46–48}.

Regulation of progenitor cell proliferation

Cerebral cortex size depends on the mode of neurogenesis and also on the proliferative activity of cortical progenitor cells. Multiple mechanisms regulating progenitor cell proliferation have recently been uncovered. Regulation of gene transcription by epigenetic mechanisms has emerged as a key factor where histone deacetylases and methyltransferases regulate the generation and position of IPCs, neuron migration, and cortical lamination^{49,50}. Similarly, regulation of chromatin accessibility and other mechanisms related to non-coding genomic regions critically determines levels and patterns of gene expression in the developing cortex, defining neuron production, cortex size, and area identity^{51,52}.

One of the most exciting findings has been the identification of mRNA species transported within the long basal process of aRGCs and locally translated at their basal endfeet, next to the pial membranes^{53–55}. This includes proteins regulating the cell cycle, like Ccnd2, and lengthening of G₁, S, and M phases of the cell cycle, which depletes progenitor cells causing premature neurogenesis or apoptosis^{56–58}.

IPCs are usually depicted as round cells extending few short processes^{3,59}, and bRGCs are usually depicted as having a single smooth and unbranched basal process^{10,60–62}. Recent observations demonstrate that the degree of process branching and elaboration in both IPCs and bRGCs is greater in ferret and human (which have a large and folded cortex) than in mouse (which has a small and smooth cortex) and this is linked to a greater proliferation rate of these cortical progenitor cells^{63,64}. Basal progenitor process growth and proliferation are related to the membrane-bound protein PALMD, enabling these cells to receive pro-proliferative signals related to integrin function^{60,64,65}. Indeed, modulation of cortical progenitor cell proliferation by cell-extrinsic signals has now been demonstrated to occur

from multiple sources, including growing axons and migrating neurons^{63,66,67}, extracellular neurotransmitters and ions^{68,69}, and Notch, Wnt, Fgf, and Shh signals⁷⁰.

Once IPCs and neurons are born from aRGCs, they must first delaminate and migrate away from the VZ toward the SVZ. This critical process is regulated by several signaling mechanisms, including Robo⁷¹, YAP⁷², Insm1 via the apical adherens protein Plekha⁷³, and the centrosome-associated protein Akna⁷⁴.

Neuronal migration

Once neurons are born, they must migrate from their place of birth in the germinal layers to their final location in the neuronal layers. Neuronal migration is a multi-step process that begins with neuronal delamination from the VZ and movement to the SVZ. (See Silva *et al.*⁶⁷ [2019] for an excellent review on neuron migration.)

Delamination

New advances in understanding cortical development have identified the microtubule-associated protein Lszts as a master regulator of cellular dynamics, promoting the delamination of neurons and bRGCs born from aRGCs by altering apical junction organization⁷⁵.

Multipolar phase transition

Once newborn neurons delaminate from the VZ, they enter the SVZ and undergo a transition phase displaying multipolar morphology⁷⁶. During this process, NeuroD1 expressed by multipolar cells represses Prdm16, which regulates mitochondrial reactive oxygen species, and this signaling axis is crucial for the regulation of the multipolar phase migration⁷⁷. Dbn1, a protein interacting with F-actin, is another key regulator of neuronal multipolar morphology, polarity, and migration by regulating the levels of N-cadherin⁷⁸.

Bipolar locomotion

Cortical neurons resume radial migration by re-acquiring polarity and extending a single leading process as they exit the SVZ. This leading process establishes intimate adhesive interactions with the basal process of RGCs for guidance in their migratory displacement. The long-held concept that the leading process of these radially migrating neurons is single and unbranched^{79,80} has been challenged by new observations, demonstrating frequent branching and thus more complex migratory behaviors than previously reported⁸¹. In fact, leading process branching is much more frequent in the developing cortex of ferret than mouse and is related to the tangential displacement of radially migrating neurons. Thus, this seems directly related to the maintenance of the radial organization of the cerebral cortex during the tangential expansion and folding of the neocortex^{81,82}. However, an excessive tangential displacement of cortical neurons is deleterious, and the horizontal tiling of the developing cerebral cortex, or regular distribution of its radial units, must be actively maintained. The microtubule stability regulator protein Memo1 plays key functions in the maintenance of RGC structure and cortical tiling by repressing the hyperbranching of the basal process of RGCs and the

excessive dispersion of radially migrating neurons⁸³. On the other hand, it was recently shown that the chromatin-modifying enzyme Prdm16 is also necessary for transcriptional silencing in RGCs and to promote the migration of late-born cortical neurons and cortical layering⁴⁹.

Evolution of cerebral cortex development

A fundamental feature of the evolution of cerebral cortex in amniotes is the phenomenal increase in neuron number and expansion in size. This process is recapitulated during embryonic development, and recent work demonstrates the importance of the balance between direct and indirect neurogenesis⁴⁴. Mechanisms regulating this critical balance, including transcriptional programs regulated by progenitor cell membrane polarity⁴⁰ and canonical signaling pathways like the unfolded protein response⁴⁶, some of which are highly conserved across amniotes like Robo and Dll1⁴⁴, are beginning to be identified.

Beyond the emergence of indirect neurogenesis, cortical evolution involved additional key mechanisms. Cell lineage labeling and single-cell transcriptional analyses have revealed a remarkable evolutionary increase in diversity of cortical progenitor cells, particularly at the genetic level. This includes, for example, multiple subtypes of aRGCs and bRGCs identified in human, macaque, or ferret but not in mouse or rat^{14,15,64,84}. Likewise, innovations in progenitor cell lineages are critical for the emergence and expansion of the OSVZ^{15,85}, the basal germinal layer typical of big and folded brains, which is absent in mice and is perturbed in human diseases that affect brain size⁸⁵.

The search for genetic mechanisms evolved in primate and human phylogeny which are likely relevant in the evolution of their neocortex, has led to the identification of primate-specific and human-specific genetic programs expressed in the developing cerebral cortex. These include whole collections of primate-specific miRNAs targeting cell cycle genes²² and also miRNA-mRNA modules in the embryonic human brain that undergo dynamic transitions during development and that identify expression networks in specific cell types^{86,87}. As for protein-coding genes, recent studies have identified genes that emerged in the recent human lineage by means of total or partial duplication, and that promote cortical progenitor cell proliferation. Other studies have identified programs of gene expression in cortical progenitor cells that are human-specific^{88–91}.

Species with large brains have a tendency of being folded, and knowledge of specific mechanisms involved in cortex folding has also increased recently. This highlights the relevance of the sodium ion channel *SCN3A* function in progenitor cells⁶⁸, of gliogenesis⁹², and of extracellular matrix proteins⁹³, in the mechanical aspect of tissue folding. Intriguingly, the organization of folds in the frontal cortex is widely conserved from Old World monkeys to hominoids⁹⁴, which offers a remarkable opportunity to study cortical evolution and the acquisition of higher brain functions in primates.

Discovering the developmental importance of microglia

Largely ignored previously in the field of neocortical development, microglia have become a new relevant component in our understanding of this process. Microglial cells are found in the developing neocortex at much earlier stages and at much greater abundance than previously considered; there, they interact intimately with progenitor cells and regulate their number, hence emerging as important players in the regulation of neurogenesis^{95–97}. Microglia may also contribute to progenitor cell delamination and expansion of ISVZ/OSVZ in primates⁹⁵. Moreover, microglia are critical regulators of brain wiring, contributing to the specification of neural circuits and relaying information from the periphery, including the microbiota^{98,99}.

Cerebral organoids as a model of human neocortical development

Although it is not possible to study the development of the human neocortex at the experimental level, major technological breakthroughs in the last few years on stem cell reprogramming and tissue culture offer possibilities that were previously unthinkable. Following the first protocol to generate cerebral organoids from human embryonic stem cells and induced pluripotent stem cells¹⁰⁰, these have become the Rosetta stone to study and manipulate human brain development¹⁰¹. Not only can we now grow human cerebral organoids for many months, recapitulating many of the early features of cortical development, but they can be grown to form functional circuits and be responsive to sensory stimuli¹⁰². Most importantly, the recent design of a culture protocol to generate highly reproducible cerebral organoids is a fundamental milestone for the consolidation of this as a faithful model of human brain development¹⁰³. Comparison between organoids grown from human and chimpanzee cells reveal human-specific features of cortical progenitor cells¹⁰⁴ and the validity of these organoids to advance our understanding of human brain evolution^{105,106}.

Conclusions

Understanding neocortical development requires working at many different levels, from single-cell transcriptomics to tissue mechanics, and this must be applied to studying histogenesis at multiple levels, from neurogenesis to connectivity and cortex folding. Understanding neocortical evolution and disease requires understanding development across relevant informative species and in the context of genetic or contextual failure. Recent technological advances offer unprecedented opportunities for conducting this research; for example, single-cell genomic analyses help elucidate molecular changes in human brain disease at the resolution of cell types¹⁰⁷, and patient-derived iPSCs are used to model and hopefully rescue the disease¹⁰⁸. Only through our ability to use and combine these amazing tools in creative ways will we decipher what makes us human and what genetic changes occurred during evolution that led to the development and emergence of the human neocortex.

Abbreviations

aRGC, apical radial glial cell; bRGC, basal radial glial cell; IPC, intermediate progenitor cell; ISVZ, inner subventricular zone; NSPC, neural stem and progenitor cell; OSVZ, outer subventricular zone; RGC, radial glial cell; SVZ, subventricular zone; VZ, ventricular zone

Acknowledgments

I am very thankful to all members of my lab and to close colleagues for thoughtful discussions and insights. I deeply apologize to all the authors whose primary work, though relevant to the topic, could not be cited here due to space limitations.

References

- Malatesta P, Hartfuss E, Götz M: **Isolation of radial glial cells by fluorescent-activated cell sorting reveals a neuronal lineage.** *Development.* 2000; **127**(24): 5253–63.
[PubMed Abstract](#)
- Nocor SC, Flint AC, Weissman TA, *et al.*: **Neurons derived from radial glial cells establish radial units in neocortex.** *Nature.* 2001; **409**(6821): 714–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Nocor SC, Martínez-Cerdeño V, Ivic L, *et al.*: **Cortical neurons arise in symmetric and asymmetric division zones and migrate through specific phases.** *Nat Neurosci.* 2004; **7**(2): 136–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Haubensak W, Attardo A, Denk W, *et al.*: **Neurons arise in the basal neuroepithelium of the early mammalian telencephalon: a major site of neurogenesis.** *Proc Natl Acad Sci U S A.* 2004; **101**(9): 3196–201.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Miyata T, Kawaguchi A, Saito K, *et al.*: **Asymmetric production of surface-dividing and non-surface-dividing cortical progenitor cells.** *Development.* 2004; **131**(13): 3133–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Attardo A, Calegari F, Haubensak W, *et al.*: **Live imaging at the onset of cortical neurogenesis reveals differential appearance of the neuronal phenotype in apical versus basal progenitor progeny.** *PLoS One.* 2008; **3**(6): e2388.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kowalczyk T, Pontious A, Englund C, *et al.*: **Intermediate neuronal progenitors (basal progenitors) produce pyramidal-projection neurons for all layers of cerebral cortex.** *Cereb Cortex.* 2009; **19**(10): 2439–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Stancik EK, Navarro-Quiroga I, Sellke R, *et al.*: **Heterogeneity in ventricular zone neural precursors contributes to neuronal fate diversity in the postnatal neocortex.** *J Neurosci.* 2010; **30**(20): 7028–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Smart IHM, Dehay C, Giroud P, *et al.*: **Unique morphological features of the proliferative zones and postmitotic compartments of the neural epithelium giving rise to striate and extrastriate cortex in the monkey.** *Cerebral Cortex.* 2002; **12**(1): 37–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Reillo I, de Juan Romero C, García-Cabezas MÁ, *et al.*: **A role for intermediate radial glia in the tangential expansion of the mammalian cerebral cortex.** *Cereb Cortex.* 2011; **21**(7): 1674–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Kelava I, Reillo I, Murayama AY, *et al.*: **Abundant occurrence of basal radial glia in the subventricular zone of embryonic neocortex of a lissencephalic primate, the common marmoset *Callithrix jacchus*.** *Cereb Cortex.* 2012; **22**(2): 469–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- García-Moreno F, Vasistha NA, Trevia N, *et al.*: **Compartmentalization of cerebral cortical germinal zones in a lissencephalic primate and gyrencephalic rodent.** *Cerebral Cortex.* 2012; **22**(2): 482–92.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Reillo I, Borrell V: **Germinal zones in the developing cerebral cortex of ferret: ontogeny, cell cycle kinetics, and diversity of progenitors.** *Cereb Cortex.* 2012; **22**(9): 2039–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Betizeau M, Cortay V, Patti D, *et al.*: **Precursor diversity and complexity of lineage relationships in the outer subventricular zone of the primate.** *Neuron.* 2013; **80**(2): 442–57.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Martínez-Martínez MÁ, De Juan Romero C, Fernández V, *et al.*: **A restricted period for formation of outer subventricular zone defined by Cdh1 and Trnp1 levels.** *Nat Commun.* 2016; **7**: 11812.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Pilz G-A, Shitamukai A, Reillo I, *et al.*: **Amplification of progenitors in the mammalian telencephalon includes a new radial glial cell type.** *Nat Commun.* 2013; **4**: 2125.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Llinares-Benadero C, Borrell V: **Deconstructing cortical folding: genetic, cellular and mechanical determinants.** *Nat Rev Neurosci.* 2019; **20**(3): 161–76.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Dehay C, Kennedy H, Kosik KS: **The outer subventricular zone and primate-specific cortical complexification.** *Neuron.* 2015; **85**(4): 683–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Johnson MB, Kawasawa YI, Mason CE, *et al.*: **Functional and evolutionary insights into human brain development through global transcriptome analysis.** *Neuron.* 2009; **62**(4): 494–509.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Miller JA, Ding SL, Sunkin SM, *et al.*: **Transcriptional landscape of the prenatal human brain.** *Nature.* 2014; **508**(7495): 199–206.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ayoub AE, Oh S, Xie Y, *et al.*: **Transcriptional programs in transient embryonic zones of the cerebral cortex defined by high-resolution mRNA sequencing.** *Proc Natl Acad Sci U S A.* 2011; **108**(36): 14950–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Arcila ML, Betizeau M, Cambronne XA, *et al.*: **Novel primate miRNAs coevolved with ancient target genes in germinal zone-specific expression patterns.** *Neuron.* 2014; **81**(6): 1255–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- de Juan Romero C, Bruder C, Tomasello U, *et al.*: **Discrete domains of gene expression in germinal layers distinguish the development of gyrencephaly.** *EMBO J.* 2015; **34**(14): 1859–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Woodworth MB, Girsakis KM, Walsh CA: **Building a lineage from single cells: genetic techniques for cell lineage tracking.** *Nat Rev Genet.* 2017; **18**(4): 230–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Pollen AA, Nowakowski TJ, Shuga J, *et al.*: **Low-coverage single-cell mRNA sequencing reveals cellular heterogeneity and activated signaling pathways in developing cerebral cortex.** *Nat Biotechnol.* 2014; **32**(10): 1053–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Miller DJ, Bhaduri A, Sestan N, *et al.*: **Shared and derived features of cellular diversity in the human cerebral cortex.** *Curr Opin Neurobiol.* 2019; **56**: 117–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Loo L, Simon JM, Xing L, *et al.*: **Single-cell transcriptomic analysis of mouse neocortical development.** *Nat Commun.* 2019; **10**(1): 134.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Telley L, Agirman G, Prados J, *et al.*: **Temporal patterning of apical progenitors and their daughter neurons in the developing neocortex.** *Science.* 2019; **364**(6400): pii: eaav2522.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Nowakowski TJ, Bhaduri A, Pollen AA, *et al.*: **Spatiotemporal gene expression trajectories reveal developmental hierarchies of the human cortex.** *Science.* 2017; **358**(6368): 1318–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Mi D, Li Z, Lim L, *et al.*: **Early emergence of cortical interneuron diversity in the mouse embryo.** *Science.* 2018; **360**(6384): 81–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mayer C, Hafemeister C, Bandler RC, *et al.*: **Developmental diversification of cortical inhibitory interneurons.** *Nature.* 2018; **555**(7697): 457–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Telley L, Govindan S, Prados J, *et al.*: **Sequential transcriptional waves direct the differentiation of newborn neurons in the mouse neocortex.** *Science.* 2016; **351**(6280): 1443–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Oberst P, Agirman G, Jabaudon D: **Principles of progenitor temporal patterning in the developing invertebrate and vertebrate nervous system.** *Curr*

- Opin Neurobiol.* 2019; **56**: 185–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
34. McConnell SK: **Constructing the cerebral cortex: neurogenesis and fate determination.** *Neuron.* 1995; **15**(4): 761–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 35. **F** Franco SJ, Gil-Sanz C, Martínez-Garay I, *et al.*: **Fate-restricted neural progenitors in the mammalian cerebral cortex.** *Science.* 2012; **337**(6095): 746–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 36. Marín O: **Brain development: The neuron family tree remodelled.** *Nature.* 2012; **490**(7419): 185–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 37. Desai AR, McConnell SK: **Progressive restriction in fate potential by neural progenitors during cerebral cortical development.** *Development.* 2000; **127**(13): 2863–72.
[PubMed Abstract](#)
 38. **F** Zahr SK, Yang G, Kazan H, *et al.*: **A Translational Repression Complex in Developing Mammalian Neural Stem Cells that Regulates Neuronal Specification.** *Neuron.* 2018; **97**(3): 520–537.e6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 39. **F** Seuntjens E, Nityanandam A, Miquelajauregui A, *et al.*: **Sip1 regulates sequential fate decisions by feedback signaling from postmitotic neurons to progenitors.** *Nat Neurosci.* 2009; **12**(11): 1373–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 40. **F** Vitali I, Fièvre S, Telley L, *et al.*: **Progenitor Hyperpolarization Regulates the Sequential Generation of Neuronal Subtypes in the Developing Neocortex.** *Cell.* 2018; **174**(5): 1264–1276.e15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 41. Fishell G, Kepecs A: **Interneuron Types as Attractors and Controllers.** *Annu Rev Neurosci.* 2019; **43**.
[PubMed Abstract](#) | [Publisher Full Text](#)
 42. Kriegstein A, Noctor S, Martínez-Cerdeño V: **Patterns of neural stem and progenitor cell division may underlie evolutionary cortical expansion.** *Nat Rev Neurosci.* 2006; **7**(11): 883–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
 43. Rakic P: **Evolution of the neocortex: a perspective from developmental biology.** *Nat Rev Neurosci.* 2009; **10**(10): 724–35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 44. **F** Cárdenas A, Villalba A, de Juan Romero C, *et al.*: **Evolution of Cortical Neurogenesis in Amniotes Controlled by Robo Signaling Levels.** *Cell.* 2018; **174**(3): 590–606.e21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 45. Cheung AF, Kondo S, Abdel-Mannan O, *et al.*: **The subventricular zone is the developmental milestone of a 6-layered neocortex: comparisons in metatherian and eutherian mammals.** *Cereb Cortex.* 2010; **20**(5): 1071–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
 46. Laguesse S, Creppe C, Nedialkova DD, *et al.*: **A Dynamic Unfolded Protein Response Contributes to the Control of Cortical Neurogenesis.** *Dev Cell.* 2015; **35**(5): 553–67.
[PubMed Abstract](#) | [Publisher Full Text](#)
 47. Alfano C, Gladwyn-Ng I, Couderc T, *et al.*: **The Unfolded Protein Response: A Key Player in Zika Virus-Associated Congenital Microcephaly.** *Front Cell Neurosci.* 2019; **13**: 94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 48. **F** Gladwyn-Ng I, Cordon-Barris L, Alfano C, *et al.*: **Stress-induced unfolded protein response contributes to Zika virus-associated microcephaly.** *Nat Neurosci.* 2018; **21**(1): 63–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 49. **F** Baizabal JM, Mistry M, García MT, *et al.*: **The Epigenetic State of PRDM16-Regulated Enhancers in Radial Glia Controls Cortical Neuron Position.** *Neuron.* 2018; **98**(5): 945–962.e8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 50. **F** Tang T, Zhang Y, Wang Y, *et al.*: **HDAC1 and HDAC2 Regulate Intermediate Progenitor Positioning to Safeguard Neocortical Development.** *Neuron.* 2019; **101**(6): 1117–1133.e5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 51. **F** de La Torre-Ubieta L, Stein JL, Won H, *et al.*: **The Dynamic Landscape of Open Chromatin during Human Cortical Neurogenesis.** *Cell.* 2018; **172**(1–2): 289–304.e18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 52. Silver DL: **Genomic divergence and brain evolution: How regulatory DNA influences development of the cerebral cortex.** *Bioessays.* 2016; **38**(2): 162–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 53. Tsunekawa Y, Britto JM, Takahashi M, *et al.*: **Cyclin D2 in the basal process of neural progenitors is linked to non-equivalent cell fates.** *EMBO J.* 2012; **31**(8): 1879–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 54. **F** Pilaz LJ, Lennox AL, Rouanet JP, *et al.*: **Dynamic mRNA Transport and Local Translation in Radial Glial Progenitors of the Developing Brain.** *Curr Biol.* 2016; **26**(24): 3383–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 55. Pilaz LJ, Silver DL: **Moving messages in the developing brain-emerging roles for mRNA transport and local translation in neural stem cells.** *FEBS Lett.* 2017; **591**(11): 1526–39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 56. **F** Pilaz LJ, McMahon JJ, Miller EE, *et al.*: **Prolonged Mitosis of Neural Progenitors Alters Cell Fate in the Developing Brain.** *Neuron.* 2016; **89**(1): 83–99.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 57. Lange C, Huttner WB, Calegari F: **Cdk4/cyclinD1 overexpression in neural stem cells shortens G1, delays neurogenesis, and promotes the generation and expansion of basal progenitors.** *Cell Stem Cell.* 2009; **5**(3): 320–31.
[PubMed Abstract](#) | [Publisher Full Text](#)
 58. Arai Y, Pulvers JN, Haffner C, *et al.*: **Neural stem and progenitor cells shorten S-phase on commitment to neuron production.** *Nat Commun.* 2011; **2**: 154.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 59. Nelson BR, Hodge RD, Bedogni F, *et al.*: **Dynamic interactions between intermediate neurogenic progenitors and radial glia in embryonic mouse neocortex: potential role in Dll1-Notch signaling.** *J Neurosci.* 2013; **33**(21): 9122–39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 60. **F** Fietz SA, Kelava I, Vogt J, *et al.*: **OSVZ progenitors of human and ferret neocortex are epithelial-like and expand by integrin signaling.** *Nat Neurosci.* 2010; **13**(6): 690–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 61. Hansen DV, Lui JH, Parker PR, *et al.*: **Neurogenic radial glia in the outer subventricular zone of human neocortex.** *Nature.* 2010; **464**(7288): 554–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
 62. Wang X, Tsai JW, LaMonica B, *et al.*: **A new subtype of progenitor cell in the mouse embryonic neocortex.** *Nat Neurosci.* 2011; **14**(5): 555–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 63. Reillo I, de Juan Romero C, Cárdenas A, *et al.*: **A Complex Code of Extrinsic Influences on Cortical Progenitor Cells of Higher Mammals.** *Cereb Cortex.* 2017; **27**(9): 4586–606.
[PubMed Abstract](#) | [Publisher Full Text](#)
 64. **F** Kalebic N, Gilardi C, Stepien B, *et al.*: **Neocortical Expansion Due to Increased Proliferation of Basal Progenitors Is Linked to Changes in Their Morphology.** *Cell Stem Cell.* 2019; **24**(4): 535–550.e9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 65. **F** Stenzel D, Wilsch-Bräuninger M, Wong FK, *et al.*: **Integrin $\alpha\beta_3$ and thyroid hormones promote expansion of progenitors in embryonic neocortex.** *Development.* 2014; **141**(4): 795–806.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 66. **F** Silva CG, Peyre E, Adhikari MH, *et al.*: **Cell-Intrinsic Control of Interneuron Migration Drives Cortical Morphogenesis.** *Cell.* 2018; **172**(5): 1063–1078.e19.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 67. **F** Silva CG, Peyre E, Nguyen L: **Cell migration promotes dynamic cellular interactions to control cerebral cortex morphogenesis.** *Nat Rev Neurosci.* 2019; **20**(6): 318–29.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 68. **F** Smith RS, Kenny CJ, Ganesh V, *et al.*: **Sodium Channel SCN3A (Na_v1.3) Regulation of Human Cerebral Cortical Folding and Oral Motor Development.** *Neuron.* 2018; **99**(5): 905–913.e7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 69. **F** Mayer S, Chen J, Velmesshev D, *et al.*: **Multimodal Single-Cell Analysis Reveals Physiological Maturation in the Developing Human Neocortex.** *Neuron.* 2019; **102**(1): 143–158.e7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 70. Bonnefont J, Tiberi L, van den Amele J, *et al.*: **Cortical Neurogenesis Requires Bcl6-Mediated Transcriptional Repression of Multiple Self-Renewal-Promoting Extrinsic Pathways.** *Neuron.* 2019; **103**(6): 1096–1108.e4.
[PubMed Abstract](#) | [Publisher Full Text](#)
 71. Borrell V, Cárdenas A, Ciceri G, *et al.*: **Slit/Robo signaling modulates the proliferation of central nervous system progenitors.** *Neuron.* 2012; **76**(2): 338–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 72. **F** Kostic M, Paridaen JTM, Long KR, *et al.*: **YAP Activity Is Necessary and Sufficient for Basal Progenitor Abundance and Proliferation in the Developing Neocortex.** *Cell Rep.* 2019; **27**(4): 1103–1118.e6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 73. **F** Tavano S, Taverna E, Kalebic N, *et al.*: **Insm1 Induces Neural Progenitor Delamination in Developing Neocortex via Downregulation of the Adherens Junction Belt-Specific Protein Plekha7.** *Neuron.* 2018; **97**(6): 1299–1314.e8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 74. Camargo Ortega G, Falk S, Johansson PA, *et al.*: **The centrosome protein AKNA regulates neurogenesis via microtubule organization.** *Nature.* 2019; **567**(7746): 113–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 75. **F** Kawaue T, Shitamukai A, Nagasaka A, *et al.*: **Lzts1 controls both neuronal delamination and outer radial glial-like cell generation during mammalian**

- cerebral development.** *Nat Commun.* 2019; **10**(1): 2780.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
76. Tabata H, Nakajima K: **Multipolar migration: the third mode of radial neuronal migration in the developing cerebral cortex.** *J Neurosci.* 2003; **23**(31): 9996–10001.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 77. Inoue M, Iwai R, Tabata H, *et al.*: **Prdm16 is crucial for progression of the multipolar phase during neural differentiation of the developing neocortex.** *Development.* 2017; **144**(3): 385–99.
[PubMed Abstract](#) | [Publisher Full Text](#)
 78. **F** Inoue S, Hayashi K, Fujita K, *et al.*: **Drebrin-like (Dbrn1) Controls Neuronal Migration via Regulating N-Cadherin Expression in the Developing Cerebral Cortex.** *J Neurosci.* 2019; **39**(4): 678–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 79. Rakic P: **Mode of cell migration to the superficial layers of fetal monkey neocortex.** *J Comp Neurol.* 1972; **145**(1): 61–83.
[PubMed Abstract](#) | [Publisher Full Text](#)
 80. Nadarajah B, Parnavelas JG: **Modes of neuronal migration in the developing cerebral cortex.** *Nat Rev Neurosci.* 2002; **3**(6): 423–32.
[PubMed Abstract](#) | [Publisher Full Text](#)
 81. Martínez-Martínez MÁ, Ciceri G, Espinós A, *et al.*: **Extensive branching of radially-migrating neurons in the mammalian cerebral cortex.** *J Comp Neurol.* 2019; **527**(10): 1558–76.
[PubMed Abstract](#) | [Publisher Full Text](#)
 82. Rakic P: **A small step for the cell, a giant leap for mankind: a hypothesis of neocortical expansion during evolution.** *Trends Neurosci.* 1995; **18**(9): 383–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 83. **F** Nakagawa N, Plestant C, Yabuno-Nakagawa K, *et al.*: **Memo1-Mediated Tiling of Radial Glial Cells Facilitates Cerebral Cortical Development.** *Neuron.* 2019; **103**(5): 836–852.e5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 84. Johnson MB, Wang PP, Atabay KD, *et al.*: **Single-cell analysis reveals transcriptional heterogeneity of neural progenitors in human cortex.** *Nat Neurosci.* 2015; **18**(5): 637–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 85. **F** Johnson MB, Sun X, Kodani A, *et al.*: **Aspm knockout ferret reveals an evolutionary mechanism governing cerebral cortical size.** *Nature.* 2018; **556**(7701): 370–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 86. Nowakowski TJ, Rani N, Golkaram M, *et al.*: **Regulation of cell-type-specific transcriptomes by microRNA networks during human brain development.** *Nat Neurosci.* 2018; **21**(12): 1784–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 87. Kosik KS, Nowakowski T: **Evolution of New miRNAs and Cerebro-Cortical Development.** *Annu Rev Neurosci.* 2018; **41**: 119–37.
[PubMed Abstract](#) | [Publisher Full Text](#)
 88. Florio M, Borrell V, Huttner WB: **Human-specific genomic signatures of neocortical expansion.** *Curr Opin Neurobiol.* 2017; **42**: 33–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
 89. **F** Florio M, Heide M, Pinson A, *et al.*: **Evolution and cell-type specificity of human-specific genes preferentially expressed in progenitors of fetal neocortex.** *eLife.* 2018; **7**: pii: e32332.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 90. **F** Fiddes IT, Lodewijk GA, Mooring M, *et al.*: **Human-Specific NOTCH2NL Genes Affect Notch Signaling and Cortical Neurogenesis.** *Cell.* 2018; **173**(6): 1356–1369.e22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 91. **F** Suzuki IK, Gacquer D, van Heurck R, *et al.*: **Human-Specific NOTCH2NL Genes Expand Cortical Neurogenesis through Delta/Notch Regulation.** *Cell.* 2018; **173**(6): 1370–1384.e16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 92. Rash BG, Duque A, Morozov YM, *et al.*: **Gliogenesis in the outer subventricular zone promotes enlargement and gyrification of the primate cerebrum.** *Proc Natl Acad Sci U S A.* 2019; **116**(14): 7089–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 93. Long KR, Newland B, Florio M, *et al.*: **Extracellular Matrix Components HAPLN1, Lumican, and Collagen I Cause Hyaluronic Acid-Dependent Folding of the Developing Human Neocortex.** *Neuron.* 2018; **99**(4): 702–719.e6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 94. **F** Amiez C, Sallet J, Hopkins WD, *et al.*: **Sulcal organization in the medial frontal cortex provides insights into primate brain evolution.** *Nat Commun.* 2019; **10**(1): 3437.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 95. **F** Barger N, Keiter J, Kreutz A, *et al.*: **Microglia: An Intrinsic Component of the Proliferative Zones in the Fetal Rhesus Monkey (Macaca mulatta) Cerebral Cortex.** *Cereb Cortex.* 2019; **29**(7): 2782–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 96. Noctor SC, Penna E, Shepherd H, *et al.*: **Periventricular microglial cells interact with dividing precursor cells in the nonhuman primate and rodent prenatal cerebral cortex.** *J Comp Neurol.* 2019; **527**(10): 1598–609.
[PubMed Abstract](#) | [Publisher Full Text](#)
 97. **F** Cunningham CL, Martinez-Cerdeno V, Noctor SC: **Microglia regulate the number of neural precursor cells in the developing cerebral cortex.** *J Neurosci.* 2013; **33**(10): 4216–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 98. Squarzone P, Oller G, Hoeffel G, *et al.*: **Microglia modulate wiring of the embryonic forebrain.** *Cell Rep.* 2014; **8**(5): 1271–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 99. Thion MS, Ginhoux F, Garel S: **Microglia and early brain development: An intimate journey.** *Science.* 2018; **362**(6411): 185–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 100. **F** Lancaster MA, Renner M, Martin CA, *et al.*: **Cerebral organoids model human brain development and microcephaly.** *Nature.* 2013; **501**(7467): 373–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 101. **F** Arlotta P, Pasca SP: **Cell diversity in the human cerebral cortex: from the embryo to brain organoids.** *Curr Opin Neurobiol.* 2019; **56**: 194–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 102. **F** Quadrato G, Nguyen T, Macosko EZ, *et al.*: **Cell diversity and network dynamics in photosensitive human brain organoids.** *Nature.* 2017; **545**(7652): 48–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 103. **F** Velasco S, Kedaigle AJ, Simmons SK, *et al.*: **Individual brain organoids reproducibly form cell diversity of the human cerebral cortex.** *Nature.* 2019; **570**(7762): 523–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 104. **F** Mora-Bermúdez F, Badsha F, Kanton S, *et al.*: **Differences and similarities between human and chimpanzee neural progenitors during cerebral cortex development.** *eLife.* 2016; **5**: pii: e18683.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 105. **F** Camp JG, Badsha F, Florio M, *et al.*: **Human cerebral organoids recapitulate gene expression programs of fetal neocortex development.** *Proc Natl Acad Sci U S A.* 2015; **112**(51): 15672–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 106. **F** Pollen AA, Bhaduri A, Andrews MG, *et al.*: **Establishing Cerebral Organoids as Models of Human-Specific Brain Evolution.** *Cell.* 2019; **176**(4): 743–756.e17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 107. **F** Velmeshev D, Schirmer L, Jung D, *et al.*: **Single-cell genomics identifies cell type-specific molecular changes in autism.** *Science.* 2019; **364**(6441): 685–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 108. **F** Klaus J, Kanton S, Kyrousi C, *et al.*: **Altered neuronal migratory trajectories in human cerebral organoids derived from individuals with neuronal heterotopia.** *Nat Med.* 2019; **25**(4): 561–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Laurent Nguyen

GIGA-Stem Cells, Molecular regulation of neurogenesis, University of Liège, Liège, Belgium

Competing Interests: Laurent Nguyen has written with Victor Borrell on one consortium paper in the last three years.

2. Simon Hippenmeyer

Institute of Science and Technology Austria (IST Austria), Klosterneuburg, Austria

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research