Check for updates

CASE REPORT

REVISED Case Report: Unprovoked venous thromboembolism in a young adult male [version 2; peer review: 2 approved]

Previously titled: Case Report: Unprovoked venous thromboembolism in a young adult

Jeyhan Dhabhar¹, Varshil Mehta ^{1,2}, Nimit Desai¹, Sameer Dawoodi¹, Sojib Bin Zaman ³¹Department of Internal Medicine, MGM Medical College, Navi Mumbai, India²Department of Internal Medicine, Dr. L. H. Hiranandani Hospital, Mumbai, India³Maternal and Child Health Division, International Centre for Diarrhoeal Disease Research, Bangladesh (icddr,b), Dhaka, Bangladesh**v2** First published: 14 Feb 2019, 8:182
<https://doi.org/10.12688/f1000research.18202.1>Latest published: 25 Mar 2019, 8:182
<https://doi.org/10.12688/f1000research.18202.2>**Abstract**

A 24-year-old male was presented to us with sudden onset of chest pain and dyspnea for the past one hour. There was no history of calf pain, trauma, surgery, prolonged immobilization, long-haul air travel, bleeding diathesis or any other co-morbidity. The patient denied any addiction history. The heart rate was 114 beats/min, and blood pressure was 106/90 mmHg. Electrocardiogram showed tachycardia with S₁Q₃T₃ pattern. The left arterio-venous Doppler study was suggestive of a thrombus in popliteal vein and sapheno-popliteal junction. The CT-Pulmonary Angiogram scan was suggestive of a massive pulmonary thromboembolism. The patient was thrombolysed with Intravenous Alteplase immediately and was put on tab Rivaroxaban for maintenance. He was later discharged after being stable.

Unprovoked venous thromboembolism (VTE) is very rare and has the potential to lead to pulmonary embolism which could be disastrous, especially in young adults. We present such a case where unprovoked VTE was diagnosed and treated. This case suggests that high clinical suspicion is the key for the diagnosis of acute pulmonary embolism, especially in the absence of history suggestive of deep vein thrombosis.

Keywords

Pulmonary Embolism, Idiopathic, Thromboembolism

Open Peer Review**Approval Status**

	1	2
version 2		
(revision)		
25 Mar 2019		
version 1		
14 Feb 2019		

1. **Shakti A. Goel** , Indian Spinal Injuries Center, New Delhi, India2. **Himel Mondal** , Fakir Mohan Medical College and Hospital, Balasore, India

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding authors: Varshil Mehta (varshilmehta@gmail.com), Sojib Bin Zaman (sojibbz@gmail.com)

Author roles: **Dhabhar J:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation; **Mehta V:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Desai N:** Writing – Review & Editing; **Dawoodi S:** Methodology, Resources, Writing – Review & Editing; **Zaman SB:** Conceptualization, Project Administration, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2019 Dhabhar J *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Dhabhar J, Mehta V, Desai N *et al.* **Case Report: Unprovoked venous thromboembolism in a young adult male [version 2; peer review: 2 approved]** F1000Research 2019, 8:182 <https://doi.org/10.12688/f1000research.18202.2>

First published: 14 Feb 2019, 8:182 <https://doi.org/10.12688/f1000research.18202.1>

REVISED Amendments from Version 1

Thanks to the reviewers for their time and helpful suggestions. These were the changes made as suggested by the reviewers:

- 1) As per the advice of the second reviewer, we have added "male" in the title.
- 2) We have also added a sentence on the heart rate and BP of the patient in the abstract as per the second reviewer's advice. The sentence reads as "The heart rate was 114/min, and blood pressure was 106/90 mmHg."
- 3) In the Introduction section, we have added the word "globally" as the second reviewer had a query whether it should be globally or regionally. So we have clarified that.
- 4) We have also added a line "Also, the detailed examination (including Homan's and Moses sign) was performed and was deemed unremarkable" in the same paragraph as it was a query asked by the first reviewer.
- 5) We also mentioned that Trop I wasn't available as a query on the same was raised by the first reviewer.
- 6) At the end of the Introduction we have added details about the patient satisfaction and follow up.

See referee reports

Introduction

Venous thromboembolism (VTE) consists of pulmonary embolism (PE) and deep vein thrombosis (DVT). It is one of the leading causes of cardiovascular disability and impaired quality of life. It also causes major long-term complications, which include recurrent VTE, chronic thromboembolic pulmonary hypertension and post-thrombotic syndrome¹.

PE is said to be the third leading cause for cardiovascular mortality (after myocardial infarction and stroke). It leads to 100 000 deaths annually globally. It is not only the leading preventable cause of death in admitted patients¹ but also the leading cause of deaths in mothers during pregnancy in developed countries².

Idiopathic or unprovoked VTE is defined as "any VTE in the absence of an identifiable predisposing factor"³. Taking into consideration that unprovoked VTE is very rare, especially in the young adults, we put forward a case report of VTE in a 24-year-old man.

Case report

A 24-year-old man was brought to the emergency department of a hospital, by his office-colleagues, complaining of sudden onset of chest pain and dyspnea at rest, for the last one hour. It was not associated with sweating, palpitations, cough, hemoptysis, syncope, giddiness, leg pain, pedal edema, fever, rash, or any bleeding manifestations.

History of calf pain, trauma, surgery, prolonged immobilization, long-haul air travel, bleeding diathesis or any other co-morbidity was not reported by the patient. The patient also denied having any addiction history. Family history was found to be insignificant.

On admission, the patient's heart rate was 114/min, and blood pressure was 106/90 mmHg. His respiratory rate was 22/min, and O₂ saturation rate was 82% at room air. BMI was 20.76 kg/m². There was no murmur or gallop on cardiovascular examination. Air entry was reduced in the left infra-axillary region. Also, the detailed examination (including Homan's and Moses sign) was performed and was deemed unremarkable. Electrocardiogram (ECG) showed tachycardia with S₁Q₃T₃ pattern, and chest X-ray was suggestive of obliteration of left costo-phrenic angle. The D-Dimer (17.31 ug/ml) was elevated, 34 times above the normal upper limit (0.5 ug/ml).

CT-Pulmonary Angiogram (Figure 1) was suggestive of a massive pulmonary thromboembolism. The pulmonary trunk was dilated to ~30 mm. There was a non-lumen occluding circumferential filling defect in the main pulmonary trunk, with maximum thickness of 4.5 mm. A large partial-lumen occluding filling defect was noted in the left main pulmonary artery, which was extending further into the hilar branch, occluding the lumen completely. Another larger complete lumen occluding filling defect was noted in the right main pulmonary artery. These filling defects were extending into the segmental and sub-segmental branches of the lateral segment of the right middle, lingual and bilateral lower lobe. The RV: LV ratio was 2:1. All four pulmonary veins were normal, and there was no evidence of mediastinal pathology.

On admission, the patient also underwent a bilateral arteriovenous Doppler study, which was suggestive of a partially-lumen- occluding thrombus in the proximal part of left popliteal vein and a completely lumen-occluding thrombus at the left saphenopopliteal junction, approximately 14 cm long. The veins of both legs showed arterialized waveforms.

On 2D echocardiography, right atrium and right ventricle was mildly dilated with grade I, tricuspid regurgitation (TR) and

Figure 1. CT Pulmonary Angiogram showing non-lumen occluding thrombus (indicated by red arrows) in the right and left main pulmonary trunk.

pulmonary arterial systolic pressure by TR jet was 55 mmHg suggestive of moderate pulmonary artery hypertension. No regional wall motion abnormality was observed and left ventricular ejection fraction was 60%.

The coagulation profile was within normal limits. All other blood investigations i.e. hemogram, serum electrolytes, renal and liver function tests were within normal range. The patient had mild hyperuricemia with serum uric acid level being 7.4 mg/dL (Normal: 3.5 – 7.2 mg/dL). Cardiac enzymes (Creatine PhosphoKinase-MB and Troponin T) were mildly elevated. Trop I levels testing was not available at our centre and hence, was not done.

Although it was a clear case of massive VTE, the underlying etiology of such an event could not be extrapolated. Since the patient was a 24-year-old man, without any risk factors or comorbidities, the final diagnosis of unprovoked VTE was made.

The patient was thrombolysed with Injection Alteplase infusion (100mg IV, over two hours) with Injection Enoxaparin 60 mg given subcutaneously every 12 hours. After which, the patient developed hypotension which was treated with inotropic support. Although he had tachycardia post-thrombolysis for the next two days, his blood pressure returned to normal on the third day. 48-hours after giving thrombolytic treatment, the left lower limb venous doppler was done which was suggestive of a partial-lumen occluding the thrombus in popliteal vein extending from saphenopopliteal junction to mid-leg approximately 10 cm long. The CT-pulmonary angiogram was not repeated (post-thrombolysis), due to financial constraints; however, the patient improved drastically. He was shifted to general ward on day 4 post admission. Tab Rivaroxaban 15 mg (12 hourly, orally) was prescribed for the next three weeks. The patient was discharged successfully on the 15th day of admission. One week after discharge, the patient was advised Tab. Rivaroxaban 20 mg, once daily. The patient was most satisfied with the treatment and the recovery he'd made, but this was not recorded in a form of any official questionnaire (but he had mentioned this verbally).

The patient had followed up in our center for the first three months, following which relocated to Uttar Pradesh, India (his hometown). He experienced no side effects of Rivaroxaban. At three months of post discharge, a lower limb doppler was repeated which showed more than 50% reduction in size of the obstructing thrombus as compared to the previous one done at admission. Post three months of discharge, the patient has not returned back yet and has been lost to follow up.

Discussion

VTE in a young male patient is life-threatening, especially if it is unprovoked. The most common and important ECG finding in a patient with PE is sinus tachycardia and the presence of S₁Q₃T₃ pattern in the ECG. Presence of these findings shortened the time for diagnosis of acute PE in our patient and the management after that⁴.

Literature suggests that the risk of early death among patients with symptomatic PE is 18-fold higher when compared with the patients having DVT alone⁵. Had this patient presented with sinus tachycardia alone, he would have undergone a battery of investigations, and prompt management (thrombolysis) would have been delayed. Furthermore, the incidence rate of VTE is about 1.5 per 1,000 person-years while appearance of DVT is twice as common⁶.

Since our patient did not have any clinical features suggestive of DVT, having a Doppler study suggestive of DVT, was more of an additional finding, confirming VTE in this patient. A young man with sudden chest pain and dyspnea without any leg pain and who is otherwise healthy, is likely to be mismanaged, as suspicion of developing pulmonary or venous thrombo-embolism is quite low.

Before attempting to use the Wells criteria, a point-score based on clinical features and the likelihood of diagnoses other than PE, the clinician must first have a suspicion of the diagnosis before attempting to apply the Wells criteria⁷. On admission, the Wells score for PE was 4.5 with a moderate probability of having a PE for the present patient based on the history and clinical features. The findings on CT Pulmonary Angiogram scan and ECG, increased the possibility of a PE, hence 3 points and 1.5 points were added for presence of tachycardia in our patient's total score. The well's score for DVT was 0, as there was no history of cancer, surgery, immobilization, calf swelling, superficial veins, lower limb swelling, tenderness, paralysis or previous history of DVT. Hence high clinical suspicion remains the key in diagnosing PE, especially in the absence of DVT.

A case of unprovoked VTE, especially acute PE, needs to be extensively worked up, to prevent recurrence. Apart from history, family members needs to be screened for coagulopathy or asymptomatic VTE, which was not done in this case. Although, coagulation profile, i.e. International Normalised Ratio (INR), prothrombin time, and activated partial thromboplastin time (APTT), was normal in our patient, it is advisable to do a thrombophilia profile. The thrombophilia profile will help to anticipate the tendency to develop pathological clotting or thrombotic disorders and consists of namely, factor V Leiden mutation, prothrombin gene mutation, protein C resistance, anti-beta 2 glycoprotein, anticardiolipin antibodies, protein S activity and serum homocysteine levels⁸. Due to the huge cost of these investigations, it becomes difficult to run such investigations while dealing with a case of acute PE in hospital. Thrombophilia profile is not indicated for the patients who are on anticoagulation therapy. Our patient refused to give consent for doing a thrombophilia profile, hence the procedure was not done.

Inj. Alteplase is indicated as a fibrinolytic agent for the lysis of the clot seen in acute massive PE⁹. Massive PE is caused by obstruction (more than 50% of the cross-sectional area) of the pulmonary arterial tree, leading to an acute and subsequently severe cardiopulmonary failure due to right ventricular overload¹⁰.

Conclusion

Silent VTE can develop into PE which may remain unrecognised for a long time before the clinical features develop. High clinical suspicion is the key, for the diagnosis of acute PE in young patients, especially in the absence of history suggestive of DVT. In the absence of hemodynamic instability, the decision of fibrinolysis, in acute massive PE, largely depends on CT-Pulmonary Angiogram.

Consent

Written informed consent was taken from the patient and hospital authority to publish this case.

Data availability

All data underlying the results are available as part of the article and no additional source data are required.

Grant information

The author(s) declare that no grants were involved in supporting this work.

Acknowledgments

Medkru Research Group.

References

- Jerjes-Sánchez C: **Mechanisms of thrombosis**. In: *Thrombolysis in Pulmonary Embolism*. Springer, Berlin. 2015; 1–17.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Konstantinides SV, Barco S, Lankeit M, *et al.*: **Management of Pulmonary Embolism: An Update**. *J Am Coll Cardiol*. 2016; 67(2): 976–990.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Iorio A, Kearon C, Filippucci E, *et al.*: **Risk of recurrence after a first episode of symptomatic venous thromboembolism provoked by a transient risk factor: a systematic review**. *Arch Intern Med*. 2010; 170(19): 1710–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tapson VF: **Acute pulmonary embolism**. *N Engl J Med*. 2008; 358(10): 1037–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Heit JA, Silverstein MD, Mohr DN, *et al.*: **Predictors of survival after deep vein thrombosis and pulmonary embolism: a population-based, cohort study**. *Arch Intern Med*. 1999; 159(5): 445–453.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Naess IA, Christiansen SC, Romundstad P, *et al.*: **Incidence and mortality of venous thrombosis: a population-based study**. *J Thromb Haemost*. 2007; 5(4): 692–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wells PS, Anderson DR, Rodger M, *et al.*: **Excluding pulmonary embolism at the bedside without diagnostic imaging: management of patients with suspected pulmonary embolism presenting to the emergency department by using a simple clinical model and d-dimer**. *Ann Intern Med*. 2001; 135(2): 98–107.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Venous thromboembolic diseases: diagnosis, management and thrombophilia testing**. Clinical guideline [CG144]. Published on 27 June 2012, Last updated on November 2015.
[Reference Source](#)
- Martin C, Sobolewski K, Bridgeman P, *et al.*: **Systemic Thrombolysis for Pulmonary Embolism: A Review**. *P T*. 2016; 41(12): 770–775.
[PubMed Abstract](#) | [Free Full Text](#)
- Sadeghi A, Brevetti GR, Kim S, *et al.*: **Acute massive pulmonary embolism: role of the cardiac surgeon**. *Tex Heart Inst J*. 2005; 32(3): 430–433.
[PubMed Abstract](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 01 April 2019

<https://doi.org/10.5256/f1000research.20440.r46157>

© 2019 Goel S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Shakti A. Goel

Department of Spine Surgery, Indian Spinal Injuries Center, New Delhi, Delhi, India

I believe that the authors have answered the queries meticulously and the article may be considered for indexing now.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Medicine, Surgery, Orthopaedics, Spine, Innovation, Technoinnovations

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 18 March 2019

<https://doi.org/10.5256/f1000research.19910.r45517>

© 2019 Mondal H. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Himel Mondal

Department of Physiology, Fakir Mohan Medical College and Hospital, Balasore, Odisha, India

Overall evaluation

This is an interesting case report of venous thromboembolism in a young adult male without any predisposing factors. The authors described the case in detail with relevant and recent literature

references. However, there is a need of a minor revision to enrich the manuscript. The author may revise the manuscript keeping in following points in mind:

Title

As this is a case of a “young adult male,” authors may add the word “male” in the title of the manuscript.

Abstract

The patient presented with chest pain and dyspnoea. The second line of the “Abstract” described that the patient did not have any “calf pain.” Authors could add some cardiological history. They could also add a glimpse of clinical examination (especially heart rate and rhythm) rather than starting with “ECG showed tachycardia...”

Keywords

Authors may add more keywords according to MeSH for wider dissemination of the article.

Introduction

PE causes “100000 deaths” annually. Is it global or regional data? Please specify.

Case report

Authors may add the time when the patient was brought to the hospital, along with the nature of his works in previous 12 hours, if the information is available.

What was the rhythm of the pulse in clinical examination? If the data is available, the authors may add it.

Discussion

Discussion is adequate and well written.

Conclusion

Written well and it is within the scope of the article.

References

References are relevant and recent.

Is the background of the case’s history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Partly

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Yes

Is the case presented with sufficient detail to be useful for other practitioners?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Cardiovascular physiology, Exercise physiology, Public health, Medical education

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 18 Mar 2019

Sojib Bin Zaman, International Centre For Diarrhoeal Disease Research, Dhaka, Bangladesh

Thank you for your reviews and approving it.

Changes we made:

In title: We have added "male" in the title.

Abstract: We have also added heart rate and bp in the abstract.

Introduction: It leads to 100 000 deaths annually globally
The heart rhythm was normal and hence we didn't add it.

Hopefully, we have made adequate changes. Once again thank you for reviewing for us.
Sojib.

Competing Interests: None

Reviewer Report 15 February 2019

<https://doi.org/10.5256/f1000research.19910.r44458>

© 2019 Goel S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Shakti A. Goel

Department of Spine Surgery, Indian Spinal Injuries Center, New Delhi, Delhi, India

This is an interesting case report which discusses a rare phenomenon of unprovoked venous thromboembolism and its management. I would congratulate the authors for their work. However, there are a few concerns:

Though the authors have mentioned that there was no history of long air travel, it will be helpful to know if the patient had any history of long car travel before the event precipitated. It has been noted that DVT is commonly encountered in patients who undergo long car travel. Such information, if added, shall make the physicians more aware.

Did the physicians/authors conduct any clinical examination to test DVT. Were Homan's or Mosses sign performed. If yes, what were the results. If not, it will be helpful to know the reason for not performing them. A little detail on clinical examination of the patient should also be added to the manuscript besides the symptoms and findings.

It will also be helpful to know the Trop I values, if they were assessed by the authors.

The authors have mentioned that patient was kept on Rivaroxaban. Were there any side effects that were encountered. It will also be helpful if the authors could mention about the follow-up period of the patient in the manuscript.

Was any assessment done in the post discharge period. Was there any change in vessel size? Any adaptive remodeling seen with the treatment using Rivaroxaban .

Was the patient satisfied with the treatment? Any satisfaction index which was assessed. Authors may add it if found to be appropriate. They could use the attached reference for the same.

References

1. Guo LW, Wang B, Goel SA, Little C, et al.: Halofuginone stimulates adaptive remodeling and preserves re-endothelialization in balloon-injured rat carotid arteries. *Circ Cardiovasc Interv.* 2014; **7** (4): 594-601 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Srivastava N, Goel S: A study of patient satisfaction level in inpatient spine department of a tertiary care multi-speciality hospital. *Journal of Medical Research and Innovation.* 2018. [Publisher Full Text](#)
3. Goel SA, Guo LW, Liu B, Kent KC: Mechanisms of post-intervention arterial remodelling. *Cardiovasc Res.* 2012; **96** (3): 363-71 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Partly

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Yes

Is the case presented with sufficient detail to be useful for other practitioners?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Medicine, Surgery, Orthopaedics, Spine, Innovation, Technoinnovations

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have

significant reservations, as outlined above.

Author Response 15 Feb 2019

Sojib Bin Zaman, International Centre For Diarrhoeal Disease Research, Dhaka, Bangladesh

Thank you Dr for reviewing our paper and taking your time out for the same.

Please find my replies to your queries.

- There was no history of prolonged air or car travel. Also, the detailed examination was performed and was deemed unremarkable; hence was not mentioned here (including Homan's and Moses sign).
- Trop I level test are not available in our center, hence was not done, However, we did Trop T and CPK MB, which were normal.
- The patient had followed up in our center for the first three months, following which relocated to Uttar Pradesh, India (his hometown). He experienced no side effects of Rivaroxaban. At three months, post discharge, a lower limb doppler was repeated which showed more than 50% reduction in size of the obstructing thrombus as compared to the previous one done at admission. Post three months of discharge, the patient has not returned back yet and has been lost to follow up.
- The patient was most satisfied with the treatment and the recovery he'd made, but this was not recorded in a form of any official questionnaire (but he had mentioned this verbally); hence not mentioned here.

Hopefully, these comments will satisfy the queries. If you have any more queries, please feel free to post and we shall be happy to reply them.

Competing Interests: No competing interests present.

Reviewer Response 19 Feb 2019

Shakti A. Goel, Indian Spinal Injuries Center, New Delhi, India

I had an opportunity to go through the review/reply and believe that the authors have addressed the comments meticulously. The article can be accepted once these details have been added to the article.

This article will certainly be very helpful for the physicians around the globe especially in low to middle income countries.

Competing Interests: None

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research