REVIEW

Recent advances in SHH medulloblastoma progression: tumor suppressor mechanisms and the tumor microenvironment

[version 1; peer review: 2 approved]

Lukas Tamayo-Orrego ¹⁻³, Frédéric Charron ^{1,2,4,5}

¹Montreal Clinical Research Institute (IRCM), Montreal, Quebec, Canada

²Integrated Program in Neuroscience, McGill University, Montreal, Quebec, Canada

³Grupo Neuroaprendizaje, Autonomous University of Manizales, Manizales, Colombia

⁴Department of Anatomy and Cell Biology, Division of Experimental Medicine, McGill University, Quebec, Canada

⁵Department of Medicine, University of Montreal, Montreal, Quebec, Canada

V1 First published: 29 Oct 2019, 8(F1000 Faculty Rev):1823

<https://doi.org/10.12688/f1000research.20013.1>

Latest published: 29 Oct 2019, 8(F1000 Faculty Rev):1823

<https://doi.org/10.12688/f1000research.20013.1>

Abstract

Medulloblastoma, the most common of the malignant pediatric brain tumors, is a group of four molecularly and clinically distinct cancers with different cells of origin. One of these medulloblastoma groups displays activation of Sonic hedgehog (SHH) signaling and originates from granule cell precursors of the developing cerebellum. Ongoing basic and clinical research efforts are tailored to discover targeted and safer therapies, which rely on the identification of the basic mechanisms regulating tumor initiation, progression, and metastasis. In SHH medulloblastoma, the mechanisms regulating neural progenitor transformation and progression to advanced tumors have been studied in some detail. The present review discusses recent advances on medulloblastoma progression derived from studies using mouse models of SHH medulloblastoma. We focus on mechanisms that regulate progression from precancerous lesions to medulloblastoma, describing novel roles played by tumor suppressor mechanisms and the tumor microenvironment.

Keywords

Medulloblastoma, Sonic hedgehog, cell senescence, tumor microenvironment, tumor progression

Open Peer Review

Approval Status

	1	2
version 1		
29 Oct 2019		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Timothy R Gershon**, University of North Carolina School of Medicine, North Carolina, USA
2. **James Manfredi**, Icahn School of Medicine at Mount Sinai, New York, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Frédéric Charron (frederic.charron@ircm.qc.ca)

Author roles: **Tamayo-Orrego L:** Conceptualization, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing;
Charron F: Conceptualization, Funding Acquisition, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the Canadian Institutes of Health Research (FDN334023) and Canada Foundation for Innovation (33768).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 Tamayo-Orrego L and Charron F. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Tamayo-Orrego L and Charron F. **Recent advances in SHH medulloblastoma progression: tumor suppressor mechanisms and the tumor microenvironment [version 1; peer review: 2 approved]** F1000Research 2019, 8(F1000 Faculty Rev):1823 <https://doi.org/10.12688/f1000research.20013.1>

First published: 29 Oct 2019, 8(F1000 Faculty Rev):1823 <https://doi.org/10.12688/f1000research.20013.1>

Introduction

The multi-stage nature of cancer

Cancer is a multi-step process where the development of high-grade tumors normally requires the occurrence of multiple and sequential genetic events¹, a process called multi-stage tumorigenesis². The paradigms of this model are adult epithelial cancers, such as pancreatic and colorectal³. Each histopathological stage of the disease is followed by a more 'malignant' form, and the transitions are governed by a genetic event or 'hit' that leads to the clonal expansion of the mutated cell; within this framework, each stage can be seen as a rate-limiting step³. In the case of adult epithelial cancers, an average of five to seven 'hits' seems to be required for the formation of a full-blown tumor^{4,5}. It is useful to introduce here the terms 'gatekeeper' and 'caretaker' mutation to distinguish, respectively, mutations that are able to *initiate* the process of tumorigenesis and those that *promote* it or allow *progression* from one stage to the next. For example, a germ-line deletion of the *APC* gene leads to the *initiation* of colorectal carcinogenesis, and therefore *APC* is a gatekeeper for the colon epithelium. On the other hand, although *TP53* mutations appear at high frequency during colon tumorigenesis, they more rarely lead to colon carcinoma when present in the germ-line; in this context, *TP53* can be considered as a caretaker gene. Gatekeepers tend to be tissue specific, as their gene products can only deregulate the growth of the specific cell populations where they play a physiological role. For example, *VHL* is a gatekeeper in renal epithelial cells, *NF1* in the peripheral nervous system, and *MEN1* in endocrine cells⁶. In the present work, we will see how the hedgehog receptor *PTCH1* functions as gatekeeper in neuronal precursors of the cerebellum.

Medulloblastomas are molecularly and clinically diverse

There has been a revolution in cancer genomics in the last two decades, and neuro-oncology is not an exception. While many different pediatric brain tumors were initially aggregated together under the denomination of 'primitive neuroectodermal tumors', new research methods enable the classification and grouping of brain tumors and the characterization of new rare diseases⁷. Medulloblastoma (MB) is one of the most common pediatric brain tumors and the most common malignant pediatric brain tumor. A tumor of the posterior fossa affecting mostly infants and children, MB is also diagnosed in adults⁸. The current knowledge based on gene expression analysis allows the identification of four molecular groups of MB⁹. These groups differ not only in terms of gene expression but also in their methylation patterns, mutational events, and clinical characteristics, such as prognosis or risk of metastasis^{8,10–14}. The four MB groups are called WNT-MB, Sonic hedgehog (SHH)-MB, group 3 MB, and group 4 MB. As the name indicates, WNT-MB and SHH-MB display deregulation in WNT signaling and SHH signaling, respectively. Group 3 MB has a photoreceptor/retinal expression signature, while group 4 MB expresses neuronal genes¹². The current idea is that each different MB group is derived from a specific cell of origin^{15,16}, which therefore determines the clinical and molecular behavior of the disease. Additionally, the four MB groups have recently been subdivided into 12 subtypes which display specific molecular and

clinical characteristics¹⁷. This section discusses the mechanisms of MB formation in SHH-MB.

Shh signaling induces granule cell precursor proliferation

Cerebellar granule neurons are the largest neuronal population of the nervous system¹⁸. These cells arise from the granule cell precursors (GCPs), a population of cells derived from the rhombic lip that populates the surface of the cerebellum and forms the external granule cell layer (EGL)¹⁹. In mice, after one week of proliferation, GCPs start to differentiate and migrate towards the internal granule cell layer (IGL) using Bergmann glia as a scaffold. The EGL disappears completely at the third postnatal week (Figure 1)²⁰. Although different molecules promote the proliferation of neuronal precursors, Shh is the most important mitogen for GCPs^{21–23}.

The essential components of the Hedgehog (Hh) pathway are the membrane proteins Patched1 (Ptc1) and Smoothened (Smo) as well as Gli transcription factors²⁴. Smo and Gli are activators of the pathway, while Ptc1 represses Smo activity. Shh binds to its receptor Ptc1, an event that relieves Smo repression by Ptc1 and allows the activation of Gli transcription factors²⁵. This signaling mechanism relying on the Ptc1–Smo–Gli axis is known as the canonical Hh signaling pathway. Shh-induced GCP proliferation is dependent on N-Myc transcription downstream of Gli²⁶ and the subsequent induction of D-type cyclins, which promote cell cycle progression and cell proliferation^{27,28}.

Mutations in Shh signaling pathway genes cause medulloblastoma

The SHH-MB group comprises 30% of all MBs¹². Mutations in *PTCH1*, *SMO*, and *SUFU* or amplification of *GLI1*, *GLI2*, and *N-MYC*⁸ cause SHH-MB. *PTCH1* mutations are the most common MB driver²⁹. Mice heterozygous for *Ptc1* (*Ptc1*^{+/-} mice), or mice bearing an active form of *Smo*, develop MBs that recapitulate the human disease, demonstrating that mutations in components of the Shh pathway are responsible for MB tumorigenesis^{30,31}. Consistent with their responsiveness to Shh signaling, GCPs have been shown through several lines of evidence to be the cell of origin of Shh-MB in mice^{32–34}. For example, activation of Smo or deletion of *Ptc1* using the GCP-specific driver *Atoh1* (also known as *Math1*) causes Shh-MB. MB can also be generated by Shh pathway activation in earlier cell progenitors (neural stem cells), such as GFAP- and Olig2-positive cells, but only after stem cells commit to a GCP lineage^{33,34}. Recent single-cell RNA-seq experiments show that among all the cerebellar cell types, GCP gene expression most closely matches SHH-MB gene expression¹⁶, supporting the conclusion that GCPs are also the cell of origin of SHH-MB in humans.

In rare cases, patients with Li-Fraumeni syndrome (germ-line *TP53* mutations) develop MB. These tumors are always SHH-MBs that display additional mutations leading to constitutive Hh pathway signaling³⁵; this requirement of Hh pathway hyperactivation for MB formation supports the idea that Hh signaling genes initiate tumorigenesis in GCPs.

Figure 1. Cerebellum development. Granule cell precursors (GCPs), the progenitors that give origin to granule neurons, proliferate in the external granule cell layer (EGL) of the cerebellum in response to Purkinje neuron-derived Shh. Granule neurons populate the internal granule cell layer after GCPs differentiate and migrate through the Purkinje cell layer using the radial processes of Bergmann glia. The peak of GCP proliferation spans the first seven postnatal days in the mouse; in humans, it extends from the second half of gestation to the sixth postnatal month³⁶. IGL, internal granule cell layer.

Hh signaling activity alone is not sufficient for the formation of advanced MB

In the case of activating *Smo* mutations, MB seems to develop very early from progenitors directly without transiting through a preneoplastic stage³¹. In contrast, the removal of one *Ptch1* allele leads to the formation of precancerous lesions, an intermediate histopathological stage³². In the specific case of the *Ptch1* model, it was initially thought that in MBs caused by *Ptch1* mutations, tumor formation is a two-step process where *Ptch1* haploinsufficiency causes preneoplasia and *Ptch1* loss of heterozygosity (LOH) causes constitutive Hh pathway activation, leading to MB progression^{37,38}. Together with the facts that pediatric tumors have been considered to display a paucity of mutations, and that *Smo* mutations seem to cause MB in one step, many early reports suggested that Hh pathway activation (through *Smo* activation or complete *Ptch1* loss) may be sufficient to drive MB formation.

However, several lines of reasoning suggested that the ‘two-hit’ model of MB formation might be incorrect. If *Ptch1* haploinsufficiency leads to preneoplasia formation, why doesn’t the whole EGL of the cerebellum become preneoplastic? By definition, a preneoplastic lesion is a discrete clonal expansion and, therefore, the cells contained within it must carry an additional genetic or epigenetic event responsible for its growth (an event not present in all other *Ptch1*^{+/-} GCPs that differentiated normally into granule neurons). Similarly, while most *Ptch1*^{+/-} mice develop various discrete preneoplastic lesions³⁸, not all of them form an advanced MB. In fact, the incidence of preneoplastic lesions is highest at P14 and decreases progressively during subsequent weeks until approaching the incidence of advanced MB³⁸. Indeed, a longitudinal MRI study of precancerous lesions resulting from *Ptch1* deletion demonstrated that some preneoplastic lesions regress

and do not become MB³⁹. In line with this reasoning, we predicted that 1) preneoplastic lesions in *Ptch1*^{+/-} mice are not *Ptch1* haploinsufficient, 2) unidentified tumor-suppressive mechanisms operate at the preneoplastic stage and limit their progression to advanced MB, and 3) additional genetic or epigenetic events regulate the progression from precancerous lesions to MB.

Cell senescence is a tumor-suppressive mechanism in SHH-MB

In addition to the two hits discussed above (*Ptch1* heterozygosity and *Ptch1* LOH), we identified additional events (‘third hits’) involved in MB formation: spontaneous *p53* mutations or *Cdkn2a* inactivation. We showed that preneoplastic lesions are the result of *Ptch1* LOH and not *Ptch1* haploinsufficiency⁴⁰. Moreover, we showed that preneoplastic lesions become senescent, therefore identifying a novel tumor suppressor mechanism for MB operating at the preneoplastic stage. Altogether, this work shows that the previous two-hit models were incomplete, since they overlooked essential components in the process of MB formation. In other words, these results demonstrate that MB molecular evolution is more complex than was previously imagined⁴¹. Additional factors support a role for cell senescence or other cooperating mechanisms in MB progression. For example, the Cdk inhibitor p27^{Kip1} regulates the rate of MB formation in *SmoA1* mice without affecting tumor incidence^{42,43}, indicating that p27^{Kip1} is an additional regulator of MB progression. In some cases, senescence responses in MB result from cell stress. Citron kinase inactivation has been shown to induce cytokinesis failure, cell death, and cell senescence in *SmoA1* MB, as demonstrated by increased p27^{Kip1}, p21, and p16^{ink4a} (*Cdkn2a*) levels⁴⁴. ATR deletion also reduces MB formation upon constitutive *Smo* activation⁴⁵.

The fact that other events and mechanisms are necessary for MB formation in addition to Hh pathway activation can change the way new MB therapies are imagined and orient research in new directions. For example, most early efforts in designing therapies were aimed at inhibiting SMO or GLI transcription factors^{46,47}. The fact that new mechanisms are at play means that additional tumor mechanisms could be targeted or exploited in order to treat MB or prevent its progression.

SHH-MB age of onset and cell senescence

Different triggers of cell senescence exist. While oncogene-induced cell senescence (OIS) is a form of senescence induced by oncogenic stress that requires and is mediated by activation of the DNA damage response (DDR)⁴⁸, proliferative or oncogenic signals can also induce telomere attrition and replicative senescence (RS)⁴⁹. This implies that different types of senescence require different mechanisms of senescence bypass. It is commonly accepted that mutations in DDR genes such as *TP53* or *ATM* evade OIS⁵⁰, while telomerase (*TERT* promoter) mutations restore telomere function and bypass RS⁵¹.

In terms of the mutation landscape, adult SHH-MBs are different compared to infant and child SHH-MBs⁵². Virtually all adult SHH-MBs with *PTCH1* mutations display *TERT* promoter mutations, while they are absent from infant/child SHH-MBs with *PTCH1* mutations²⁹. This strongly suggests that adult SHH-MBs undergo RS, consistent with the idea that those tumors develop over a longer period of time (Figure 2); this idea is also supported by the higher mutational load of adult SHH-MBs^{52,53}, indicative of longer tumor latency. In contrast, SHH-MBs in children are enriched in *TP53* mutations¹⁷, consistent with the idea that SHH-MBs in younger patients undergo OIS.

The existence of cell senescence in MB preneoplasia has potential implications and opens new avenues of research.

Although in our experiments cell senescence operates as a tumor-suppressive mechanism, the functions and potential roles of senescent cells in SHH-MB are presently unknown. Whether preneoplastic senescent cells contribute or respond to environmental changes in other preneoplastic cells is not known; additionally, whether senescent and resident stem-like cells interact remains an open question. These questions are discussed below (Figure 3).

Emerging players in MB progression

Tumor microenvironment

Recent work has investigated the role of the tumor microenvironment in MB progression. Brain tumors are composed of not only tumor and stem-like cells but also tumor-associated components (stroma) such as vascular and immune cells, astrocytes, microglia, and extracellular matrix⁵⁴. This has potential clinical implications, since tumor-associated cells could be the focus of targeted therapies.

One of the key features distinguishing SHH-MB among other MB groups is its frequent desmoplastic histology⁸. Desmoplasia is defined as the growth of fibrous or connective tissues and therefore reflects the presence of tumor stroma. Surprisingly, little is known about the causes of this tumor histological phenotype in SHH-MB. However, some interesting insights can be drawn from the study of other cancer types in which Shh signaling induces desmoplasia, such as prostate and pancreatic cancers⁵⁵. In these epithelial cancers, tumor cells produce Shh, which signals to the stroma and promotes tumor growth^{56,57}. These studies highlighted the crosstalk mediated by Shh between cancer cells and other cell components within tumors.

A recent report studied gene expression signatures and designed algorithms to extrapolate the relative contribution of cells from the tumor microenvironment in groups of MB and relative to

Figure 2. Different cell senescence mechanisms could explain tumor latency differences in SHH medulloblastomas (SHH-MBs). While *TP53* mutations are enriched in child SHH-MB, *TERT* promoter mutations exclusively affect SHH-MB diagnosed in adults. We speculate that different cell senescence mechanisms underlie differences in MB latency. One possibility is that slow-growing precancerous lesions undergo replicative senescence (RS) that needs to be evaded by *TERT* promoter mutations in adult cases. In children, oncogene activation causes fast proliferation and oncogene-induced senescence (OIS), driving the acquisition of *TP53* mutations or *p16^{ink4}* inactivation.

Figure 3. Medulloblastoma formation in *Ptch1*^{+/-} mice and components of the tumor microenvironment. Besides cancer cells, preneoplastic lesions and advanced medulloblastoma contain multiple cell types that form the tumor stroma. Preneoplasia contains senescent cells and recruits the formation of blood vessels and immune cells, including macrophages. The cell types composing this tumor microenvironment regulate preneoplasia progression to advanced tumors. GCP, granule cell precursor.

other brain tumor entities. This study found that MBs display low expression of immune markers (including PD1, a cell-surface immune checkpoint receptor) compared to other brain tumors⁵⁸. Additionally, it was found that SHH-MBs display signatures predictive of fibroblast, T cell, and macrophage infiltration⁵⁸, findings corroborated by flow cytometry studies⁵⁹. Interestingly, studies are starting to show functional roles for these cell types in MB tumorigenesis and are summarized in this section.

Macrophages are the most abundant type of immune cells in brain tumors⁵⁴. Among the MB groups, SHH-MB displays the highest number of macrophages⁶⁰. Tumor-associated macrophages (TAMs) have been shown to promote or restrain tumor growth depending on the context. In astrocytomas, TAMs have been shown to release cytokines that facilitate tumor cell proliferation and survival and therefore display tumor-promoting roles⁶¹. Two kinds of macrophages, M1 and M2, have traditionally been described. A first report indicated that human SHH-MBs displaying high numbers of M1 macrophages display worse prognosis⁶². However, a recent study in mice using genetic and cell biology approaches demonstrated that TAMs in Shh-MB caused by SmoA1 activation display tumor cell killing properties and therefore have a tumor suppressive role⁶³; TAMs in Shh-MB corresponded predominantly to bone marrow-derived monocytes that invaded the MB tissue during tumor growth⁶³.

Additional studies will be required to determine specific functional roles of macrophage subtypes in MB as well as the relative contribution of microglia to tumor formation.

T lymphocytes also infiltrate primary MB⁶⁴. In orthotopic transplantation experiments, it was shown that MB cells trigger the secretion of the T cell chemoattractant RANTES from the endothelium, leading to T cell migration⁶⁴. Other studies using transgenic mouse models have shown that Shh-MB tumor cells produce TGF- β , which leads to reduced CD8 T lymphocyte expansion and activation, limiting anti-tumor cytotoxic activity⁶⁵. MB-bearing mice also display low numbers of peripheral T cells and smaller spleens, indicating that MB affects immune function systemically⁶⁵. Consistent with these findings, high numbers of T helper lymphocytes (TH1) as well as high levels of IFN γ and TNF α correlate with good prognosis in human MB patients undergoing chemotherapy and hematopoietic stem cell transplantation⁶⁶. While these findings suggest that MB cells modulate and perhaps inhibit T cell-dependent responses, further work is required to determine the specific functions of T cell subpopulations and NK (natural killer) cells in MB pathogenesis and whether they could be exploited for therapies.

Astrocytes are abundant in human SHH-MBs and in various mouse models of Shh-MB. Tumor astrocyte-derived Shh

was shown to induce the proliferation of advanced MB tumor cells from *Atoh1-Cre; Ptch1^{fl/fl}* mice, preventing their differentiation and leading to increased tumor growth⁶⁷. In this case, it seems that tumor cells exploit the normal Shh-secreting function of astrocytes to their advantage. It is unknown whether other astrocyte functions, such as maintaining tissue homeostasis, also play a role in MB tumorigenesis. Additionally, whether tumor-associated astrocytes are recruited from other central nervous system (CNS) regions and whether they are related to Bergmann glia or are perhaps derived from MB stem cells remain to be determined.

Role of the vasculature and extracellular matrix

The presence of the blood brain barrier (BBB) distinguishes the vasculature of the CNS. Composed of endothelial cells, pericytes, and astrocyte processes, the BBB isolates the brain parenchyma from toxic circulating compounds, including chemotherapeutics³⁴. In contrast to WNT-MB, SHH-MB displays a BBB, which seems to be induced by tumor cells⁶⁸. This process is controlled by Wnt signaling in vascular endothelial cells. Importantly, it has been shown that Norrin/Frizzled signaling regulates angiogenic remodeling and acts as a tumor-suppressive mechanism; in the absence of Fzd4 signaling, angiogenic changes led to increased cell proliferation, reduced apoptosis, and increased rates of *Ptch1* LOH and progression to advanced tumors in *Ptch1*^{+/-} mice⁶⁹.

In vitro studies of endothelial cells show that Shh promotes blood vessel formation via cytoskeletal remodeling⁷⁰. In the healthy brain, Hh signaling promotes BBB formation and integrity: astrocyte-derived Shh signals to endothelial cells, where it decreases trans-endothelial permeability and promotes immune quiescence by decreasing inflammatory mediators and leukocyte migration⁷¹. This role of Shh perhaps explains the relatively low levels of inflammatory markers in SHH-MB compared to other molecular MB groups⁵⁸.

SHH-MB also displays expression of coagulation and angiogenesis signaling genes compared to other MB groups. Tissue factor and protease-activated receptor 1 (thrombin receptor), essential coagulation and angiogenesis factors, are upregulated in SHH-MB⁷². It has been shown that the extracellular protease Serpine2 regulates MB progression in *Ptch1* mice by maintaining preneoplastic lesions in a proliferative state⁷³, although the mechanism is unknown. Finally, MB therapies also have an impact on tumor vasculature. For example, irradiation induces angiogenesis in a matrix metalloproteinase 9-dependent manner in MB⁷⁴.

An interesting avenue of research is to determine the potential cell-cell interactions in the perivascular niche and the cross-talk between vascular structures and MB stem cells. MB stem cells (Nestin⁺, Prominin⁺) are closely associated with capillaries in the perivascular niche, where endothelial cells from the tumor vasculature secrete factors that promote stem cell self-renewal and propagation *in vitro* and *in vivo*⁷⁵. The Akt/Pten signaling pathway controls self-renewal of neural stem cells (NSCs)⁷⁶. Additionally, NSCs of the perivascular niche are resistant to radiation therapy, a response mediated by

activation of the Akt/mTOR pathway⁷⁷ and which might also involve YAP1⁷⁸. Pten inactivation in Nestin⁺ cells in early post-natal periods causes the expansion of perivascular cells and, in combination with *p53* inactivation, leads to Shh-MB⁷⁹. Together, these studies highlight a critical role for the perivascular niche in MB initiation, growth, and resistance. The facts that Wnt signaling is essential for stem cell function and that Wnt ligands are produced by endothelial cells might explain why MB stem cells reside in the perivascular niche.

Microenvironment invasion and metastasis

Three essential steps regulate metastasis: initiation, dispersal, and colonization. MBs most often display meningeal metastases, a process called leptomeningeal dissemination⁸⁰. While the cerebrospinal fluid may be the most favorable route of dissemination in MB, a hematogenous route has been demonstrated for MB⁸¹. Hepatocyte growth factor (HGF), a potent inducer of epithelial-to-mesenchymal transition (EMT), induces aggressive MB in collaboration with Shh. The chemokine CCL2 and its receptor CCR2 were expressed at higher levels in MB metastases compared to primary MBs. Expression of either CCL2 or CCR2 dramatically increased leptomeningeal metastasis in a model of Shh-MB⁸¹. The kinase PERK has been shown to enhance MB cell migration and invasion *in vitro* in a VEGF signaling-dependent manner⁸². These two studies show that regulators of the tissue microenvironment can contribute to metastasis in MB.

Cell senescence, inflammation, and stem cells

In other cancers or tissue contexts, an interesting link between senescence and cell reprogramming has been discovered⁸³. Transient exposure to senescence-associated secretory phenotype (SASP) factors, chemokines produced by senescent cells, can induce the expression of stem cell genes (such as *Nestin*, *Lgr6*, and *CD34* and cell plasticity in keratinocytes and pancreatic epithelial cells⁸⁴). This response promotes tissue regeneration. Conversely, *in vivo* expression of reprogramming factors induces both senescence and reprogramming; tissue damage and the ability to undergo senescence foster the reprogramming potential⁸⁵. Moreover, cells induced to senesce exhibit plasticity and a greater tumor-initiating potential when they escape senescence than cells that were never senescent⁸⁶. These results raise the possibility that senescent and inflammatory cells found within MBs could impact MB tumor-initiating cells (stem cells). In this context, increased Wnt signaling in senescent cells due to epigenetic changes is the main regulator of stemness⁸⁶. Given that the Wnt pathway is a major physiological regulator of stem cell self-renewal during development in many different tissue contexts such as intestinal crypts, hematopoietic stem cells, hair follicle buds, and neuroepithelial cells^{87,88}, this suggests that senescent cancer cells might co-opt the Wnt pathway to reactivate stemness and evade senescence⁸³. This phenomenon of Wnt re-activation has been documented in cases where progenitor cells (not stem cells) become cancer cells upon oncogene activation⁸⁸.

Several stem cell/progenitor populations have been described in SHH-MB. Besides the EGL containing GCPs, the cerebellum displays a ventricular zone containing NSCs that

give origin to all cerebellar cell types other than glutamatergic neurons⁸⁹. The classical marker expressed by NSCs is the intermediate filament Nestin. Additionally, NSCs are positive for GFAP and subsets of them seem to express Sox2 and Olig2, and constitutive Hh activation in this population causes SHH-MB that first transit through a GCP cell fate^{33,90}. The early postnatal cerebellum also contains a population of Proliferating (CD133)-positive stem cells⁹¹ located close to the white matter that seem to be Nestin⁺. Deletion of *p53* and *Retinoblastoma* in these Proliferating⁺ stem cells leads to MB⁹², although with low efficiency. Moreover, a subset of Atoh1⁺ and CD133⁻ GCPs express high levels of the glycoprotein CD15 and have been shown to be tumor-propagating cells in *Ptch1*^{+/-} MB^{93,94}. Together, these results indicate that ventricular zone NSCs and GCPs from the EGL are the main cells with the potential to give origin to SHH-MB. Additionally, a subpopulation of Nestin⁺ GCPs located in the inner layer of the EGL has been shown to display increased tumorigenic potential compared to Atoh1⁺ GCPs⁹⁵; these cells are Shh responsive, and it was later proposed that Nestin has the ability to bind Gli3 and prevent its conversion to a repressive form, leading to higher Hh pathway activity⁹⁶.

Furthermore, a population of SHH-MB tumor-propagating cells expressing Sox2⁺ was described as the source of MB repopulating cells subsequent to therapy⁹⁷. While Sox2⁺ cells were insensitive to Hh pathway inhibition, they displayed Wnt pathway activation, and modulation of Wnt signaling regulated their tumor-propagating abilities⁹⁸. In this way, Sox2⁺ MB cancer cells are similar to other cancer stem cells and NSCs of the cerebellar ventricular zone, which proliferate in response to Wnt signaling⁹⁹. Although Sox2⁺ MB cells are tumor-propagating cells present in primary Shh-MBs, their origin and whether they are derived from ventricular NSCs are unknown. Because Bergmann glia express Sox2, this cell type is a potential source of Sox2⁺ MB cancer-propagating cells.

Although insensitive to Hh pathway inhibition, Sox2⁺ cells seem to be specific to SHH-MB; moreover, activation of Hh signaling genes (such as *SmoM2* mutation) increased the number of Sox2⁺ cells in the cerebellum⁹⁰, suggesting that some Hh-dependent process underlies the generation of Sox2⁺ MB cells. The polycomb protein Bmi1 is a key regulator of stem cell function. Hh signaling activation induces Bmi1 expression¹⁰⁰ and Bmi1 is required for normal GCP expansion¹⁰¹. While Bmi1 overexpression is insufficient to cause MB¹⁰², its deletion

completely prevents the formation of MB in the *SmoA1* model¹⁰³; this effect was due to impaired progression of tumor lesions (preneoplasia), which displayed increased numbers of apoptotic and p19^{Arf} cells and reduced proliferation. Whether Bmi1 is expressed or plays a role in Sox2 MB cells is unknown, but it is interesting to speculate that Hh signaling-dependent regulation of Bmi1 and other potential stem cell regulators could be the source of MB Sox2⁺ cells.

Conclusion

Because of the histological similarities between tumors and stromal components undergoing tissue remodeling, pathologists since the times of Virchow have traditionally described tumors as “wounds that do not heal”¹⁰⁴; the recent recognition of the multiple roles of the tumor microenvironment in cancer confirms this idea and enables new potential ways of understanding tumors. Relative to other cancers, little is known about how the tumor microenvironment regulates MB formation. However, the identification of critical steps regulating tumor formation, such as progression, offers an interesting framework to address how the recently identified tumor cell components regulate MB formation.

Among these microenvironment cell components, clear roles are currently being established for some cell types. Macrophages seem to have a tumor-suppressive role in Shh-MB, although the specific function of macrophage subtypes, and the relative contribution of microglia, needs better understanding. Less is known about the roles of lymphocytes, although initial studies suggest that MB cells inhibit T-cell function. Besides immune cells, glia play an important role in MB formation in which astrocytes secrete Shh, promoting MB cell proliferation.

Tumor and stromal cells are regulated by signaling molecules in different ways. Shh promotes the formation and maintains the integrity of the BBB, therefore influencing factors like vascular permeability and immune cell infiltration. Wnt signaling has also been shown to regulate the formation of the BBB. The interplay between Hh and Wnt signaling in the perivascular niche, and the potential ways it regulates perivascular tumor-propagating cells, is an exciting topic of research. Finally, important research has been carried out showing cell populations other than GCPs with tumor-initiating or tumor-propagating abilities. How these cell populations arise within the tumor, whether they are sensitive to inflammatory signals in the microenvironment, and how they respond to Hh and Wnt signaling are current topics of MB research.

References

1. Fearon ER, Vogelstein B: **A genetic model for colorectal tumorigenesis.** *Cell.* 1990; **61**(5): 759-67.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Vogelstein B, Kinzler KW: **The multistep nature of cancer.** *Trends Genet.* 1993; **9**(4): 138-41.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Frank SA: **Dynamics of Cancer: Incidence, Inheritance, and Evolution.** (Princeton (NJ)). 2007.
[PubMed Abstract](#)
4. Knudson AG: **Two genetic hits (more or less) to cancer.** *Nat Rev Cancer.* 2001; **1**(2): 157-62.
[PubMed Abstract](#) | [Publisher Full Text](#)

5. Vogelstein B, Fearon ER, Hamilton SR, *et al.*: Genetic alterations during colorectal-tumor development. *N Engl J Med.* 1988; 319(9): 525-32.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Weinberg RA: **The biology of cancer.** Second edition. edn (New York: Garland Science, Taylor & Francis Group). 2014.
[Reference Source](#)
7. Sturm D, Orr BA, Toprak UH, *et al.*: New Brain Tumor Entities Emerge from Molecular Classification of CNS-PNETs. *Cell.* 2016; 164(5): 1060-72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. **F** Kool M, Korshunov A, Remke M, *et al.*: Molecular subgroups of medulloblastoma: An international meta-analysis of transcriptome, genetic aberrations, and clinical data of WNT, SHH, Group 3, and Group 4 medulloblastomas. *Acta Neuropathol.* 2012; 123(4): 473-84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
9. **F** Taylor MD, Northcott PA, Korshunov A, *et al.*: Molecular subgroups of medulloblastoma: the current consensus. *Acta Neuropathol.* 2012; 123(4): 465-72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
10. **F** Hovestadt V, Jones DT, Picelli S, *et al.*: Decoding the regulatory landscape of medulloblastoma using DNA methylation sequencing. *Nature.* 2014; 510(7506): 537-41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
11. Kool M, Koster J, Bunt J, *et al.*: Integrated genomics identifies five medulloblastoma subtypes with distinct genetic profiles, pathway signatures and clinicopathological features. *PLoS One.* 2008; 3(8): e3088.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Northcott PA, Jones DT, Kool M, *et al.*: Medulloblastomics: the end of the beginning. *Nat Rev Cancer.* 2012; 12(12): 818-34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. **F** Northcott PA, Shih DJ, Peacock J, *et al.*: Subgroup-specific structural variation across 1,000 medulloblastoma genomes. *Nature.* 2012; 488(7409): 49-56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
14. **F** Pugh TJ, Weeraratne SD, Archer TC, *et al.*: Medulloblastoma exome sequencing uncovers subtype-specific somatic mutations. *Nature.* 2012; 488(7409): 106-10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
15. Gibson P, Tong Y, Robinson G, *et al.*: Subtypes of medulloblastoma have distinct developmental origins. *Nature.* 2010; 468(7327): 1095-9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. **F** Vladoiu MC, El-Hamamy I, Donovan LK, *et al.*: Childhood cerebellar tumours mirror conserved fetal transcriptional programs. *Nature.* 2019; 572(7767): 67-73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
17. Cavalli FMG, Remke M, Rampasek L, *et al.*: Intertumoral Heterogeneity within Medulloblastoma Subgroups. *Cancer Cell.* 2017; 31(6): 737-754.e6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. **F** Azevedo FA, Carvalho LR, Grinberg LT, *et al.*: Equal numbers of neuronal and nonneuronal cells make the human brain an isometrically scaled-up primate brain. *J Comp Neurol.* 2009; 513(5): 532-41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
19. Hatten ME, Heintz N: Mechanisms of neural patterning and specification in the developing cerebellum. *Annu Rev Neurosci.* 1995; 18: 385-408.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. Sillitoe RV, Joyner AL: Morphology, molecular codes, and circuitry produce the three-dimensional complexity of the cerebellum. *Annu Rev Cell Dev Biol.* 2007; 23: 549-77.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Dahmane N, Ruiz i Altaba A: Sonic hedgehog regulates the growth and patterning of the cerebellum. *Development.* 1999; 126(14): 3089-100.
[PubMed Abstract](#)
22. Wallace VA: Purkinje-cell-derived Sonic hedgehog regulates granule neuron precursor cell proliferation in the developing mouse cerebellum. *Curr Biol.* 1999; 9(8): 445-8.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Wechsler-Reya RJ, Scott MP: Control of neuronal precursor proliferation in the cerebellum by Sonic Hedgehog. *Neuron.* 1999; 22(1): 103-14.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Jiang J, Hui CC: Hedgehog signaling in development and cancer. *Dev Cell.* 2008; 15(6): 801-12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Briscoe J, Théron PP: The mechanisms of Hedgehog signalling and its roles in development and disease. *Nat Rev Mol Cell Biol.* 2013; 14(7): 416-29.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Kenney AM, Cole MD, Rowitch DH: *Nmyc* upregulation by sonic hedgehog signaling promotes proliferation in developing cerebellar granule neuron precursors. *Development.* 2003; 130(1): 15-28.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Kenney AM, Rowitch DH: Sonic hedgehog promotes G₁ cyclin expression and sustained cell cycle progression in mammalian neuronal precursors. *Mol Cell Biol.* 2000; 20(3): 9055-67.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Oliver TG, Grasdeder LL, Carroll AL, *et al.*: Transcriptional profiling of the Sonic hedgehog response: A critical role for N-myc in proliferation of neuronal precursors. *Proc Natl Acad Sci U S A.* 2003; 100(12): 7331-6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Northcott PA, Buchhalter I, Morrissy AS, *et al.*: The whole-genome landscape of medulloblastoma subtypes. *Nature.* 2017; 547(7663): 311-317.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Goodrich LV, Milenkovic L, Higgins KM, *et al.*: Altered neural cell fates and medulloblastoma in mouse patched mutants. *Science.* 1997; 277(5329): 1109-13.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Mao J, Ligon KL, Rakhlin EY, *et al.*: A novel somatic mouse model to survey tumorigenic potential applied to the Hedgehog pathway. *Cancer Res.* 2006; 66(20): 10171-8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. **F** Oliver TG, Read TA, Kessler JD, *et al.*: Loss of *patched* and disruption of granule cell development in a pre-neoplastic stage of medulloblastoma. *Development.* 2005; 132(10): 2425-39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
33. Schüller U, Heine VM, Mao J, *et al.*: Acquisition of granule neuron precursor identity is a critical determinant of progenitor cell competence to form Shh-induced medulloblastoma. *Cancer Cell.* 2008; 14(2): 123-34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Yang ZJ, Ellis T, Markant SL, *et al.*: Medulloblastoma can be initiated by deletion of *Patched* in lineage-restricted progenitors or stem cells. *Cancer Cell.* 2008; 14(2): 135-45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. Rausch T, Jones DT, Zapotka M, *et al.*: Genome sequencing of pediatric medulloblastoma links catastrophic DNA rearrangements with *TP53* mutations. *Cell.* 2012; 148(1-2): 59-71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Bayer SA, Altman J, Russo RJ, *et al.*: Timetables of neurogenesis in the human brain based on experimentally determined patterns in the rat. *Neurotoxicology.* 1993; 14(1): 83-144.
[PubMed Abstract](#)
37. Ayrault O, Zindy F, Reh J, *et al.*: Two tumor suppressors, *p27^{Kip1}* and *patched-1*, collaborate to prevent medulloblastoma. *Mol Cancer Res.* 2009; 7(1): 33-40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Pazzaglia S, Tanori M, Mancuso M, *et al.*: Two-hit model for progression of medulloblastoma preneoplasia in Patched heterozygous mice. *Oncogene.* 2006; 25(40): 5575-80.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Suero-Abreu GA, Praveen Raju G, Aristizábal O, *et al.*: *In vivo* Mn-enhanced MRI for early tumor detection and growth rate analysis in a mouse medulloblastoma model. *Neoplasia.* 2014; 16(12): 993-1006.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Tamayo-Orrego L, Wu CL, Bouchard N, *et al.*: Evasion of Cell Senescence Leads to Medulloblastoma Progression. *Cell Rep.* 2016; 14(12): 2925-37.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Tamayo-Orrego L, Swikert SM, Charron F: Evasion of cell senescence in SHH medulloblastoma. *Cell Cycle.* 2016; 15(16): 2102-2107.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Bhatia B, Malik A, Fernandez-L A, *et al.*: *p27^{Kip1}*, a double-edged sword in Shh-mediated medulloblastoma: Tumor accelerator and suppressor. *Cell Cycle.* 2010; 9(21): 4307-14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. Hatton BA, Ellison DW, Gajjar A, *et al.*: Prognostic value and functional consequences of cell cycle inhibitor *p27^{Kip1}* loss in medulloblastoma. *Biomark Res.* 2013; 1(1): 14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
44. Pallavicini G, Sgrò F, Garello F, *et al.*: Inactivation of Citron Kinase Inhibits Medulloblastoma Progression by Inducing Apoptosis and Cell Senescence. *Cancer Res.* 2018; 78(16): 4599-4612.
[PubMed Abstract](#) | [Publisher Full Text](#)
45. Lang PY, Nanjangud GJ, Sokolsky-Papkov M, *et al.*: ATR maintains chromosomal integrity during postnatal cerebellar neurogenesis and is required for medulloblastoma formation. *Development.* 2016; 143(21): 4038-4052.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. Kim J, Aftab BT, Tang JY, *et al.*: Itraconazole and arsenic trioxide inhibit Hedgehog pathway activation and tumor growth associated with acquired resistance to smoothened antagonists. *Cancer Cell.* 2013; 23(1): 23-34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
47. **F** Romer JT, Kimura H, Magdaleno S, *et al.*: Suppression of the Shh pathway using a small molecule inhibitor eliminates medulloblastoma in *Ptc1^{flx}p53^{flx}* mice. *Cancer Cell.* 2004; 6(3): 229-40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
48. **F** Di Micco R, Fumagalli M, Cicalese A, *et al.*: Oncogene-induced senescence is a DNA damage response triggered by DNA hyper-replication. *Nature.* 2006; 444(7119): 638-42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

49. Campisi J: **Aging, cellular senescence, and cancer.** *Annu Rev Physiol.* 2013; **75**: 685–705.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Negrini S, Gorgoulis VG, Halazonetis TD: **Genomic instability—an evolving hallmark of cancer.** *Nat Rev Mol Cell Biol.* 2010; **11**(3): 220–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
51. Shay JW: **Role of Telomeres and Telomerase in Aging and Cancer.** *Cancer Discov.* 2016; **6**(6): 584–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
52. Kool M, Jones DTW, Jäger N, et al.: **Genome sequencing of SHH medulloblastoma predicts genotype-related response to smoothened inhibition.** *Cancer Cell.* 2014; **25**(3): 393–405.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
53. Jones DT, Jäger N, Kool M, et al.: **Dissecting the genomic complexity underlying medulloblastoma.** *Nature.* 2012; **488**(7409): 100–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
54. Quail DF, Joyce JA: **The Microenvironmental Landscape of Brain Tumors.** *Cancer Cell.* 2017; **31**(3): 326–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
55. Bailey JM, Swanson BJ, Hamada T, et al.: **Sonic hedgehog promotes desmoplasia in pancreatic cancer.** *Clin Cancer Res.* 2008; **14**(19): 5995–6004.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. Fan L, Pepicelli CV, Dibble CC, et al.: **Hedgehog signaling promotes prostate xenograft tumor growth.** *Endocrinology.* 2004; **145**(8): 3961–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
57. Yauch RL, Gould SE, Scales SJ, et al.: **A paracrine requirement for hedgehog signalling in cancer.** *Nature.* 2008; **455**(7211): 406–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
58. Bockmayr M, Mohme M, Klauschen F, et al.: **Subgroup-specific immune and stromal microenvironment in medulloblastoma.** *Oncoimmunology.* 2018; **7**(9): e1462430.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
59. Pham CD, Flores C, Yang C, et al.: **Differential Immune Microenvironments and Response to Immune Checkpoint Blockade among Molecular Subtypes of Murine Medulloblastoma.** *Clin Cancer Res.* 2016; **22**(3): 582–95.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
60. Margol AS, Robison NJ, Gnanachandran J, et al.: **Tumor-associated macrophages in SHH subgroup of medulloblastomas.** *Clin Cancer Res.* 2015; **21**(6): 1457–65.
[PubMed Abstract](#) | [Publisher Full Text](#)
61. Hambardzumyan D, Gutmann DH, Kettenmann H: **The role of microglia and macrophages in glioma maintenance and progression.** *Nat Neurosci.* 2016; **19**(1): 20–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
62. Lee C, Lee J, Choi SA, et al.: **M1 macrophage recruitment correlates with worse outcome in SHH Medulloblastomas.** *BMC Cancer.* 2018; **18**(1): 535.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
63. **F** Maximov V, Chen Z, Wei Y, et al.: **Tumour-associated macrophages exhibit anti-tumoural properties in Sonic Hedgehog medulloblastoma.** *Nat Commun.* 2019; **10**(1): 2410.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
64. Salsman VS, Chow KK, Shaffer DR, et al.: **Crosstalk between medulloblastoma cells and endothelium triggers a strong chemotactic signal recruiting T lymphocytes to the tumor microenvironment.** *PLoS One.* 2011; **6**(5): e20267.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
65. Gate D, Danielpour M, Rodriguez J, et al.: **T-cell TGF- β signaling abrogation restricts medulloblastoma progression.** *Proc Natl Acad Sci U S A.* 2014; **111**(33): E3458–E3466.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
66. Wiegner V, Eyrych M, Rutkowski S, et al.: **TH1 predominance is associated with improved survival in pediatric medulloblastoma patients.** *Cancer Immunol Immunother.* 2011; **60**(5): 693–703.
[PubMed Abstract](#) | [Publisher Full Text](#)
67. **F** Liu Y, Yuelling LW, Wang Y, et al.: **Astrocytes Promote Medulloblastoma Progression through Hedgehog Secretion.** *Cancer Res.* 2017; **77**(23): 6692–703.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
68. Phoenix TN, Patmore DM, Boop S, et al.: **Medulloblastoma Genotype Dictates Blood Brain Barrier Phenotype.** *Cancer Cell.* 2016; **29**(4): 508–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
69. Bassett EA, Tokarew N, Allemanno EA, et al.: **Norrin/FRizzled4 signalling in the preneoplastic niche blocks medulloblastoma initiation.** *eLife.* 2016; **5**: pii: e16764.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
70. Chinchilla P, Xiao L, Kazanietz MG, et al.: **Hedgehog proteins activate pro-angiogenic responses in endothelial cells through non-canonical signaling pathways.** *Cell Cycle.* 2010; **9**(3): 570–79.
[PubMed Abstract](#) | [Publisher Full Text](#)
71. **F** Alvarez JI, Dodelet-Devillers A, Kebir H, et al.: **The Hedgehog pathway promotes blood-brain barrier integrity and CNS immune quiescence.** *Science.* 2011; **334**(6063): 1727–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
72. D'Asti E, Kool M, Pfister SM, et al.: **Coagulation and angiogenic gene expression profiles are defined by molecular subgroups of medulloblastoma: evidence for growth factor-thrombin cross-talk.** *J Thromb Haemost.* 2014; **12**(11): 1838–49.
[PubMed Abstract](#) | [Publisher Full Text](#)
73. Vaillant C, Valdivieso P, Nuciforo S, et al.: **Serpine2/PN-1 Is Required for Proliferative Expansion of Pre-Neoplastic Lesions and Malignant Progression to Medulloblastoma.** *PLoS One.* 2015; **10**(4): e0124870.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
74. Asuthkar S, Velpula KK, Nalla AK, et al.: **Irradiation-induced angiogenesis is associated with an MMP-9-miR-494-syndecan-1 regulatory loop in medulloblastoma cells.** *Oncogene.* 2014; **33**(15): 1922–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
75. **F** Calabrese C, Poppleton H, Kocak M, et al.: **A perivascular niche for brain tumor stem cells.** *Cancer Cell.* 2007; **11**(1): 69–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
76. Groszer M, Erickson R, Scripture-Adams DD, et al.: **PTEN negatively regulates neural stem cell self-renewal by modulating G0-G1 cell cycle entry.** *Proc Natl Acad Sci U S A.* 2006; **103**(1): 111–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
77. Hambardzumyan D, Becher OJ, Rosenblum MK, et al.: **PI3K pathway regulates survival of cancer stem cells residing in the perivascular niche following radiation in medulloblastoma in vivo.** *Genes Dev.* 2008; **22**(4): 436–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
78. Fernandez-L A, Northcott PA, Dalton J, et al.: **YAP1 is amplified and up-regulated in hedgehog-associated medulloblastomas and mediates Sonic hedgehog-driven neural precursor proliferation.** *Genes Dev.* 2009; **23**(23): 2729–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
79. Zhu G, Rankin SL, Larson JD, et al.: **PTEN Signaling in the Postnatal Perivascular Progenitor Niche Drives Medulloblastoma Formation.** *Cancer Res.* 2017; **77**(1): 123–133.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
80. Fults DW, Taylor MD, Garzia L: **Leptomeningeal dissemination: a sinister pattern of medulloblastoma growth.** *J Neurosurg Pediatr.* 2019; **23**(5): 613–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
81. Garzia L, Kijima N, Morrissy AS, et al.: **A Hematogenous Route for Medulloblastoma Leptomeningeal Metastases.** *Cell.* 2018; **173**(6): 1549.
[PubMed Abstract](#) | [Publisher Full Text](#)
82. Jamison S, Lin Y, Lin W: **Pancreatic endoplasmic reticulum kinase activation promotes medulloblastoma cell migration and invasion through induction of vascular endothelial growth factor A.** *PLoS One.* 2015; **10**(3): e0120252.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
83. Lee S, Schmitt CA: **The dynamic nature of senescence in cancer.** *Nat Cell Biol.* 2019; **21**(1): 94–101.
[PubMed Abstract](#) | [Publisher Full Text](#)
84. Ritschka B, Storer M, Mas A, et al.: **The senescence-associated secretory phenotype induces cellular plasticity and tissue regeneration.** *Genes Dev.* 2017; **31**(2): 172–183.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
85. Mosteiro L, Pantoja C, Alcazar N, et al.: **Tissue damage and senescence provide critical signals for cellular reprogramming in vivo.** *Science.* 2016; **354**(6315): pii: aaf4445
[PubMed Abstract](#) | [Publisher Full Text](#)
86. **F** Milanovic M, Fan DNY, Belenki D, et al.: **Senescence-associated reprogramming promotes cancer stemness.** *Nature.* 2018; **553**(7686): 96–100.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
87. **F** Ouspenskaia T, Matos I, Mertz AF, et al.: **WNT-SHH Antagonism Specifies and Expands Stem Cells prior to Niche Formation.** *Cell.* 2016; **164**(1–2): 156–169.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
88. **F** Wang Y, Krivtsov AV, Sinha AU, et al.: **The Wnt/beta-catenin pathway is required for the development of leukemia stem cells in AML.** *Science.* 2010; **327**(5973): 1650–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
89. Wang J, Wechsler-Reya RJ: **The role of stem cells and progenitors in the genesis of medulloblastoma.** *Exp Neurol.* 2014; **260**: 69–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
90. Ahlfeld J, Favaro R, Pagella P, et al.: **Sox2 requirement in sonic hedgehog-associated medulloblastoma.** *Cancer Res.* 2013; **73**(12): 3796–807.
[PubMed Abstract](#) | [Publisher Full Text](#)
91. Lee A, Kessler JD, Read TA, et al.: **Isolation of neural stem cells from the postnatal cerebellum.** *Nat Neurosci.* 2005; **8**(6): 723–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
92. Sutter R, Shakhova O, Bhagat H, et al.: **Cerebellar stem cells act as medulloblastoma-initiating cells in a mouse model and a neural stem cell signature characterizes a subset of human medulloblastomas.** *Oncogene.* 2010; **29**(12): 1845–56.
[PubMed Abstract](#) | [Publisher Full Text](#)
93. Read T-A, Fogarty MP, Markant SL, et al.: **Identification of CD15 as a marker for tumor-propagating cells in a mouse model of medulloblastoma.** *Cancer Cell.* 2009; **15**(2): 135–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

94. Ward RJ, Lee L, Graham K, *et al.*: **Multipotent CD15⁺ cancer stem cells in patched-1-deficient mouse medulloblastoma.** *Cancer Res.* 2009; **69**(11): 4682–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
95. Li P, Du F, Yuelling LW, *et al.*: **A population of Nestin-expressing progenitors in the cerebellum exhibits increased tumorigenicity.** *Nat Neurosci.* 2013; **16**(12): 1737–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
96. Li P, Lee EH, Du F, *et al.*: **Nestin Mediates Hedgehog Pathway Tumorigenesis.** *Cancer Res.* 2016; **76**(18): 5573–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
97. Vanner RJ, Remke M, Gallo M, *et al.*: **Quiescent sox2⁺ cells drive hierarchical growth and relapse in sonic hedgehog subgroup medulloblastoma.** *Cancer Cell.* 2014; **26**(1): 33–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
98. Rodriguez-Blanco J, Pednekar L, Penas C, *et al.*: **Inhibition of WNT signaling attenuates self-renewal of SHH-subgroup medulloblastoma.** *Oncogene.* 2017; **36**(45): 6306–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
99. Pei Y, Brun SN, Markant SL, *et al.*: **WNT signaling increases proliferation and impairs differentiation of stem cells in the developing cerebellum.** *Development.* 2012; **139**(10): 1724–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
100. Wang X, Venugopal C, Manoranjan B, *et al.*: **Sonic hedgehog regulates Bmi1 in human medulloblastoma brain tumor-initiating cells.** *Oncogene.* 2012; **31**(2): 187–99.
[PubMed Abstract](#) | [Publisher Full Text](#)
101. Leung C, Lingbeek M, Shakhova O, *et al.*: **Bmi1 is essential for cerebellar development and is overexpressed in human medulloblastomas.** *Nature.* 2004; **428**(6980): 337–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
102. Behesti H, Bhagat H, Dubuc AM, *et al.*: **Bmi1 overexpression in the cerebellar granule cell lineage of mice affects cell proliferation and survival without initiating medulloblastoma formation.** *Dis Model Mech.* 2013; **6**(1): 49–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
103. Michael LE, Westerman BA, Ermilov AN, *et al.*: **Bmi1 is required for Hedgehog pathway-driven medulloblastoma expansion.** *Neoplasia.* 2008; **10**(12): 1343–9, 5p following 1349.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
104. Dvorak HF: **Tumors: Wounds that do not heal. Similarities between tumor stroma generation and wound healing.** *N Engl J Med.* 1986; **315**(26): 1650–9.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **James Manfredi**

Department of Oncological Sciences and Graduate School of Biomedical Sciences, Icahn School of Medicine at Mount Sinai, New York, USA

Competing Interests: No competing interests were disclosed.

2. **Timothy R Gershon**

Department of Neurology, University of North Carolina School of Medicine, North Carolina, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research