REVIEW

Recent advances in signal integration mechanisms in the unfolded protein response [version 1; peer review: 2 approved]

G. Elif Karagöz ¹, Tomás Aragón², Diego Acosta-Alvear ³

¹Max Perutz Labs Vienna, Medical University of Vienna, Vienna, Austria

²Department of Gene Therapy and Regulation of Gene Expression, University of Navarra, Pamplona, Spain

³Department of Molecular, Cellular, and Developmental Biology, University of California, Santa Barbara, Santa Barbara, Santa Barbara, CA, USA

V1 First published: 01 Nov 2019, 8(F1000 Faculty Rev):1840
<https://doi.org/10.12688/f1000research.19848.1>

Latest published: 01 Nov 2019, 8(F1000 Faculty Rev):1840
<https://doi.org/10.12688/f1000research.19848.1>

Abstract

Since its discovery more than 25 years ago, great progress has been made in our understanding of the unfolded protein response (UPR), a homeostatic mechanism that adjusts endoplasmic reticulum (ER) function to satisfy the physiological demands of the cell. However, if ER homeostasis is unattainable, the UPR switches to drive cell death to remove defective cells in an effort to protect the health of the organism. This functional dichotomy places the UPR at the crossroads of the adaptation versus apoptosis decision. Here, we focus on new developments in UPR signaling mechanisms, in the interconnectivity among the signaling pathways that make up the UPR in higher eukaryotes, and in the coordination between the UPR and other fundamental cellular processes.

Keywords

ER stress, unfolded protein response, signal transduction pathway interconnectivity

Open Peer Review

Approval Status

	1	2
version 1		
01 Nov 2019		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Neil Bulleid**, College of Medical Veterinary and Life Sciences, University of Glasgow, Glasgow, UK
2. **Jeffrey L. Brodsky**, University of Pittsburgh, Pittsburgh, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Diego Acosta-Alvear (daa@lifesci.ucsb.edu)

Author roles: Karagöz GE: Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; Aragón T: Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; Acosta-Alvear D: Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2019 Karagöz GE *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Karagöz GE, Aragón T and Acosta-Alvear D. **Recent advances in signal integration mechanisms in the unfolded protein response [version 1; peer review: 2 approved]** F1000Research 2019, 8(F1000 Faculty Rev):1840 <https://doi.org/10.12688/f1000research.19848.1>

First published: 01 Nov 2019, 8(F1000 Faculty Rev):1840 <https://doi.org/10.12688/f1000research.19848.1>

Introduction

The unfolded protein response (UPR) comprises a collection of evolutionarily conserved signaling pathways in eukaryotes that monitor endoplasmic reticulum (ER) functions. In its most fundamental form, the UPR maintains the health of secreted and transmembrane proteins. Most of these proteins, which comprise roughly one third of the proteome, translocate into the ER for their folding and maturation. Because these proteins establish communication networks among cells and between cells and the extracellular environment, the UPR is essential for normal cell and organismal physiology. Indeed, the UPR has long been known to be essential for the differentiation, development, and maintenance of professional secretory cells and secretory tissues^{1–7}. However, UPR functions are not restricted to these tissues, as all key metazoan UPR sensor/transducers are ubiquitously expressed and sustain diverse processes in different tissues^{8–15}.

ER-resident sensors that detect perturbations of ER composition and function, commonly known as “ER stress”, initiate the UPR (reviewed in 16). ER stress occurs when the folding or degradative capacities of the ER are exceeded (for example, upon expression of mutant proteins that cannot be properly folded or by incapacitation of ER quality-control systems by mutations or pathogens). The UPR transmits information about the ER status to the cell nucleus by inducing the expression of transcription factors^{17–21}. The UPR also exerts post-transcriptional control by regulating protein synthesis and mRNA stability^{22–24}. The coordination of transcriptional and post-transcriptional UPR mechanisms determines the cellular decision of whether to adapt or to die if ER stress is unmitigable.

Thousands of articles published to date (a PubMed search for “unfolded protein response” shows more than 7,800 articles, including more than 3,800 in the past five years alone) have revealed the inner workings of the UPR and have portrayed it as a complex network of interconnected signaling pathways. In this review, we discuss new developments in UPR signaling mechanisms, connections among metazoan UPR signaling pathways, and UPR signaling beyond the ER.

Signal transduction in the UPR: early models and new insights

Three ER–transmembrane stress sensors control the UPR: the kinase/endoribonuclease IRE1, the kinase PERK, and the membrane-tethered transcription factor ATF6 (reviewed in 16). Evolutionary conservation provided the first clues about specialization in UPR signal transduction, which perhaps arose with the advent of multicellularity. IRE1 is ubiquitously found from yeasts to metazoans, whereas PERK and ATF6 appear only in animals²⁵. With regard to ER stress detection, the classic view posits that IRE1, PERK, and ATF6 detect protein-folding perturbations in the ER lumen. Lately, all of them have also been suggested to sense imbalances in lipid composition of the ER membrane^{26–31}. It has increasingly been recognized that ER deficits leading to proteo- or lipo-toxicity are dealt with by the UPR through different regulatory mechanisms (reviewed in 32). These findings coalesce into a model in which UPR sensors work

together holistically by integrating information about distinct/diverse ER deficiencies to launch homeostatic or apoptotic programs.

IRE1 and PERK detect ER stress by direct binding of unfolded protein ligands in the ER lumen, and the reversible dissociation of IRE1 and PERK from the ER luminal chaperone BiP regulates their activation/deactivation dynamics^{33–41}. Even though the interaction of BiP with the UPR sensors is long known, other interactions between UPR sensors and ER chaperones have only recently been revealed. For example, the co-chaperone ERDJ4 specifically facilitates BiP’s interaction with IRE1³⁸, and the activities of both IRE1 and PERK are fine-tuned by various ER chaperones, providing tissue- and context-specific UPR regulation. The protein disulfide isomerase PDIA6 modulates the deactivation of IRE1 and PERK^{42,43}, and the ER chaperone HSP47 associates with IRE1 but not PERK to regulate its activity⁴⁴. By contrast, the ER luminal protein canopy homolog 2 (CNPY2) binds only to PERK during ER stress to regulate downstream signaling⁴⁵. Moreover, in skeletal muscle cells, the ER oxidoreductase calsequestrin binds IRE1 to inhibit its activity⁴⁶. These findings support a view wherein the signaling capacity of UPR sensors can be refined to accommodate specific outputs according to the specific needs of cells and tissues. Most likely, the aforementioned interactions of UPR sensors with chaperones also influence the temporal dynamics of UPR signaling activation and deactivation, thereby providing an additional level of regulatory control leading to the life versus death decision in cells^{47–50}. In this way, modulation of the UPR at the level of UPR sensor–chaperone interactions serves as an initial point for information integration to regulate complex downstream UPR outputs.

How ATF6, by comparison with IRE1 and PERK, detects ER stress is less well understood. The established model posits that ER stress leads to ATF6 export to the Golgi apparatus, where resident proteases cleave it to liberate a soluble transcription factor, ATF6N^{51,52}. Accumulating evidence suggests that ATF6 may act as a redox sensor or that it is coupled to one, as the reduction of intra- and inter-molecular disulfide bonds in its ER luminal domain appears to be required for trafficking to the Golgi apparatus⁵³. The recent observation of aberrant and attenuated ATF6 signaling in cells deficient in the oxidoreductase ERp18, which associates with ATF6 during ER stress, supports this view⁵⁴.

Self-association/dissociation of UPR sensors is another canonical feature of the UPR. ER stress–induced oligomerization of IRE1 and PERK in the plane of the ER membrane drives the *trans*-autophosphorylation of their respective cytosolic kinase domains (reviewed in 16). Active PERK phosphorylates eIF2 α (the α subunit the eukaryotic translation initiation factor 2), causing a decrease in global protein synthesis²⁴. Paradoxically, a subset of mRNAs bearing small upstream open reading frames (uORFs) on their 5’ untranslated regions (5’ UTRs) are preferentially translated in these conditions, including those encoding the transcription factors ATF4 and CHOP^{21,55–58}. By increasing the cellular metabolic capacity, ATF4 effects are mostly cytoprotective⁵⁹ whereas CHOP is largely regarded as pro-apoptotic^{60–63}. It

has recently been shown that, in addition to translational control by uORFs, N6-methyladenosine (m6A) methylation of the *ATF4* mRNA also contributes to the control of its translation. Demethylation of the 5' UTR of the *ATF4* mRNA by the demethylase ALKBH5 enhances its translational re-initiation, promoting the synthesis of ATF4 during ER stress⁶⁴. This mechanism highlights the exquisite fine-tuning of protein synthesis re-programming during stress.

Unlike PERK, IRE1 has no known kinase substrates besides itself, and its active kinase domain licenses the allosteric activation of its C-terminal RNase domain^{65,66}. Active IRE1 excises a small unconventional intron from the mRNA encoding the transcription factor XBP1^{18,19}. The resulting exons are joined by the tRNA ligase RTCB to produce a new mRNA encoding the transcription factor XBP1S ("S", for spliced)^{67–69}. Even though *XBP1* mRNA splicing was discovered almost 20 years ago, some of the salient features of this mechanism have been elucidated recently. Five years ago, RTCB was independently identified by three groups as the *XBP1* mRNA splicing ligase^{67–69}. A conformational change in the *XBP1* mRNA, dubbed an RNA "zipper", which is required to eject the intron and hold the exons together after cleavage, was described shortly after⁷⁰. Additional recent work has shown that an intact 2'-3' cyclic phosphate—long known to be left on the RNA ends after cleavage by IRE1⁷¹—is essential for completion of the *XBP1* mRNA splicing reaction⁷². The opposing activities of the cyclic phosphodiesterase CNP and the RNA cyclase RTCA control the availability of the cyclic phosphate⁷². Targeted quantitative proteomics analyses revealed that IRE1 is found in complex with RTCB in cells⁷³, suggesting that the *XBP1* mRNA splicing can be completed immediately after mRNA cleavage. This newly described multi-step regulation of *XBP1* mRNA splicing could also provide regulatory layers controlling a tunable UPR output.

IRE1 signaling is not confined to *XBP1* mRNA splicing. IRE1 also cleaves ER-bound mRNAs in a process known as regulated IRE1-dependent decay (RIDD)^{22,23}. When first discovered, RIDD was thought to protect the ER by lowering ER load through the selective cleavage of mRNAs²³; however, a recent finding challenges this view. RIDD of a single mRNA encoding the lysosome trafficking factor BLOS1 has been shown to protect cells from proteotoxicity by enhancing their capacity to degrade protein aggregates by microautophagy⁷⁴. The precise molecular mechanism that determines the fate of an mRNA encountering IRE1—splicing or RIDD—appears to hinge on the aforementioned *XBP1* mRNA zipper, which is absent in RIDD targets studied to date; implanting this mRNA zipper structure into RIDD target mRNAs results in their splicing⁷⁵.

Our understanding of IRE1 signaling mechanisms has also expanded lately. Several lines of evidence support the notion that IRE1 is an integrating node linking the UPR and the ER protein co-translational targeting machinery. IRE1 has been shown to bind to the Sec61 translocon⁷⁶, and impairing this interaction resulted in dysregulated IRE1 activity⁷⁷. The recently developed Perturb-seq method, which combines single-cell RNA-seq with CRISPR-based genetic screens, further substantiated these observations

by showing that depletion of translocon subunits resulted in exclusive IRE1 activation without impact on other UPR signaling pathways⁷⁸. More recently, RNA–protein cross-linking and mass spectrometry-based approaches revealed that IRE1 associates with the signal recognition particle, tRNAs, mRNAs, and ribosomes in living cells⁷³. All of these observations converge on a model in which IRE1 oversees the health and availability of translocons while it monitors the co-translational targeting machinery at the ER surface.

Cross-talk between UPR signaling pathways controls adaptation and death

A basic level of pathway interconnectivity in the UPR comprises the coordinated actions of transcription factors. In the adaptive phase of the UPR, ATF6N and XBP1S increase the synthesis of chaperones, protein-folding enzymes, and proteins that take part in ER protein turnover mechanisms, and they physically enlarge the ER by upregulating endomembrane biosynthesis^{9,79–83}. In parallel, ATF4 upregulates the biosynthetic capacity of the cell by controlling genes required for antioxidant responses and amino acid import⁸⁹. Adaptive transcriptional signals further integrate at the level of combinatorial regulation. For example, ATF6N and XBP1S can form heterodimers⁸⁴, thereby expanding the repertoire of UPR *cis*-regulatory elements^{9,20,83,85,86}. Post-transcriptional regulatory control by XBP1U ("U" for unspliced), encoded by the *XBP1* precursor mRNA, adjusts the transcriptional responses; XBP1U regulates the turnover of XBP1S and ATF6N, setting a molecular timer for the duration of the adaptive phase of the UPR^{87,88}.

Not all UPR transcriptional outputs are adaptive. ATF4, ATF6N, and XBP1S converge on the induction of CHOP^{9,21,83,85}, which negatively impacts the ER protein-folding capacity, downregulates the anti-apoptotic protein BCL2, and upregulates the pro-apoptotic signaling protein death receptor 5 (DR5)^{50,61,63,89,90}. However, recent data showing that genetic depletion of ATF6—but not IRE1 or PERK—significantly impaired the growth of HeLa cells engineered to adopt a plasma cell-like secretory phenotype that predisposes them to ER stress suggest that ATF6 is mostly cytoprotective⁹¹. The homeostatic role of the IRE1–XBP1 axis has recently been challenged by the discovery of cytotoxic XBP1-driven responses⁹². These data show that surpassing a critical ER stress threshold allows XBP1S to indirectly induce the expression of the ER calcium channels TMEM38B and ITPR1, leading to depletion of ER calcium stores, thereby providing a positive feedback loop to aggravate ER stress⁹². In this model, cellular demise arises from collateral damage caused by the transcription factor KLF9 downstream of XBP1S and not by the direct control of *bona fide* pro-apoptotic genes by XBP1S.

XBP1-independent roles with regard to cytoprotection or apoptosis have also been proposed for IRE1. One model suggests that IRE1 promotes cell death by degrading microRNAs (miRNAs) targeting mRNAs encoding the pro-apoptotic proteins TXNIP and caspase-2^{93,94}. This model is somewhat disputed, as caspase-2 appears to be dispensable for eliciting cell death in response to ER stress^{50,95}. Another view posits that RIDD could

lead to cell death through pervasive RNA cleavage⁹⁶. The notion of antagonistic IRE1 roles has been further substantiated by a recent finding showing that XBP1S promotes glioblastoma tumor progression while RIDD obstructs it⁹⁷.

Antagonism among UPR sensors also provides a model for UPR-dependent cell fate control. This model considers a dynamic cross-talk between cytoprotective and pro-apoptotic signals emanating from different UPR sensors. IRE1 promotes cytoprotection through *XBP1* mRNA splicing and RIDD, while PERK promotes apoptosis by inducing the ATF4–CHOP axis and the gene encoding *DR5* downstream of it⁵⁰. *DR5* signaling instructs the cells to die in response to unmitigable ER stress through the extrinsic (caspase-8-dependent) apoptotic pathway; IRE1 combats the apoptotic signal by degrading the *DR5* mRNA⁵⁰. Concomitantly, the phosphatase RPAP2, downstream of PERK, dephosphorylates IRE1 attenuating IRE1 signaling⁹⁸. Along with RPAP2, caspase-mediated IRE1 turnover can also play a role in terminating signaling and enforcing the apoptotic program⁹⁹. Surprisingly, cleavage of IRE1 by caspases also generates a proteolytic fragment that antagonizes BAX-driven pro-apoptotic signaling from the mitochondria⁹⁹, further substantiating the notion of extensive functional cross-talk between the UPR and other fundamental cellular processes.

PERK also stimulates the production of GADD34, a regulatory subunit of PP1 (protein phosphatase 1), which dephosphorylates eIF2 α , thereby establishing a negative feedback loop controlling responses downstream of PERK^{100,101}. The cross-connectivity between UPR signaling pathways has been further substantiated by recent RNA-seq and ribosome profiling experiments that show that PERK has the potential to repress—both transcriptionally and post-transcriptionally—the expression of a subset of cytoprotective genes induced by XBP1S and ATF6N¹⁰². Together, these observations support a model in which a molecular timer, set off by opposing signals downstream of IRE1 and PERK, coordinates the survival versus death decision. In this model, the activities of IRE1 and PERK are coordinated, ensuring that specific responses kick in as the response to ER stress progresses. At the beginning of the response, homeostatic mechanisms are enforced while apoptotic ones are suppressed (for example, induction of *XBP1* mRNA splicing, RIDD of the *DR5* mRNA). If stress persists, apoptosis mechanisms take over (for example, shutdown of IRE1 signaling, suppression of cytoprotective genes downstream of XBP1S and ATF6).

Apart from the mechanisms discussed above, protein quality-control mechanisms of UPR sensors have been implicated in UPR signaling; the protein levels of IRE1, PERK, and ATF6 are controlled by ER-associated degradation (ERAD)^{103–105}. These observations suggest that ERAD engages negative feedback loops that could enforce the survival versus death decision. Moreover, recent data suggest that post-translational modification of IRE1 and PERK by ufmylation, which regulates their stability, plays an important role in regulating apoptosis and plasma cell development^{106,107}. Importantly, post-transcriptional mechanisms operate alongside protein turnover devices to regulate the UPR. It has increasingly been recognized that non-coding RNAs—mostly

miRNAs but also long non-coding RNAs—act as important regulators of the UPR (reviewed in 108). This type of regulatory control adds a layer of complexity to the homeostatic or apoptotic programs controlled by the UPR.

Beyond the ER: non-canonical UPR mechanisms and interorganellar communication

As we move forward in our understanding of the UPR, we have come to recognize that UPR signaling extends beyond protecting ER physiology. Recent observations support an integral role of the UPR as a hub for interorganellar communication. At ER–mitochondria membrane contact sites, IRE1 coordinates mitochondrial physiological bioenergetics by engaging the calcium channel ITPR¹⁰⁹. IRE1 regulation at ER–mitochondria junctions also tunes biomedically relevant processes, such as the regulation of T-cell responses¹¹⁰. In parallel, PERK stimulates the assembly of respiratory chain supercomplexes in a nutritional and ER stress-dependent manner^{111,112}. The UPR–mitochondria interplay is bidirectional: a genomic high-content screen in yeast demonstrated that mitochondrial heme biosynthesis enables optimal UPR signaling¹¹³. More recently, the mitochondrial ubiquitin ligase MITOL was shown to ubiquitylate IRE1 at ER–mitochondria contact sites to suppress its activity and prevent apoptosis¹¹⁴. These studies showcase the conserved role of the UPR as an integration node for interorganellar communication.

IRE1 and PERK are also found at ER–plasma membrane (ER–PM) contact sites through their association with the cytoskeletal scaffold filamin A (FLNA). PERK-driven FLNA recruitment expands ER–PM contacts and replenishes ER calcium stores¹¹⁵, while IRE1 engages FLNA to facilitate cell motility¹¹⁶. These UPR mechanisms are independent of the enzymatic activities of IRE1 or PERK, relying instead on dynamic UPR sensor clustering. Unexpectedly, the role of UPR–organelle cross-talk in modulating UPR signaling was further illustrated by the observation that ceapins, recently discovered ATF6 signaling inhibitors^{117,118}, work by providing a neomorphic artificial tether between the ER and peroxisomes¹¹⁹.

In spite of the compelling evidence supporting non-canonical roles for IRE1 signaling, a recent report demonstrated that *XBP1* mRNA splicing is the only IRE1 activity required for medaka fish development and growth¹²⁰. However, this observation does not negate the relevance of cross-connectivity between the UPR and other cellular processes in maintaining organismal homeostasis. Far from behaving as a self-contained transcriptional program, the UPR drives the expression of distinct sets of genes in a cell type- and stimulus-dependent manner that involves metabolic, inflammatory, or developmental cues^{9,121,122}. Moreover, out of hundreds of UPR regulated genes, many have no direct functions in maintaining ER homeostasis. For example, some UPR target genes coordinate the DNA damage and repair response, hinting at a key role for UPR in maintaining genome integrity^{9,122,123}. Together, canonical and non-canonical mechanisms emerging from different UPR sensors—at least for IRE1 and PERK—assemble into a multipronged signaling relay that most likely determines the robustness of the UPR while coordinating the activities of other organelles and functions within the cell.

UPR signal integration at the organism level

The interconnected nature of UPR signaling is not restricted to single cells, as it has been increasingly clear that a cell non-autonomous UPR plays a role in maintaining organismal health. Work in *Caenorhabditis elegans* showed that IRE1 regulates stress resistance and longevity through an unidentified diffusible factor, which is a downstream target of XBP1¹²⁴. Moreover, ectopic expression of XBP1S in mouse neurons leads to stress responses in liver and improves hepatic insulin sensitivity¹²⁵. More recently, a role for UPR in the maintenance (or disruption) of circadian rhythms in animals has been elucidated. Through the combined action of transcriptional and translational mechanisms, the UPR regulates the expression of circadian factors to ensure the oscillatory control of metabolic rhythms^{126,127}. On the flip side, under ER stress conditions, PERK inhibits the circadian heterodimeric components BMAL1 and CLOCK, abrogating cellular rhythms in cells and animals^{128,129}.

The fundamental roles of the UPR in coordinating organism-level homeostasis are further accentuated by the potential for therapeutic utility of recently discovered small-molecule modulators of the UPR. In some instances, it may be desirable to suppress UPR signaling for a favorable outcome, as has been recently demonstrated for pharmacologic inhibition of IRE1 in orthotopic models of multiple myeloma¹³⁰ and atherosclerosis¹³¹ or by blocking IRE1 functions as a promising strategy for pain management¹³². In other instances, it may be desirable to enhance UPR signaling to increase the protein processing capacity of the ER. Recently discovered selective activators of ATF6 show promise in this regard as they alleviate amyloidogenic protein aggregation and proteotoxicity in various disease models, including myocardial ischemia^{133–135}. The positive effects of the small-molecule ISRIB, which renders cells insensitive to the effects of eIF2 α phosphorylation¹³⁶, on models of neurological disorders and brain injury further substantiate this notion. For example, ISRIB rescues the effects of genetic mutations observed in vanishing white matter disease¹³⁷, reverses cognitive deficits in mouse models of traumatic brain injury¹³⁸, alleviates social behavioral defects and anxiety-like symptoms in a genetic mouse model¹³⁹, and protects neural cells in mouse models of prion disease¹⁴⁰. Importantly, a small molecule called Sephin1, which has been proposed to obstruct eIF2 α dephosphorylation, also provides therapeutic benefit in mouse models of protein misfolding neuropathologies^{141–143}. These findings highlight the potential for controlling eIF2 α phosphorylation in different pathologies. All of these observations underscore the notion that the pharmacological manipulation of the UPR is an attractive opportunity for targeted therapeutic intervention in multiple diseases.

Taken together, the aforementioned findings portray a bottom-up view of the UPR homeostat that extends beyond the scope of maintaining ER homeostasis. In the UPR hierarchy, at its most basic level, the UPR maintains ER physiology. In the next level, the UPR interdigitates with cellular physiology through dynamic connections between the ER and other cellular components and processes. In a higher plane, the UPR integrates signals at the multicellular level, facilitating communication about specific states between cells and tissues and modulating the oscillatory

dynamics of organismal rhythms. By interlocking fundamental biological processes at multiple levels, the UPR emerges as a main regulatory hub that maintains the health of the whole organism.

What does the future hold for the UPR signaling mechanisms?

The past five years have seen great progress in our understanding of the UPR. These developments raise new questions. For example, although we know that clustering is essential for UPR sensor activation, we do not understand the mechanisms that would drive their spatiotemporal organization in specific domains of the ER membrane. We also know very little about the mechanistic details that underlie the functional partitioning of each UPR signaling pathway. Is the main function of IRE1 to survey translocon integrity and availability while other homeostatic functions are delegated to ATF6? Are pro-apoptotic functions limited mostly to PERK? The significance of other molecular mechanisms also remains obscure. Are there other RIDD substrates that, akin to BLOS1, control specific cytoprotective or pro-apoptotic responses? Is the ribosome a signaling hub for IRE1 or PERK? Do non-canonical UPR mechanisms impact the differentiation of cells and tissues, or do they exist simply to fine-tune cellular responses to specific physiological states? These questions will provide fertile ground for research for years to come.

Another recent observation that begets new questions is the finding that not every cell in a population responds in the same way to the same ER stress input⁷⁸. Is it possible that, akin to the innate antiviral type-I interferon response, alerting neighboring cells about a potential threat to ER functions allows coordinated responses that maintain tissue and organism homeostasis? Is it possible that cells exhibit different UPR types depending on whether they are actively cycling, have exited the cell cycle irreversibly in preparation for terminal differentiation, or are terminally differentiated? Future work addressing these questions could reveal new UPR signaling paradigms in multicellular organisms.

The differences in the response to ER stress of individual cells in an asynchronous population strongly hint at a fundamental connection between UPR signaling and the cell cycle. Substantiating this notion, the yeast XBP1 homolog Hac1 has been implicated in cytokinesis¹⁴⁴, and blocking IRE1 signaling delayed cell cycle progression in helper T-cells¹⁴⁵. In addition, PERK signaling has been proposed to negatively impact cell cycle progression^{146,147}. A putative role of XBP1U in controlling cell proliferation has recently been discovered: XBP1U acts as a negative regulator of the p53/p21 tumor suppressor axis, revealing a potential oncogenic role for this protein¹⁴⁸. Even though this role is independent of IRE1, it is possible that it remains linked to the UPR since both ATF6N and XBP1S induce *XBP1* mRNA expression^{9,19}. Future work is required to uncover the mechanistic details behind a putative “ER health checkpoint” in the metazoan cell cycle.

The developments discussed here highlight that the UPR story is far from complete. They portray a far more complex molecular circuitry for the UPR than previously anticipated.

This new knowledge on basic UPR mechanisms has the potential to dramatically change our outlook on how to manipulate the UPR for therapeutic intervention. It is becoming clear that the UPR is much more than the sum of its parts and that this Rube Goldberg stress signaling device will continue to gift significant discoveries for years to come.

Abbreviations

5' UTR, 5' untranslated region; ALKBH5, AlkB homolog 5 RNA demethylase; ATF4, activating transcription factor 4; ATF6, activating transcription factor 6; BAX, BCL2-associated X protein; BCL2, apoptosis regulator BCL2/B-cell CLL/lymphoma 2; BiP, binding-immunoglobulin protein; BLOS1, biogenesis of lysosome-related organelles complex 1 subunit 1; BMAL1, brain and muscle ARNT-like 1; CHOP, CCAAT/enhancer-binding protein homologous protein; CLOCK, circadian locomotor output cycles protein kaput; CNP, 2',3' cyclic nucleotide 3' phosphodiesterase; CNPY2, protein canopy homolog 2; CRISPR, clustered regularly interspaced short palindromic repeats; DR5, death receptor 5; eIF2 α , eukaryotic translation initiation factor 2, subunit alpha; ER, endoplasmic reticulum; ERAD, ER-associated degradation; ERDJ4, ER DnaJ homolog

4; ERp18, endoplasmic reticulum resident protein of 18 kDa; FLNA, filamin A; GADD34, growth arrest and DNA damage-inducible 34/PP1 regulatory subunit 15A; HSP47, heat shock protein 47; IRE1, inositol-requiring enzyme 1; ISRIB, integrated stress response inhibitor; ITPR1, inositol 1,4,5-trisphosphate receptor type 1; KLF9, Krueppel-like factor 9; m6A, N6-methyladenosine; miRNA, microRNA; MITOL, mitochondrial ubiquitin ligase; PDIA6, protein disulfide isomerase family A member 6; PERK, PKR-like endoplasmic reticulum kinase; PM, plasma membrane; PP1, protein phosphatase 1; RIDD, regulated IRE1-dependent decay; RPAP2, RNA polymerase II-associated protein 2; RTCA, RNA 3'-terminal phosphate cyclase; RTCB, tRNA-splicing ligase RtcB homolog; TMEM38B, transmembrane protein 38B; TXNIP, thioredoxin-interacting protein; uORF, upstream open reading frame; UPR, unfolded protein response; XBP1, X-box binding protein

Acknowledgments

We thank Carolina Arias (University of California, Santa Barbara) and the members of the Acosta-Alvear laboratory for their suggestions and critical revision of the manuscript.

References

- Reimold AM, Iwakoshi NN, Manis J, *et al.*: **Plasma cell differentiation requires the transcription factor XBP-1.** *Nature*. 2001; **412**(6844): 300–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Iwakoshi NN, Lee AH, Vallabhajosyula P, *et al.*: **Plasma cell differentiation and the unfolded protein response intersect at the transcription factor XBP-1.** *Nat Immunol*. 2003; **4**(4): 321–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Lee AH, Chu GC, Iwakoshi NN, *et al.*: **XBP-1 is required for biogenesis of cellular secretory machinery of exocrine glands.** *EMBO J*. 2005; **24**(24): 4368–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gass JN, Gifford NM, Brewer JW: **Activation of an unfolded protein response during differentiation of antibody-secreting B cells.** *J Biol Chem*. 2002; **277**(50): 49047–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Zhang K, Wong HN, Song B, *et al.*: **The unfolded protein response sensor IRE1 α is required at 2 distinct steps in B cell lymphopoiesis.** *J Clin Invest*. 2005; **115**(2): 268–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gao Y, Sartori DJ, Li C, *et al.*: **PERK is required in the adult pancreas and is essential for maintenance of glucose homeostasis.** *Mol Cell Biol*. 2012; **32**(24): 5129–39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Zhang W, Feng D, Li Y, *et al.*: **PERK EIF2AK3 control of pancreatic beta cell differentiation and proliferation is required for postnatal glucose homeostasis.** *Cell Metab*. 2006; **4**(6): 491–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cornejo VH, Pihán P, Vidal RL, *et al.*: **Role of the unfolded protein response in organ physiology: lessons from mouse models.** *IUBMB Life*. 2013; **65**(12): 962–75.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Acosta-Alvear D, Zhou Y, Blais A, *et al.*: **XBP1 controls diverse cell type- and condition-specific transcriptional regulatory networks.** *Mol Cell*. 2007; **27**(1): 53–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Dalton RP, Lyons DB, Lomvardas S: **Co-opting the unfolded protein response to elicit olfactory receptor feedback.** *Cell*. 2013; **155**(2): 321–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Cho YM, Jang YS, Jang YM, *et al.*: **Induction of unfolded protein response during neuronal induction of rat bone marrow stromal cells and mouse embryonic stem cells.** *Exp Mol Med*. 2009; **41**(6): 440.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kawada K, Iekumo T, Saito R, *et al.*: **Aberrant neuronal differentiation and inhibition of dendrite outgrowth resulting from endoplasmic reticulum stress.** *J Neurosci Res*. 2014; **92**(9): 1122–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Tekko T, Lilleväli K, Luuk H, *et al.*: **Initiation and developmental dynamics of *Wfs1* expression in the context of neural differentiation and ER stress in mouse forebrain.** *Int J Dev Neurosci*. 2014; **35**: 80–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Martínez G, Vidal RL, Mardones P, *et al.*: **Regulation of Memory Formation by the Transcription Factor XBP1.** *Cell Rep*. 2016; **14**(6): 1382–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Murao N, Nishitoh H: **Role of the unfolded protein response in the development of central nervous system.** *J Biochem*. 2017; **162**(3): 155–62.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Karagöz GE, Acosta-Alvear D, Walter P: **The Unfolded Protein Response: Detecting and Responding to Fluctuations in the Protein-Folding Capacity of the Endoplasmic Reticulum.** *Cold Spring Harb Perspect Biol*. 2019; **11**(9): pii: a033886.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Cox JS, Walter P: **A novel mechanism for regulating activity of a transcription factor that controls the unfolded protein response.** *Cell*. 1996; **87**(3): 391–404.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Calton M, Zeng H, Urano F, *et al.*: **IRE1 couples endoplasmic reticulum load to secretory capacity by processing the *XBP-1* mRNA.** *Nature*. 2002; **415**(6867): 92–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Yoshida H, Matsui T, Yamamoto A, *et al.*: **XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription factor.** *Cell*. 2001; **107**(7): 881–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Yoshida H, Haze K, Yanagi H, *et al.*: **Identification of the *cis*-acting endoplasmic reticulum stress response element responsible for transcriptional induction of mammalian glucose-regulated proteins. Involvement of basic leucine zipper transcription factors.** *J Biol Chem*. 1998; **273**(50): 33741–9.
[PubMed Abstract](#) | [Publisher Full Text](#)

21. Harding HP, Novoa I, Zhang Y, *et al.*: Regulated translation initiation controls stress-induced gene expression in mammalian cells. *Mol Cell*. 2000; 6(5): 1099–108. [PubMed Abstract](#) | [Publisher Full Text](#)
22. Hollien J, Lin JH, Li H, *et al.*: Regulated Ire1-dependent decay of messenger RNAs in mammalian cells. *J Cell Biol*. 2009; 186(3): 323–31. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. **F** Hollien J, Weissman JS: Decay of endoplasmic reticulum-localized mRNAs during the unfolded protein response. *Science*. 2006; 313(5783): 104–7. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
24. Harding HP, Zhang Y, Bertolotti A, *et al.*: Perk is essential for translational regulation and cell survival during the unfolded protein response. *Mol Cell*. 2000; 5(5): 897–904. [PubMed Abstract](#) | [Publisher Full Text](#)
25. Mori K: Signalling pathways in the unfolded protein response: development from yeast to mammals. *J Biochem*. 2009; 146(6): 743–50. [PubMed Abstract](#) | [Publisher Full Text](#)
26. **F** Halbleib K, Pesek K, Covino R, *et al.*: Activation of the Unfolded Protein Response by Lipid Bilayer Stress. *Mol Cell*. 2017; 67(4): 673–684.e8. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
27. Koh JH, Wang L, Beaudoin-Chabot C, *et al.*: Lipid bilayer stress-activated IRE-1 modulates autophagy during endoplasmic reticulum stress. *J Cell Sci*. 2018; 131(22): pii: jcs217992. [PubMed Abstract](#) | [Publisher Full Text](#)
28. Promlek T, Ishiwata-Kimata Y, Shido M, *et al.*: Membrane aberrancy and unfolded proteins activate the endoplasmic reticulum stress sensor Ire1 in different ways. *Mol Biol Cell*. 2011; 22(18): 3520–32. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. **F** Volmer R, van der Ploeg K, Ron D: Membrane lipid saturation activates endoplasmic reticulum unfolded protein response transducers through their transmembrane domains. *Proc Natl Acad Sci U S A*. 2013; 110(12): 4628–33. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
30. Kono N, Amin-Wetzel N, Ron D: Generic membrane-spanning features endow IRE1 α with responsiveness to membrane aberrancy. *Mol Biol Cell*. 2017; 28(17): 2318–32. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. **F** Tam AB, Roberts LS, Chandra V, *et al.*: The UPR Activator ATF6 Responds to Proteotoxic and Lipotoxic Stress by Distinct Mechanisms. *Dev Cell*. 2018; 46(3): 327–343.e7. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
32. Fun XH, Thibault G: Lipid bilayer stress and proteotoxic stress-induced unfolded protein response deploy divergent transcriptional and non-transcriptional programmes. *Biochim Biophys Acta Mol Cell Biol Lipids*. 2019; pii: S1388-1981(19)30062-9. [PubMed Abstract](#) | [Publisher Full Text](#)
33. Karagöz GE, Acosta-Alvear D, Nguyen HT, *et al.*: An unfolded protein-induced conformational switch activates mammalian IRE1. *eLife*. 2017; 6: pii: e30700. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. **F** Wang P, Li J, Tao J, *et al.*: The luminal domain of the ER stress sensor protein PERK binds misfolded proteins and thereby triggers PERK oligomerization. *J Biol Chem*. 2018; 293(11): 4110–21. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
35. Dalton RP, Karagöz GE, Kahiapo G, *et al.*: Olfactory and Vomeronasal Receptor Feedback Employ Divergent Mechanisms of PERK Activation. *Biorxiv*. 2018. [Publisher Full Text](#)
36. **F** Carrara M, Prischi F, Nowak PR, *et al.*: Noncanonical binding of BIP ATPase domain to Ire1 and Perk is dissociated by unfolded protein C α 1 to initiate ER stress signaling. *eLife*. 2015; 4. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
37. **F** Carrara M, Prischi F, Nowak PR, *et al.*: Crystal structures reveal transient PERK luminal domain tetramerization in endoplasmic reticulum stress signaling. *EMBO J*. 2015; 34(11): 1589–600. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
38. **F** Amin-Wetzel N, Saunders RA, Kamphuis MJ, *et al.*: A J-Protein Co-chaperone Recruits BIP to Monomerize IRE1 and Repress the Unfolded Protein Response. *Cell*. 2017; 171(7): 1625–1637.e13. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
39. Bertolotti A, Zhang Y, Hendershot LM, *et al.*: Dynamic interaction of BIP and ER stress transducers in the unfolded-protein response. *Nat Cell Biol*. 2000; 2(6): 326–32. [PubMed Abstract](#) | [Publisher Full Text](#)
40. **F** Pincus D, Chevalier MW, Aragón T, *et al.*: BIP binding to the ER-stress sensor Ire1 tunes the homeostatic behavior of the unfolded protein response. *PLoS Biol*. 2010; 8(7): e1000415. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
41. Okamura K, Kimata Y, Higashio H, *et al.*: Dissociation of Kar2p/BIP from an ER sensory molecule, Ire1p, triggers the unfolded protein response in yeast. *Biochem Biophys Res Commun*. 2000; 279(2): 445–50. [PubMed Abstract](#) | [Publisher Full Text](#)
42. **F** Eletto D, Eletto D, Dersh D, *et al.*: Protein disulfide isomerase A6 controls the decay of IRE1 α signaling via disulfide-dependent association. *Mol Cell*. 2014; 53(4): 562–76. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
43. **F** Eletto D, Eletto D, Boyle S, *et al.*: PDIA6 regulates insulin secretion by selectively inhibiting the RIDD activity of IRE1. *FASEB J*. 2016; 30(2): 653–65. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
44. Sepulveda D, Rojas-Rivera D, Rodríguez DA, *et al.*: Interactome Screening Identifies the ER Luminal Chaperone Hsp47 as a Regulator of the Unfolded Protein Response Transducer IRE1 α . *Mol Cell*. 2018; 69(2): 238–252.e7. [PubMed Abstract](#) | [Publisher Full Text](#)
45. Hong F, Liu B, Wu BX, *et al.*: CNPY2 is a key initiator of the PERK-CHOP pathway of the unfolded protein response. *Nat Struct Mol Biol*. 2017; 24(10): 834–9. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. Wang Q, Groenendyk J, Paskevicius T, *et al.*: Two pools of IRE1 α in cardiac and skeletal muscle cells. *FASEB J*. 2019; 33(8): 8892–904. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
47. Yoshida H, Matsui T, Hosokawa N, *et al.*: A time-dependent phase shift in the mammalian unfolded protein response. *Dev Cell*. 2003; 4(2): 265–71. [PubMed Abstract](#)
48. **F** Lin JH, Li H, Yasumura D, *et al.*: IRE1 signaling affects cell fate during the unfolded protein response. *Science*. 2007; 318(5852): 944–9. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
49. Li H, Korenykh AV, Behrman SL, *et al.*: Mammalian endoplasmic reticulum stress sensor IRE1 signals by dynamic clustering. *Proc Natl Acad Sci U S A*. 2010; 107(37): 16113–8. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. **F** Lu M, Lawrence DA, Marsters S, *et al.*: Opposing unfolded-protein-response signals converge on death receptor 5 to control apoptosis. *Science*. 2014; 345(6192): 98–101. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
51. Haze K, Yoshida H, Yanagi H, *et al.*: Mammalian transcription factor ATF6 is synthesized as a transmembrane protein and activated by proteolysis in response to endoplasmic reticulum stress. *Mol Biol Cell*. 1999; 10(11): 3787–99. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
52. Ye J, Rawson RB, Komuro R, *et al.*: ER stress induces cleavage of membrane-bound ATF6 by the same proteases that process SREBPs. *Mol Cell*. 2000; 6(6): 1355–64. [PubMed Abstract](#) | [Publisher Full Text](#)
53. Nakanaka S, Okada T, Yoshida H, *et al.*: Role of disulfide bridges formed in the luminal domain of ATF6 in sensing endoplasmic reticulum stress. *Mol Cell Biol*. 2007; 27(3): 1027–43. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
54. **F** Oka OBV, Lith M, Rudolf J, *et al.*: ERp18 regulates activation of ATF6 α during unfolded protein response. *EMBO J*. 2019; 38(15): [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
55. Palam LR, Baird TD, Wek RC: Phosphorylation of eIF2 facilitates ribosomal bypass of an inhibitory upstream ORF to enhance CHOP translation. *J Biol Chem*. 2011; 286(13): 10939–49. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. Vattem KM, Wek RC: Reinitiation involving upstream ORFs regulates ATF4 mRNA translation in mammalian cells. *Proc Natl Acad Sci U S A*. 2004; 101(31): 11269–74. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
57. Lu PD, Harding HP, Ron D: Translation reinitiation at alternative open reading frames regulates gene expression in an integrated stress response. *J Cell Biol*. 2004; 167(1): 27–33. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
58. Jousse C, Bruhat A, Carraro V, *et al.*: Inhibition of CHOP translation by a peptide encoded by an open reading frame localized in the chop 5'UTR. *Nucleic Acids Res*. 2001; 29(21): 4341–51. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
59. Harding HP, Zhang Y, Zeng H, *et al.*: An integrated stress response regulates amino acid metabolism and resistance to oxidative stress. *Mol Cell*. 2003; 11(3): 619–33. [PubMed Abstract](#) | [Publisher Full Text](#)
60. Zinszner H, Kuroda M, Wang X, *et al.*: CHOP is implicated in programmed cell death in response to impaired function of the endoplasmic reticulum. *Genes Dev*. 1998; 12(7): 982–95. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
61. McCullough KD, Martindale JL, Klotz LO, *et al.*: Gadd153 sensitizes cells to endoplasmic reticulum stress by down-regulating Bcl2 and perturbing the cellular redox state. *Mol Cell Biol*. 2001; 21(4): 1249–59. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
62. Maytin EV, Ubeda M, Lin JC, *et al.*: Stress-inducible transcription factor CHOP/gadd153 induces apoptosis in mammalian cells via p38 kinase-dependent and -independent mechanisms. *Exp Cell Res*. 2001; 267(2): 193–204. [PubMed Abstract](#) | [Publisher Full Text](#)
63. **F** Marciniak SJ, Yun CY, Oyadomari S, *et al.*: CHOP induces death by promoting protein synthesis and oxidation in the stressed endoplasmic reticulum. *Genes Dev*. 2004; 18(24): 3066–77. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

64. **F** Zhou J, Wan J, Shu XE, *et al.*: **N⁶-Methyladenosine Guides mRNA Alternative Translation during Integrated Stress Response.** *Mol Cell.* 2018; 69(4): 636–647.e7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
65. Mendez AS, Alfaro J, Morales-Soto MA, *et al.*: **Endoplasmic reticulum stress-independent activation of unfolded protein response kinases by a small molecule ATP-mimic.** *eLife.* 2015; 4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
66. **F** Korennykh AV, Egea PF, Korostelev AA, *et al.*: **The unfolded protein response signals through high-order assembly of Ire1.** *Nature.* 2009; 457(7230): 687–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
67. **F** Lu Y, Liang F-X, Wang X: **A synthetic biology approach identifies the mammalian UPR RNA ligase RtcB.** *Mol Cell.* 2014; 55(5): 758–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
68. **F** Kosmaczewski SG, Edwards TJ, Han SM, *et al.*: **The RtcB RNA ligase is an essential component of the metazoan unfolded protein response.** *EMBO Rep.* 2014; 15(12): 1278–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
69. **F** Jurkin J, Henkel T, Nielsen AF, *et al.*: **The mammalian tRNA ligase complex mediates splicing of XBP1 mRNA and controls antibody secretion in plasma cells.** *EMBO J.* 2014; 33(24): 2922–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
70. Peschek J, Acosta-Alvear D, Mendez AS, *et al.*: **A conformational RNA zipper promotes intron ejection during non-conventional XBP1 mRNA splicing.** *EMBO Rep.* 2015; 16(12): 1688–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
71. Gonzalez TN, Sidrauski C, Dörfler S, *et al.*: **Mechanism of non-spliceosomal mRNA splicing in the unfolded protein response pathway.** *EMBO J.* 1999; 18(11): 3119–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
72. **F** Unlu I, Lu Y, Wang X: **The cyclic phosphodiesterase CNP and RNA cyclase RtcA fine-tune noncanonical XBP1 splicing during ER stress.** *J Biol Chem.* 2018; 293(50): 19365–19376.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
73. **F** Acosta-Alvear D, Karagöz GE, Fröhlich F, *et al.*: **The unfolded protein response and endoplasmic reticulum protein targeting machineries converge on the stress sensor IRE1.** *eLife.* 2018; 7: pii: e43036.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
74. **F** Bae D, Moore KA, Mella JM, *et al.*: **Degradation of Blos1 mRNA by IRE1 repositions lysosomes and protects cells from stress.** *J Cell Biol.* 2019; 218(4): 1118–1127.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
75. Li W, Okreglak V, Peschek J, *et al.*: **Engineering ER-stress dependent non-conventional mRNA splicing.** *eLife.* 2018; 7: pii: e35388.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
76. **F** Plumb R, Zhang ZR, Appathurai S, *et al.*: **A functional link between the co-translational protein translocation pathway and the UPR.** *eLife.* 2015; 4: e07426.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
77. **F** Sundaram A, Plumb R, Appathurai S, *et al.*: **The Sec61 translocon limits IRE1 α signaling during the unfolded protein response.** *eLife.* 2017; 6: pii: e27187.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
78. **F** Adamson B, Norman TM, Jost M, *et al.*: **A Multiplexed Single-Cell CRISPR Screening Platform Enables Systematic Dissection of the Unfolded Protein Response.** *Cell.* 2016; 167(7): 1867–1882.e21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
79. Adachi Y, Yamamoto K, Okada T, *et al.*: **ATF6 is a transcription factor specializing in the regulation of quality control proteins in the endoplasmic reticulum.** *Cell Struct Funct.* 2008; 33(1): 75–89.
[PubMed Abstract](#) | [Publisher Full Text](#)
80. Bommasamy H, Back SH, Fagone P, *et al.*: **ATF6 α induces XBP1-independent expansion of the endoplasmic reticulum.** *J Cell Sci.* 2009; 122(Pt 10): 1626–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
81. **F** Sriburi R, Jackowski S, Mori K, *et al.*: **XBP1: a link between the unfolded protein response, lipid biosynthesis, and biogenesis of the endoplasmic reticulum.** *J Cell Biol.* 2004; 167(1): 35–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
82. Lee AH, Iwakoshi NN, Glimcher LH: **XBP-1 regulates a subset of endoplasmic reticulum resident chaperone genes in the unfolded protein response.** *Mol Cell Biol.* 2003; 23(21): 7448–59.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
83. Okada T, Yoshida H, Akazawa R, *et al.*: **Distinct roles of activating transcription factor 6 (ATF6) and double-stranded RNA-activated protein kinase-like endoplasmic reticulum kinase (PERK) in transcription during the mammalian unfolded protein response.** *Biochem J.* 2002; 366(Pt 2): 585–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
84. Yamamoto K, Sato T, Matsui T, *et al.*: **Transcriptional induction of mammalian ER quality control proteins is mediated by single or combined action of ATF6 α and XBP1.** *Dev Cell.* 2007; 13(3): 365–76.
[PubMed Abstract](#) | [Publisher Full Text](#)
85. Yoshida H, Okada T, Haze K, *et al.*: **ATF6 activated by proteolysis binds in the presence of NF-Y (CBF) directly to the cis-acting element responsible for the mammalian unfolded protein response.** *Mol Cell Biol.* 2000; 20(18): 6755–67.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
86. Yoshida H, Okada T, Haze K, *et al.*: **Endoplasmic reticulum stress-induced formation of transcription factor complex ERSF including NF-Y (CBF) and activating transcription factors 6 α and 6 β that activates the mammalian unfolded protein response.** *Mol Cell Biol.* 2001; 21(4): 1239–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
87. Yoshida H, Oku M, Suzuki M, *et al.*: **pXBP1(U) encoded in XBP1 pre-mRNA negatively regulates unfolded protein response activator pXBP1(S) in mammalian ER stress response.** *J Cell Biol.* 2006; 172(4): 565–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
88. Yoshida H, Uemura A, Mori K: **pXBP1(U), a negative regulator of the unfolded protein response activator pXBP1(S), targets ATF6 but not ATF4 in proteasome-mediated degradation.** *Cell Struct Funct.* 2009; 34(1): 1–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
89. Yamaguchi H, Wang HG: **CHOP is involved in endoplasmic reticulum stress-induced apoptosis by enhancing DR5 expression in human carcinoma cells.** *J Biol Chem.* 2004; 279(44): 45495–502.
[PubMed Abstract](#) | [Publisher Full Text](#)
90. Abdelrahim M, Newman K, Vanderlaag K, *et al.*: **3,3'-diindolylmethane (DIM) and its derivatives induce apoptosis in pancreatic cancer cells through endoplasmic reticulum stress-dependent upregulation of DR5.** *Carcinogenesis.* 2006; 27: 717–28.
[PubMed Abstract](#) | [Publisher Full Text](#)
91. **F** Vitale M, Bakunts A, Orsi A, *et al.*: **Inadequate BIP availability defines endoplasmic reticulum stress.** *eLife.* 2019; 8: pii: e41168.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
92. **F** Fink EE, Moparthy S, Bagati A, *et al.*: **XBP1-KLF9 Axis Acts as a Molecular Rheostat to Control the Transition from Adaptive to Cytotoxic Unfolded Protein Response.** *Cell Rep.* 2018; 25(1): 212–223.e4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
93. Lerner AG, Upton JP, Praveen PVK, *et al.*: **IRE1 α induces thioredoxin-interacting protein to activate the NLRP3 inflammasome and promote programmed cell death under irremediable ER stress.** *Cell Metab.* 2012; 16(2): 250–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
94. **F** Upton JP, Wang L, Han D, *et al.*: **IRE1 α cleaves select microRNAs during ER stress to derepress translation of proapoptotic Caspase-2.** *Science.* 2012; 338(6108): 818–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
95. **F** Sandow JJ, Dorstyn L, O'Reilly LA, *et al.*: **ER stress does not cause upregulation and activation of caspase-2 to initiate apoptosis.** *Cell Death Differ.* 2014; 21(3): 475–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
96. **F** Han D, Lerner AG, Vande Walle L, *et al.*: **IRE1 α kinase activation modes control alternate endoribonuclease outputs to determine divergent cell fates.** *Cell.* 2009; 138(3): 562–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
97. **F** Lhomond S, Avril T, Dejeans N, *et al.*: **Dual IRE1 RNase functions dictate glioblastoma development.** *EMBO Mol Med.* 2018; 10(3): e7929.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
98. **F** Chang T-K, Lawrence DA, Lu M, *et al.*: **Coordination between Two Branches of the Unfolded Protein Response Determines Apoptotic Cell Fate.** *Mol Cell.* 2018; 71(4): 629–636.e5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
99. **F** Shemorry A, Harnoss JM, Guttman O, *et al.*: **Caspase-mediated cleavage of IRE1 controls apoptotic cell commitment during endoplasmic reticulum stress.** *eLife.* 2019; 8: e47084.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
100. Novoa I, Zeng H, Harding HP, *et al.*: **Feedback Inhibition of the Unfolded Protein Response by GADD34-Mediated Dephosphorylation of eIF2 α .** *J Cell Biol.* 2001; 153(5): 1011–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
101. Ma Y, Hendershot LM: **Delineation of a negative feedback regulatory loop that controls protein translation during endoplasmic reticulum stress.** *J Biol Chem.* 2003; 278(37): 34864–73.
[PubMed Abstract](#) | [Publisher Full Text](#)
102. **F** Gonen N, Sabath N, Burge CB, *et al.*: **Widespread PERK-dependent repression of ER targets in response to ER stress.** *Sci Rep.* 2019; 9(1): 4330.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
103. **F** Sun S, Shi G, Sha H, *et al.*: **IRE1 α is an endogenous substrate of endoplasmic-reticulum-associated degradation.** *Nat Cell Biol.* 2015; 17(12): 1546–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
104. Horimoto S, Ninagawa S, Okada T, *et al.*: **The Unfolded Protein Response Transducer ATF6 Represents a Novel Transmembrane-type Endoplasmic**

- Reticulum-associated Degradation Substrate Requiring Both Mannose Trimming and SEL1L Protein.** *J Biol Chem.* 2013; 288(44): 31517–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
105. Jeon YJ, Kim T, Park D, *et al.*: miRNA-mediated TUSC3 deficiency enhances UPR and ERAD to promote metastatic potential of NSCLC. *Nat Commun.* 2018; 9(1): 5110.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 106. **F** Zhu H, Bhatt B, Sivaprakasam S, *et al.*: Ufbp1 promotes plasma cell development and ER expansion by modulating distinct branches of UPR. *Nat Commun.* 2019; 10(1): 1084.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 107. Liu J, Wang Y, Song L, *et al.*: A critical role of DDGK1 in endoplasmic reticulum homeostasis via regulation of IRE1 α stability. *Nat Commun.* 2017; 8: 263.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 108. McMahon M, Samali A, Chevet E: Regulation of the unfolded protein response by noncoding RNA. *Am J Physiol Cell Physiol.* 2017; 313(3): C243–C254.
[PubMed Abstract](#) | [Publisher Full Text](#)
 109. **F** Carreras-Sureda A, Jaña F, Urra H, *et al.*: Non-canonical function of IRE1 α determines mitochondria-associated endoplasmic reticulum composition to control calcium transfer and bioenergetics. *Nat Cell Biol.* 2019; 21(6): 755–67.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 110. **F** Song M, Sandoval TA, Chae CS, *et al.*: IRE1 α -XBP1 controls T cell function in ovarian cancer by regulating mitochondrial activity. *Nature.* 2018; 562(7727): 423–428.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 111. **F** Balsa E, Soustek MS, Thomas A, *et al.*: ER and Nutrient Stress Promote Assembly of Respiratory Chain Supercomplexes through the PERK-eIF2 α Axis. *Mol Cell.* 2019; 74(5): 877–890.e6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 112. Lebeau J, Saunders JM, Moraes VWR, *et al.*: The PERK Arm of the Unfolded Protein Response Regulates Mitochondrial Morphology during Acute Endoplasmic Reticulum Stress. *Cell Rep.* 2018; 22(11): 2827–2836.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 113. Cohen N, Breker M, Bakunts A, *et al.*: Iron affects Ire1 clustering propensity and the amplitude of endoplasmic reticulum stress signaling. *J Cell Sci.* 2017; 130(19): 3222–3233.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 114. **F** Takeda K, Nagashima S, Shiba I, *et al.*: MITOL prevents ER stress-induced apoptosis by IRE 1 α ubiquitylation at ER-mitochondria contact sites. *EMBO J.* 2019; 38(15): e100999.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 115. van Vliet AR, Giordano F, Gerlo S, *et al.*: The ER Stress Sensor PERK Coordinates ER-Plasma Membrane Contact Site Formation through Interaction with Filamin-A and F-Actin Remodeling. *Mol Cell.* 2017; 65(5): 885–899.e6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 116. **F** Urra H, Henríquez DR, Cánovas J, *et al.*: IRE1 α governs cytoskeleton remodelling and cell migration through a direct interaction with filamin A. *Nat Cell Biol.* 2018; 20(8): 942–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 117. Gallagher CM, Garri C, Cain EL, *et al.*: Ceapins are a new class of unfolded protein response inhibitors, selectively targeting the ATF6 α branch. *eLife.* 2016; 5: pii: e11878.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 118. Gallagher CM, Walter P: Ceapins inhibit ATF6 α signaling by selectively preventing transport of ATF6 α to the Golgi apparatus during ER stress. *eLife.* 2016; 5: pii: e11880.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 119. **F** Torres SE, Gallagher CM, Plate L, *et al.*: Ceapins block the unfolded protein response sensor ATF6 α by inducing a neomorphic inter-organelle tether. *eLife.* 2019; 8: pii: e46595.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 120. **F** Ishikawa T, Kashima M, Nagano AJ, *et al.*: Unfolded protein response transducer IRE1-mediated signaling independent of XBP1 mRNA splicing is not required for growth and development of medaka fish. *eLife.* 2017; 6: pii: e26845.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 121. **F** Chen X, Iliopoulos D, Zhang Q, *et al.*: XBP1 promotes triple-negative breast cancer by controlling the HIF1 α pathway. *Nature.* 2014; 508(7494): 103–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 122. Argemí J, Kress TR, Chang HCY, *et al.*: X-box Binding Protein 1 Regulates Unfolded Protein, Acute-Phase, and DNA Damage Responses During Regeneration of Mouse Liver. *Gastroenterology.* 2017; 152(5): 1203–1216.e15.
[PubMed Abstract](#) | [Publisher Full Text](#)
 123. Rendleman J, Cheng Z, Maity S, *et al.*: New insights into the cellular temporal response to proteostatic stress. *eLife.* 2018; 7: pii: e39054.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 124. **F** Taylor RC, Dillin A: XBP-1 is a cell-nonautonomous regulator of stress resistance and longevity. *Cell.* 2013; 153(7): 1435–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 125. **F** Williams KW, Liu T, Kong X, *et al.*: Xbp1s in Pomc neurons connects ER stress with energy balance and glucose homeostasis. *Cell Metab.* 2014; 20(3): 471–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 126. Maillou C, Martín J, Sebastián D, *et al.*: Circadian- and UPR-dependent control of CPEB4 mediates a translational response to counteract hepatic steatosis under ER stress. *Nat Cell Biol.* 2017; 19(2): 94–105.
[PubMed Abstract](#) | [Publisher Full Text](#)
 127. Zhu B, Zhang Q, Pan Y, *et al.*: A Cell-Autonomous Mammalian 12 hr Clock Coordinates Metabolic and Stress Rhythms. *Cell Metab.* 2017; 25(6): 1305–1319.e9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 128. **F** Bu Y, Yoshida A, Chitnis N, *et al.*: A PERK-miR-211 axis suppresses circadian regulators and protein synthesis to promote cancer cell survival. *Nat Cell Biol.* 2018; 20(1): 104–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 129. **F** Gao L, Chen H, Li C, *et al.*: ER stress activation impairs the expression of circadian clock and clock-controlled genes in NIH3T3 cells via an ATF4-dependent mechanism. *Cell Signal.* 2019; 57: 89–101.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 130. Harnoss JM, Le Thomas A, Shemorry A, *et al.*: Disruption of IRE1 α through its kinase domain attenuates multiple myeloma. *Proc Natl Acad Sci U S A.* 2019; 116(33): 16420–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 131. **F** Tufanli O, Telkoparan Akililar P, Acosta-Alvear D, *et al.*: Targeting IRE1 with small molecules counteracts progression of atherosclerosis. *Proc Natl Acad Sci U S A.* 2017; 114(8): E1395–E1404.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 132. **F** Chopra S, Giovanelli P, Alvarado-Vazquez PA, *et al.*: IRE1 α -XBP1 signaling in leukocytes controls prostaglandin biosynthesis and pain. *Science.* 2019; 365(6450): pii: eaau6499.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 133. **F** Paxman R, Plate L, Blackwood EA, *et al.*: Pharmacologic ATF6 activating compounds are metabolically activated to selectively modify endoplasmic reticulum proteins. *eLife.* 2018; 7: pii: e37168.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 134. Plate L, Cooley CB, Chen JJ, *et al.*: Small molecule proteostasis regulators that reprogram the ER to reduce extracellular protein aggregation. *eLife.* 2016; 5: pii: e15550.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 135. **F** Blackwood EA, Azizi K, Thuerauf DJ, *et al.*: Pharmacologic ATF6 activation confers global protection in widespread disease models by reprogramming cellular proteostasis. *Nat Commun.* 2019; 10(1): 187.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 136. Sidrauski C, Acosta-Alvear D, Khoutorsky A, *et al.*: Pharmacological brake-release of mRNA translation enhances cognitive memory. *eLife.* 2013; 2: e00498.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 137. **F** Wong YL, LeBon L, Edalji R, *et al.*: The small molecule ISRIB rescues the stability and activity of Vanishing White Matter Disease eIF2B mutant complexes. *eLife.* 2018; 7: pii: e32733.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 138. Chou A, Krukowski K, Jopson T, *et al.*: Inhibition of the integrated stress response reverses cognitive deficits after traumatic brain injury. *Proc Natl Acad Sci U S A.* 2017; 114(31): E6420–E6426.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 139. Kabir ZD, Che A, Fischer DK, *et al.*: Rescue of impaired sociability and anxiety-like behavior in adult cacna1c-deficient mice by pharmacologically targeting eIF2 α . *Mol Psychiatry.* 2017; 22(8): 1096–109.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 140. Halliday M, Radford H, Sekine Y, *et al.*: Partial restoration of protein synthesis rates by the small molecule ISRIB prevents neurodegeneration without pancreatic toxicity. *Cell Death Dis.* 2015; 6: e1672.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 141. **F** Das I, Krzyzosiak A, Schneider K, *et al.*: Preventing proteostasis diseases by selective inhibition of a phosphatase regulatory subunit. *Science.* 2015; 348(6231): 239–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 142. **F** Chen Y, Podojil JR, Kunjamma RB, *et al.*: Sephin1, which prolongs the integrated stress response, is a promising therapeutic for multiple sclerosis. *Brain.* 2019; 142(2): 344–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 143. **F** Carrara M, Sigurdardottir A, Bertolotti A: Decoding the selectivity of eIF2 α holophosphatases and PPP1R15A inhibitors. *Nat Struct Mol Biol.* 2017; 24(9): 708–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 144. Bicknell AA, Babour A, Federovitch CM, *et al.*: A novel role in cytokinesis reveals a housekeeping function for the unfolded protein response. *J Cell Biol.* 2007; 177(6): 1017–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

145. Pramanik J, Chen X, Kar G, *et al.*: **Genome-wide analyses reveal the IRE1a-XBP1 pathway promotes T helper cell differentiation by resolving secretory stress and accelerating proliferation.** *Genome Med.* 2018; **10**(1): 76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
146. Brewer JW, Diehl JA: **PERK mediates cell-cycle exit during the mammalian unfolded protein response.** *Proc Natl Acad Sci U S A.* 2000; **97**(23): 12625–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
147. Hamanaka RB, Bennett BS, Cullinan SB, *et al.*: **PERK and GCN2 contribute to eIF2alpha phosphorylation and cell cycle arrest after activation of the unfolded protein response pathway.** *Mol Biol Cell.* 2005; **16**(12): 5493–501.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
148. Huang C, Wu S, Ji H, *et al.*: **Identification of XBP1-u as a novel regulator of the MDM2/p53 axis using an shRNA library.** *Sci Adv.* 2017; **3**(10): e1701383.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Jeffrey L. Brodsky**

Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA, USA

Competing Interests: No competing interests were disclosed.

2. **Neil Bulleid**

Institute of Molecular, Cell and Systems Biology, College of Medical Veterinary and Life Sciences, University of Glasgow, Glasgow, UK

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research