REVIEW

Recent advances in understanding extremophiles [version 1; peer review: 2 approved]

James A Coker

Department of Sciences, University of Maryland Global Campus, Adelphi, MD, USA

V1 First published: 13 Nov 2019, 8(F1000 Faculty Rev):1917
<https://doi.org/10.12688/f1000research.20765.1>Latest published: 13 Nov 2019, 8(F1000 Faculty Rev):1917
<https://doi.org/10.12688/f1000research.20765.1>

Abstract

Despite the typical human notion that the Earth is a habitable planet, over three quarters of our planet is uninhabitable by us without assistance. The organisms that live and thrive in these “inhospitable” environments are known by the name extremophiles and are found in all Domains of Life. Despite our general lack of knowledge about them, they have already assisted humans in many ways and still have much more to give. In this review, I describe how they have adapted to live/thrive/survive in their niches, helped scientists unlock major scientific discoveries, advance the field of biotechnology, and inform us about the boundaries of Life and where we might find it in the Universe.

Keywords

extremophile, acidophile, alkaliphile, halophile, psychrophile, thermophile, piezophile, astrobiology, biotechnology, radiophile, model organism, replication, TBP, TFB, TATA, 16S rRNA, PCB, PHA

Open Peer Review

Approval Status

	1	2
version 1
13 Nov 2019		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Noah Mesbah**, Suez Canal University, Ismalia, Egypt
2. **Linda DeVeaux**, New Mexico Institute of Mining and Technology, Socorro, USA

Any comments on the article can be found at the end of the article.

Corresponding author: James A Coker (James.Coker@umuc.edu)

Author roles: Coker JA: Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2019 Coker JA. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Coker JA. **Recent advances in understanding extremophiles [version 1; peer review: 2 approved]** F1000Research 2019, 8(F1000 Faculty Rev):1917 <https://doi.org/10.12688/f1000research.20765.1>

First published: 13 Nov 2019, 8(F1000 Faculty Rev):1917 <https://doi.org/10.12688/f1000research.20765.1>

Introduction

As human beings, we tend to look at the world from our perspective and in turn the conditions under which our species thrives as “normal” and “typical” for the planet Earth; however, this is quite far from the truth. When looked at as a whole, the Earth is actually quite a cold place since 90% of the world’s oceans are not more than 5 °C¹. When the polar and alpine regions are factored in, cold environments account for roughly three quarters of the planet Earth. Therefore, in regard to extremophiles, it is important to understand that they are “extreme” only because they are found in regions where human beings typically cannot live unassisted. It is also important to think of them as growing and thriving in these conditions and not merely eking out an existence as the “last organism standing” that was able to survive just a bit longer than the rest. After all, they have evolved a set of adaptations that have made these “extreme” environments their preferred niche.

All three Domains of Life are represented in all of Earth’s extreme environments. However, a vast majority are prokaryotes, as they are the most numerous organisms on the planet, so it should come as no surprise that they have been isolated or detected literally all over the Earth. Just looking at the known limits of growth — minus 12 °C (*Psychromonas ingrahamii*) to 129 °C (*Geogemma barossii*), pH of less than 0 (*Picrophilus torridus*) to 13 (*Plectonema nostocorum* and *Hydrogenophaga* sp.), pressures of more than 100 Mpa (*Shewanella benthica*), beyond saturating conditions of NaCl and KCl (*Haloflex volcanii*), and high levels of ultraviolet (UV) (>100 J/m²) and gamma (>12 kGy) radiation (*Halobacterium* sp. NRC-1 and *Deinococcus radiodurans*)^{2–8}—should more than confirm this. In addition to these gradients, viable microbes have been found from the mesosphere (48 to 77 km above the Earth’s surface)⁹ to the Mariana Trench/Challenger Deep (11 km below the ocean surface)⁶ to several kilometers below the Earth’s surface (for example, the South African gold mines)¹⁰.

In the following review, I will catalog the adaptations that the organisms that live in these environments have evolved to persist and thrive. I will also describe a few major contributions made to the body of scientific knowledge that were made by using these organisms as model systems. I will finish by discussing two topics most often associated with extremophiles: biotechnology applications and astrobiology.

Adaptations to environmental conditions

pH: acidophiles

With few exceptions, these organisms maintain a cellular pH near neutral, which requires maintaining a pH gradient that is several orders of magnitude different on either side of the plasma membrane¹¹. One benefit of this is that there is always a ready supply of protons to power the proton motive force and form adenosine triphosphate (ATP) via the electron transport chain. However, the flip side is that the unusually high differences in concentration of protons (compared with neutrophiles) mean that, if left unchecked, the incoming protons have a greater capacity to dramatically change the internal cellular pH, which would lead to cell death.

To counteract these conditions, acidophiles have evolved several mechanisms. One of these is a cell membrane that is fairly impermeable to protons¹². In the Archaea branch of Life, this impermeability has been shown via tetraether lipids, differences in lipid head-group structures, a bulky isoprenoid core, and the fact that its ether linkages are less sensitive to acid hydrolysis compared with the ester linkages in the Bacteria and Eukarya Domains¹³. Another mechanism is the reduced pore size of membrane channels, which has been shown for *Acidithiobacillus ferrooxidans*¹⁴. Acidophiles also have a net positive potential charge inside the cell, which can counteract the high concentration of H⁺ ions in their surroundings. They also employ active proton pumping as has been observed in *Bacillus* and *Thermoplasma* species¹⁵. It has not been shown that there is an identifiable acid response signature in the genomes of acidophiles. However, it is interesting to note that the genomes of acidophiles are predominately smaller than neutrophiles¹³. It is currently unclear why this should be and whether it confers any evolutionary advantage to these organisms.

pH: alkaliphiles

When adaptations are discussed, alkaliphiles are often grouped with halophiles, as they are typically found in saline environments¹⁶. However, the response to high pH is specific to these organisms and is worth discussing. The cytoplasm of alkaliphiles, like that of acidophiles, is typically near neutral pH¹⁷; therefore, alkaliphiles also have to overcome an imbalance of H⁺ ions¹⁸. Where acidophiles are “swimming” in H⁺ ions, alkaliphiles are in a relative desert by comparison.

In response to this challenge, alkaliphiles have developed a negatively charged cell wall, which lowers the pH of the environment just outside the cell. They also produce an acidic secondary cell wall composed of teichurono-peptide and teichuronic acid or polyglutamic acid. These acids attract H⁺ and repel OH⁻, possibly helping to generate the proton motive force needed to drive ATP synthesis. In several alkaliphilic *Bacillus* species, the proton motive force for ATP synthesis is driven by Na⁺ or K⁺ antiporters that catalyse an electrogenic exchange of outwardly moving ions (Na⁺ or K⁺) and an increased number of entering H⁺¹⁹. More generally, alkaliphiles are able to use these antiporters (Na⁺/H⁺ and K⁺/H⁺)²⁰ and also produce acids to reduce the internal pH when it is too high for metabolism to occur⁷.

Salinity: halophiles

Organisms that require a saline environment to grow (also known as halophiles) can be found along a continuum of salinity. These organisms have adopted either a “salt in” or “salt out” approach as the main adaptation for their ability to thrive in these conditions²¹.

As suggested by the name, the salt-in approach means that these organisms have salinity/ion concentrations (up to 4 or 5 M) that are similar both inside and outside the cell membrane. As such, the cytosol of these organisms presents a significant challenge for the regular biochemistry of life. High-salinity conditions typically strip water molecules from proteins, resulting in aggregation and precipitation, and often this is due to exposed

hydrophobic patches binding to one another. To counteract this, these organisms have evolved a proteome that is composed of primarily acidic proteins²², and the acidic residues (aspartic and glutamic acid) are typically found on the surface of most of their proteins. These acidic residues have been shown to coordinate water molecules (that is, H^+ of water interacts with the COO^- of the acidic side chain) around the proteins forming a “water cage” that protects the proteins from being dehydrated and precipitating out of solution^{23,24}. As a result of the large-scale evolutionary changes needed for this survival (that is, changes to the proteome), these organisms tend to live mainly in environments where salinity does not dramatically fluctuate frequently. Thus far, only prokaryotes (bacteria and archaea) have been shown to adopt this strategy²⁵.

In contrast to salt-in organisms, the salt-out organisms have differing concentrations of salt/ions inside and outside the cell membrane (similar to H^+ with acido- and alkaliphiles). This strategy is more energy-intensive than the salt-in strategy¹⁶, as these organisms actively accumulate ions and organic osmolytes (for example, glycine betaine, ectoine/5-hydroxyectoine, glucosylglycerol, and dimethylsulfoniopropionate²⁶) to maintain turgor pressure. The accumulated compatible solutes eventually can be released into the environment via mechanosensitive channels or used as an energy source during times of lower external salinity²⁵. As the salt-out approach requires fewer large-scale evolutionary changes, organisms that have adopted it are able to grow over a wider range of salinity.

Radiation: radiophiles

Organisms’ responses to primarily two main types of radiation—ionizing (gamma) radiation and UV radiation—have been studied. Although on the surface it may seem that the same mechanisms of adaptation should be involved in both, there are, in fact, quite a few differences, which are probably due to the types of damage caused by each.

Ionizing radiation is responsible mainly for double-stranded breaks in the genome of organisms. However, it has also been shown to damage both proteins and lipids and induce persistent oxidative stress²⁷. Therefore, ionizing radioresistant organisms have developed all, or a combination of, the following strategies: novel and adaptive DNA repair mechanisms, antioxidant and enzymatic defense systems, and a condensed nucleoid. Fast and accurate repair of genomes is essential in surviving doses of ionizing radiation. This has been shown to be accomplished through the use of the nucleotide excision repair pathway (*uvrA1B*), base excision repair pathway (*ung* and *mutY*), and homologous recombination pathway (*recA*, *ruvA*, *ddrA*, and *pprA*) in *D. radiodurans*²⁸ and single-stranded binding proteins in *Halobacterium* sp. NRC-1 (Rfa-like genes)⁴ and *D. radiodurans* (DdrB and SSB)^{29,30}. In some especially sensitive species, protein damage causes death before double-stranded breaks start to form. It has been suggested that, especially for bacterial species, the role of protein damage in cell death due to ionizing radiation is underestimated. For example, *D. radiodurans* cells contain

several oxidative stress prevention and tolerance mechanisms: cell cleaning through elimination of oxidized macromolecules, selective protection of proteins against oxidative damage, and suppression of reactive oxygen species production. A condensed nucleoid has also been shown to promote the efficiency/accuracy of DNA repair³¹ and to limit the diffusion³² of radiation-generated DNA fragments.

Unlike ionizing radiation where DNA damage is primarily double-stranded breaks, UV radiation damages DNA in more subtle ways through the formation of cyclobutene pyrimidine dimers (that is, thymine dimers) and pyrimidine (6-4) pyrimidone photoproducts (that is, 6-4 photo products). These two types of damage account for roughly 80% of photolesions induced by UV radiation³³. However, cyclobutene pyrimidine dimers are far more numerous and typically outnumber pyrimidine (6-4) pyrimidone photoproducts 3 to 1. To repair these DNA lesions, organisms use a combination of photoreactivation (*phr*) genes, nucleotide excision repair (*uvrABCD*, *xpf*, and *rad*), base excision repair (*mutY* and *nth*), and homologous recombination (*recA* and *radA/51*)³³. Additionally, organisms have evolved a suite of photoprotection devices to protect themselves from continual exposure to UV radiation. These include carotenoids, superoxide dismutases and hydroperoxidases, gene duplication via polyploidy, and genome composition (that is, reduction in the number of bipyrimidine sequences)³³. However, as in ionizing radiation, reactive oxygen species interference with normal metabolic processes is a more typical cause of cell death³⁴.

Pressure: piezophiles

These organisms are typically found in deep lakes such as Baikal (1.6 km) and Tanganyika (1.5 km), the ocean, and sub-surface communities. To have an idea of how much pressure is involved, one must remember that hydrostatic pressure increases roughly 10.5 kPa per meter depth while lithostatic (overburden) pressure increases about 22.6 kPa per meter. This means that microbes growing at the bottom of the Mariana Trench (10.9 km below the ocean surface) are subjected to 114.4 kPa of pressure but that those in the South African gold mines (3.5 km below the Earth’s surface) face 79.1 kPa.

With increasing pressures, membranes lose fluidity and permeability as lipids pack more tightly and enter a gel phase similar to what happens at cold temperatures. To counteract this, bacterial piezophiles have been shown to incorporate polyunsaturated and monounsaturated fatty acids¹¹ or phosphatidylglycerol and phosphatidylcholine instead of phosphatidylethanolamine³⁵. Little is known about the adaptations in their archaeal counterparts, but alterations in the amount of certain archaeols seem to be important³⁶. Other mechanisms of adaptation are generally not well known, but a few reports suggest that changes are not due to a specific group of genes/enzymes but rather are the result of an overall change in metabolism^{37–39} — similar to how the salt-in halophiles have evolved an acidic proteome to keep proteins soluble and active at high salinities rather than altering the expression of a few genes.

Temperature: psychrophiles

As mentioned above, the Earth and the Universe are both predominately cold environments. At first glance, psychrophile should be easy to define (that is, as something that grows in the cold) and it has often been defined as an organism with an optimum temperature of less than 15 °C⁴⁰. However, this definition has multiple problems: it is arbitrary, does not account for eukaryotes, and treats cells as mere thermal units⁴¹. Others have adopted the terms eurypsychrophile and stenopsychrophile to refer to a “broad” or “narrow” range of growth at low temperatures⁴². Still, some believe that the use of these terms does not “push” researchers enough to search for “true” psychrophiles – those organisms able to grow well below 0 °C. As such, there have been attempts to classify psychrophiles as those organisms that grow below 5 °C and to introduce the term cryophile, defined as organisms that can grow below 0 °C⁴³. So, whichever term you prefer, it is important to remember that in our efforts to communicate, scientists often impose strict restrictions that Nature does not create or recognize.

Psychrophiles have been isolated from a variety of natural (for example, polar regions, frozen lakes, and winter soils) and man-made (for example, refrigerators and cooling vents) environments. Microbes are generally subject to the temperature of their environment and as a consequence must find ways to adapt to the limitation placed on them by temperature.

To compensate for the negative effects of cold temperatures, organisms have developed several physiological adaptations, including regulating the fluidity of their membranes, synthesizing temperature-related chaperones, and producing antifreeze molecules⁴⁴. Psychrophiles regulate membrane fluidity through an increase in the number of branched-chain or unsaturated fatty acids or a shortening of the length of the fatty-acyl chains or both. Molecular chaperones are used to aid in the refolding of proteins and affect the levels of protein synthesis⁴⁵. Compatible solutes are used as cryoprotectors to lower the freezing point of the cytoplasm⁴⁶ and possibly prevent aggregation/denaturation of proteins, stabilize membranes, and scavenge free radicals in cold conditions⁴⁷. To reduce the damage caused by forming ice crystals, they may also use antifreeze or ice nucleation proteins^{48,49}. Antifreeze proteins act by binding seed ice crystals, thereby inhibiting their growth while ice nucleation proteins prevent the supercooling of water by ice crystal formation.

Structural proteins are also affected by temperature. Enzymes must overcome at least two obstacles in order to maintain activity at low temperatures: cold denaturation and slower reaction rates. Cold denaturation occurs at low temperatures because decreasing temperature results in more ordered water molecules surrounding a protein's surface, which results in their being less associated with the protein and pushing the system equilibrium toward the unfolded state⁴⁴. The second problem for enzymes at low temperatures is slower reaction rates. According to the Boltzmann equation, reaction rates increase with increasing temperature and decrease two- to three-fold for every 10 °C decrease. Therefore, if cold-active enzymes are to have activities on par with their mesophilic counterparts, they must have developed structural changes to overcome these thermodynamic barriers⁵⁰.

Temperature: thermophiles

All portions of thermophiles, as with psychrophiles, are constantly exposed to temperature; therefore, they have adapted all macromolecules (DNA, lipids, and proteins) and complexes (cell surface, ribosomes, RNA polymerase, and so on) to remain functional. Among the most studied aspects of adaptation for thermophiles are those found in proteins. Additional networks of hydrogen bonds, decreased length of surface loops, enhanced secondary structure propensity, higher core hydrophobicity, increased van der Waals interactions, ionic interactions, and increased packing density have all been shown to contribute to protein thermostability²². It was more recently shown that, in addition to using the above mechanisms, archaeal cells use a structure-stabilization approach (that is, proteins are more compact than their mesophilic homologs) but that bacterial cells use a sequence-stabilization approach (that is, a small number of strong interactions)⁵¹. Another well-studied aspect of adaptation is the lipid composition of thermophilic membranes. Certain organisms, like *Thermatoga maritima*, have novel/specific lipids (15, 16-dimethyl-30-glycerylox-triacontanedioic acid)¹¹. The ether-based lipids of archaea have also been shown to be resistant to hydrolysis at high temperatures. These are also found in meso- and psychrophiles and so are not a specific thermal adaptation. However, some thermophilic archaeal cells do contain a monolayer composed of a “fused lipid bi-layer” that has also been shown to resist hydrolysis at higher temperatures²⁴.

The DNA of thermophiles also has a thermal resistance in that it has positive supertwists added by reverse gyrase⁵². Additionally, an increase in GC base pairs in specific regions (stem-loops) has been shown to stabilize DNA. Archaeal thermophiles also have histones that are closely related to the H2A/B, H3, and H4 core histone of eukaryotes. The binding of these histones has been shown to increase the melting temperature of DNA⁵³.

Model organisms and major discoveries

Until very recently, a major drag on extremophile research was a lack of model organisms. Typically, after isolation, the genes of extremophilic organisms would be cloned and transformed into well-established model organisms like *Escherichia coli*. This was a necessary step to keep a forward momentum in research; however, it was also limiting as it allowed studies targeting only a small number of genes/proteins at a time and not the whole organism grown under *in situ* conditions. It also kept researchers from pushing limits as it meant that in-depth studies would always be relegated to genes/proteins from organisms that could “conform” to being studied in mesophilic model systems. Omics experiments have alleviated some of these problems; however, they cannot truly replace studies that need to be carried out *in vivo*. For these experiments, true model organisms that are extremophiles are needed. Fortunately, there are now model organisms for all extremophile groups mentioned in this review; they include *Leptospirillum ferriphilum* (acidophile)^{54,55}, *Sulfolobus solfataricus* (acidophile and thermophile)^{56,57}, *Natronomonas pharaonis* (alkaliphile and halophile)^{58,59}, *Bacillus halodurans* (halophile)^{60,61}, *H. volcanii* (halophile)^{62,63}, *Halobacterium* sp. NRC-1 (halophile and radiophile)^{64,65}, *Walleimia ichthyophaga* (halophile)^{66,67}, *D. radiodurans* (radiophile)^{68,69}, *Thermococcus barophilus*

(piezophile)^{70,71}, *Halorubrum lacusprofundi* (halophile and psychrophile)^{72,73}, *Pseudoalteromonas haloplanktis* (psychrophile)^{74,75}, *Thermococcus kodakarensis* (thermophile)^{76,77}, and *Thermus thermophilus* (thermophile)^{78,79}.

Other than detailed knowledge about the mechanisms that allow these organisms to thrive, studies of these model organisms have also led to some significant discoveries. One of these was the finding of a novel gene regulation mechanism found in *Halobacterium* sp. NRC-1⁸⁰ that was also shown to exist in other archaeal cells with multiple TATA-binding proteins (TBPs) or transcription factor B (TFBs) or both^{81,82}. This mechanism uses pairs of TBPs and TFBs to express/regulate specific sets of genes and is similar to a mechanism found in the Metazoa^{83,84}. Another is the primacy of biological function (that is, proteome/lipidome) over information (that is, genome). Studies performed using *D. radiodurans* showed that cells can function for some time after losing their genome (that is, complete fragmentation of the genome by gamma irradiation); however, the same is not true for a cell that loses its proteome⁸⁵. Just recently, there were reports using *T. thermophilus* and other organisms suggesting that the 16S genes of prokaryotes, once thought to be species-specific and so used for decades for phylogenetic analysis, are promiscuous and horizontally transferred⁷⁹.

Biotechnology applications

There have been four great success stories for the application of extremophiles and their products: the application of DNA polymerases from thermophiles to polymerase chain reaction, the use of thermophilic organisms/enzymes to produce biofuels, the use of acidophilic and halophilic organisms/enzymes in biomineralization, and the use of halophilic organisms/enzymes to produce carotenoids⁸⁵. Extremophilic lipases/proteases have also been used extensively in detergents, specifically designed for cold water washing, and fine chemical synthesis⁸⁶.

In addition to the current applications, several other possibilities have been suggested over the years. These include using cold-active beta-galactosidase to hydrolyze lactose in dairy products⁸⁷ and using thermophilic and alkaliphilic starch-hydrolyzing enzymes (for example, alpha-amylase, glucoamylase, and pullulanase) to make a range of products like ethanol and high-fructose corn syrup^{85,88}. Additionally, the products of extremophiles like sugars (for example, trehalose) and amino acid derivatives (for example, ectoine) as stabilizers for antibodies and vaccines or as skin care products^{24,89} have been suggested. Several extremophiles produce polyhydroxyalkanoates (PHAs),

which are a heterogeneous group of polyesters and can be used to generate bioplastics⁹⁰. Finally, several extremophiles have been tested and shown to be helpful, at least on a small scale, in bioremediation efforts against heavy metals, radioactive isotopes, hydrocarbons, and polychlorobiphenyls (PCBs) to name a few⁹¹.

Astrobiology and origin-of-life theories

In addition to teaching us about the limits of life on Earth, extremophiles can tell us about the limits of life in the Universe. We should not expect that life, if found, in other parts of the Universe will necessarily be the same as on Earth; however, the laws of chemistry and physics we know suggest that life requires building blocks (for example, nucleotides and amino acids), an energy source (for example, solar radiation or redox reactions), and a liquid solvent (for example, water)⁹². The known limits of life on Earth are mentioned in the introduction above; however, scientists keep finding exceptions and it is likely that the absolute limits, if they exist, have not been discovered. Indeed, the true limits of life are possibly the availability of water, building blocks, and an energy source as opposed to a specific temperature, pH, and so on⁹³.

Although extremophiles are often discussed by the predominant environmental pressure, in reality there are typically multiple extremes (for example, cold and high salinity or heat and acid). When considering these environments on Earth and other extraplanetary bodies, we must look increasingly toward polyextremophiles as the true model organisms for astrobiology⁹⁴.

At first blush, it might seem that the ever-extending boundary of life is a boon to detecting life on extraplanetary bodies. Indeed, that is the case but only to a limited extent. It is true that, as knowledge expands, we extend the types of places we expect to find life. However, one should remember that, with the exception of the Oceans, the environments where extremophiles thrive are quite far from a planetary mean⁹⁵. Additionally, organisms may be “active” (that is, metabolizing, dividing, and so on) only for brief moments in transient environments (for example, seasonal rains/water)⁹⁶. As such, many extraplanetary bodies that could support life are likely to be overlooked. It also means that “one-off” observations of extraplanetary bodies are highly unlikely to find signatures of life.

Abbreviations

TBP, TATA-binding protein; TFB, transcription factor B; UV, ultraviolet

References

1. Cavicchioli R, Charlton T, Ertan H, et al.: **Biotechnological uses of enzymes from psychrophiles**. *Microb Biotechnol*. 2011; 4(4): 449–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
2. Auman AJ, Breeze JL, Gosink JJ, et al.: ***Psychromonas ingrahamii* sp. nov., a novel gas vacuolate, psychrophilic bacterium isolated from Arctic polar sea**

ice. *Int J Syst Evol Microbiol*. 2006; 56(Pt 5): 1001–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

3. Johnson DB, Kanao T, Hedrich S: **Redox Transformations of Iron at Extremely Low pH: Fundamental and Applied Aspects**. *Front Microbiol*. 2012; 3: 96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

4. DeVeaux LC, Müller JA, Smith J, *et al.*: **Extremely radiation-resistant mutants of a halophilic archaeon with increased single-stranded DNA-binding protein (RPA) gene expression.** *Radiat Res.* 2007; **168**(4): 507–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Horikoshi K: **Alkaliphiles: some applications of their products for biotechnology.** *Microbiol Mol Biol Rev.* 1999; **63**(4): 735–50, table of contents.
[PubMed Abstract](#) | [Free Full Text](#)
6. Kato C, Li L, Nogi Y, *et al.*: **Extremely barophilic bacteria isolated from the Mariana Trench, Challenger Deep, at a depth of 11,000 meters.** *Appl Environ Microbiol.* 1998; **64**(4): 1510–3.
[PubMed Abstract](#) | [Free Full Text](#)
7. Moran-Reyna A, Coker JA: **The effects of extremes of pH on the growth and transcriptomic profiles of three haloarchaea [version 2; peer review: 3 approved].** *F1000Res.* 2014; **3**: 168.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. **F** Takai K, Nakamura K, Toki T, *et al.*: **Cell proliferation at 122 degrees C and isotopically heavy CH₄ production by a hyperthermophilic methanogen under high-pressure cultivation.** *Proc Natl Acad Sci U S A.* 2008; **105**(31): 10949–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
9. Imshenetsky AA, Lysenko SV, Kazakov GA: **Upper boundary of the biosphere.** *Appl Environ Microbiol.* 1978; **35**(1): 1–5.
[PubMed Abstract](#) | [Free Full Text](#)
10. Takai K, Moser DP, DeFlaun M, *et al.*: **Archaeal diversity in waters from deep South African gold mines.** *Appl Environ Microbiol.* 2001; **67**(12): 5750–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. **F** Siliakus MF, van der Oost J, Kengen SWM: **Adaptations of archaeal and bacterial membranes to variations in temperature, pH and pressure.** *Extremophiles.* 2017; **21**(4): 651–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
12. Konings WN, Albers SV, Konings S, *et al.*: **The cell membrane plays a crucial role in survival of bacteria and archaea in extreme environments.** *Antonie Van Leeuwenhoek.* 2002; **81**(1–4): 61–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Baker-Austin C, Dopson M: **Life in acid: pH homeostasis in acidophiles.** *Trends Microbiol.* 2007; **15**(4): 165–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Amaro AM, Chamorro D, Seeger M, *et al.*: **Effect of external pH perturbations on *in vivo* protein synthesis by the acidophilic bacterium *Thiobacillus ferrooxidans*.** *J Bacteriol.* 1991; **173**(2): 910–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Michels M, Bakker EP: **Generation of a large, protonophore-sensitive proton motive force and pH difference in the acidophilic bacteria *Thermoplasma acidophilum* and *Bacillus acidocaldarius*.** *J Bacteriol.* 1985; **161**(1): 231–7.
[PubMed Abstract](#) | [Free Full Text](#)
16. Pikuta EV, Hoover RB, Tang J: **Microbial extremophiles at the limits of life.** *Crit Rev Microbiol.* 2008; **33**(3): 183–209.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Slonczewski JL, Fujisawa M, Dopson M, *et al.*: **Cytoplasmic pH measurement and homeostasis in bacteria and archaea.** *Adv Microb Physiol.* 2009; **55**: 1–79, 317.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. **F** Mamo G: **Challenges and Adaptations of Life in Alkaline Habitats.** *Adv Biochem Eng Biotechnol.* 2019.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
19. Preiss L, Hicks DB, Suzuki S, *et al.*: **Alkaliphilic Bacteria with Impact on Industrial Applications, Concepts of Early Life Forms, and Bioenergetics of ATP Synthesis.** *Front Bioeng Biotechnol.* 2015; **3**: 75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Krulwich TA, Sachs G, Padan E: **Molecular aspects of bacterial pH sensing and homeostasis.** *Nat Rev Microbiol.* 2011; **9**(5): 330–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Yin J, Chen JC, Wu Q, *et al.*: **Halophiles, coming stars for industrial biotechnology.** *Biotechnol Adv.* 2015; **33**(7): 1433–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. **F** Brininger C, Spradlin S, Cobani L, *et al.*: **The more adaptive to change, the more likely you are to survive: Protein adaptation in extremophiles.** *Semin Cell Dev Biol.* 2018; **84**: 158–69.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
23. DasSarma S, DasSarma P: **Halophiles and their enzymes: negativity put to good use.** *Curr Opin Microbiol.* 2015; **25**: 120–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. DasSarma S, Coker JA, DasSarma P: **“Archaea”.** In *Encyclopedia of Microbiology*. Oxford Academic Press, 2009; 1–23.
[Publisher Full Text](#)
25. Oren A: **Microbial life at high salt concentrations: phylogenetic and metabolic diversity.** *Saline Systems.* 2008; **4**: 2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Roberts MF: **Organic compatible solutes of halotolerant and halophilic microorganisms.** *Saline Systems.* 2005; **1**: 5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Slade D, Radman M: **Oxidative stress resistance in *Deinococcus radiodurans*.** *Microbiol Mol Biol Rev.* 2011; **75**(1): 133–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Pavlopoulou A, Savva GD, Louka M, *et al.*: **Unraveling the mechanisms of extreme radioresistance in prokaryotes: Lessons from nature.** *Mutat Res Rev Mutat Res.* 2016; **767**: 92–107.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Cox MM, Keck JL, Battista JR: **Rising from the Ashes: DNA Repair in *Deinococcus radiodurans*.** *PLoS Genet.* 2010; **6**(1): e1000815.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Lockhart JS, DeVeaux LC: **The essential role of the *Deinococcus radiodurans* *ssb* gene in cell survival and radiation tolerance.** *PLoS One.* 2013; **8**(8): e71651.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Minsky A, Shimoni E, Englander J: **Ring-like nucleoids and DNA repair through error-free nonhomologous end joining in *Deinococcus radiodurans*.** *J Bacteriol.* 2006; **188**(17): 6047–51; discussion 6052.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. **F** Daly MJ, Gaidamakova EK, Matrosovaya VY, *et al.*: **Protein oxidation implicated as the primary determinant of bacterial radioresistance.** *PLoS Biol.* 2007; **5**(4): e92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
33. **F** Jones DL, Baxter BK: **DNA Repair and Photoprotection: Mechanisms of Overcoming Environmental Ultraviolet Radiation Exposure in Halophilic Archaea.** *Front Microbiol.* 2017; **8**: 1882.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
34. Quintana-Cabrera R, Fernandez-Fernandez S, Bobo-Jimenez V, *et al.*: **γ -Glutamylcysteine detoxifies reactive oxygen species by acting as glutathione peroxidase-1 cofactor.** *Nat Commun.* 2012; **3**: 718.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. Jebbar M, Franzetti B, Girard E, *et al.*: **Microbial diversity and adaptation to high hydrostatic pressure in deep-sea hydrothermal vents prokaryotes.** *Extremophiles.* 2015; **19**(4): 721–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Kaneshiro SM, Clark DS: **Pressure effects on the composition and thermal behavior of lipids from the deep-sea thermophile *Methanococcus jannaschii*.** *J Bacteriol.* 1995; **177**(13): 3668–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Michoud G, Jebbar M: **High hydrostatic pressure adaptive strategies in an obligate piezophile *Pyrococcus yayanosii*.** *Sci Rep.* 2016; **6**: 27289.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Vannier P, Michoud G, Oger P, *et al.*: **Genome expression of *Thermococcus barophilus* and *Thermococcus kodakarensis* in response to different hydrostatic pressure conditions.** *Res Microbiol.* 2015; **166**(9): 717–25.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Amrani A, Bergon A, Holota H, *et al.*: **Transcriptomics reveal several gene expression patterns in the piezophile *Desulfovibrio hydrothermalis* in response to hydrostatic pressure.** *PLoS One.* 2014; **9**(9): e106831.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Morita RY: **Psychrophilic bacteria.** *Bacteriol Rev.* 1975; **39**(2): 144–67.
[PubMed Abstract](#) | [Free Full Text](#)
41. Neidhardt FC, Ingraham JL, Schaechter M: **Physiology of the bacterial cell: a molecular approach.** Sunderland: Sinauer Associates. 1990.
[Reference Source](#)
42. **F** Raymond-Bouchard I, Tremblay J, Altschuler I, *et al.*: **Comparative Transcriptomics of Cold Growth and Adaptive Features of a Eury- and Steno-Psychrophile.** *Front Microbiol.* 2018; **9**: 1565.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
43. Coker JA: **Structure and function relationships in the cold-active beta-galactosidase, BgaS, examining theories of enzyme cold-adaptation.** 2004.
[Reference Source](#)
44. de Maayer P, Anderson D, Cary C, *et al.*: **Some like it cold: understanding the survival strategies of psychrophiles.** *EMBO Rep.* 2014; **15**(5): 508–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
45. Math RK, Jin HM, Kim JM, *et al.*: **Comparative genomics reveals adaptation by *Alteromonas* sp. SN2 to marine tidal-flat conditions: cold tolerance and aromatic hydrocarbon metabolism.** *PLoS One.* 2012; **7**(4): e35784.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. Casanueva A, Tuffin M, Cary C, *et al.*: **Molecular adaptations to psychrophily: the impact of ‘omic’ technologies.** *Trends Microbiol.* 2010; **18**(8): 374–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
47. **F** Kandror O, DeLeon A, Goldberg AL: **Trehalose synthesis is induced upon exposure of *Escherichia coli* to cold and is essential for viability at low temperatures.** *Proc Natl Acad Sci U S A.* 2002; **99**(15): 9727–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
48. Celik Y, Drori R, Pertaya-Braun N, *et al.*: **Microfluidic experiments reveal that antifreeze proteins bound to ice crystals suffice to prevent their growth.** *Proc Natl Acad Sci U S A.* 2013; **110**(4): 1309–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
49. Kawahara H: **The structures and functions of ice crystal-controlling proteins from bacteria.** *J Biosci Bioeng.* 2002; **94**(6): 492–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
50. Santiago M, Ramirez-Sarmiento CA, Zamora RA, *et al.*: **Discovery, Molecular Mechanisms, and Industrial Applications of Cold-Active Enzymes.** *Front Microbiol.* 2016; **7**: 1408.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
51. Berezovsky IN, Shakhnovich EI: **Physics and evolution of thermophilic adaptation.** *Proc Natl Acad Sci U S A.* 2005; **102**(36): 12742–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
52. Jamroze A, Perugini G, Valenti A, *et al.*: **The Reverse Gyrase from *Pyrobaculum***

- calidifontis**, a Novel Extremely Thermophilic DNA Topoisomerase Endowed with DNA Unwinding and Annealing Activities. *J Biol Chem.* 2014; 289(6): 3231–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
53. Stetter KO: **Extremophiles and their adaptation to hot environments.** *FEBS Lett.* 1999; 452(1–2): 22–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
 54. Christel S, Herold M, Bellenberg S, *et al.*: **Multi-omics Reveals the Lifestyle of the Acidophilic, Mineral-Oxidizing Model Species *Leptospirillum ferriphilum*.** *Appl Environ Microbiol.* 2018; 84(3): pii: e02091-17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 55. Cardenas JP, Quatrini R, Homes DS: **Progress in acidophile genomics.** In *Acidophiles: life in extremely acidic environments.* Norfolk, Caister Academic Press, 2016; 179–197.
[Publisher Full Text](#)
 56. Quehenberger J, Shen L, Albers SV, *et al.*: **Sulfolobus - A Potential Key Organism in Future Biotechnology.** *Front Microbiol.* 2017; 8: 2474.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 57. Figueiredo AS, Kouril T, Esser D, *et al.*: **Systems biology of the modified branched Entner-Doudoroff pathway in *Sulfolobus solfataricus*.** *PLoS One.* 2017; 12(7): e0180331.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 58. Falb M, Pfeiffer F, Palm P, *et al.*: **Living with two extremes: Conclusions from the genome sequence of *Natronomonas pharaonis*.** *Genome Res.* 2005; 15(10): 1336–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 59. Oren: **Two centuries of microbiological research in the Wadi Natrun, Egypt: a model system for the study of the ecology, physiology, and taxonomy of the haloalkaliphilic microorganisms.** In *Polymicrobiology.* Springer, 2013; 101–119.
[Publisher Full Text](#)
 60. Van-Thuoc D, Hashim SO, Hatti-Kaul R, *et al.*: **Ecto-enzyme-mediated protection of enzyme from the effect of pH and temperature stress: A study using *Bacillus halodurans* xylanase as a model.** *Appl Microbiol Biotechnol.* 2013; 97(14): 6271–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 61. Takami H, Nakasone K, Takaki Y, *et al.*: **Complete genome sequence of the alkaliphilic bacterium *Bacillus halodurans* and genomic sequence comparison with *Bacillus subtilis*.** *Nucleic Acids Res.* 2000; 28(21): 4317–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 62. Hartman AL, Norais C, Badger JH, *et al.*: **The Complete Genome Sequence of *Haloferax volcanii* DS2, a Model Archaeon.** *PLoS One.* 2010; 5(3): e9605.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 63. Allers T, Barak S, Liddell S, *et al.*: **Improved Strains and Plasmid Vectors for Conditional Overexpression of His-Tagged Proteins in *Haloferax volcanii*.** *Appl Environ Microbiol.* 2010; 76(6): 1759–69.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 64. Kennedy SP, Ng WV, Salzberg SL, *et al.*: **Understanding the adaptation of *Halobacterium* species NRC-1 to its extreme environment through computational analysis of its genome sequence.** *Genome Res.* 2001; 11(10): 1641–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 65. Coker JA, DasSarma P, Capes M, *et al.*: **Multiple Replication Origins of *Halobacterium* sp. Strain NRC-1: Properties of the Conserved *orc7*-Dependent *oriC1*.** *J Bacteriol.* 2009; 191(16): 5253–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 66. Zajc J, Liu Y, Dai W, *et al.*: **Genome and transcriptome sequencing of the halophilic fungus *Wallemia ichthyophaga*: haloadaptations present and absent.** *BMC Genomics.* 2013; 14: 617.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 67. Plemenitaš A, Konte T, Gostinčar C, *et al.*: **Transport Systems in Halophilic Fungi.** *Adv Exp Med Biol.* 2016; 892: 307–25.
[PubMed Abstract](#) | [Publisher Full Text](#)
 68. Krisko A, Radman M: **Biology of Extreme Radiation Resistance: The Way of *Deinococcus radiodurans*.** *Cold Spring Harb Perspect Biol.* 2013; 5(7): pii: a012765.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 69. Zhou Y, Shen P, Lan Q, *et al.*: **High-coverage proteomics reveals methionine auxotrophy in *Deinococcus radiodurans*.** *Proteomics.* 2017; 17(13–14): 1700072.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 70. Birien T, Thiel A, Henneke G, *et al.*: **Development of an Effective 6-Methylpurine Counterselection Marker for Genetic Manipulation in *Thermococcus barophilus*.** *Genes (Basel).* 2018; 9(2): pii: E77.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 71. Oger P, Sokolova TG, Kozhevnikova DA, *et al.*: **Complete Genome Sequence of the Hyperthermophilic and Piezophilic Archaeon *Thermococcus barophilus* Ch5, Capable of Growth at the Expense of Hydrogenogenesis from Carbon Monoxide and Formate.** *Genome Announc.* 2016; 4(1): pii: e01534-15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 72. Liao Y, Williams TJ, Walsh JC, *et al.*: **Developing a genetic manipulation system for the Antarctic archaeon, *Halorubrum lacusprofundi*: investigating acetamidase gene function.** *Sci Rep.* 2016; 6: 34639.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 73. Tschitschko B, Williams TJ, Allen MA, *et al.*: **Ecophysiological Distinctions of Haloarchaea from a Hypersaline Antarctic Lake as Determined by Metaproteomics.** *Appl Environ Microbiol.* 2016; 82(11): 3165–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 74. Parrilli E, Tedesco P, Fondi M, *et al.*: **The art of adapting to extreme environments: The model system *Pseudoalteromonas*.** *Phys Life Rev.* 2019; pii: S1571-0645(19)30066-1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 75. Mocali S, Chiellini C, Fabiani A, *et al.*: **Ecology of cold environments: new insights of bacterial metabolic adaptation through an integrated genomic-phenomic approach.** *Sci Rep.* 2017; 7(1): 839.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 76. Aslam M, Horiuchi A, Simons JR, *et al.*: **Engineering of the Hyperthermophilic Archaeon *Thermococcus kodakarensis* for Chitin-Dependent Hydrogen Production.** *Appl Environ Microbiol.* 2017; 83(15): pii: e00280-17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 77. Atomi H, Reeve J: **Microbe Profile: *Thermococcus kodakarensis*: the model hyperthermophilic archaeon.** *Microbiology.* 2019; 165(11): 1166–1168.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 78. Alvarez L, Bricio C, Blesa A, *et al.*: **Transferable denitrification capability of *Thermus thermophilus*.** *Appl Environ Microbiol.* 2014; 80(1): 19–28.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 79. Miyazaki K, Tomariguchi N: **Occurrence of randomly recombined functional 16S rRNA genes in *Thermus thermophilus* suggests genetic interoperability and promiscuity of bacterial 16S rRNAs.** *Sci Rep.* 2019; 9(1): 11233.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 80. Coker JA, DasSarma S: **Genetic and transcriptomic analysis of transcription factor genes in the model halophilic Archaeon: coordinate action of TbpD and TfbA.** *BMC Genet.* 2007; 8: 61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 81. Kramm K, Engel C, Grohmann D: **Transcription initiation factor TBP: old friend new questions.** *Biochem Soc Trans.* 2019; 47(1): 411–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 82. Reichlen MJ, Murakami KS, Ferry JG: **Functional analysis of the three TATA binding protein homologs in *Methanosarcina acetivorans*.** *J Bacteriol.* 2010; 192(6): 1511–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 83. Wang YL, Duttkie SH, Chen K, *et al.*: **TRF2, but not TBP, mediates the transcription of ribosomal protein genes.** *Genes Dev.* 2014; 28(14): 1550–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 84. D'Alessio JA, Wright KJ, Tjian R: **Shifting players and paradigms in cell-specific transcription.** *Mol Cell.* 2009; 36(6): 924–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 85. Coker JA: **Extremophiles and biotechnology: current uses and prospects [version 1; peer review: 2 approved].** *F1000Res.* 2016; 5: pii: F1000 Faculty Rev-396.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 86. Kavitha M: **Cold active lipases – an update.** *Front Life Sci.* 2016; 9(3): 226–38.
[Publisher Full Text](#)
 87. Coker JA, Brenchley JE: **Protein engineering of a cold-active beta-galactosidase from *Arthrobacter* sp. SB to increase lactose hydrolysis reveals new sites affecting low temperature activity.** *Extremophiles.* 2006; 10(6): 515–24.
[PubMed Abstract](#) | [Publisher Full Text](#)
 88. Schiraldi C, Giuliano M, De Rosa M: **Perspectives on biotechnological applications of archaea.** *Archaea.* 2002; 1(2): 75–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 89. Dalmaso GZ, Ferreira D, Vermelho AB: **Marine extremophiles: a source of hydrolases for biotechnological applications.** *Mar Drugs.* 2015; 13(4): 1925–65.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 90. DasSarma P, Coker JA, Huse V, *et al.*: **"Halophiles, industrial applications".** In: *Encyclopedia of Industrial Biotechnology: Bioprocess, Bioseparation, and Cell Technology.* John Wiley & Sons, Inc. 2010.
[Publisher Full Text](#)
 91. Orellana R, Macaya C, Bravo G, *et al.*: **Living at the Frontiers of Life: Extremophiles in Chile and Their Potential for Bioremediation.** *Front Microbiol.* 2018; 9: 2309.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 92. Schwieterman EW, Kiang NY, Parenteau MN, *et al.*: **Exoplanet Biosignatures: A Review of Remotely Detectable Signs of Life.** *Astrobiology.* 2018; 18(6): 663–708.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 93. Merino N, Aronson HS, Bojanova DP, *et al.*: **Living at the Extremes: Extremophiles and the Limits of Life in a Planetary Context.** *Front Microbiol.* 2019; 10: 780.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 94. DasSarma S, DasSarma P, Laye VJ, *et al.*: **Extremophilic models for astrobiology: haloarchaeal survival strategies and pigments for remote sensing.** *Extremophiles.* 2019; 1–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 95. Reid IN, Sparks WB, Lubow S, *et al.*: **Terrestrial models for extraterrestrial life: Methanogens and halophiles at Martian temperatures.** *Int J Astrobiol.* 2006; 5(2): 89–97.
[Publisher Full Text](#)
 96. Davila AF, Gómez-Silva B, de los Rios A, *et al.*: **Facilitation of endolithic microbial survival in the hyperarid core of the Atacama Desert by mineral deliquescence.** *J Geophys Res.* 2008; 113(G1): n/a-n/a.
[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Linda DeVeaux**

Department of Biology, New Mexico Institute of Mining and Technology, Socorro, NM, USA

Competing Interests: No competing interests were disclosed.

2. **Noah Mesbah**

Department of Biochemistry, Faculty of Pharmacy, Suez Canal University, Ismailia, Egypt

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research