RESEARCH ARTICLE

REVISED Predictive statistical model for the factors associated with prenatal depression among pregnant adolescents in Maha Sarakham province, Thailand [version 2; peer review: 1 approved, 1 not approved]

Bhanwasa Jantasin¹ , Witaya Yoosook¹, Supawadee Thaewpia²

¹Faculty of Public Health, Mahasarakham University, Maha Sarakham, 44160, Thailand

²Maternal and Child Department, Boromarajonani College of Nursing Khonkaen, Khon Kaen, 40000, Thailand

V2 First published: 14 Nov 2019, 8:1921
<https://doi.org/10.12688/f1000research.21007.1>
 Latest published: 29 Apr 2020, 8:1921
<https://doi.org/10.12688/f1000research.21007.2>

Abstract

Background: The rate of adolescent pregnancy has steadily increased in Thailand over the years and depression is considered an important psychological problem in many pregnant adolescents. The aims of this study were to determine the proportion of pregnant adolescents with depressive symptoms and to identify a predictive statistical model for the factors related to prenatal depression among adolescents at antenatal care clinics in 10 community health centers in Maha Sarakham Province.

Methods: A cross-sectional study was conducted in 230 randomly selected participants who were pregnant adolescents who volunteered to participate in the research, those aged ≤ 19 years, and had an estimated gestational age of ≤ 18 weeks. Those with severe obstetrics complications or a psychiatric disorder were excluded. The subjects were asked to complete a demographic questionnaire and the Thai version of the Edinburgh Postnatal Depression Scale. The mean age of participants was 17.0 (± 1.4) years.

Results: Overall, 43.9% of the adolescents had a cut off score of ≥ 11 , suggestive of prenatal depressive symptoms. Multivariate analysis identified that pregnant adolescents at a primary school education level were 6.5 times more likely to experience depression symptoms than those with a diploma. Those who were emotionally supported by their husband or partner compared to relatives had a five times higher chance of presenting depressive symptoms, and those with a perceived bad relationship with their husband or partner had seven times higher. The model was well-calibrated (goodness-of-fit test, $p = 0.675$) and had high discriminative power (ROC curve = 0.73 (95%CI = 0.67 to 0.79)).

Conclusions: More attention is needed to reduce the known problems with pregnant adolescents and to ensure they continue with their

Open Peer Review

Approval Status

	1	2
version 2		
(revision)		
29 Apr 2020	view	view
version 1		
14 Nov 2019	view	

1. **Felix Emeka Anyiam** , University of Port Harcourt, Port Harcourt, Nigeria
2. **Caroline Trumpff**, Columbia University Medical Center, New York, USA

Any reports and responses or comments on the article can be found at the end of the article.

education. Lastly, public education to support mental health initiatives for individuals and families should be encouraged.

Keywords

Adolescent pregnancy, Community hospital, Prenatal depression, Thai Edinburgh Postnatal Depression Scale

This article is included in the **Developmental Psychology and Cognition** gateway.

Corresponding author: Bhanwasa Jantasin (bhanwasa@smnc.ac.th)

Author roles: **Jantasin B:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Resources, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Yoosook W:** Supervision; **Thaewpia S:** Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Jantasin B *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Jantasin B, Yoosook W and Thaewpia S. **Predictive statistical model for the factors associated with prenatal depression among pregnant adolescents in Maha Sarakham province, Thailand [version 2; peer review: 1 approved, 1 not approved]** F1000Research 2020, 8:1921 <https://doi.org/10.12688/f1000research.21007.2>

First published: 14 Nov 2019, 8:1921 <https://doi.org/10.12688/f1000research.21007.1>

REVISED Amendments from Version 1

We have revised Table 2 in the result section and added some missing words according to suggestion from the reviewer.

Any further responses from the reviewers can be found at the end of the article

Introduction

Adolescents can develop health problems during pregnancy that adversely affect both their physical and psychological health (UNFPA, 2014a). Pregnant adolescents commonly experience depressive symptoms (Lancaster *et al.*, 2010); the prevalence has been estimated to be approximately 20% (Abajobir *et al.*, 2016; Nasreen *et al.*, 2011; Roomruangwong & Epperson, 2011). Pregnant adolescents with depression and their infants are at a higher risk of increased morbidity (Tzilos *et al.*, 2012), resulting in a greater risk for preterm labor and intrauterine fetal growth retardation (Field, 2017; Grote *et al.*, 2010; Orr *et al.*, 2007). Depressed pregnant adolescents also have a greater risk of suicidal ideations and suicide attempts than those without depression (Hodgkinson *et al.*, 2010).

Risk factors predisposing adult pregnant women to depressive symptoms include daily maternal stress, loss of self-confidence, and inadequate social support and emotional coping (Da Costa *et al.*, 2000; Dennis *et al.*, 2007). Risks factors for depression among pregnant adolescents include not only those listed for adult pregnant women but also include unplanned pregnancy, low education, non-marital status, alcohol and drug use, inadequate income, and a history of emotional, sexual, and physical abuse (Koniak-Griffin *et al.*, 1996; Meltzer-Brody *et al.*, 2013; Siegel & Brandon, 2014; Uthaisaisanwong *et al.*, 2015). Because prenatal depression in adolescence is a significant factor for predicting postpartum depression (Dennis *et al.*, 2007), the ability to identify prenatal depression at the earliest stage of pregnancy is particularly important for the health and well-being of mothers and infants and for public health in general.

Thailand has a high rate of adolescent pregnancy (Ministry of Public Health, 2017). Studies by Limlomwongse & Liabsuetrakul (2006), and Uthaisaisanwong *et al.* (2015) in Bangkok and the South of Thailand have shown that the incidence of prenatal depression among adolescents is 46% and 20.5%, respectively. Little research, however, has been conducted in northeastern Thailand, especially in Maha Sarakham province, where the rate of adolescent pregnancy is reported to be high (Ministry of Public Health, 2017). Therefore, the aim of this study was to estimate the proportion of prenatal depressive symptoms among adolescents in the Maha Sarakham Province, Thailand, and to identify factors associated with prenatal depression symptoms in adolescents.

Methods

Study subjects and data collection

This cross-sectional study was conducted from July to October 2017 with 230 pregnant adolescents. The participants

were selected using simple randomization in order to avoid sampling bias. Every week, two to three pregnant adolescents were recruited from each antenatal clinic in 10 community hospitals in the Maha Sarakham province, Thailand by randomizing HN numbers (patient identification numbers were drawn as lots) among all of the pregnant adolescents who met the inclusion criteria. Inclusion criteria were 1) pregnant adolescents who volunteered to participate, 2) those ≤ 19 years of age, 3) and those with an estimated gestational age of ≤ 18 weeks. The exclusion criteria were pregnant adolescents with obstetric complications or those who had been diagnosed with a psychiatric disorder. Initially, a sample size of 209 adolescents was calculated (Hsieh *et al.*, 1998) assuming a prenatal depression prevalence of 46% (Uthaisaisanwong *et al.*, 2015) with an alpha error of 0.05 and a power of 80%. An extra 21 adolescents (10%) were added to compensate for possible incomplete data, giving a total sample of 230 participants. All participants were recruited in person during their first appointment at the antenatal clinic. After receiving informed consent, participants provided their demographic and socio-economic information on a written questionnaire (available as *Extended data*; Jantasin *et al.*, 2019b; Jantasin *et al.*, 2019c) then completed the Thai version of the Edinburgh Postnatal Depression Scale (Pitanupong *et al.*, 2007).

The Edinburgh Postnatal Depression Scale (EPDS) is an assessment tool available in Thailand to screen for depressive symptoms. Originally developed to detect possible depressive symptoms in the postpartum period (Cox *et al.*, 1987), the EPDS has been successfully used to identify depressive symptoms in prenatal women (Eberhard-Gran *et al.*, 2001; Lee *et al.*, 2004). In 2003, the EPDS was translated to Thai by Pitanupong *et al.* (2007). Three doctorally prepared Thai experts in midwifery reviewed the translation for content validity and approved all items without recommendations. Published psychometric testing supports the 10-item self-assessment questionnaire's validity and reliability among Thai pregnant women (Pitanupong *et al.*, 2007; Vacharaporn *et al.*, 2003). Preliminary testing for internal reliability using 30 people, not part of the study, showed a Cronbach's alpha = 0.76.

Each of the ten EPDS items has four response options ranging from 0 to 3. The response descriptors vary somewhat depending on the item's wording. The participants choose the descriptor most closely indicating their emotions during the prior seven days. Seven items require reverse scoring so that higher total scores are more suggestive of depressive symptoms. A total score can range between 0 and 30. The cutoff score for the EPDS that indicates the presence of depression symptoms has been reported between 10 and 14 (Bolton *et al.*, 1998; Evans *et al.*, 2001). We defined an EPDS score of 11 or greater as a positive screen for depressive symptoms (Uthaisaisanwong *et al.*, 2015).

Statistical analysis

Data analysis were conducted with descriptive statistics to present demographic data. Multivariable analysis through unconditional multiple logistic regression was used to identify factors associated with prenatal depression among pregnant

adolescents and to determine potential associations of socio-demographic information with prenatal depressive symptoms (EPDS $\geq 11=1$ and $<11=0$). Crude analysis through simple logistic regression analysis was used. After that, all variables with p -values less than 0.25 (Hosmer & Lemeshow, 2000) were selected to include an initial regression model for the assessment of multicollinearity. Then, backward elimination for a fitted regression model was performed. The model was developed and its goodness-of-fit was evaluated by determining the predictive accuracy and cut-off that was suitable for clustering towards depressive symptoms among pregnant adolescents using a receiver operating characteristics (ROC) curve. Statistical significance was set at $p < 0.05$. Analyses were done with Statistical Package for the Social Sciences (SPSS) software, version 18.0 (SPSS®, Chicago, USA).

Ethical considerations

The study was approved by the Institutional Ethics Committee at Mahasarakham University, Thailand. (Ref No. 054/2560; 21 July 2017). Adolescents of legal age offered their informed consent volunteering to participate in the study. Parents of adolescents younger than 18 years of age provided written informed consent prior to their child's participation.

Results

Association between depressive symptoms and participant variables

Of the 230 participants, 101 adolescents (43.9%) met the EPDS cutoff score of ≥ 11 suggestive of prenatal depressive symptoms. The remaining 129 adolescents (56.1%) scored below the cutoff score (Table 1).

The mean age of the 101 adolescents was 17.0 (SD $\pm$ 1.4) years old. The majority of those were >15 years old ($n=80$, 80.0%) and 21 (20.0%) were ≤ 15 of age. In bivariate analysis, depressive symptoms were significantly correlated with receiving less education, being a single mother with one supporter, having a husband as a supporter, and having a poor relationship with a husband, their family and friends. Pregnant adolescents who only finished primary school were 7.3 times more likely to have prenatal depressive symptoms than those with education to diploma level. In terms of marital status, the adolescents who were single mothers were at 9.4 times higher risk of depressive symptoms than those living together with their partner, with or without being married. Pregnant adolescents who had one supporter were 2.8 times more likely to be at risk of depressive symptoms compared to those with three supporters. The adolescents who had a husband as a supporter tended to be 3.7 times higher at risk of depressive symptoms when compared to those who were supported by relatives. Moreover, pregnant adolescents who reported a bad relationship with their husband, family, and friends were 5.5 times, 2.2 times, 2.8 times higher at risk of depressive symptoms than those with good relationship, respectively (Table 1). Responses to each question from each participant are available as *Underlying data* (Jantasin *et al.*, 2019a).

Linear regression analysis

According to Table 1, eight variables with a p -value of less than 0.25 were selected and included in the initial regression model. From the multiple logistic regression analysis, there were only three factors that were significantly associated with prenatal depressive symptoms: level of education, the perceived support from a husband or partner, and the perceived relationship with a husband or partner (Table 2). Those pregnant adolescents with a primary school education level were 6.5 times more likely to develop prenatal depressive symptoms compared to those with a diploma degree [AOR=6.5 (95% CI: 1.5 to 28.7)]. The pregnant adolescents who perceived support from their husband or partner were five times more likely to have prenatal depressive symptoms compared to those with relatives' support [AOR=5.1 (95% CI: 1.9 to 14.5)]. The chance of developing prenatal depressive symptoms in those with a bad relationship with the husband was approximately seven times higher than pregnant adolescents who reported a good relationship [AOR=7.4 (95% CI: 3.1 to 17.8)].

Three variables in the final model influencing with prenatal depressive symptoms significantly were as follows:

$P(Y=1/X) = 1/[1+\text{EXP}(-1.99+ 1.88 \text{ Education; Primary} + 0.85 \text{ Education; Secondary} + 1.65 \text{ Supporter; Husband/partner} + 0.45 \text{ Supporter; Their family} + 2.00 \text{ Relationship with a husband/partner}]$

To validate how well the model fits a set of observations, Hosmer-Lemeshow's goodness-of-fit test was used and the results showed that the model was well-fit data (p -value = 0.675). The results of testing the model's predictive power by creating the relational graph of sensitivity and 1-specificity showed that the model had a good predictive power with ROC of 0.73 (95% CI 0.67 to 0.79) (Figure 1) at the cut point of -0.11, a sensitivity of 57.43, a specificity of 81.40, and an accuracy of 71.30.

Discussion

There was an unexpectedly high proportion of pregnant adolescents with prenatal depression symptoms in the ten community hospitals in the Maha Sarakham Province, at 43.9%. This proportion is twice the number reported worldwide (Abajobir *et al.*, 2016; Nasreen *et al.*, 2011; Roomruangwong & Epperson, 2011). However, the findings are similar to the studies reported by Uthapaisanwong *et al.* (2015) and Nirattharadorn *et al.* (2005), with the risk for prenatal depression from EPDS screening at 46% and 47.1%, respectively. Findings indicate that the underlying reasons are associated with the adolescents' low level of education, the husband or partner who supports her, and the relationship she has with him. Bunevicius *et al.* (2009) and Lydsdottir *et al.* (2014) also found that pregnant adolescents with a low level of education are more at risk of having depressive symptoms (see also Field (2017) and Osok (2016)). In Thailand, pregnancy in adolescents often leads to dropping out from school.

Table 1. Simple logistic regression assessing whether factors among participants were associated with depressive symptoms.

Factors	Depressive symptoms (n = 101) n (%)	Not depressive symptoms (n = 129) n (%)	Crude OR (95%CI)	p- value
Age (years)				0.231
≤15 years	21 (52.5)	19 (47.5)	1.5 (0.8, 3.0)	
>15 years	80 (42.1)	110 (57.9)	1	
Education				0.006
Primary	17 (70.8)	7 (29.2)	7.3 (1.9, 27.9)	
Secondary	79 (42.5)	107 (57.5)	2.2 (0.8, 6.3)	
Diploma	5 (25.0)	15 (75.0)	1	
Return to study				0.886
No return to study	58 (45.3)	70 (54.7)	1.2 (0.6, 2.4)	
Return to formal education	26 (42.6)	35 (57.4)	1.1 (0.5, 2.3)	
Return to non-formal education	17 (41.5)	24 (58.5)	1	
Occupation				.545
Unemployed	86 (44.8)	106 (55.2)	1.2 (0.6, 2.5)	
Employee	15 (39.5)	23 (60.5)	1	
Personal income				0.545
No	86 (44.8)	11 (52.4)	1.2 (0.6, 2.5)	
Yes	15 (39.5)	118 (56.5)	1	
Marital status				<0.001
Single mom	13 (86.7)	2 (13.3)	9.4 (2.1, 42.6)	
Living together with or without marriage	88 (40.9)	127 (59.1)	1	
Living with				0.553
Husband/partner	13 (50.0)	13 (50.0)	1.4 (0.6, 3.3)	
Adolescent's family	26 (48.2)	28 (51.9)	1.3 (0.7, 2.5)	
Husband/partner's family	62 (41.3)	88 (58.7)	1	
Pregnancy planning				0.720
Yes	10 (47.6)	11 (52.4)	1.2 (0.5, 2.9)	
No	91 (43.5)	118 (56.5)	1	
Number of supporters				0.087
1	41 (51.9)	38 (48.1)	2.8 (1.1, 7.4)	
2	53 (42.1)	73 (57.9)	1.9 (0.7, 4.8)	
≥3	7 (28.0)	18 (72.0)	1	
Supporters				0.013
Husband/partner	23 (65.7)	12 (34.3)	3.7 (1.4, 9.6)	
Adolescent's family	64 (41.6)	90 (58.4)	1.4 (0.7, 2.8)	
Adolescent's relatives	14 (34.2)	27 (65.8)	1	
Relationship with a husband/partner			<0.001	
Bad	27 (77.1)	8 (22.9)	5.5 (2.4, 12.8)	
Good	74 (37.9)	121 (62.5)	1	
Relationship with adolescent's family			0.016	
Bad	28 (59.6)	19 (40.4)	2.2 (1.2, 4.3)	
Good	73 (39.9)	110 (60.1)	1	
Relationship with adolescent's friends			<0.001	
Bad	64 (56.6)	49 (43.4)	2.8 (1.6, 4.8)	
Good	37 (31.6)	80 (68.4)	1	

Table 2. Multiple logistic regression of the factors associated with prenatal depressive symptoms during adolescent pregnancy.

Factors	Co-efficient	SE	Adjusted OR	95% CI	p-value
Education (ref=diploma)					0.013
Primary	1.88	0.75	6.5	1.5-28.7	
Secondary	0.85	0.60		0.7-7.6	
Supporter (ref=relative)					0.002
Husband/partner	1.65	0.52	5.1	1.9-14.5	
Their family	0.45	0.41		0.7 – 3.5	
Relationship with a husband/partner (ref=good)	2.00	0.45	7.4	3.1-17.8	<0.001
Constants	-1.99	0.68	-	-	-

Diagonal segments are produced by ties.

Figure 1. Graph shows associations between sensitivity with the 1-specificity of the model. Area under ROC curve = 0.73 (95%CI 0.67 to 0.79).

Consequently, a low level of education limits the opportunities for better employment and earning a higher income, which results in increased stress levels of these pregnant adolescents (Kaewjanta, 2012).

We found that not only was the emotional support coming from a husband or partner (compared to the pregnant adolescent's family) a risk factor for depressive symptoms, but also a poor relationship with that person contributes to her depression.

This finding seemingly contradicts Stapleton *et al.* (2012) who found that a husband's support reduces stress and positively affects pregnant women emotions. However, the perceived emotional support experienced by a pregnant adolescent from her husband or partner needs to be interpreted simultaneously with the finding of the poor relationship she experiences with him. The perceived adverse support from a husband or partner and the negative relationship with him likely increases the adolescents' depressive symptoms.

From the predictive model of prenatal depression symptoms by determining ROC, it was found that level of education, the perceived support from a husband or partner, and the perceived relationship with a husband or partner could predict prenatal depression symptoms well.

The 95% CI of the narrow ROC reflects that the model had predictive accuracy. Although the sensitivity was not adequate, specificity was at a high level. The model of three variables could be applied in operational planning to prevent and solve the problem of pregnant adolescents with prenatal depression symptoms.

The current Thai cultural norm is not to provide support to adolescents who become pregnant at a young age; society does not consider them to be ready for motherhood or a marital relationship. Adolescents who become pregnant may prefer to hide their condition from the family. Some may need to leave or transfer to a new school. They may lose their job and income as a result of the pregnancy or because of the change in living arrangements. Frequently, adolescent couples lack the maturity and emotional intelligence to begin a marital relationship, which might lead to domestic violence, although the latter was not part of the study's investigation or findings. In more agreement with our findings are the reports (Muangpin *et al.*, 2010; UNFPA, 2014a) that pregnant adolescents with depressive symptoms and other mental health problems are adversely affected by the lack of emotional support and a negative relationship with a husband.

This cross-sectional study does not permit a closer understanding of the long-term trajectory of adolescents who become pregnant and develop depressive symptoms, nor can it examine how their pregnancy might affect their adult lives. This information could come from longitudinal and qualitative studies. In addition, the sample in this study was confined to only one region of the country; therefore, the findings may not be generalizable to other parts of Thailand. Future research might be conducted in other areas, which should consider adult women who became pregnant in adolescence and have shown depressive symptoms and what meaning that held for them in their later lives. Similarly, the husbands or partners of the adolescents should be studied as to their perceptions of the experiences related to unexpected fatherhood with adolescent women.

In conclusion, the adolescent pregnancy rate in Thailand is increasing and adolescents are vulnerable to suffer from depressive

symptoms during pregnancy. As a matter of public health concern, more attention is needed to help reduce known problems with pregnant adolescents. Efforts to keep adolescents in school or ensure they return to school after delivery of the newborn may be an important step in improving the mental health of adolescent mothers. Public advocacy and education to support mental health initiatives for individuals and couples should be encouraged as a matter of health policy. If couples have become isolated from their families, attempts to reconcile and reunite them by public health personnel may promote better mental health.

Data availability

Underlying data

Figshare: Raw data - factors associated with depression. <http://doi.org/10.6084/m9.figshare.10072898.v2> (Jantasin *et al.*, 2019a).

This project contains the following underlying data:

- Raw-data-factors-associated-with-depression-edit.csv (raw data for each participant extracted in this study).
- Raw data Dictionary.docx (data dictionary defining all abbreviations and values)

Extended data

Figshare: Demographic questionnaire and EPDS in English. <http://doi.org/10.6084/m9.figshare.10073012> (Jantasin *et al.*, 2019b).

Figshare: Demographic questionnaire and EPDS in Thai. <http://doi.org/10.6084/m9.figshare.10073009> (Jantasin *et al.*, 2019c).

Data are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

Acknowledgements

The authors would like to thank Dr. Sutin Chanabun for providing guidance and support for statistical analysis, further thanks for Asst. Prof. Niruan Turnbull for the consultant of manuscript, the registered nurses who work at antenatal care clinics in 10 community hospitals in the Maha Sarakham province, and the pregnant adolescents who volunteered for this study. This study could not have been achievable without the dedicated support of all of them.

References

- Abajobir AA, Maravilla JC, Alati R, *et al.*: **A systematic review and meta-analysis of the association between unintended pregnancy and perinatal depression.** *J Affect Disord.* 2016; **192**: 56–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bolton HL, Hughes PM, Turton P, *et al.*: **Incidence and demographic correlates of**

- depressive symptoms during pregnancy in an inner London population.** *J Psychosom Obstet Gynaecol.* 1998; **19**(4): 202–209.
[PubMed Abstract](#) | [Publisher Full Text](#)

- Bunevicius A, Kusminskas L, Pop VJ, *et al.*: **Screening for antenatal depression with the Edinburgh Depression Scale.** *J Psychosom Obstet Gynaecol.* 2009;

30(4): 238–243.

[PubMed Abstract](#) | [Publisher Full Text](#)

Cox JL, Holden JM, Sagovsky R: **Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale.** *Br J Psychiatry.* 1987; **150**(6): 782–786.

[PubMed Abstract](#) | [Publisher Full Text](#)

Da Costa D, Larouche J, Dritsa M, *et al.*: **Psychosocial correlates of prepartum and postpartum depressed mood.** *J Affect Disord.* 2000; **59**(1): 31–40.

[PubMed Abstract](#) | [Publisher Full Text](#)

Dennis CL, Ross LE, Grigoriadis S: **Psychosocial and psychological interventions for treating antenatal depression.** *Cochrane Database Syst Rev.* 2007; (3): CD006309.

[PubMed Abstract](#) | [Publisher Full Text](#)

Eberhard-Gran M, Eskild A, Tambs K, *et al.*: **Review of validation studies of the Edinburgh Postnatal Depression Scale.** *Acta Psychiatr Scand.* 2001; **104**(4): 243–249.

[PubMed Abstract](#) | [Publisher Full Text](#)

Evans J, Heron J, Francomb H, *et al.*: **Cohort study of depressed mood during pregnancy and after childbirth.** *BMJ.* 2001; **323**(7307): 257–260.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Field T: **Prenatal Depression Risk Factors, Developmental Effects and Interventions: A Review.** *J Pregnancy Child Health.* 2017; **4**(1): pii: 301.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Grote NK, Bridge JA, Gavin AR, *et al.*: **A meta-analysis of depression during pregnancy and the risk of preterm birth, low birth weight, and intrauterine growth restriction.** *Arch Gen Psychiatry.* 2010; **67**(10): 1012–1024.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Hodgkinson SC, Colantuoni E, Roberts D, *et al.*: **Depressive symptoms and birth outcomes among pregnant teenagers.** *J Pediatr Adolesc Gynecol.* 2010; **23**(1): 16–22.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Hosmer DW Jr, Lemeshow S: **Applied Logistic Regression.** 2nd ed. New York: A Wiley Interscience Publication. 2000.

[Reference Source](#)

Hsieh FY, Bloch DA, Larsen MD: **A simple method of sample size calculation for linear and logistic regression.** *Stat Med.* 1998; **17**(14): 1623–1634.

[PubMed Abstract](#) | [Publisher Full Text](#)

Jantasin B, Yoosook W, Thaewpia S: **Raw data - factors associated with depression.** *figshare.* Dataset. 2019a.

<http://www.doi.org/10.6084/m9.figshare.10072898.v2>

Jantasin B, Yoosook W, Thaewpia S: **Demographic questionnaire and EPDS in English.** *figshare.* Dataset. 2019b.

<http://www.doi.org/10.6084/m9.figshare.10073012.v1>

Jantasin B, Yoosook W, Thaewpia S: **Demographic questionnaire and EPDS in Thai.** *figshare.* Dataset. 2019c.

<http://www.doi.org/10.6084/m9.figshare.10073009.v1>

Kaewjanta N: **Depression in teenage pregnancy: factors, affect and prevention.** *Journal of Nursing Science & Health.* 2012; **35**(1): 82–88.

[Reference Source](#)

Koniak-Griffin D, Walker DS, de Traversay J: **Predictors of depression symptoms in pregnant adolescents.** *J Perinatol.* 1996; **16**(1): 69–76.

[PubMed Abstract](#)

Lancaster CA, Gold KJ, Flynn HA, *et al.*: **Risk factors for depressive symptoms during pregnancy: a systematic review.** *Am J Obstet Gynecol.* 2010; **202**(1): 5–14.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Lee DT, Chan SS, Sahota DS, *et al.*: **A prevalence study of antenatal depression among Chinese women.** *J Affect Disord.* 2004; **82**(1): 93–99.

[PubMed Abstract](#) | [Publisher Full Text](#)

Limlomwongse N, Liabsuetrakul T: **Cohort study of depressive moods in Thai women during late pregnancy and 6–8 weeks of postpartum using the Edinburgh Postnatal Depression Scale (EPDS).** *Arch Womens Ment Health.* 2006; **9**(3): 131–138.

[PubMed Abstract](#) | [Publisher Full Text](#)

Lydsdottir LB, Howard LM, Olafsdottir H, *et al.*: **The mental health characteristics of pregnant women with depressive symptoms identified by the Edinburgh Postnatal Depression Scale.** *J Clin Psychiatry.* 2014; **75**(4): 393–8.

[PubMed Abstract](#) | [Publisher Full Text](#)

Meltzer-Brody S, Bledsoe-Mansori SE, Johnson N, *et al.*: **A prospective study of perinatal depression and trauma history in pregnant minority adolescents.** *Am J Obstet Gynecol.* 2013; **208**(3): 211.e1–7.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Ministry of Public Health: **Statistics on Adolescent Births, Thailand 2015.**

Department of Health, Ministry of Public Health. 2017; **1**: 11–146.

[Reference Source](#)

Muangpin S, Tiansawad S, Kantaruksa K, *et al.*: **Northeastern Thai adolescents' perceptions of being unmarried and pregnant.** *Pacific Rim Int J Nurs Res.* 2010; **14**(2): 149–161.

[Reference Source](#)

Nasreen HE, Kabir ZN, Forsell Y, *et al.*: **Prevalence and associated factors of depressive and anxiety symptoms during pregnancy: a population based study in rural Bangladesh.** *BMC Womens Health.* 2011; **11**: 22.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Niratttharadorn MK, Gennaro S, Vorapongsathorn T, *et al.*: **Self-esteem, social support and depression in Thai adolescent mothers.** *Thai J Nurs Res.* 2005; **9**(1): 63–75.

[Reference Source](#)

Orr ST, Blazer DG, James SA, *et al.*: **Depressive symptoms and indicators of maternal health status during pregnancy.** *J Womens Health (Larchmt).* 2007; **16**(4): 535–542.

[PubMed Abstract](#) | [Publisher Full Text](#)

Osok OJ: **Depression and psychosocial risk factors associated with pregnant adolescent mixed method study based at Kangemi health centre.** Nairobi University. 2016; [Cite 2018 september 15].

[Reference Source](#)

Pitanupong J, Liabsuetrakul T, Vittayanont A: **Validation of the Thai Edinburgh Postnatal Depression Scale for screening postpartum depression.** *Psychiatry Res.* 2007; **149**(1–3): 253–259.

[PubMed Abstract](#) | [Publisher Full Text](#)

Roomruangwong C, Epperson CN: **Perinatal depression in Asian women: prevalence, associated factors, and cultural aspects.** *Asian Biomedicine.* 2011; **5**(2): 179–193.

[Publisher Full Text](#)

Siegel RS, Brandon AR: **Adolescents, pregnancy, and mental health.** *J Pediatr Adolesc Gynecol.* 2014; **27**(3): 138–150.

[PubMed Abstract](#) | [Publisher Full Text](#)

Stapleton LR, Schetter CD, Westling E, *et al.*: **Perceived partner support in pregnancy predicts lower maternal and infant distress.** NIH public access. *J Fam Psychol.* 2012; **26**(3): 453–463.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Tzilos GK, Zlotnick C, Raker C, *et al.*: **Psychosocial factors associated with depression severity in pregnant adolescents.** *Arch Womens Ment Health.* 2012; **15**(5): 397–401.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

UNFPA: **The state of Thailand's population 2013: Motherhood in childhood facing the challenge of adolescent pregnancy.** Thailand: Prang Priyatrak. 2014a.

[Reference Source](#)

UNFPA: **The state of Thailand's population Report 2013.** 2014b; [Cite 2018 April 19].

[Reference Source](#)

Uthaisaisanwong A, Rungruxsirivorn T, Roomruangwong C, *et al.*: **Associated factors of prenatal depression among teenage pregnant women at King Chulalongkorn Memorial Hospital.** *J Med Assoc Thai.* 2015; **98**(5): 437–443.

[PubMed Abstract](#)

Vacharaporn K, Pitanupong J, Samangsi N: **Development of the Edinburgh Postnatal Depression Scale Thai version.** *J Mental Health Thai.* 2003; **11**(3): 164–169.

[Reference Source](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 13 November 2020

<https://doi.org/10.5256/f1000research.26093.r72522>

© 2020 Trumpff C. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Caroline Trumpff

Division of Behavioral Medicine, Department of Psychiatry, Columbia University Medical Center, New York, NY, USA

General comments:

This study assessed the proportion and risk factors of pregnant adolescent depression in in Maha Sarakham Province of Thailand. Description of the methods to assess risk factors is lacking. The title: "Predictive statistical model for the factors associated with prenatal depression among pregnant adolescents in Maha Sarakham province, Thailand" is misleading; the author should clarify which type of factors were assessed, here it seems that the investigation was limited to socio-economic factors (education) and social support. Spelling and grammar mistakes were found throughout. The language should be edited by an English native speaker.

Major comments:

Abstract:

- What about the odd of depression in pregnant adolescents supported by both relatives as well as the father?

Introduction:

- Can the other add recent references on the risk factors for depression in pregnancy.

Method:

- Why did you exclude pregnant adolescents with psychiatric disorders? Did you excluded adolescents who previously experienced depression?
- "Initially, a sample size of 209 adolescents was calculated (Hsieh et al., 1998) assuming a prenatal depression prevalence of 46% (Uthaipaisanwong et al., 2015) with an alpha error of 0.05 and a power of 80%." Could you provide more information on the way you performed sample size calculation? Based on which effect did you compute the sample size calculation?

Finding predictors of depression or finding a certain prevalence of depression? Is the study powered to define predictors of depression?

- Could you describe how did you measure factors associated with prenatal depression? Can you list the factors and give the reference of the tools used (e.g. was perceived social support measured with a validated tool)?

Results:

- "The adolescents who had a husband as a supporter tended to be 3.7 times higher at risk of depressive symptoms when compared to those who were supported by relatives." Do you mean pregnant adolescents who had only their husband as a supporter?
- "The pregnant adolescents who perceived support from their husband or partner were five times more likely to have prenatal depressive symptoms compared to those with relatives' support [AOR=5.1 (95% CI:1.9 to 14.5)]." I really wonder how this was measured.

Discussion:

- "We found that not only was the emotional support coming from a husband or partner (compared to the pregnant adolescent's family) a risk factor for depressive symptoms, but also a poor relationship with that person contributes to her depression." This sentence should be revised for clarity.
- "Frequently, adolescent couples lack the maturity and emotional intelligence to begin a marital relationship, which might lead to domestic violence, although the latter was not part of the study's investigation or findings." I would be more cautious in the language used (e.g. adolescent couples may lack...) or completely delete this sentence.
- "In addition, the sample in this study was confined to only one region of the country; therefore, the findings may not be generalizable to other parts of Thailand." Please modify accordingly.

Figure and tables:

- Define the factors in the legend.

Minors comments:

Methods:

- "the EPDS has been successfully used to identify depressive. symptoms in prenatal women". "Symptoms" should be "symptoms".
- "Data analysis were conducted with descriptive statistics to present demographic data." This sentence could be modified as follows: "Descriptive statistics were used to present demographic data".
- "to determine potential associations **of** sociodemographic information with prenatal depressive symptoms (EPDS $\geq 11=1$ and $<11=0$)." This sentence could be modified as follows "to determine potential associations **between** sociodemographic information and prenatal depressive symptoms (EPDS $\geq 11=1$ and $<11=0$)."

Results:

- "According to Table 1, eight variables with a p-value of less than 0.25 were selected and included in the initial regression model." This sentence should be corrected. Suggestion: "The initial regression model included eight variables selected with a p-value of less than 0.25 (see. Table 1)."

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

No

If applicable, is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: psychopathology, psychobiology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Reviewer Report 29 April 2020

<https://doi.org/10.5256/f1000research.26093.r62876>

© 2020 Anyiam F. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Felix Emeka Anyiam

University of Port Harcourt , Port Harcourt, Nigeria

I have no further comments to make as the revised article is well done.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Public Health, Population Health, Global Health, Epidemiological Research, Biostatistics, Open Research Data, Open Science, Data Science, Non-communicable diseases, Maternal and Child Health, Vulnerable Populations, Population at Risk, Reproductive Health, Data Management, Community-Based Research, HIV/AIDS Prevention, TB Prevention, Health Management and Health Promotion

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 10 February 2020

<https://doi.org/10.5256/f1000research.23117.r58099>

© 2020 Anyiam F. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Felix Emeka Anyiam

University of Port Harcourt , Port Harcourt, Nigeria

Summary:

The article titled "Predictive statistical model for the factors associated with prenatal depression

among pregnant adolescents in Maha Sarakham province, Thailand", is a cross-sectional study focused on adolescent pregnant mothers aged ≤ 19 years, and have an estimated gestational age of ≤ 18 weeks. The study is relevant, as it highlighted several risk factors associated with prenatal depression and some of which are: level of education, perceived support from husband and perceived relationship with husband. The study found a high prevalence of depression among this group of people which I think is a public health concern needing an urgent governmental intervention.

Report:

A. Title:

1. The title for this study is not appropriate as I did not see a **predictive statistical modeling**. Predictive modeling in statistics is more complex than what you have here. Sometimes, data mining and probability are applied in order to forecast outcomes. I think you should just leave the title as:
Factors associated with prenatal depression among pregnant adolescents in Maha Sarakham province, Thailand

B. Abstract:

1. The results section, the sentence with the words: "... and those with a perceived bad relationship with their husband or partner seven times higher." should read: "...and those with a perceived bad relationship with their husband or partner **had** seven times higher."

C. Introduction:

1. The statement, "Adolescents can develop health problems during pregnancy that adversely affect both physical and psychological health (UNFPA, 2014b)." should read, "Adolescents can develop health problems during pregnancy that adversely affect both **their** physical and psychological health (UNFPA, 2014b).
2. I do not think your introduction reference should begin with UNFPA, 2014b but instead UNFPA, 2014a. "a" should come before "b." Please do switch the reference.
3. I am not sure what the phrase "...adult women" means in the sentence: "Risks factors for depression among pregnant adolescents include not only those listed for adult women..." as you did not list any risk factors for adult women.
4. The statement, "...and a history of emotion, sexual, and physical abuse.." should read, "...and a history of emotional, sexual, and physical abuse.
5. "Little research, however, has been conducted in northeastern Thailand, especially in Maha Sarakham province, where the rate of adolescent pregnancy is reported to be high." is a very strong statement and should be backed-up with literature.

D. Results:

1. Your Table 2 has 4 p-values but you only have 3 variables here you are associating. Please confirm and correct.

2. In Table 2, you also have 5 Adjusted OR. I think this should be just the 3 you mentioned in your text: 6.5, 5.1 and 7.4.

E. Discussion:

1. "As a matter of public health..." should read, "As a matter of public health **concern**".

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Public Health, Population Health, Global Health, Epidemiological Research, Biostatistics, Open Research Data, Open Science, Data Science, Non-communicable diseases, Maternal and Child Health, Vulnerable Populations, Population at Risk, Reproductive Health, Data Management, Community-Based Research, HIV/AIDS Prevention, TB Prevention, Health Management and Health Promotion

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 24 Apr 2020

Bhanwasa Jantasin, Mahasarakham University, Maha Sarakham, Thailand

Report:

A. Title:

1. The title for this study is not appropriate as I did not see a **predictive statistical modeling**. Predictive modeling in statistics is more complex than what you have here. Sometimes, data mining and probability are applied in order to forecast outcomes. I think you should just leave the title as: **Factors associated with prenatal depression**

among pregnant adolescents in Maha Sarakham province, Thailand

Answer:

We discussed the title and prefer to preserve it as our statistical analysis (ROC = 73%) is plausible to forecast future outcomes.

B. Abstract:

1. The results section, the sentence with the words: "... and those with a perceived bad relationship with their husband or partner seven times higher." should read: "...and those with a perceived bad relationship with their husband or partner **had** seven times higher."

Answer: we added the missing word according to the comment.

C. Introduction:

1. The statement, "Adolescents can develop health problems during pregnancy that adversely affect both physical and psychological health (UNFPA, 2014b)." should read, "Adolescents can develop health problems during pregnancy that adversely affect both **their** physical and psychological health (UNFPA, 2014b).

Answer: we added the missing word according to the comment.

1. I do not think your introduction reference should begin with UNFPA, 2014b but instead UNFPA, 2014a. "a" should come before "b." Please do switch the reference.

Answer: we revised the reference according to the comment.

1. I am not sure what the phrase "...adult women" means in the sentence: "Risks factors for depression among pregnant adolescents include not only those listed for adult women..." as you did not list any risk factors for adult women

Answer: We mean "pregnant adult women" and the risks factors are listed in the first sentence. We added the word 'pregnant' in front of 'adult women' to make it more clear.

1. The statement, "...and a history of emotion, sexual, and physical abuse.." should read, "...and a history of **emotional**, sexual, and physical abuse.

Answer: we added the missing word according to the comment.

1. "Little research, however, has been conducted in northeastern Thailand, especially in Maha Sarakham province, where the rate of adolescent pregnancy is reported to be high." is a very strong statement and should be backed-up with literature.

Answer: we added the reference according to the comment.

D. Results:

1. Your Table 2 has 4 p-values but you only have 3 variables here you are associating. Please confirm and correct.
2. In Table 2, you also have 5 Adjusted OR. I think this should be just the 3 you mentioned in your text: 6.5, 5.1 and 7.4.

Answer: we revised the table according to the comment.

E. Discussion:

1. "As a matter of public health..." should read, "As a matter of public health **concern**".

Answer: we added the missing word according to the comment.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research