REVIEW

Recent advances in taste transduction and signaling [version 1; peer review: 2 approved]

Sue C. Kinnamon, Thomas E. Finger

Rocky Mountain Taste & Smell Center, Department of Otolaryngology and Department of Cell & Developmental Biology, University of Colorado School of Medicine, Aurora, CO, 80045, USA

V1 First published: 17 Dec 2019, 8(F1000 Faculty Rev):2117
<https://doi.org/10.12688/f1000research.21099.1>

Latest published: 17 Dec 2019, 8(F1000 Faculty Rev):2117
<https://doi.org/10.12688/f1000research.21099.1>

Abstract

In the last few years, single-cell profiling of taste cells and ganglion cells has advanced our understanding of transduction, encoding, and transmission of information from taste buds as relayed to the central nervous system. This review focuses on new knowledge from these molecular approaches and attempts to place this in the context of previous questions and findings in the field. The individual taste cells within a taste bud are molecularly specialized for detection of one of the primary taste qualities: salt, sour, sweet, umami, and bitter. Transduction and transmitter release mechanisms differ substantially for taste cells transducing sour (Type III cells) compared with those transducing the qualities of sweet, umami, or bitter (Type II cells), although ultimately all transmission of taste relies on activation of purinergic P2X receptors on the afferent nerves. The ganglion cells providing innervation to the taste buds also appear divisible into functional and molecular subtypes, and each ganglion cell is primarily but not exclusively responsive to one taste quality.

Keywords

taste bud, transduction, geniculate ganglion, purinergic transmission, serotonin, taste, signalling, nervous system

Open Peer Review

Approval Status

	1	2
version 1
17 Dec 2019		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Susan Travers**, Ohio State University, Columbus, USA
2. **Stephen Roper**, University of Miami Miller School of Medicine, Miami, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Thomas E. Finger (tom.finger@cuanschutz.edu)

Author roles: Kinnamon SC: Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; Finger TE: Conceptualization, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported in part by grants from the National Institute on Deafness and Other Communication Disorders: 1R01DC012931 and 1R01DC014728 (TEF), 1R01DC017679 and 1R01DC012555 (SCK).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 Kinnamon SC and Finger TE. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Kinnamon SC and Finger TE. **Recent advances in taste transduction and signaling [version 1; peer review: 2 approved]** F1000Research 2019, 8(F1000 Faculty Rev):2117 <https://doi.org/10.12688/f1000research.21099.1>

First published: 17 Dec 2019, 8(F1000 Faculty Rev):2117 <https://doi.org/10.12688/f1000research.21099.1>

This review will focus on progress in the last few years in understanding transduction, coding, and specificity of the peripheral gustatory system starting with the cellular components of taste buds. Taste buds comprise 50 to 100 taste cells, which are specialized epithelial cells including signal transducing cells and glial-like supporting cells. The taste cells respond to taste substances in the saliva to generate a biological signal that then must be transmitted to the taste sensory nerves conveying the message to the nucleus of the solitary tract in the brain stem. This review will focus on recent discoveries in the processes of taste transduction and transmission, which significantly impact our understanding of this system.

In mammals, the sense of taste detects a wide variety of compounds which canonically fall into only five main sensory qualities (or modalities): sweet, umami (savory), bitter, sour, and salty. Each taste cell is most responsive to a single taste quality¹, and each taste bud contains one or more cells capable of responding to each of the taste qualities; that is, taste cells are “chemically tuned” whereas taste buds are not.

Methodological advances

Two relatively new methodologies have been put to good effect in analyzing taste buds and their associated ganglia. Functional imaging of ganglion cells expressing genetically encoded activity sensors has allowed direct analysis of the breadth of tuning of inputs to the central nervous system (CNS)^{2,3}, whereas single-cell transcriptomics has permitted molecular and functional classification of both taste cells and ganglion cells⁴⁻⁸. The implications of these and other recent studies are described below and summarized in Table 1.

Transduction

The taste cells are divisible into four types characterized by both morphological and molecular features and given the names Type I, Type II, Type III and Type IV (Figure 1). Type I cells are similar in many ways to astrocytes and Type IV cells are immature cells, whereas Type II and Type III cells serve as the transducing elements for different taste qualities.

Type II cells use G protein–coupled receptors for sweet (T1R2 + T1R3), umami (T1R1 + T1R3), or bitter (T2Rs) to

Figure 1. Cell types in taste buds. Four different morphological and molecularly distinct types of cells populate taste buds. Types II and III transduce different classes of tastes, whereas Type I cells are more glial-like. Type IV cells are the immature population, which develop into the other cell types over the span of a few days. Figure generated from data in 12.

initiate a transduction cascade, whereas Type III cells rely on ion channels for transduction of the ionic tastes of salty and sour. The receptors and downstream signaling cascade for the Type II cells (sweet, umami, or bitter) have been well described since the early part of this century¹³ and involve a phospholipase C (PLC)-mediated cascade culminating in the activation of the Ca⁺⁺-responsive channels TRPM5 and TRPM4¹⁴ to depolarize the cell sufficiently to generate an action potential via voltage-gated Na⁺ channels (SCN2A, SCN3A, and SCN9A¹⁵). Why axonless receptor cells should generate action potentials is of interest and is likely related to the release mechanism for neurotransmitter from Type II taste cells as described below.

Whereas early studies suggested that a single sweet taste receptor (T1R2 + T2R3) mediates all responses to sugars and sweeteners¹⁶, recent studies suggest that other mechanisms also play a role for glucose-containing sugars but not for artificial sweeteners. Glucose transporters and the K_{ATP} channel, which are expressed in sweet-responsive (T1R3-expressing) taste cells¹⁷, are involved in cephalic phase insulin release independent of the neural signal for sweet transmitted to the nervous system¹⁸. The exact mechanism by which activation of the taste cells evokes insulin release is unclear but may involve humoral rather than neural signals.

Sour

In 2006, Huang *et al.*¹⁹ showed that sour detection depended on cells expressing PKD2L1—cells subsequently identified as a subset of Type III cells²⁰. In 2011, Horio *et al.* showed that PKD2L1 itself was not necessary for transduction of protons²¹.

Table 1. Reported molecular characteristics of taste cells and ganglion cells.

Taste quality	Sweet	Umami	Bitter	Sour	Salty
Ganglion cells	<i>Spon1</i>	<i>Cdh4?</i> ^a	<i>Cdh13 ?</i> ^a	<i>Penk, Htr3a</i>	<i>Egr2?</i> ^a
Taste cells	<i>Sema7A</i>		<i>Sema3A</i>	<i>Pkd2L1, OTOP1</i>	

Data are from 6,8–11. ^aThe meta-analysis by Anderson and Larson¹¹ does not support *Cdh4*, *Cdh13*, and *Egr2* as marking unique clusters. *Cdh13* marked two clusters that also express *Olfm3*. *Olfm3* is associated with several ganglion cell clusters but is never associated with *Penk* and so may mark cells innervating Type II cells⁶ but not a particular subset of Type II cells. Factors in question are indicated by “?”.

Rather, transduction of sour involves permeation of H^+ through an apical ion channel²² subsequently identified as OTOPI⁶. Using PKD2L1 as a molecular identifier for sour-responsive taste cells, Liman *and co-workers*⁷ and Zhang *et al.*⁸ went on to confirm OTOPI as the necessary transduction channel underlying sour taste. The entering H^+ ions not only directly depolarize the taste cells but also block Kir2.1 K^+ channels²³, thereby amplifying the depolarization of the entering H^+ ions. The resulting depolarization triggers voltage-gated Na^+ channels (SCN2A¹⁵) to generate action potentials that activate voltage-gated Ca^{++} channels triggering the release of synaptic vesicles²⁴.

In keeping with the PKD2L1 cells being the sour-transducing cells, optogenetic driving of these cells evokes an aversive response²⁵. Curiously, another study²⁶ reported that optogenetic driving of the PKD2L1 population drives drinking behavior in thirsty mice. Why the mice should respond with drinking to a sensation of sour is still unresolved, although Zocchi *et al.*²⁶ suggest that a subset of PKD2L1-expressing cells may convey a specific “water taste” as is known for insects^{27–29}.

Salt

Historically, responsiveness to salt has been separated into amiloride-sensitive (AS) and amiloride-insensitive (AI) modalities³⁰. Confounding our understanding of salt taste is that low concentrations of salt are appetitive whereas high concentrations are aversive. Furthermore, while Na^+ is important perceptually for salt, other substances, not containing Na^+ , also are salty-tasting. The multiple perceptual and chemical properties suggest that more than one transduction mechanism may be involved. Supporting this concept are the results from^{31,32}. These studies argue that the molecular correlate of AS salt, the epithelial sodium channel (ENaC), underlies the appetitive qualities of Na^+ but that AI salt detection of high concentrations of Na^+ relies on a subpopulation of the bitter-responsive Type II cells and a subset of the sour-responsive Type III cells^{5,32,33}. Furthermore, a recent study³⁴ suggests that the AI salt transduction mechanism may directly involve Cl^- , but the actual mechanism remains elusive since known Cl^- channel and transporter blockers have no effect on salt taste. Two recent works confirm a previous study³⁵ suggesting that AS-responsive taste cells fall into a unique class of taste cells not identified by the canonical reporters (TRPM5 for Type II and CAR4 for Type III³² or PIRT for Types II and III³⁴). Further confounding the interpretation of these studies on Na^+ transduction is the finding that, compared with rodents, humans—who enjoy low salt and avoid high salt—do not have a large AS component of salt taste³⁶, although chimpanzees do have a minor AS component³⁷.

Peripheral neurotransmission

Whatever means are used for transduction, the Type II and Type III cells ultimately must release one or more neurotransmitters to activate the afferent nerves. Activity-dependent release of ATP from Type II cells^{38,39} and 5-HT from Type III cells⁴⁰ was clearly demonstrated by several investigators, but whether other transmitters (for example, glutamate or acetylcholine) also may be involved remains unclear. Although ATP acting on neural P2X receptors was identified as a crucial transmitter for

all tastes over a decade ago⁴¹, only recently has the contribution of 5-HT acting on neural 5-HT₃ receptors been elucidated in terms of transmission of sour taste⁴². In addition, taste cells may directly release peptides such as glucagon-like peptide 1 (GLP-1)^{43,44}. These additional transmitters may act to modulate adjacent taste cells⁴⁵ and activate afferent nerve fibers.

Although both Type II and Type III cells require action potentials for transmitter release, the mechanisms of release for these two cell types are quite different (Figure 2). Type III cells use a conventional synapse involving voltage-gated Ca^{++} channels and SNARE mechanisms to effect release of synaptic vesicles^{24,46,47}. In contrast, Type II taste cells (transducing bitter, sweet, or umami) rely on action potentials to trigger the voltage-gated large-pore channel CALHM1 to release ATP^{48–50}. The pore size of this channel is sufficient to permit passage of ATP which serves as an obligatory transmitter in this system^{41,51,52}. The biophysical properties of CALHM1 as described in the seminal article on this channel⁴⁸ did not exactly match the properties of the release channel in taste buds. A more recent report⁴⁹ showed that the channel in taste buds consists of two subunits, CALHM1 and CALHM3, which together form a channel matching the properties of the Type II cell release channel.

Whereas the mechanism of release for ATP is well established for Type II cells, the source of ATP for transmission of sour information from Type III cells remains enigmatic. No one has yet demonstrated direct release of ATP from Type III cells, yet transmission of Type III cell taste qualities (for example, sour) is dependent on intact purinergic signaling to P2X receptors⁴¹. Type III cells do not express CALHM1⁴⁸ and so do not use CALHM1/3-mediated release. Another possible means of release of ATP is via synaptic vesicles, but Type III cells are reported to lack the vesicular ATP-transporter, VNUT⁵³, presumed to be necessary for loading of synaptic vesicles with ATP. So, what other possible sources exist for release of ATP? One suggestion is that Type III cells may trigger ATP release via interaction with other taste cell types. If so, this interaction does not require participation of Type II cells since mice lacking Type II cells (Skn1A-KO)⁵⁴ and CALHM1 KO⁴⁸ mice show nearly normal responses to sour. Hence, the transmission of sour information to taste nerves does not require the presence of Type II taste cells nor the function of CALHM1 to release ATP. Furthermore, a recent study showing high-resolution reconstructions of taste buds shows that Type III cells seldom directly contact Type II cells since processes from Type I cells intervene⁵⁵, suggesting that any interactions between Type II and Type III cells may be indirect.

Tuning specificity of taste cells and nerve fibers

One of the classic discussions in taste research concerns the specificity of chemical tuning in the peripheral taste system, that is, whether individual taste cells or nerve fibers respond to single classes of taste stimuli or respond more broadly across several modalities. The former, known as “labeled line” coding, suggests that taste information (for example, for sweet taste) is transduced by a unique subset of taste cells and transmitted over dedicated nerve fibers that signal that particular quality (sweet)

Figure 2. Taste transduction cascades. Transduction pathways for the two different types of taste-transducing cells: Type III for sour and Type II for sweet, bitter, or umami. The different responsiveness of Type II cells is dictated by the type of receptor each cell expresses, not by downstream members of the transduction cascade. AtpMito, atypical mitochondria; ER, endoplasmic reticulum; G α , alpha subunit of G protein; G $\beta\gamma$, beta-gamma subunits of G protein; IP $_3$, inositol trisphosphate; IP $_3$ R3, inositol trisphosphate receptor isoform 3; PLC β 2, phospholipase C isoform β 2; TRPM5, transient receptor potential cation channel subfamily M member 5; VGCC, voltage-gated calcium channel; VG-Na $^+$, voltage-gated sodium channel.

to the brain. Conversely, a broadly tuned peripheral system, utilizing “cross-fiber pattern” coding, relies on nerve fibers that are more broadly responsive but that respond maximally to particular qualities, necessitating comparison of activity across the fiber population in order for the brain to extract quality information. Receptor expression data strongly indicate that taste receptor cells largely express receptors for a single taste quality and this suggests that labeled line coding is plausible⁵⁵.

So, the question then is how specifically the taste cells activate particular nerve fibers. Two articles published in 2015 address this question but arrive at somewhat different conclusions, although detailed comparison of the data in the two studies shows similar results. In both works, the investigators used optical recording methods to assess activity in geniculate ganglion cells that innervate taste buds in fungiform papillae on the front of the tongue. Barretto *et al.*², using relatively low concentrations of taste stimuli, report that taste ganglion cells largely respond to a single stimulus class, although many multiply responsive cells are evident in the data. Conversely, Wu *et al.*³ found that ganglion cell specificity is dominant only for low concentrations of tastants. As concentrations increase, more and more ganglion cells are recruited so that the majority of cells will respond to more than one class of tastant at moderate - high stimulus levels. This finding tends to support a cross-fiber model of decoding afferent input, particularly at higher stimulus concentrations. It should be noted that loss of specificity at high stimulus levels is

similar to encoding in other sensory modalities; for example, a dim blue light of 505 nm will selectively activate the blue-sensitive, short-wavelength cones in the retina. But when the same wavelength of light is more intense, it will also activate the green, medium-wavelength cones and ultimately the red, long-wavelength cones. So absolute color cannot be determined by reading the output of only one class of cones but rather requires comparisons of activity levels across the different types of cones. Similarly, taste quality identification may require comparison of activity levels across incoming lines of information which preferentially, but not absolutely, encode particular qualities. In other words, while the afferent input is “tuned” as would be necessary for a labeled line system, absolute quality information can be extracted only by comparing the pattern and level of activity across fiber classes.

Another requirement for labeled line coding is connectional specificity between taste cells and nerve fibers; that is, sweet receptor-expressing taste cells should connect to nerve fibers receiving synaptic input only from other sweet receptor cells and so forth. Such specificity has been suggested recently for both Type III cells and Type II cells. Taste nerve fibers highly expressing the serotonin receptor 5-HT $_3$ show preferential connectivity with Type III taste cells (sour) that accumulate and release serotonin⁹. Argument for Type II cell connectional specification is based on transcriptome profiling of geniculate ganglion cells. Like dorsal root ganglion cells⁵⁶, geniculate ganglion cells

fall into distinct molecular classes⁴. Those ganglion cells that innervate taste buds express Phox2b, whereas those providing somatosensory innervation to the ear do not⁵⁷, so this transcription factor can be used to identify ganglion cells that innervate taste buds. Lee *et al.*¹⁰ showed that connectional specificity for Type II cells is maintained by molecular encoding of receptor cell and nerve fiber identities by SEMA family guidance molecules. Likewise, Zhang *et al.*⁸ found that genetic deletion of particular gene products correlates with loss of behavioral response to particular tastants—*Cdh4* for umami, *Cdh13* for bitter, and *Egr2* for salty—suggesting that these factors may serve to identify particular classes of gustatory ganglion cells (see Table 1). Since these proteins are expressed widely in the CNS (including in taste-processing areas), it is unclear whether the reported behavioral changes are attributable to changes in ganglion cell functionality or changes higher in the neuraxis. Furthermore, a more recent meta-analysis of these and other transcriptome data on ganglion cell subclasses fails to support the segregation of geniculate ganglion cell subtypes according to expression of these cadherins¹¹. Whether this all equates to absolute functional specificity of the taste neurons remains open to question. Substantial evidence exists for the possibility of cell-to-cell communication in taste buds^{1,58}, and side-band activity in the taste nerves might result from such interactions between taste cells rather than direct convergence of input onto single fibers.

Another interesting outcome of the molecular profiling of the geniculate ganglion neurons innervating taste buds is the identification of a population expressing *Piezo2*, a marker of touch-sensitive ganglion cells⁴. Such touch sensitivity of a subpopulation of gustatory ganglion cells would explain the residual tactile responsiveness of the gustatory nerves following genetic deletion of P2X receptors necessary for transmission from taste cells to nerve fibers⁴¹. Thus, the nerve fibers themselves may be touch-sensitive and require no activation from the taste buds for tactile activation.

Is “fatty” a taste?

A classic question in the field of taste research is “How many primary tastes are there?” In the 20th century, the debate focused on whether umami (savory) was a primary taste distinct from the classic qualities of salty, sour, sweet, and bitter. The

discovery of a distinct molecular receptor for glutamate, coupled with the identification of glutamate (umami)-specific fibers in gustatory nerves and behavioral experiments showing the discriminability of glutamate from other tastes, clinched the case that umami is indeed a distinct primary taste. In the last few decades, the question of whether the taste of fat is a primary taste quality has been debated⁵⁹. Potential fat receptors GPR120 and CD36 were identified molecularly in taste buds about a decade ago^{60,61}. But the mere presence of a receptor does not necessarily equate to the existence of a separate coding channel as would be expected of a primary taste quality. This year, Ninomiya *et al.*⁶² identified, for the first time, a population of nerve fibers (F-fibers) in the chorda tympani nerve that respond uniquely to a fatty acid, linoleic acid, lending further credence to the idea that fat is a unique taste quality. Complicating the situation is that some of the F-fibers also show responses to glutamate, and although mice can be trained to recognize the taste of linoleic acid, they confound it with glutamate (see also 63), suggesting that fatty taste sensations may be intermingled with those for umami. Thus, the question remains as to whether fatty acid taste is a distinct primary taste or more a modulator of other taste qualities.

Remaining unanswered questions

- What mechanisms and cells underlie transduction of AI salt taste, and what cells are responsible for AS salt taste?
- What is the source of ATP required for transmission of information from Type III (sour) cells to the afferent nerves?
- How much intercellular communication occurs within taste buds, and how does this affect the nature of the signal transmitted to the nerve fibers?
- Is fatty a primary taste quality? And how many other unrecognized primary taste qualities may exist?

Acknowledgments

The authors thank Brigit High and Eric Larson (Univ. Colorado Sch. Medicine) for comments on this work, and Yannick K. Dzowo (Univ. Colorado Sch. Medicine) for assistance with preparing Figure 1.

References

1. Roper SD: **Signal transduction and information processing in mammalian taste buds.** *Pflügers Arch.* 2007; **454**(5): 759–76. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
2. Barretto RPJ, Gillis-Smith S, Chandrashekar J, *et al.*: **The neural representation of taste quality at the periphery.** *Nature.* 2015; **517**(7534): 373–6. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
3. Wu A, Dvoryanchikov G, Pereira E, *et al.*: **Breadth of tuning in taste afferent neurons varies with stimulus strength.** *Nat Commun.* 2015; **6**: 8171. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
4. Dvoryanchikov G, Hernandez D, Roebber JK, *et al.*: **Transcriptomes and neurotransmitter profiles of classes of gustatory and somatosensory neurons in the geniculate ganglion.** *Nat Commun.* 2017; **8**(1): 760. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
5. Sukumaran SK, Lewandowski BC, Qin Y, *et al.*: **Whole transcriptome profiling of taste bud cells.** *Sci Rep.* 2017; **7**(1): 7595. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
6. Tu YH, Cooper AJ, Teng B, *et al.*: **An evolutionarily conserved gene family**

- encodes proton-selective ion channels. *Science*. 2018; 359(6379): 1047–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
7. Teng B, Wilson CE, Tu YH, *et al.*: Cellular and Neural Responses to Sour Stimuli Require the Proton Channel Otop1. *Curr Biol*. 2019; 29(21): 3647–3656.e5.
[PubMed Abstract](#) | [Publisher Full Text](#)
 8. **F** Zhang J, Jin H, Zhang W, *et al.*: Sour Sensing from the Tongue to the Brain. *Cell*. 2019; 179(2): 392–402.e15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 9. Stratford JM, Larson ED, Yang R, *et al.*: 5-HT_{3A}-driven green fluorescent protein delineates gustatory fibers innervating sour-responsive taste cells: A labeled line for sour taste? *J Comp Neurol*. 2017; 525(10): 2358–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 10. Lee H, Macpherson LJ, Parada CA, *et al.*: Rewiring the taste system. *Nature*. 2017; 548(7667): 330–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 11. Anderson CB, Larson ED: Single Cell Transcriptional Profiling of Phox2b-Expressing Geniculate Ganglion Neurons. *bioRxiv*, 2019; 812578.
[Publisher Full Text](#)
 12. Yang R, Dzowo YK, Wilson CE, *et al.*: Three-dimensional reconstructions of mouse circumvallate taste buds using serial blockface scanning electron microscopy: I. Cell types and the apical region of the taste bud. *J Comp Neurol*. 2019; 25: 205.
[PubMed Abstract](#) | [Publisher Full Text](#)
 13. Zhang Y, Hoon MA, Chandrashekar J, *et al.*: Coding of sweet, bitter, and umami tastes: different receptor cells sharing similar signaling pathways. *Cell*. 2003; 112(3): 293–301.
[PubMed Abstract](#) | [Publisher Full Text](#)
 14. **F** Dutta Banik D, Martin LE, Freichel M, *et al.*: TRPM4 and TRPM5 are both required for normal signaling in taste receptor cells. *Proc Natl Acad Sci U S A*. 2018; 115(4): E772–E781.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 15. Gao N, Lu M, Echeverri F, *et al.*: Voltage-gated sodium channels in taste bud cells. *BMC Neurosci*. 2009; 10: 20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 16. **F** Nelson G, Hoon MA, Chandrashekar J, *et al.*: Mammalian sweet taste receptors. *Cell*. 2001; 106(3): 381–390.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 17. **F** Yee KK, Sukumaran SK, Kotha R, *et al.*: Glucose transporters and ATP-gated K⁺ (KATP) metabolic sensors are present in type 1 taste receptor 3 (T1r3)-expressing taste cells. *Proc Natl Acad Sci U S A*. 2011; 108(13): 5431–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 18. **F** Glendinning JL, Frim YG, Hochman A, *et al.*: Glucose elicits cephalic-phase insulin release in mice by activating K_{ATP} channels in taste cells. *Am J Physiol Regul Integr Comp Physiol*. 2017; 312(4): R597–R610.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 19. **F** Huang AL, Chen X, Hoon MA, *et al.*: The cells and logic for mammalian sour taste detection. *Nature*. 2006; 442(7105): 934–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 20. Kataoka S, Yang R, Ishimaru Y, *et al.*: The candidate sour taste receptor, PKD2L1, is expressed by type III taste cells in the mouse. *Chem Senses*. 2008; 33(3): 243–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 21. Horio N, Yoshida R, Yasumatsu K, *et al.*: Sour taste responses in mice lacking PKD channels. *PLoS One*. 2011; 6(5): e20007.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 22. **F** Chang RB, Waters H, Liman ER: A proton current drives action potentials in genetically identified sour taste cells. *Proc Natl Acad Sci U S A*. 2010; 107(51): 22320–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 23. Ye W, Chang RB, Bushman JD, *et al.*: The K⁺ channel K_{ir}2.1 functions in tandem with proton influx to mediate sour taste transduction. *Proc Natl Acad Sci U S A*. 2016; 113(2): E229–E238.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 24. Vandenbeuch A, Zorec R, Kinnamon SC: Capacitance measurements of regulated exocytosis in mouse taste cells. *J Neurosci*. 2010; 30(44): 14695–701.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 25. Wilson CE, Vandenbeuch A, Kinnamon SC: Physiological and Behavioral Responses to Otopotent Stimulation of PKD2L1⁺ Type III Taste Cells. *eNeuro*. 2019; 6(2): pii: ENEURO.0107–19.2019.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 26. Zocchi D, Wennemuth G, Oka Y: The cellular mechanism for water detection in the mammalian taste system. *Nat Neurosci*. 2017; 20(7): 927–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
 27. **F** Cameron P, Hiroi M, Ngai J, *et al.*: The molecular basis for water taste in *Drosophila*. *Nature*. 2010; 465(7294): 91–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 28. Chen Z, Wang Q, Wang Z: The amiloride-sensitive epithelial Na⁺ channel PPK28 is essential for *drosophila* gustatory water reception. *J Neurosci*. 2010; 30(18): 6247–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 29. **F** Matthews BJ, Younger MA, Vossell LB: The ion channel *ppk301* controls freshwater egg-laying in the mosquito *Aedes aegypti*. *eLife*. 2019; 8: pii: e43963.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 30. Heck GL, Miesner S, DeSimone JA: Salt taste transduction occurs through an amiloride-sensitive sodium transport pathway. *Science*. 1984; 223(4634): 403–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
 31. Chandrashekar J, Hoon MA, Ryba NJ, *et al.*: The receptors and cells for mammalian taste. *Nature*. 2006; 444(7117): 288–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
 32. **F** Chandrashekar J, Kuhn C, Oka Y, *et al.*: The cells and peripheral representation of sodium taste in mice. *Nature*. 2010; 464(7286): 297–301.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 33. **F** Lewandowski BC, Sukumaran SK, Margolske RF, *et al.*: Amiloride-Insensitive Salt Taste Is Mediated by Two Populations of Type III Taste Cells with Distinct Transduction Mechanisms. *J Neurosci*. 2016; 36(6): 1942–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 34. **F** Roeber JK, Roper SD, Chaudhari N: The Role of the Anion in Salt (NaCl) Detection by Mouse Taste Buds. *J Neurosci*. 2019; 39(32): 6224–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 35. Vandenbeuch A, Clapp TR, Kinnamon SC: Amiloride-sensitive channels in type I fungiform taste cells in mouse. *BMC Neurosci*. 2008; 9: 1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 36. Desor JA, Finn J: Effects of amiloride on salt taste in humans. *Chem Senses*. 1989; 14(6): 793–803.
[Publisher Full Text](#)
 37. Hellekant G, Ninomiya Y, Danilova V: Taste in chimpanzees II: single chorda tympani fibers. *Physiol Behav*. 1997; 61(6): 829–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
 38. **F** Huang YJ, Maruyama Y, Dvoryanchikov G, *et al.*: The role of pannexin 1 hemichannels in ATP release and cell-cell communication in mouse taste buds. *Proc Natl Acad Sci U S A*. 2007; 104(15): 6436–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 39. **F** Murata Y, Yasuo T, Yoshida R, *et al.*: Action potential-enhanced ATP release from taste cells through hemichannels. *J Neurophysiol*. 2010; 104(2): 896–901.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 40. **F** Huang YJ, Maruyama Y, Lu KS, *et al.*: Mouse taste buds use serotonin as a neurotransmitter. *J Neurosci*. 2005; 25(4): 843–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 41. **F** Finger TE, Danilova V, Barrows J, *et al.*: ATP signaling is crucial for communication from taste buds to gustatory nerves. *Science*. 2005; 310(5753): 1495–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 42. **F** Larson ED, Vandenbeuch A, Voigt A, *et al.*: The Role of 5-HT₃ Receptors in Signaling from Taste Buds to Nerves. *J Neurosci*. 2015; 35(48): 15984–95.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 43. **F** Shin YK, Martin B, Golden E, *et al.*: Modulation of taste sensitivity by GLP-1 signaling. *J Neurochem*. 2008; 106(1): 455–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 44. **F** Takai S, Yasumatsu K, Inoue M, *et al.*: Glucagon-like peptide-1 is specifically involved in sweet taste transmission. *FASEB J*. 2015; 29(6): 2268–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 45. Huang YA, Dando R, Roper SD: Autocrine and paracrine roles for ATP and serotonin in mouse taste buds. *J Neurosci*. 2009; 29(44): 13909–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 46. Yang R, Crowley HH, Rock ME, *et al.*: Taste cells with synapses in rat circumvallate papillae display SNAP-25-like immunoreactivity. *J Comp Neurol*. 2000; 424(2): 205–15.
[PubMed Abstract](#) | [Publisher Full Text](#)
 47. DeFazio RA, Dvoryanchikov G, Maruyama Y, *et al.*: Separate populations of receptor cells and presynaptic cells in mouse taste buds. *J Neurosci*. 2006; 26(15): 3971–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 48. **F** Taruno A, Vingtdex V, Ohmoto M, *et al.*: CALHM1 ion channel mediates purinergic neurotransmission of sweet, bitter and umami tastes. *Nature*. 2013; 495(7440): 223–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 49. **F** Ma Z, Taruno A, Ohmoto M, *et al.*: CALHM3 Is Essential for Rapid Ion Channel-Mediated Purinergic Neurotransmission of GPCR-Mediated Tastes. *Neuron*. 2018; 98(3): 547–561.e10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 50. **F** Romanov RA, Lasher RS, High B, *et al.*: Chemical synapses without synaptic vesicles: Purinergic neurotransmission through a CALHM1 channel-mitochondrial signaling complex. *Sci Signal*. 2018; 11(529): pii: eaao1815.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 51. Kinnamon SC, Finger TE: A taste for ATP: neurotransmission in taste buds. *Front Cell Neurosci*. 2013; 7: 264.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

52. Vandenbeuch A, Anderson CB, Ford AP, *et al.*: **A selective P2X3, P2X2/3 receptor antagonist abolishes responses to all taste stimuli in mice.** *Chem Senses*. 2013; **38**: 86.
[PubMed Abstract](#) | [Publisher Full Text](#)
53. Iwatsuki K, Ichikawa R, Hiasa M, *et al.*: **Identification of the vesicular nucleotide transporter (VNUT) in taste cells.** *Biochem Biophys Res Commun*. 2009; **388**(1): 1–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
54. Matsumoto I, Ohmoto M, Narukawa M, *et al.*: **Skn-1a (Pou2f3) specifies taste receptor cell lineage.** *Nat Neurosci*. 2011; **14**(6): 685–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
55. Yarmolinsky DA, Zuker CS, Ryba NJ: **Common sense about taste: from mammals to insects.** *Cell*. 2009; **139**(2): 234–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. Usoskin D, Furlan A, Islam S, *et al.*: **Unbiased classification of sensory neuron types by large-scale single-cell RNA sequencing.** *Nat Neurosci*. 2015; **18**(1): 145–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
57. Ohman-Gault L, Huang T, Krimm R: **The transcription factor Phox2b distinguishes between oral and non-oral sensory neurons in the geniculate ganglion.** *J Comp Neurol*. 2017; **525**(18): 3935–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
58. Roper SD: **Taste buds as peripheral chemosensory processors.** *Semin Cell Dev Biol*. 2013; **24**(1): 71–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
59. Besnard P, Passilly-Degrace P, Khan NA: **Taste of Fat: A Sixth Taste Modality?** *Physiol Rev*. 2016; **96**(1): 151–76.
[PubMed Abstract](#) | [Publisher Full Text](#)
60. Matsumura S, Mizushige T, Yoneda T, *et al.*: **GPR expression in the rat taste bud relating to fatty acid sensing.** *Biomed Res*. 2007; **28**(1): 49–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
61. Gaillard D, Laugerette F, Darcel N, *et al.*: **The gustatory pathway is involved in CD36-mediated orosensory perception of long-chain fatty acids in the mouse.** *FASEB J*. 2008; **22**(5): 1458–68.
[PubMed Abstract](#) | [Publisher Full Text](#)
62. Yasumatsu K, Iwata S, Inoue M, *et al.*: **Fatty acid taste quality information via GPR120 in the anterior tongue of mice.** *Acta Physiol (Oxf)*. 2019; **226**(1): e13215.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
63. Stratford JM, Curtis KS, Contreras RJ: **Linoleic acid increases chorda tympani nerve responses to and behavioral preferences for monosodium glutamate by male and female rats.** *Am J Physiol Regul Integr Comp Physiol*. 2008; **295**(3): R764–R772.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Stephen Roper**

Department of Physiology & Biophysics and Department of Otolaryngology, University of Miami Miller School of Medicine, Miami, FL, 33136, USA

Competing Interests: No competing interests were disclosed.

2. **Susan Travers**

Division of Biosciences, College of Dentistry, Ohio State University, Columbus, OH, USA

Competing Interests: No competing interests were disclosed.

Comments on this article

Version 1

Reader Comment 23 Dec 2019

Anthony Sclafani, Brooklyn College of the City University of New York, Brooklyn, New York, USA

In addition to fatty taste, there is substantial evidence for the existence of a maltodextrin, or more specifically a maltooligosaccharide taste in various species including rats, mice and humans (Sclafani, 1987; Sclafani, 2004; Spector and Schier, 2018; Lim and Pullicin, 2019). This was first suggested by the attraction displayed by rats to maltodextrin solutions (e.g., Polycose) in brief-access or sham-feeding tests that minimize postoral nutrient feedback and by the weak cross-generalization of taste aversions conditioned to maltodextrin and sucrose (Sclafani, 1987). Subsequent studies revealed that deletion of the T1R2 and/or T1R3 sweet receptor elements substantially reduced the behavioral response of mice to sugars but had minimal effects on their response to maltodextrin (Treesukosol *et al.*, 2009; Zukerman *et al.*, 2009; Treesukosol *et al.*, 2011). Yet deletion of downstream taste signaling elements (α -gustducin, TRPM5, CALHM1, P2X2/P2X3) greatly attenuates the behavioral responses of mice to maltodextrin as well as sugar solutions (Sclafani *et al.*, 2007; Taruno *et al.*, 2013; Sclafani and Ackroff, 2014). Consistent with the mouse findings, blocking the sweet taste receptor with lactisole blocks sugar but not maltooligosaccharide detection in humans (Lim and Pullicin, 2019). Humans differ from rodents in that they do not

display a strong behavioral attraction to maltodextrin taste. However, the oral detection of maltodextrins is reported to improve human exercise performance (Jeukendrup and Chambers, 2010; Bortolotti *et al.*, 2014). Interestingly, there are no published reports on maltodextrin taste effects on exercise performance in rodents for comparison. Importantly, the identity of the putative maltooligosacchride taste receptor is unknown, which represents a major gap in our understanding of this taste modality.

Bortolotti, H., Altimari, L.R., Vitor-Costa, M. and Cyrino, E.S. (2014) *Mouth rinse with carbohydrate drink and performance: hypothesis of its central action: review article*. International SportMed Journal, 15, 84-95.

Jeukendrup, A.E. and Chambers, E.S. (2010) *Oral carbohydrate sensing and exercise performance*. Curr. Opin. Clin. Nutr. Metab. Care, 13, 447-451.

Lim, J. and Pullicin, A.J. (2019) *Oral carbohydrate sensing: Beyond sweet taste*. Physiol. Behav., 202, 14-25.

Sclafani, A. (1987) *Carbohydrate taste, appetite, and obesity: An overview*. Neurosci. Biobehav. Rev., 11, 131-153.

Sclafani, A. (2004) *The sixth taste*. Appetite, 43, 1-3.

Sclafani, A. and Ackroff, K. (2014) *Maltodextrin and fat preference deficits in "taste-blind" P2X2/P2X3 knockout mice*. Chem. Senses, 39, 507-514.

Sclafani, A., Zukerman, S., Glendinning, J.I. and Margolskee, R.F. (2007) *Fat and carbohydrate preferences in mice: The contribution of α -gustducin and Trpm5 taste signaling proteins*. Am. J. Physiol. Regul. Integr. Comp. Physiol., 293, R1504-R1513.

Spector, A.C. and Schier, L.A. (2018) *Behavioral evidence that select carbohydrate stimuli activate T1R-independent receptor mechanisms*. Appetite, 122, 26-31.

Taruno, A., Vingtdeux, V., Ohmoto, M., Ma, Z., Dvoryanchikov, G., Li, A., Adrien, L., Zhao, H., Leung, S., Abernethy, M., Koppel, J., Davies, P., Civan, M.M., Chaudhari, N., Matsumoto, I., Hellekant, G., Tordoff, M.G., Marambaud, P. and Foskett, J.K. (2013) *CALHM1 ion channel mediates purinergic neurotransmission of sweet, bitter and umami tastes*. Nature, 495, 223-226.

Treesukosol, Y., Blonde, G. and Spector, A.C. (2009) *The T1R2 and T1R3 subunits are individually unnecessary for normal affective licking responses to Polycose: Implications for saccharide taste receptors in mice*. Am. J. Physiol. Regul. Integr. Comp. Physiol., 296, R855-R865.

Competing Interests: No competing interests

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research