REVIEW

Sec translocon has an insertase-like function in addition to polypeptide conduction through the channel [version 1; peer review: 4 approved]

Koreaki Ito , Naomi Shimokawa-Chiba, Shinobu Chiba

Faculty of Life Sciences and Institute for Protein Dynamics, Kyoto Sangyo University, Kyoto, Japan

V1 First published: 20 Dec 2019, 8(F1000 Faculty Rev):2126
<https://doi.org/10.12688/f1000research.21065.1>
 Latest published: 20 Dec 2019, 8(F1000 Faculty Rev):2126
<https://doi.org/10.12688/f1000research.21065.1>

Abstract

The Sec translocon provides a polypeptide-conducting channel, which is insulated from the hydrophobic lipidic environment of the membrane, for translocation of hydrophilic passenger polypeptides. Its lateral gate allows a downstream hydrophobic segment (stop-transfer sequence) to exit the channel laterally for integration into the lipid phase. We note that this channel model only partly accounts for the translocon function. The other essential role of translocon is to facilitate *de novo* insertion of the N-terminal topogenic segment of a substrate polypeptide into the membrane. Recent structural studies suggest that *de novo* insertion does not use the polypeptide-conducting channel; instead, it takes place directly at the lateral gate, which is prone to opening. We propose that the *de novo* insertion process, in concept, is similar to that of insertases (such as YidC in bacteria and EMC3 in eukaryotes), in which an intramembrane surface of the machinery provides the halfway point of insertion.

Keywords

sec translocon, insertase, SecY, Sec61, membrane protein, YidC

Open Peer Review

Approval Status

	1	2	3	4
version 1
20 Dec 2019				

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Ross Dalbey**, The Ohio State University, Ohio, USA
2. **Gunnar von Heijne**, Stockholm University, Stockholm, Sweden
3. **Richard Zimmermann**, Saarland University, Homburg, Germany
4. **Arnold J M Driessen**, University of Groningen, Groningen, The Netherlands

Any comments on the article can be found at the end of the article.

Corresponding author: Koreaki Ito (kito@cc.kyoto-su.ac.jp)

Author roles: **Ito K:** Conceptualization, Investigation, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Shimokawa-Chiba N:** Investigation, Resources; **Chiba S:** Conceptualization, Investigation, Project Administration, Resources

Competing interests: No competing interests were disclosed.

Grant information: Work from the authors' laboratories was supported by grants from MEXT and JSPS Grant-in-Aid for Scientific Research (Grant No. 16H04788 to SC, 26116008 to SC and KI).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 Ito K *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Ito K, Shimokawa-Chiba N and Chiba S. **Sec translocon has an insertase-like function in addition to polypeptide conduction through the channel [version 1; peer review: 4 approved]** F1000Research 2019, 8(F1000 Faculty Rev):2126 <https://doi.org/10.12688/f1000research.21065.1>

First published: 20 Dec 2019, 8(F1000 Faculty Rev):2126 <https://doi.org/10.12688/f1000research.21065.1>

Introduction

In protein localization, hydrophobic segments of polypeptides play a central role by their ability to partition into the hydrophobic core of the membrane¹⁻³. At the same time, proteins that are integrated into the membrane or localized in extra-cytosolic locations require facilitation by specific cellular mechanisms in reaching their destinations and achieving their topographical relationships to the membrane. The difficulties arise because the cytosol, where the translation of genetic messages takes place, is compartmentalized by membranes to make it discrete from other cellular compartments and the surroundings. Consequently, newly synthesized polypeptides must follow a proper pathway typically involving bio-machinery to integrate into or cross the membrane. Such pathways alleviate the difficulty of overcoming specific energy barriers that the moving polypeptide encounters. Hydrophilic parts of polypeptides must overcome the hydrophobic barrier of the lipid hydrocarbon to cross the membrane, whereas hydrophobic polypeptide segments should overcome the hydrophilic barrier of the phospholipid head groups to partition into the lipidic membrane interior.

In this review, we attempt to clarify the role of the Sec translocon, a principal and conserved cellular machine, in assisting protein translocation across and insertion into the membrane, corresponding to the cytoplasmic membrane in the case of

bacteria and the endoplasmic reticulum (ER) membrane in the case of eukaryotic cells⁴. Specifically, we illuminate the importance of *de novo* insertion of a hydrophobic polypeptide stretch into the membrane.

Topogenic insertion sequences

Topogenic sequences responsible for *de novo* membrane insertion are called signal sequences in the case of secretory proteins and signal-anchor sequences in the case of integral membrane proteins. For this section, readers are advised to refer to [Figure 1](#), which is discussed later in this article in conjunction with the translocon functions. Topogenic sequences invariably contain core sequences enriched in hydrophobic residues, which insert into the membrane to span it with a specific orientation. The $C_{in}-N_{out}$ orientation is called type I and $N_{in}-C_{out}$ type II (“in” indicates the cytosolic side and “out” the *trans* side of the membrane [see [Figure 1](#)]). The orientation is determined by charge characters and length of the flanking hydrophilic regions¹.

A canonical signal sequence assumes the type II ($N_{in}-C_{out}$) orientation, hence inducing translocation of the mature region that follows it. The signal sequence then receives proteolytic cleavage on the *trans* side by the processing enzyme, signal peptidase, which liberates the mature domain from the membrane. Membrane anchor sequences do not undergo cleavage.

Figure 1. Model diagrams of the Sec-mediated *de novo* insertion of hydrophobic domains. The Sec translocon is schematically depicted to show the polypeptide-conducting channel in the center, the plug helix in blue, and the open lateral gate in yellow. The substrate polypeptide is shown in red, with the thick part representing a hydrophobic segment. R shows the resting (quiescent) state. N1 and N2 show insertion with the $C_{in}-N_{out}$ orientation. C1 through C6 shows insertion with the $N_{in}-C_{out}$ orientation. Targeting events before insertion could differ depending on the substrates or the organisms. The figure is intended to show the co-translational process with the ribosomal exit tunnel in gray and the tRNA at the growing end of the nascent polypeptide as green rectangles. The timings of plug dislocation and polypeptide enclosure within the channel are shown arbitrarily, as neither has been defined precisely. At least in an early stage of insertion, the plug still occludes the central pore⁵. Depending on the substrate proteins, the proteolytic cleavage of the signal peptide may or may not take place. Finally, if the second hydrophobic segment (stop-transfer sequence shown in purple) follows, the translocation step halts, and the hydrophobic segment exits the channel laterally to become a type I transmembrane domain, as shown in C6 (the orientation of the stop-transfer is type I, but, here, we classify different proteins by the mode of *de novo* insertion of their N-terminal regions). It should be noted that the above diagrams do not take into account the proposal that integration initially proceeds with the N_{out} orientation and reorientation events, later on, accompany the determination of type I versus type II routes⁶. ER, endoplasmic reticulum.

Although signal sequences and type II signal-anchor sequences are similar in their membrane topologies, they have different features, such as the length of the hydrophobic core⁷ and the predicted behaviors in lipidic environments⁸. Nevertheless, it is useful to assume that these two categories of topogenic sequences share the fundamental principle that underlies their membrane insertion. We point out two observations that support this view. First, leucine-based model sequences can use the common translocon (see below) and function like either a signal sequence or a signal-anchor sequence depending on the length of the leucine stretch and the presence or the absence of the signal peptidase cleavage motif⁷. Second, in engineered fusion proteins, transmembrane sequences can direct export of the mature domain of alkaline phosphatase⁹, a bacterial periplasmic protein, even with accompanying cleavage by signal peptidase¹⁰.

The $C_{in}-N_{out}$ (type I) transmembrane state can be generated by an N-terminal type I signal-anchor sequence as well as by an internal “stop-transfer” sequence that follows a *de novo* insertion signal (see Figure 1). In our view (see below), the stop-transfer sequence does not fall into the category of *de novo* insertion sequence; rather, it depends on the function of the preceding insertion sequence in the upstream region of the polypeptide. Type II *de novo* insertion signals can reside at internal sites such that multiple insertion events in combination with stop-transfer lead to the biogenesis of multi-spanning membrane proteins⁶. For insertion, the topogenic region of the polypeptide must be targeted to the insertion machinery of the membrane. A typical mechanism is signal recognition particle-mediated co-translational targeting, but we do not discuss targeting mechanisms¹¹ further in this review. Finally, a class of proteins called tail-anchored membrane proteins contains an insertion sequence near the C-terminal end, which must integrate into the membrane with the type II orientation post-translationally¹².

The Sec translocon paradigm must be expanded to account for *de novo* insertion

The determination of the Sec translocon structure provided the first look at a polypeptide-conducting channel that allows a substrate polypeptide to traverse the membrane without directly meeting the phospholipid constituent of the membrane^{5,13–16}. The conduit is gated by a plug-like helix that seals the channel in the quiescent state (Figure 2, panel 1), whereas it accommodates a translocating polypeptide in the active, open state with the dislocated plug (Figure 2, panel 2). The channel is shaped like an hourglass, with a gasket-like constriction at the center, which is formed by hydrophobic sidechains and prevents the leakage of small molecules during ongoing translocation (Figure 2, panel 3). The translocon is used not only for the complete export of proteins from the cytosol but also for the biogenesis of integral membrane proteins¹³. For this purpose, the translocon contains the lateral gate, which consists of a few of its transmembrane helices. In a well-known mode of translocon-facilitated integration, called stop-transfer^{6,17}, a hydrophobic segment of the polypeptide exits the channel laterally to reach the lipid phase of the membrane (Figure 2, panel 4). This is made possible because the lateral gate of the translocon opens in response to the entry of the hydrophobic segment into the channel, allowing its lateral partition into the lipid phase and acquisition of the $C_{in}-N_{out}$ transmembrane configuration^{13,18}.

This beautiful model of the polypeptide-conducting channel, however, leaves a critical question unanswered: how is the translocating polypeptide accommodated in the channel? *De novo* insertion of an N-terminal polypeptide segment into the membrane will bring the rest of the nascent/newly synthesized polypeptide into the channel interior. *De novo* insertion, which must be induced by the upstream topogenic signal, is a prerequisite for the insertion of a stop-transfer-type hydrophobic segment through the lateral gate into the lipid phase.

Figure 2. Simplified views of the Sec translocon. Panels 1 and 2 show vertical cutaway images, and panels 3 and 4 show horizontal cutaway images viewed from the trans side. The pore ring, shown in green, is at the constriction of the channel and consists of hydrophobic amino acids that surround the translocating polypeptide (panel 3). The plug, shown in blue, keeps the vertical gate closed in the resting state (panel 1) and is dislocated to open the gate in the working state (panel 2). The translocating polypeptide is shown by the red line (panel 2) or cutaway disc (panel 3). The signal peptide is shown by the thick line (panel 2) in the state already disengaged from the translocon and proteolytically processed. Earlier events of signal peptide insertion into the membrane are shown in Figure 1. The purple line (panel 2) and purple disc (panel 4) show a downstream hydrophobic segment that exits the channel laterally via the open lateral gate (panel 4). This figure was prepared by referring to the structural depiction presented as Figure 1 in Rapoport *et al.*¹⁹.

In the original publication of the translocon structure, the authors explained that the signal peptide may use a lipid-translocon interface at the lateral gate region to integrate into the membrane¹³, and recent structural studies support this notion (see below). Probably because the model of the polypeptide-conducting channel is so compelling, however, discussion of membrane integration processes of signal peptides and signal-anchor peptides sometimes starts with an implicit assumption that they reside initially within the polypeptide-conducting channel of the translocon. Recent studies disfavor this assumption (see below). It is of vital importance to elucidate the actual pathways and mechanisms used by Sec translocon for *de novo* membrane insertion of hydrophobic stretches. Before looking into this function of the translocon, we will summarize the importance and the mechanism of actions of the conserved membrane protein biogenesis factors called insertases.

The YidC insertase provides an intramembrane platform that facilitates membrane insertion of a class of membrane protein

We point out that the *de novo* integration function of translocon can be viewed as similar in concept to that of “insertases”, which occur in bacteria (the plasma membrane), mitochondria, chloroplasts, and eukaryotes (the ER membrane). Among them, YidC in bacteria is best characterized in its high-resolution structures^{20–23} and its insertase function executed independently of the Sec machinery^{24–26}. YidC facilitates the insertion of a class of simple membrane proteins with one or two transmembrane segment(s) and a small periplasmic (trans-side) domain. Structural, genetic, and biochemical studies suggest that the functional unit of YidC is a monomer^{20,21,23,27,28}. Its transmembrane helices form an intramembrane cavity that is open toward the cytosol and the membrane interior (Figure 3); the cavity is embraced by horizontal helices on the cytosolic and the

periplasmic sides^{20–22}. Notably, the cavity is highly hydrophilic and water-accessible, and these features proved to be functionally important^{29,30}. The presence of an arginine deep in the cavity wall is required for the insertion of a class of substrate proteins with negative charges on the periplasmic tail (Figure 3). The transient charge attraction is one of the strategies that YidC adopts for the insertase function^{20,31–33}, although it is not the exclusive mechanism of insertion³⁴.

In contrast to the translocon channel, which is optimized to accommodate a long and moving hydrophilic polypeptide chain, the insertase does not enclose its substrate (Figure 3). The current model of the YidC insertase suggests that it provides an intramembrane platform that a substrate polypeptide uses as the halfway point of insertion^{20,29}. The hydrophilic cavity that is exposed to the lipid phase could cause hydrophobic mismatch problems, which may be alleviated in part by the flexible cytoplasmic loop that covers it²², but could nevertheless disturb the bilayer structure, which might contribute to the insertion mechanism³⁰. The cavity would lower the energy cost of translocation of the short and hydrophilic periplasmic tail by binding to it transiently and circumventing the hydrophobic barrier. This event may (further) disturb the phospholipid organization and lessens the head group barrier against the hydrophobic segment partitioning into the lipidic phase, which will, in turn, drive concomitant translocation of the hydrophilic tail from the halfway point to the periplasmic side (Figure 3). Circumvention of one barrier or a chain of barriers sequentially could trigger the forward movement of the substrate to the energetically stable, membrane-integrated state. This principle could work for *de novo* insertion of a variety of transmembrane sequences if amphipathic arrangements of the intramembrane platforms are adequately tuned to make them usable as the halfway point. It should be noted that such a platform can be formed dynamically upon access by a substrate protein.

Figure 3. A model diagram of the insertase function. The YidC insertase provides a hydrophilic cavity within the membrane, which a substrate polypeptide uses as the halfway point of insertion²⁰. The figure depicts a possible intermediate state of a class of substrates that have a negatively charged periplasmic tail, which may thereafter be translocated to the periplasmic side, as shown by the arrow, in coordination with the hydrophobic segment partition into the lipid phase.

Insertases as ubiquitous factors in biological kingdoms

Recent studies reveal that eukaryotic cells also possess insertases in the ER membrane. They include Get1/2 and the ER membrane complex 3 (EMC3) subunit of the EMC^{35,36}. Archaea also have a YidC-like protein³⁷. Eukaryotic Get1/2 mediates the insertion of tail-anchored proteins as an insertase^{12,38}. The bacterial YidC insertase also facilitates the biogenesis of tail-anchored proteins^{39–41}. The EMC also functions as an insertase for a subset of tail-anchored membrane proteins⁴². The EMC is involved in the biogenesis of other types of membrane proteins as well, including multi-spanning membrane proteins with moderately hydrophobic transmembrane regions^{36,43,44}. At least in one analyzed case, the EMC acts as an insertase for the insertion of the first transmembrane segment of a multi-spanning membrane protein⁴⁵.

Both YidC and the EMC have been reported to cooperate with the Sec translocon in assisting the correct biogenesis of membrane proteins that have multiple transmembrane regions. They may contribute to this mode of function by fulfilling chaperone-like roles that facilitate the “folding” of membrane

proteins^{36,43,46}. However, the division of labor between insertase and Sec in inserting different transmembrane domains could also explain the apparent chaperone function of the EMC/YidC. Specifically, the EMC's ability to facilitate the insertion of the first transmembrane segment of a multi-spanning membrane protein substrate in the correct orientation allows for the correct insertion of the remaining transmembrane regions by the Sec machinery⁴⁵. The general applicability of such a relay mechanism for insertase–Sec-mediated membrane protein biogenesis remains to be investigated.

The translocon lateral gate is dynamic in handling hydrophobic topogenic segments of substrates

High-resolution structural information on the Sec translocon in its working state started to accumulate recently through both X-ray crystallography and electron microscopy-based approaches, the latter of which is undergoing remarkable technical progress. Also, efforts are being made to determine structures of translocon that is membrane integrated, rather than after isolation in detergent-solubilized states, by use of lipidic cubic phase crystals¹⁵, reconstitution into the nanodisc bilayer^{5,47,48}, and cryo-electron tomography of native membranes from cells⁴⁹. A central point in understanding the roles of the Sec translocon in mediating *de novo* membrane insertion of hydrophobic domains is the roles played by the lateral gate region. A current consensus is that the lateral gate region can accommodate (transiently) the signal sequence or the membrane anchor sequence of either orientation, in its “open” form with a ~22° relative rotation of the N- and C-terminal halves of SecY/SecE61α^{5,47,50–53}, which is also the case for the lateral exit of a stop-transfer-type hydrophobic segment⁵⁴. The substrate helix enters this region in a manner replacing a “placeholder” helix of the translocon⁵¹ and opening the lateral gate toward the lipid phase and also to the trans-side (or the cytosolic side depending on the stages of insertion⁵) of the membrane^{5,47,48,52,55–57}.

While the *de novo* inserting helix should end up in the lipid-embedded state⁵⁸, the snapshot pictures do not reveal a temporal order of the events in the sequential insertion process. There are crucial questions: when and how is the lateral gate open partially or fully, and when is the plug helix displaced to create a trans-membrane channel? The literature on the timing of lateral gate opening is disparate at a glance, but we can extract one crucial characteristic of the lateral gate: it is mobile and prone to opening to varying extents under different conditions. Structural studies detected a crack on the cytosolic side of the translocon, where some polypeptide moiety from an adjacent molecule resided, possibly as a signal peptide mimic. Substrate polypeptides may use such a crack for the initial recognition and entry step in both the co- and post-translational pathways^{14–16,51,57} (Figure 1, C1).

On the other hand, the targeting/driving partners of the translocon, the ribosome or the SecA motor, can induce partial opening of the lateral gate without involving any substrate polypeptide^{16,59}. Strikingly, cryo-electron tomography studies of the membrane-embedded Sec61–ribosome complex show that the

translocon even assumes a fully open conformation in the native membrane without involving any polypeptide substrate⁴⁹. Also, Sec62/63, the Sec61 partner in the post-translational mode of translocation, causes large lateral gate opening in the absence of substrate^{60,61}.

A recent study revealed an early intermediate structure of the signal peptide insertion process⁵, in which nanodisc-integrated SecYEG was complexed with the nascent chain ribosome complex that has a 48-amino-acid-long N-terminal part exposed outside the ribosome. In this structure, the space created by lateral gate helices is open toward the cytosol, while the plug helix still closes the channel. The early intermediate structure differs from the structure determined for the later stage of insertion with the periplasmic-side-open (cytosolic-side-closed) lateral gate accompanied by a ~22° oblique rotation of a set of helices⁵¹. Thus, dynamic conformational changes of translocon range from a crack formation on the cytosolic side, through the cytoplasmic opening of the lateral gate, to the full opening of the lateral gate (Figure 1, C1 to C3). As already mentioned, the dynamic nature of the lateral gate region even allows its full opening before the substrate's entry^{49,60,61}.

The Sec translocon function to mediate *de novo* membrane insertion can be viewed as insertase-like

With the current knowledge of the insertion process discussed above, it is still unclear how the “passenger” hydrophilic polypeptide following the N-terminal signal is accommodated within the polypeptide-conducting channel (Figure 1, C4). The plug displacement is a relatively late event, as several structures have been reported in which the plug is not yet displaced in translocon complexes that are already engaged with a substrate or a substrate mimic^{5,15,50,54}. These observations indicate that early events of insertion proceed before the opening of the polypeptide-conducting channel (Figure 1, N1 and C1–C2). Probably, in *de novo* insertion, substrate polypeptides interact directly with the lateral gate region, which undergoes dynamic remodeling of the constituent transmembrane helices. Although the signal sequence transiently intercalates between some transmembrane helices of the translocon⁵¹, its residues also face the lipid phase in the early and later steps of insertion^{5,47,48,52,55,56}. We envision that the rearranging lateral gate region provides intramembrane platforms, whose hydrophobic properties lower the energy cost of insertion by interacting with the incoming polypeptide. In an extreme case, the strongly hydrophobic substrate might slide along the lipid–translocon interface, as proposed by Cymer *et al.*¹, which is supported by thermodynamic considerations¹.

The Sec translocon and YidC provide the client polypeptide with thermodynamically similar environments that support membrane protein partition⁶², suggesting that they share the underlying principles. While the initial crack on the cytosolic side of the translocon forms a hydrophobic patch^{15,51}, the “primed” lateral gate has a hydrophilic seam⁵¹. Upon further opening, the lateral gate cavity will have hydrophobic helices in front⁵¹, but it could also have some hydrophilic parts on the wall deep inside⁵⁰, potentially akin to the YidC cavity. Substrate polypeptides

may use the changing hydrophobic characters of the lateral gate in their membrane insertion steps. A recent report by He *et al.*⁶³ proposes that two transmembrane segments of *Escherichia coli* YidC provide a “greasy slide” for a hydrophobic core of a client polypeptide to interact transiently during the insertion process. The insertion of the hydrophobic segment in a loop-like conformation appears to precede the hydrophilic N-tail interaction with the hydrophilic groove of YidC. Thus, an insertase can have multiple intramembrane surfaces of different amphipathic characters, in line with our proposal that the Sec lateral gate could function in similar fashions.

The concept of the polypeptide-conducting channel goes back to the proposal by Blobel and Dobberstein in 1975⁶⁴, followed by genetic identification of SecY⁶⁵ and SecE⁶⁶, biochemical demonstration of their translocase functions^{67–69}, and, finally, the realization of its structural entity 29 years later by van den Berg *et al.*¹³. These studies have been influential for our understanding of living cells. We now learn that the translocon must

integrate its insertase-like functions and the polypeptide-conduction function to execute the biological task. The thermodynamic principle of membrane insertion suggests the importance of the lipidic constituents of the membrane in *de novo* polypeptide insertion. Indeed, Nishiyama and coworkers have identified a glycolipid molecule, termed MPIase, that is dedicated to protein integration into the membrane in *Escherichia coli*^{70–72}. The responsible enzyme is conserved in prokaryotic and eukaryotic cells. Now, we are at a stage where studies from various directions merge to enable us to better understand the biogenesis of membrane proteins.

Acknowledgments

We thank Tomoya Tsukazaki of Nara Institute of Science and Technology, and Hiroyuki Mori and Yoshinori Akiyama of Kyoto University for helpful comments and discussion on this review article.

References

1. Cymer F, von Heijne G, White SH: **Mechanisms of integral membrane protein insertion and folding.** *J Mol Biol.* 2015; **427**(5): 999–1022.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
2. Hessa T, Kim H, Bihlmaier K, *et al.*: **Recognition of transmembrane helices by the endoplasmic reticulum translocon.** *Nature.* 2005; **433**(7024): 377–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
3. Gumbart JC, Chipot C: **Decrypting protein insertion through the translocon with free-energy calculations.** *Biochim Biophys Acta.* 2016; **1858**(7 Pt B): 1663–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Rapoport TA: **Protein translocation across the eukaryotic endoplasmic reticulum and bacterial plasma membranes.** *Nature.* 2007; **450**(7170): 663–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Kater L, Frieg B, Berninghausen O, *et al.*: **Partially inserted nascent chain unzips the lateral gate of the Sec translocon.** *EMBO Rep.* 2019; **20**(10): e48191.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
6. Spiess M, Junne T, Janoschke M: **Membrane Protein Integration and Topogenesis at the ER.** *Protein J.* 2019; **38**(3): 306–316.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
7. Nilsson I, Whitley P, von Heijne G: **The COOH-terminal ends of internal signal and signal-anchor sequences are positioned differently in the ER translocase.** *J Cell Biol.* 1994; **126**(5): 1127–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. de Vrije GJ, Batenburg AM, Killian JA, *et al.*: **Lipid involvement in protein translocation in *Escherichia coli*.** *Mol Microbiol.* 1990; **4**(1): 143–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Manoil C, Beckwith J: **A genetic approach to analyzing membrane protein topology.** *Science.* 1986; **233**(4771): 1403–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Akiyama Y, Inada T, Nakamura Y, *et al.*: **SecY, a multispanning integral membrane protein, contains a potential leader peptidase cleavage site.** *J Bacteriol.* 1990; **172**(6): 2888–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Guna A, Hegde RS: **Transmembrane Domain Recognition during Membrane Protein Biogenesis and Quality Control.** *Curr Biol.* 2018; **28**(8): R498–R511.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
12. Wang F, Chan C, Weir NR, *et al.*: **The Get1/2 transmembrane complex is an endoplasmic-reticulum membrane protein insertase.** *Nature.* 2014; **512**(7515): 441–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
13. Van den Berg B, Clemons WM Jr, Collinson I, *et al.*: **X-ray structure of a protein-conducting channel.** *Nature.* 2004; **427**(6969): 36–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
14. Tsukazaki T, Mori H, Fukai S, *et al.*: **Conformational transition of Sec machinery inferred from bacterial SecYE structures.** *Nature.* 2008; **455**(7215): 988–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Tanaka Y, Sugano Y, Takemoto M, *et al.*: **Crystal Structures of SecYEG in Lipidic Cubic Phase Elucidate a Precise Resting and a Peptide-Bound State.** *Cell Rep.* 2015; **13**(8): 1561–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Voorhees RM, Fernández IS, Scheres SH, *et al.*: **Structure of the mammalian ribosome-Sec61 complex to 3.4 Å resolution.** *Cell.* 2014; **157**(7): 1632–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
17. Yost CS, Hedgpeth J, Lingappa VR: **A stop transfer sequence confers predictable transmembrane orientation to a previously secreted protein in cell-free systems.** *Cell.* 1983; **34**(3): 759–66.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Rapoport TA, Goder V, Heinrich SU, *et al.*: **Membrane-protein integration and the role of the translocation channel.** *Trends Cell Biol.* 2004; **14**(10): 568–75.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Rapoport TA, Li L, Park E: **Structural and Mechanistic Insights into Protein Translocation.** *Annu Rev Cell Dev Biol.* 2017; **33**: 369–390.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. Kumazaki K, Chiba S, Takemoto M, *et al.*: **Structural basis of Sec-independent membrane protein insertion by YidC.** *Nature.* 2014; **509**(7501): 516–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
21. Kumazaki K, Kishimoto T, Furukawa A, *et al.*: **Crystal structure of *Escherichia coli* YidC, a membrane protein chaperone and insertase.** *Sci Rep.* 2014; **4**: 7299.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Tanaka Y, Izumioka A, Abdul Hamid A, *et al.*: **2.8-Å crystal structure of *Escherichia coli* YidC revealing all core regions, including flexible C2 loop.** *Biochem Biophys Res Commun.* 2018; **505**(1): 141–145.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
23. Xin Y, Zhao Y, Zheng J, *et al.*: **Structure of YidC from *Thermotoga maritima* and its implications for YidC-mediated membrane protein insertion.** *FASEB J.* 2018; **32**(5): 2411–2421.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Samuelson JC, Chen M, Jiang F, *et al.*: **YidC mediates membrane protein insertion in bacteria.** *Nature.* 2000; **406**(6796): 637–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Kiefer D, Kuhn A: **YidC-mediated membrane insertion.** *FEMS Microbiol Lett.* 2018; **365**(12).
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
26. Shanmugam SK, Dalbey RE: **The Conserved Role of YidC in Membrane Protein Biogenesis.** In: Sandkvist, Cascales, Christie, editors. *Protein Secretion in Bacteria.* American Society of Microbiology. 2019; **7**(1): 43–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Spann D, Pross E, Chen Y, *et al.*: **Each protomer of a dimeric YidC functions as a single membrane insertase.** *Sci Rep.* 2018; **8**(1): 589.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

28. Kedrov A, Wickles S, Crevenna AH, *et al.*: **Structural Dynamics of the YidC: Ribosome Complex during Membrane Protein Biogenesis.** *Cell Rep.* 2016; 17(11): 2943–2954.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Shimokawa-Chiba N, Kumazaki K, Tsukazaki T, *et al.*: **Hydrophilic microenvironment required for the channel-independent insertase function of YidC protein.** *Proc Natl Acad Sci U S A.* 2015; 112(16): 5063–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. **F** Chen Y, Capponi S, Zhu L, *et al.*: **YidC Insertase of *Escherichia coli*: Water Accessibility and Membrane Shaping.** *Structure.* 2017; 25(9): 1403–1414.e3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
31. Price CE, Driessen AJ: **Conserved negative charges in the transmembrane segments of subunit K of the NADH:ubiquinone oxidoreductase determine its dependence on YidC for membrane insertion.** *J Biol Chem.* 2010; 285(6): 3575–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. **F** Gray AN, Henderson-Frost JM, Boyd D, *et al.*: **Unbalanced charge distribution as a determinant for dependence of a subset of *Escherichia coli* membrane proteins on the membrane insertase YidC.** *mBio.* 2011; 2(6): pii: e00238–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
33. **F** Zhu L, Wasey A, White SH, *et al.*: **Charge composition features of model single-span membrane proteins that determine selection of YidC and SecYEG translocase pathways in *Escherichia coli*.** *J Biol Chem.* 2013; 288(11): 7704–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
34. Chen Y, Soman R, Shanmugam SK, *et al.*: **The role of the strictly conserved positively charged residue differs among the Gram-positive, Gram-negative, and chloroplast YidC homologs.** *J Biol Chem.* 2014; 289(51): 35656–67.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. **F** Anghel SA, McGilvray PT, Hegde RS, *et al.*: **Identification of Oxa1 Homologs Operating in the Eukaryotic Endoplasmic Reticulum.** *Cell Rep.* 2017; 21(13): 3708–3716.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
36. **F** Chitwood PJ, Hegde RS: **The Role of EMC during Membrane Protein Biogenesis.** *Trends Cell Biol.* 2019; 29(5): 371–384.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
37. **F** Borowska MT, Dominik PK, Anghel SA, *et al.*: **A YidC-like Protein in the Archaeal Plasma Membrane.** *Structure.* 2015; 23(9): 1715–1724.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
38. Zalisko BE, Chan C, Denic V, *et al.*: **Tail-Anchored Protein Insertion by a Single Get1/2 Heterodimer.** *Cell Rep.* 2017; 20(10): 2287–2293.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. **F** Pross E, Soussoula L, Seil I, *et al.*: **Membrane Targeting and Insertion of the C-Tail Protein SciP.** *J Mol Biol.* 2016; 428(20): 4218–4227.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
40. Peschke M, Le Goff M, Konings GM, *et al.*: **SRP, FtsY, DnaK and YidC Are Required for the Biogenesis of the *E. coli* Tail-Anchored Membrane Proteins DJC and Fik.** *J Mol Biol.* 2018; 430(3): 389–403.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Aschtgen MS, Zoued A, Llobès R, *et al.*: **The C-tail anchored TssL subunit, an essential protein of the enteroglycocalyx *Escherichia coli* Sci-1 Type VI secretion system, is inserted by YidC.** *MicrobiologyOpen.* 2012; 1(1): 71–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. **F** Guna A, Volkmar N, Christianson JC, *et al.*: **The ER membrane protein complex is a transmembrane domain insertase.** *Science.* 2018; 359(6374): 470–473.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
43. **F** Shurtleff MJ, Itzhak DN, Hussmann JA, *et al.*: **The ER membrane protein complex interacts cotranslationally to enable biogenesis of multipass membrane proteins.** *eLife.* 2018; 7: pii: e37018.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
44. Tian S, Wu Q, Zhou B, *et al.*: **Proteomic Analysis Identifies Membrane Proteins Dependent on the ER Membrane Protein Complex.** *Cell Rep.* 2019; 28(10): 2517–2526. e5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
45. **F** Chitwood PJ, Juszkievicz S, Guna A, *et al.*: **EMC Is Required to Initiate Accurate Membrane Protein Topogenesis.** *Cell.* 2018; 175(6): 1507–1519.e16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
46. Nagamori S, Smirnova IN, Kaback HR: **Role of YidC in folding of polytopic membrane proteins.** *J Cell Biol.* 2004; 165(1): 53–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
47. **F** Ma C, Wu X, Sun D, *et al.*: **Structure of the substrate-engaged SecA-SecY protein translocation machine.** *Nat Commun.* 2019; 10(1): 2872.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
48. **F** Frauenfeld J, Gumbart J, Sluis EO, *et al.*: **Cryo-EM structure of the ribosome-SecY complex in the membrane environment.** *Nat Struct Mol Biol.* 2011; 18(5): 614–21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
49. Pfeffer S, Burbaum L, Unverdorben P, *et al.*: **Structure of the native Sec61 protein-conducting channel.** *Nat Commun.* 2015; 6: 8403.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Egea PF, Stroud RM: **Lateral opening of a translocon upon entry of protein suggests the mechanism of insertion into membranes.** *Proc Natl Acad Sci U S A.* 2010; 107(40): 17182–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
51. **F** Voorhees RM, Hegde RS: **Structure of the Sec61 channel opened by a signal sequence.** *Science.* 2015; 351(6268): 88–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
52. **F** Park E, Ménétret JF, Gumbart JC, *et al.*: **Structure of the SecY channel during initiation of protein translocation.** *Nature.* 2014; 506(7486): 102–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
53. **F** Jomaa A, Boehringer D, Leibundgut M, *et al.*: **Structures of the *E. coli* translating ribosome with SRP and its receptor and with the translocon.** *Nat Commun.* 2016; 7: 10471.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
54. **F** Gogala M, Becker T, Beatrix B, *et al.*: **Structures of the Sec61 complex engaged in nascent peptide translocation or membrane insertion.** *Nature.* 2014; 506(7486): 107–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
55. Plath K, Mothes W, Wilkinson BM, *et al.*: **Signal sequence recognition in posttranslational protein transport across the yeast ER membrane.** *Cell.* 1998; 94(6): 795–807.
[PubMed Abstract](#) | [Publisher Full Text](#)
56. Urbanus ML, Scotti PA, Froderberg L, *et al.*: **Sec-dependent membrane protein insertion: sequential interaction of nascent FtsQ with SecY and YidC.** *EMBO Rep.* 2001; 2(6): 524–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
57. MacKinnon AL, Paavilainen VO, Sharma A, *et al.*: **An allosteric Sec61 inhibitor traps nascent transmembrane helices at the lateral gate.** *eLife.* 2014; 3: e01483.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
58. Bischoff L, Wickles S, Berninghausen O, *et al.*: **Visualization of a polytopic membrane protein during SecY-mediated membrane insertion.** *Nat Commun.* 2014; 5: 4103.
[PubMed Abstract](#) | [Publisher Full Text](#)
59. **F** Zimmer J, Nam Y, Rapoport TA: **Structure of a complex of the ATPase SecA and the protein-translocation channel.** *Nature.* 2008; 455(7215): 936–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
60. **F** Wu X, Cabanos C, Rapoport TA: **Structure of the post-translational protein translocation machinery of the ER membrane.** *Nature.* 2019; 566(7742): 136–139.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
61. **F** Itskanov S, Park E: **Structure of the posttranslational Sec protein-translocation channel complex from yeast.** *Science.* 2019; 363(6422): 84–87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
62. Ojemalm K, Botelho SC, Stüdle C, *et al.*: **Quantitative analysis of SecYEG-mediated insertion of transmembrane α -helices into the bacterial inner membrane.** *J Mol Biol.* 2013; 425(15): 2813–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
63. He H, Kuhn A, Dalbey RE: **Tracking the Stepwise Movement of a Membrane-inserting Protein *In Vivo*.** *J Mol Biol.* 2019; pii: S0022-2836(19)30606-0.
[PubMed Abstract](#) | [Publisher Full Text](#)
64. Blobel G, Dobberstein B: **Transfer of proteins across membranes. I. Presence of proteolytically processed and unprocessed nascent immunoglobulin light chains on membrane-bound ribosomes of murine myeloma.** *J Cell Biol.* 1975; 67(3): 835–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
65. Ito K, Wittekind M, Nomura M, *et al.*: **A temperature-sensitive mutant of *E. coli* exhibiting slow processing of exported proteins.** *Cell.* 1983; 32(3): 789–97.
[PubMed Abstract](#) | [Publisher Full Text](#)
66. Deshaies RJ, Schekman R: **A yeast mutant defective at an early stage in import of secretory protein precursors into the endoplasmic reticulum.** *J Cell Biol.* 1987; 105(2): 633–45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
67. Brundage L, Hendrick JP, Schiebel E, *et al.*: **The purified *E. coli* integral membrane protein SecY/E is sufficient for reconstitution of SecA-dependent precursor protein translocation.** *Cell.* 1990; 62(4): 649–57.
[PubMed Abstract](#) | [Publisher Full Text](#)
68. Akimaru J, Matsuyama S, Tokuda H, *et al.*: **Reconstitution of a protein translocation system containing purified SecY, SecE, and SecA from *Escherichia coli*.** *Proc Natl Acad Sci U S A.* 1991; 88(15): 6545–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
69. Görlich D, Rapoport TA: **Protein translocation into proteoliposomes reconstituted from purified components of the endoplasmic reticulum membrane.** *Cell.* 1993; 75(4): 615–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
70. Nishiyama K, Maeda M, Yanagisawa K, *et al.*: **MPlase is a glycolipozyme essential for membrane protein integration.** *Nat Commun.* 2012; 3: 1260.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
71. **F** Sawasato K, Sato R, Nishikawa H, *et al.*: **CdsA is involved in biosynthesis of glycolipid MPlase essential for membrane protein integration *in vivo*.** *Sci Rep.* 2019; 9(1): 1372.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
72. Sasaki M, Nishikawa H, Suzuki S, *et al.*: **The bacterial protein YidC accelerates MPlase-dependent integration of membrane proteins.** *J Biol Chem.* 2019; 294(49): 18898–18908.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

- 1. Arnold J M Driessen**
Molecular Microbiology, Groningen Biomolecular Sciences and Biotechnology Institute, and the Zernike Institute of Advanced Materials, University of Groningen, Groningen, The Netherlands
Competing Interests: No competing interests were disclosed.
- 2. Richard Zimmermann**
Medical Biochemistry and Molecular Biology, Saarland University, Homburg, Germany
Competing Interests: No competing interests were disclosed.
- 3. Gunnar von Heijne**
Department of Biochemistry and Biophysics, Stockholm University, Stockholm, Sweden
Competing Interests: No competing interests were disclosed.
- 4. Ross Dalbey**
Department of Chemistry and Biochemistry, The Ohio State University, Ohio, USA
Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research