REVIEW

Stress as a potential moderator of ovarian hormone influences on binge eating in women [version 1; peer review: 2 approved]

Natasha Fowler¹, Phuong T. Vo¹, Cheryl L. Sisk², Kelly L. Klump¹

¹Department of Psychology, Michigan State University, 316 Physics Road, East Lansing, MI, 48824-1116, USA

²Neuroscience Program, Michigan State University, 293 Farm Lane, East Lansing, MI, 48824-1116, USA

V1 First published: 27 Feb 2019, 8(F1000 Faculty Rev):222
<https://doi.org/10.12688/f1000research.16895.1>

Latest published: 27 Feb 2019, 8(F1000 Faculty Rev):222
<https://doi.org/10.12688/f1000research.16895.1>

Abstract

Previous research has demonstrated significant associations between increased levels of ovarian hormones and increased rates of binge eating (BE) in women. However, whereas all women experience fluctuations in ovarian hormones across the menstrual cycle, not all women binge eat in response to these fluctuations, suggesting that other factors must contribute. Stress is one potential contributing factor. Specifically, it may be that hormone-BE associations are stronger in women who experience high levels of stress, particularly as stress has been shown to be a precipitant to BE episodes in women. To date, no studies have directly examined stress as a moderator of hormone-BE associations, but indirect data (that is, associations between BE and stress and between ovarian hormones and stress) could provide initial clues about moderating effects. Given the above, the purpose of this narrative review was to evaluate these indirect data and their promise for understanding the role of stress in hormone-BE associations. Studies examining associations between all three phenotypes (that is, ovarian hormones, stress, and BE) in animals and humans were reviewed to provide the most thorough and up-to-date review of the literature on the potential moderating effects of stress on ovarian hormone-BE associations. Overall, current evidence suggests that associations between hormones and BE may be stronger in women with high stress levels, possibly via altered hypothalamic-pituitary-adrenal axis response to stress and increased sensitivity to and altered effects of ovarian hormones during stress. Additional studies are necessary to directly examine stress as a moderator of ovarian hormone-BE associations and identify the mechanisms underlying these effects.

Keywords

stress, ovarian hormones, binge eating, bulimia nervosa, binge eating disorder, estrogen, progesterone, women

Open Peer Review

Approval Status

	1	2
version 1
27 Feb 2019		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Jessica Baker**, University of North Carolina, Chapel Hill, USA
2. **Yong Xu**, Baylor College of Medicine, Houston, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Kelly L. Klump (klump@msu.edu)

Author roles: **Fowler N:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Vo PT:** Writing – Review & Editing; **Sisk CL:** Writing – Review & Editing; **Klump KL:** Conceptualization, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by grants awarded to KKL and CLS by the National Institute of Mental Health (NIMH) (MH 111715-01A1S1). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIMH. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2019 Fowler N *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Fowler N, Vo PT, Sisk CL and Klump KL. **Stress as a potential moderator of ovarian hormone influences on binge eating in women [version 1; peer review: 2 approved]** F1000Research 2019, 8(F1000 Faculty Rev):222 <https://doi.org/10.12688/f1000research.16895.1>

First published: 27 Feb 2019, 8(F1000 Faculty Rev):222 <https://doi.org/10.12688/f1000research.16895.1>

Introduction

Binge eating (BE) is a core symptom of many eating disorders, such as anorexia nervosa (AN), bulimia nervosa (BN), and binge eating disorder (BED)¹. BE is defined as repeated overconsumption of an objectively large amount of food (for example, >1000 calories) with a loss of control over the eating².

Although the etiology of BE remains largely unknown, emerging data suggest a role for ovarian hormones^{3–7}. Indeed, both animal^{8–11} and human^{3–7} studies suggest that changing levels of both estrogen and progesterone likely contribute to risk for BE and possibly its predominance in females. Interestingly, whereas all women experience fluctuations in ovarian hormones, not all women develop BE. The prevalence of BE in the population ranges from 1.5%¹² to 26%¹³, suggesting that there are likely moderators of the hormone-BE association that explain why some women binge eat in response to hormone fluctuations whereas others do not.

This review aimed to explore one such moderator—stress—defined in this review as the psychological perception of a stressful life event or the physiological response to a stressful life event, which is measured via hypothalamic–pituitary–adrenal (HPA) axis responsivity—and its impact on hormone-BE associations in women. Stress is an ideal candidate as a moderator of these associations. BE increases with increasing levels of stress^{14–17}, and stress responsivity varies across the menstrual cycle in women¹⁸. Thus, ovarian hormone–BE associations may be particularly strong in women with higher stress levels. If this is the case, then stress may account for why some, but not all, women binge eat in response to hormonal risk.

This review briefly describes the literature linking ovarian hormones to BE in women and reviews the potential role of stress responses in ovarian hormone–BE associations. Whenever possible, data from animal and human studies are explored to provide the most comprehensive picture of stress effects. Owing to the effects of starvation on ovarian hormones (for example, lack of cyclicity and loss of monthly cycles), most studies have not examined ovarian hormone effects on AN. For the purpose of this review, studies were limited to human and animal models of BE, BED, and BN. We end by suggesting areas for future research with the overarching goal of informing etiological models to understand why some, but not all, women binge eat during high-risk hormonal periods.

Ovarian hormones and binge eating

Animal studies

Animal studies provide some of the strongest support for ovarian hormone influences on eating behavior. Findings dating back more than 40 years show that estrogen has anorexic effects on food intake, with higher levels decreasing food intake and lower levels increasing food consumption⁸. These data come from strong experimental studies in which the ovaries are removed—via ovariectomy (OVX)—and then hormones are re-administered exogenously. Following OVX, rats, mice, and monkeys^{19–23} significantly increase chow (that is, nutritive food) intake and body weight, which normalize following exogenous estradiol treatment^{8,11,24}. Treatment with progesterone

alone does not directly affect food intake^{25–28}, but progesterone antagonizes the effects of estradiol and increases food intake in OVX rats^{11,24}. Fluctuations in food intake across the estrous cycle are also present in rodents, such that food consumption increases in non-ovulatory phases (for example, diestrous) and decreases during ovulation (that is, estrous)^{8,29–34}. Together, these data suggest that estrogen exerts direct, inhibitory effects on food intake while progesterone has stimulatory effects in the presence of estrogen.

Unfortunately, comparatively fewer animal studies have examined ovarian hormone effects on BE per se. Results thus far replicate those for chow intake^{9–11,24}. BE is typically assessed in these studies by exposing animals to intermittent (for example, 3 days per week) access to palatable food (PF) (that is, high-fat/-sugar food). Using these models, studies have shown that rats undergoing OVX in adulthood significantly increase binge-like eating^{9,11} that is reversed by estradiol treatment^{11,35} (and combination estradiol/progesterone treatment in some studies²⁴). Binge-like eating also increases in intact animals across the estrous cycle when estradiol levels are low (that is, diestrous) and decreases when estradiol levels are high (that is, estrous)¹⁰.

Human studies

Human research generally corroborates findings from animal studies. In women, it is difficult to directly manipulate hormone levels, but changes in BE across the menstrual cycle have been examined as a quasi-experimental design for understanding hormone effects. In these studies, changes in food intake and BE that are predicted by natural (and monthly) changes in hormones are used as an indication of longitudinal and predictive associations between ovarian hormones and BE. As in animal findings, food intake in women decreases with increasing estradiol levels and low progesterone levels (that is, follicular phase)^{36–38} and increases when both hormone levels are high (that is, mid-luteal phase)^{36–38}. Nearly identical patterns of effects have been observed for BE^{4–7}. When estradiol levels are high and progesterone levels are low (for example, follicular phase and ovulation), rates of BE and emotional eating (EE) (that is, a significant correlate and predictor of BE, defined as overeating in response to negative emotions) are low^{3–5,39}. Conversely, when both estradiol and progesterone levels are high (for example, mid-luteal phase), rates of BE/EE significantly increase^{3–5,39}. Natural fluctuations in ovarian hormones across the menstrual cycle seem to account for these effects, as higher levels of both estradiol and progesterone predict increased levels of BE and EE across the cycle^{4,39,40}. These associations between hormones and BE/EE are present in community^{5,39–41} and clinical^{4,6} samples of women, although effects appear stronger in clinical samples^{4,6}.

Overall, animal and human studies have significant predictive (and, in animal studies, causal) associations between ovarian hormones and changes in food intake and BE in females^{3–11}. Although the factors contributing to these associations remain unknown, it seems likely that changes in gene transcription are major contributors. Ovarian hormones are steroid hormones whose primary function is to regulate gene transcription/expression (and, thereby, protein synthesis) in the neurobiological systems

(for example, serotonin and dopamine) known to be disrupted in eating disorders^{42,43}. Thus, it is possible that during risky hormonal milieus (for example, non-estrous or mid-luteal phases) there is differential regulation of risky or protective genes for BE in key neurobiological systems⁴⁴. Data from twin studies indirectly support this possibility, as the heritability of eating disorder symptoms varies dramatically across the menstrual cycle⁴⁰ and other hormonal “events” (for example, puberty)^{45–47}, and suggest that ovarian hormone activation and de-activation are associated with changes in genetic risk. This is likely one set of moderators that contribute to hormone-BE associations, whereby associations between hormones and behavioral phenotypes vary by the presence/absence of risk/protective genes. Nonetheless, other moderators likely also contribute and may also interact with ovarian hormones to influence behavior. The psychological and physiological responses to stress are strong candidates in this regard.

Stress and binge eating

Animal studies

Stress is a common precipitant to BE^{14–17}. Animal models repeatedly demonstrate that rats, mice, and non-human primates subjected to stress exhibit hyperphagia of general food intake^{48–51} and binge-like eating^{48,52–54}. Results are consistent across various forms of stress tasks (for example, shock stress, restriction/refeeding, and chronic social defeat)⁵⁰. For example, following repeated bouts of foot shock, female BE-prone (BEP) rats consumed significantly larger amounts of sucrose than BE-resistant (BER) rats⁵³. BEP rats also consumed sucrose more rapidly and demonstrated compulsive-like intake of sucrose following stress compared with BER rats⁵³. Mice exhibit similar increases in food intake following chronic subordination stress (that is, repeatedly exposing a submissive animal to an aggressive, dominant animal) that becomes exacerbated in response to food restriction (a known precipitant to overeating)^{48,52,55} or acute social defeat⁴⁹. Restriction/refeeding stress (that is, access to PF diet followed by food deprivation and then refeeding of PF diet) also induces a hyperphagic rebound upon refeeding of PF⁵⁶.

Stress effects on eating behavior appear dependent upon BE phenotype. Prior to stress exposure and intermittent PF access, both BEP and BER animals exhibit increases in plasma corticosterone levels⁵³. However, following repeated exposure to stress, BEP animals begin to display a blunted corticosterone response to stress^{53,57}, indicative of a hypoactive HPA-axis response, compared with BER animals that display the typical post-stress increase in plasma corticosterone⁵⁸. Because differences in stress response between the phenotypes do not appear until stress and PF access induction, it is possible that there are inherent physiological and neurobiological differences between the two BE phenotypes that become exacerbated upon exposure to stress.

One potential neurobiological difference that may contribute to the development of the two phenotypes is the corticotropin-releasing hormone (CRH) system within hypo- and extra-thalamic brain regions. Chronic stress increases mRNA expression of CRH

and CRH receptors in brain regions associated with food intake (for example, paraventricular nucleus) and stress response (for example, central amygdala)^{57,58}. However, increases in CRH levels are significantly higher in BEP compared with BER rats following stress⁵⁷. CRH triggers release of adrenocorticotrophic hormone from the pituitary, which stimulates release of glucocorticoids (for example, corticosterone) from the adrenal glands¹⁸. Therefore, elevated CRH levels increase corticosterone levels in animals. Increased corticosterone levels are associated with increased anxiety-like behaviors^{59,60} and concordant increases in food intake^{58,61}. These results suggest increased susceptibility to stress in BEP versus BER animals that may alter underlying neural circuitry involved in stress and food intake regulation in order to perpetuate the BE phenotype.

Human studies

In humans, the majority of studies examining acute stress effects on BE in the laboratory use physical (for example, cold pressor test)⁶², psychosocial (for example, Trier Social Stress Test [TSST])⁶³, or cognitive (for example, Stroop test)⁶⁴ stress tasks. A smaller number of studies also examine daily experiences of stress by using self-report questionnaires (for example, Perceived Stress Scale)⁶⁵ or biological measures of stress responses (for example, salivary or urinary cortisol). Finally, chronic stress has been examined by studying samples with known or self-reported exposure to chronic stress (for example, trauma and abuse) measured via self-report questionnaires (for example, Childhood Trauma Questionnaire)⁶⁶ or longitudinal biomarkers of stress (for example, daily salivary cortisol assays).

Consistent with the effects found in animal studies, effects of acute stress on eating behavior exist in women^{67–69} and appear dependent upon BE status. Following acute psychosocial (that is, TSST) or physical (that is, cold pressor task) stress, women who binge eat rated foods as more palatable⁶⁷, displayed increased motivation toward food^{68,70,71}, and showed an increased desire to binge eat⁷¹ compared with non-stressed BED and control women. Food intake^{72–74} and BE^{75,76} also increase following exposure to acute stress in women who binge eat compared with control women. Elevated basal cortisol levels^{69,77–80} and a blunted acute stress response^{81,82}, indicative of chronic stress exposure and altered HPA-axis responses, are also observed in women who binge eat¹⁸. Interestingly, women who binge eat are more likely to have experienced chronic life stress (for example, abuse and trauma)^{77–80,83,84} and often exhibit a hypoactive HPA-axis response¹⁷ compared with non-BE women, indicating potentially important associations between chronic stress or altered stress responses (or both) and BE in women.

However, not all women binge eat in response to stress. Currently, it is unknown why stress increases the risk for BE in some, but not all, women. Studies in humans have yet to test the effects of CRH mRNA and receptor expression observed in animals (described above). However, studies suggest that differential neural activation in women who binge eat may partially explain this discrepancy. Following the presentation of food cues from pre- to post-stress induction, women with BN exhibited decreased activation in brain regions associated with

emotion regulation (for example, anterior cingulate cortex, amygdala, and ventromedial prefrontal cortex)^{85,86} and self-focus (for example, precuneus)⁶⁸ compared with healthy controls. Among the women who binge eat, those with greater de-activation in these brain regions reported increased stress levels prior to BE^{85,86}. This differential response to stress on eating behavior suggests that, as in animal studies, inherent differences in neural circuitry following stress may increase susceptibility to BE in a subset of women. These findings highlight the need for additional studies that elaborate upon the mechanisms through which stress might influence BE.

Stress and ovarian hormones

Animal studies

Ovarian hormones play a critical role in modulating the stress response in animals and humans. Specifically, ovarian hormones regulate HPA-axis functioning via antagonistic effects on glucocorticoids⁸⁷. At high levels of estradiol, as in ovarian hormone regulation of eating behavior, glucocorticoid receptor levels are decreased⁸⁸, HPA-axis responsivity is enhanced¹⁸, and animals exhibit fewer stress-like (for example, anxiety) behaviors^{88,89}. Conversely, when either progesterone alone or both ovarian hormones are elevated, glucocorticoid levels increase^{90–92}, HPA-axis activity decreases¹⁸, and animals exhibit more stress-like behaviors⁸⁸.

It should be noted that when estradiol is administered to OVX animals without progesterone, some studies have reported the expected decreases in glucocorticoid levels^{90,93,94} whereas others have reported increases in glucocorticoid levels^{91,95}. Differing results may be due to different responses based on estradiol dose, such that the administration of high physiological levels of estradiol produces protective effects against stress⁹⁴, whereas the administration of lower physiological levels may not produce protective effects^{91,96}. Additional research is needed to confirm these impressions and identify the potential mechanisms underlying these dose-response relationships. However, initial results^{18,87–89,97–99} are promising in supporting a protective role for estrogen against stress and an antagonistic role for progesterone⁹⁶ against estrogen, similar to the effects observed with eating behavior. Notably, during more severe and chronic levels of stress, ovarian hormone functioning can shut down because of antagonistic effects of elevated glucocorticoid (for example, cortisol) levels on ovarian hormones¹⁰⁰. In these cases, the estrous/menstrual cycle becomes irregular and/or cycling stops. Because the hypothalamic–pituitary–gonadal axis is completely shut down, these effects cannot account for the hormone-BE associations that are observed in regularly cycling animals and humans and that are the focus of this review. Nonetheless, these extreme stress–hormone interactions would be important to examine in populations of women with BE who are no longer menstruating and who have experienced high levels of chronic stress.

Interestingly, ovarian hormone and glucocorticoid (for example, cortisol) receptors colocalize in brain networks¹⁰¹ and binding sites of promotor regions of similar genes responsible for regulating stress and food intake^{87,102,103}. This colocalization creates competition for binding sites and leads to increased

susceptibility for stress–ovarian hormone interactions in females across ovarian hormone levels, such that at high estradiol and low progesterone levels, glucocorticoid levels are low compared with when both ovarian hormones are elevated¹⁰⁴.

Human studies

Findings from human studies corroborate data from animal studies suggesting associations between ovarian hormones and stress responses in women. In women, ovarian hormones modulate HPA-axis functioning via antagonistic effects on glucocorticoids¹⁸. For example, women with¹⁰⁵ and without^{104,106–108} BE perceive stimuli (for example, giving a speech) as less stressful when estradiol levels are high and progesterone levels are low, compared with when both hormone levels are elevated. Results from community samples of women also suggest that basal plasma cortisol levels and adrenal production of cortisol post-stress are higher post-ovulation versus pre-ovulation¹⁰⁴.

Ovarian hormones also regulate neural activity in key subcortical regions of stress circuitry (for example, amygdala, hippocampus, and hypothalamus)^{109–111} such that when both ovarian hormones are elevated (for example, during the mid-luteal phase), activation of stress circuitry increases compared with when estradiol is elevated and progesterone levels are low (for example, during the follicular phase)^{112,113}. Studies examining the effects of progesterone on stress are limited, but current theories suggest an antagonistic role for progesterone against the protective benefits of estrogen, particularly at peak progesterone levels^{92,114,115}. Taken together, these findings suggest a protective role for estrogen against stress and a possible antagonistic role for progesterone toward estrogen.

Stress as a moderator of ovarian hormone–binge eating associations

As previously stated, not all women binge eat in response to ovarian hormones or stress. Given the associations between ovarian hormones, stress, and BE (described above), it is possible that stress moderates ovarian hormone–BE associations, strengthening these associations in highly stressed women. In these women, risky hormonal milieus (that is, high levels of estradiol and progesterone) may enhance psychological (for example, perceived levels of stress) and biological (for example, elevated glucocorticoid levels) stress responses and lead to even higher levels of BE compared with women with lower stress levels.

Unfortunately, to date, no studies have examined whether stress moderates hormone-BE associations in humans. Studies have examined this association in animals and found significant moderating effects of stress on ovarian hormone–BE associations^{10,116–118}. Acute stress (that is, frustrative non-reward) significantly increased PF consumption (by 30% to 40%) in BEP rats studied during low estradiol phases (for example, diestrous) as compared with non-stressed rats and stressed rats studied during high estradiol phases (for example, estrous)¹⁰. Stress (for example, food-restriction stress) also significantly increases PF consumption in OVX rats compared with non-stressed OVX rats¹¹. Treatment with exogenous estrogen and progesterone attenuates these stress-induced increases in PF consumption in

OVX animals¹¹, although the magnitude of attenuation seems dependent on the intensity of the stressor^{11,116–118}. Together, these results suggest that ovarian hormones exert protective effects against increased BE at low stress (for example, restriction) conditions. However, under high stress/chronic stress conditions, the protective effects of estrogen may be decreased or even reversed.

Conclusions

The etiology of BE remains to be fully elucidated; however, previous studies have repeatedly established ovarian hormones^{4,6,8} and stress^{14–17} as critical contributing factors to BE in animals and women. Studies also demonstrate substantial regulation of the stress response by ovarian hormones with patterns mimicking those found for ovarian hormone influences on BE (that is, more significant stress effects at elevated versus low ovarian hormone levels)^{101–104,119}. Stress may also initiate different physiological and neural responses in animals^{53,57} and women^{68,85,86} who do and do not binge eat, which reinforce the BE phenotype. These differences may become exacerbated at varying ovarian hormone levels. Lastly, stress increases PF consumption in female rats with lower estradiol levels or who have undergone OVX. Overall, the current literature provides strong support for individual associations among stress, ovarian hormones, and BE in both women and animals, and emerging data from animal studies indirectly support stress as a moderator of ovarian hormone influence over general food intake.

Clearly, more evidence is needed to more firmly establish stress as a moderator of hormone-BE associations. For animal studies, a key starting place would be to examine stress and hormone effects on binge-like eating rather than general food intake. OVX animal models should also be used as strong experimental controls to examine binge-like eating and PF consumption before and after both stress exposure and exogenous hormone administration. Lastly, studies should examine effects of multiple stressor types (for example, foot shock and restraint stress) to determine the moderating effect of different stressor types on ovarian hormone-BE associations.

Human studies need similarly strong quasi-experimental designs. First, studies should examine daily stress, hormone, and BE levels to examine stress by hormone interactions on BE. The

menstrual cycle is one model in which to test these effects due to natural fluctuations in ovarian hormones. Oral contraceptive (OC) use can also be used to examine stress by hormone interactions on BE. Because OCs consist of active and inactive pill phases, human studies can mimic hormonal manipulation paradigms seen in animal literature (for example, OVX and hormone administration). In addition to self-reported stress, lab-based stressors (for example, TSST) should be used to assess subjective and objective responses to stress as they relate to hormone-BE associations.

Although this review focused on women of reproductive age, data suggest that other developmental periods (for example, puberty^{45–47} and menopause^{120–123}) may be important for BE risk. Puberty and menopause are critical developmental periods during which dramatic increases or decreases in ovarian hormone levels can activate or de-activate the protective effects of estrogen on stress responsivity^{124–128} and BE^{45–47,120–122}. Future studies are needed to better understand the role of stress and ovarian hormone interactions on BE risk during these risky developmental hormonal milieus.

Finally, relatively little is known about the neurobiological mechanisms underlying the independent or interactive effects of stress and ovarian hormones on BE. Initial studies suggest that increased expression of CRH in brain regions associated with food intake and stress response^{57,58} and decreased activation in limbic and frontal regions following stress^{68,85,86} may be important. However, more studies using neuroimaging (for example, functional magnetic resonance imaging) in humans and brain processing (for example, optogenetics) in animals are needed to explore these possibilities.

Grant information

This work was supported by grants awarded to KKL and CLS by the National Institute of Mental Health (NIMH) (MH 111715-01A1S1). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIMH.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Hoek HW: **Incidence, prevalence and mortality of anorexia nervosa and other eating disorders.** *Curr Opin Psychiatry*. 2006; 19(4): 389–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
- American Psychiatric Association: **Diagnostic and Statistical Manual of Mental Disorders.** 5th ed. Arlington, VA: American Psychiatric Publishing; 2013.
[Publisher Full Text](#)
- Culbert KM, Racine SE, Klump KL: **Hormonal Factors and Disturbances in Eating Disorders.** *Curr Psychiatry Rep*. 2016; 18(7): 65.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Edler C, Lipson SF, Keel PK: **Ovarian hormones and binge eating in bulimia nervosa.** *Psychol Med*. 2007; 37(1): 131–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Klump KL, Keel PK, Culbert KM, *et al.*: **Ovarian hormones and binge eating: exploring associations in community samples.** *Psychol Med*. 2008; 38(12): 1749–57.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Klump KL, Racine SE, Hildebrandt B, *et al.*: **Ovarian Hormone Influences on Dysregulated Eating: A Comparison of Associations in Women with versus without Binge Episodes.** *Clin Psychol Sci*. 2014; 2(4): 545–59.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Racine SE, Culbert KM, Keel PK, *et al.*: **Differential associations between ovarian hormones and disordered eating symptoms across the menstrual cycle in women.** *Int J Eat Disord*. 2012; 45(3): 333–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

8. Asarian L, Geary N: **Sex differences in the physiology of eating.** *Am J Physiol Regul Integr Comp Physiol.* 2013; **305**(11): R1215–67.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. Klump KL, Suisman JL, Culbert KM, *et al.*: **The effects of ovariectomy on binge eating proneness in adult female rats.** *Horm Behav.* 2011; **59**(4): 585–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. **F** Micioni Di Bonaventura MV, Lutz TA, Romano A, *et al.*: **Estrogenic suppression of binge-like eating elicited by cyclic food restriction and frustrative-nonreward stress in female rats.** *Int J Eat Disord.* 2017; **50**(6): 624–35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
11. Yu Z, Geary N, Corwin RL: **Ovarian hormones inhibit fat intake under binge-type conditions in ovariectomized rats.** *Physiol Behav.* 2008; **95**(3): 501–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. The National Eating Disorders Coalition: **Eating disorders prevention, treatment, & management: An evidence review.** 2010.
[Reference Source](#)
13. **F** Marzilli E, Cerniglia L, Cimino S: **A narrative review of binge eating disorder in adolescence: prevalence, impact, and psychological treatment strategies.** *Adolesc Health Med Ther.* 2018; **9**: 17–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
14. Levine MD, Marcus MD: **Eating behavior following stress in women with and without bulimic symptoms.** *Ann Behav Med.* 1997; **19**(2): 132–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Pike KM, Wilfley D, Hilbert A, *et al.*: **Antecedent life events of binge-eating disorder.** *Psychiatry Res.* 2006; **142**(1): 19–29.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Copeland WE, Bulik CM, Zucker N, *et al.*: **Does childhood bullying predict eating disorder symptoms? A prospective, longitudinal analysis.** *Int J Eat Disord.* 2015; **48**(8): 1141–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Lo Sauro C, Raval di C, Cabras PL, *et al.*: **Stress, hypothalamic-pituitary-adrenal axis and eating disorders.** *Neuropsychobiology.* 2008; **57**(3): 95–115.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. **F** Oyola MG, Handa RJ: **Hypothalamic-pituitary-adrenal and hypothalamic-pituitary-gonadal axes: sex differences in regulation of stress responsivity.** *Stress.* 2017; **20**(5): 476–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
19. Blaustein JD, Gentry RT, Roy EJ, *et al.*: **Effects of ovariectomy and estradiol on body weight and food intake in gold thioglucose-treated mice.** *Physiol Behav.* 1976; **17**(6): 1027–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. Drewett RF: **Oestrous and dioestrous components of the ovarian inhibition on hunger in the rat.** *Anim Behav.* 1973; **21**(4): 772–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Gentry RT, Wade GN: **Sex differences in sensitivity of food intake, body weight, and running-wheel activity to ovarian steroids in rats.** *J Comp Physiol Psychol.* 1976; **90**(8): 747–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Sullivan E, Daniels A, Cameron J: **Ovariectomy leads to rapid changes in food intake, body weight, and metabolic regulation in female rhesus monkeys.** *Soc Neurosci Abstr.* 2003; **803**: 19.
23. Witte MM, Resuehr D, Chandler AR, *et al.*: **Female mice and rats exhibit species-specific metabolic and behavioral responses to ovariectomy.** *Gen Comp Endocrinol.* 2010; **166**(3): 520–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Yu Z, Geary N, Corwin RL: **Individual effects of estradiol and progesterone on food intake and body weight in ovariectomized binge rats.** *Physiol Behav.* 2011; **104**(5): 687–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Wade GN: **Some effects of ovarian hormones on food intake and body weight in female rats.** *J Comp Physiol Psychol.* 1975; **88**(1): 183–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Wade GN: **Gonadal hormones and behavioral regulation of body weight.** *Physiol Behav.* 1972; **8**(3): 523–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Geary N, Trace D, McEwen B, *et al.*: **Cyclic estradiol replacement increases the satiety effect of CCK-8 in ovariectomized rats.** *Physiol Behav.* 1994; **56**(2): 281–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
28. Grueso E, Rocha M, Puerta M: **Plasma and cerebrospinal fluid leptin levels are maintained despite enhanced food intake in progesterone-treated rats.** *Eur J Endocrinol.* 2001; **144**(6): 659–65.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Butera PC: **Estradiol and the control of food intake.** *Physiol Behav.* 2010; **99**(2): 175–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Geary N: **The Estrogenic Inhibition of Eating.** *Neurobiology of Food and Fluid Intake.* 2nd ed. (E.M. Stricker & S.C. Woods, ed.). New York, NY: Kluwer; 2004; **14**: 307–345.
[Publisher Full Text](#)
31. Geary N: **Endocrine controls of eating: CCK, leptin, and ghrelin.** *Physiol Behav.* 2004; **81**(5): 719–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Rosenblatt H, Dyrenfurth I, Ferin M, *et al.*: **Food intake and the menstrual cycle in rhesus monkeys.** *Physiol Behav.* 1980; **24**(3): 447–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
33. Kemnitz JW, Houser WD, Eisele SG, *et al.*: **Pregnancy and Fetal Development in the Rhesus Monkey.** (P.W. Nathanielsz, J.T. Parer, eds.). Ithaca, New York: Perinatology Press; 1984.
34. Kemnitz JW, Gibber JR, Eisele SG, *et al.*: **Relationship of Reproductive Condition to Food Intake and Sucrose Consumption of Female Rhesus Monkeys.** (D.M. Taub & F.A. King, ed.). New York: Van Nostrand Reinhold; 1986.
[Reference Source](#)
35. Cao X, Xu P, Oyola MG, *et al.*: **Estrogens stimulate serotonin neurons to inhibit binge-like eating in mice.** *J Clin Invest.* 2014; **124**(10): 4351–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Fong AK, Kretsch MJ: **Changes in dietary intake, urinary nitrogen, and urinary volume across the menstrual cycle.** *Am J Clin Nutr.* 1993; **57**(1): 43–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Gong EJ, Garrel D, Calloway DH: **Menstrual cycle and voluntary food intake.** *Am J Clin Nutr.* 1989; **49**(2): 252–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Lyons PM, Truswell AS, Mira M, *et al.*: **Reduction of food intake in the ovulatory phase of the menstrual cycle.** *Am J Clin Nutr.* 1989; **49**(6): 1164–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Klump KL, Keel PK, Racine SE, *et al.*: **The interactive effects of estrogen and progesterone on changes in emotional eating across the menstrual cycle.** *J Abnorm Psychol.* 2013; **122**(1): 131–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Klump KL, Hildebrandt BA, O'Connor SM, *et al.*: **Changes in genetic risk for emotional eating across the menstrual cycle: a longitudinal study.** *Psychol Med.* 2015; **45**(15): 3227–37.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Klump KL, O'Connor SM, Hildebrandt BA, *et al.*: **Differential Effects of Estrogen and Progesterone on Genetic and Environmental Risk for Emotional Eating in Women.** *Clin Psychol Sci.* 2016; **4**(5): 895–908.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Wilson JD, Foster DW, Kronenberg HM, *et al.*: **Williams Textbook of Endocrinology.** 9th ed. Philadelphia, PA: W.B. Saunders Company; 1998.
[Reference Source](#)
43. Ostlund H, Keller E, Hurd YL: **Estrogen receptor gene expression in relation to neuropsychiatric disorders.** *Ann N Y Acad Sci.* 2003; **1007**(1): 54–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Klump KL, Culbert KM, Sisk CL: **Sex Differences in Binge Eating: Gonadal Hormone Effects Across Development.** *Annu Rev Clin Psychol.* 2017; **13**: 183–207.
[PubMed Abstract](#) | [Publisher Full Text](#)
45. **F** Klump KL, Burt SA, McGue M, *et al.*: **Changes in genetic and environmental influences on disordered eating across adolescence: a longitudinal twin study.** *Arch Gen Psychiatry.* 2007; **64**(12): 1409–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
46. Klump KL: **Puberty as a critical risk period for eating disorders: a review of human and animal studies.** *Horm Behav.* 2013; **64**(2): 399–410.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
47. Klump KL, Culbert KM, O'Connor S, *et al.*: **The significant effects of puberty on the genetic diathesis of binge eating in girls.** *Int J Eat Disord.* 2017; **50**(8): 984–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. Corwin RL, Avena NM, Boggiano MM: **Feeding and reward: perspectives from three rat models of binge eating.** *Physiol Behav.* 2011; **104**(1): 87–97.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
49. Razzoli M, Sanghez V, Bartolomucci A: **Chronic subordination stress induces hyperphagia and disrupts eating behavior in mice modeling binge-eating-like disorder.** *Front Nutr.* 2015; **1**(30): pii: 00030.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. **F** Razzoli M, Pearson C, Crow S, *et al.*: **Stress, overeating, and obesity: Insights from human studies and preclinical models.** *Neurosci Biobehav Rev.* 2017; **76**(Pt A): 154–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
51. Michopoulos V, Toufexis D, Wilson ME: **Social stress interacts with diet history to promote emotional feeding in females.** *Psychoneuroendocrinology.* 2012; **37**(9): 1479–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
52. Boggiano MM, Chandler PC: **Binge eating in rats produced by combining dieting with stress.** *Curr Protoc Neurosci.* 2006; **Chapter 9**(Unit9): 23A.
[PubMed Abstract](#) | [Publisher Full Text](#)
53. **F** Calvez J, Timofeeva E: **Behavioral and hormonal responses to stress in binge-like eating prone female rats.** *Physiol Behav.* 2016; **157**: 28–38.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
54. Hagan MM, Chandler PC, Wauford PK, *et al.*: **The role of palatable food and hunger as trigger factors in an animal model of stress induced binge eating.** *Int J Eat Disord.* 2003; **34**(2): 183–97.
[PubMed Abstract](#) | [Publisher Full Text](#)

55. Hagan MM, Wauford PK, Chandler PC, *et al.*: **A new animal model of binge eating: key synergistic role of past caloric restriction and stress.** *Physiol Behav.* 2002; **77**(1): 45–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
56. Hagan MM, Moss DE: **Persistence of binge-eating patterns after a history of restriction with intermittent bouts of refeeding on palatable food in rats: implications for bulimia nervosa.** *Int J Eat Disord.* 1997; **22**(4): 411–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
57. **F** Calvez J, de Ávila C, Guèvremont G, *et al.*: **Stress differentially regulates brain expression of corticotropin-releasing factor in binge-like eating prone and resistant female rats.** *Appetite.* 2016; **107**: 585–95.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
58. **F** Alcántara-Alonso V, Amaya MI, Matamoros-Trejo G, *et al.*: **Altered functionality of the corticotropin-releasing hormone receptor-2 in the hypothalamic paraventricular nucleus of hyperphagic maternally separated rats.** *Neuropeptides.* 2017; **63**: 75–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
59. Machado TD, Dalle Molle R, Laureano DP, *et al.*: **Early life stress is associated with anxiety, increased stress responsivity and preference for “comfort foods” in adult female rats.** *Stress.* 2013; **16**(5): 549–56.
[PubMed Abstract](#) | [Publisher Full Text](#)
60. Ortolani D, Garcia MC, Melo-Thomas L, *et al.*: **Stress-induced endocrine response and anxiety: the effects of comfort food in rats.** *Stress.* 2014; **17**(2): 211–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
61. **F** Sefton C, Harno E, Davies A, *et al.*: **Elevated Hypothalamic Glucocorticoid Levels Are Associated With Obesity and Hyperphagia in Male Mice.** *Endocrinology.* 2016; **157**(11): 4257–65.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
62. Lovallo W: **The cold pressor test and autonomic function: a review and integration.** *Psychophysiology.* 1975; **12**(3): 268–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
63. Kirschbaum C, Pirke KM, Hellhammer DH: **The ‘Trier Social Stress Test’—a tool for investigating psychobiological stress responses in a laboratory setting.** *Neuropsychobiology.* 1993; **28**(1–2): 76–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
64. Stroop JR: **Studies of interference in serial verbal reactions.** *J Exp Psychol.* 1935; **18**(6): 643–62.
[Publisher Full Text](#)
65. Cohen S, Kamarck T, Mermelstein R: **A global measure of perceived stress.** *J Health Soc Behav.* 1983; **24**(4): 385–96.
[PubMed Abstract](#) | [Publisher Full Text](#)
66. Bernstein D, Fink L: **Childhood Trauma Questionnaire: A Retrospective Self-Report Manual.** San Antonio, TX: The Psychological Corporation; 1998.
[Reference Source](#)
67. **F** Lyu Z, Jackson T: **Acute Stressors Reduce Neural Inhibition to Food Cues and Increase Eating Among Binge Eating Disorder Symptomatic Women.** *Front Behav Neurosci.* 2016; **10**: 188.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
68. **F** Collins B, Breithaupt L, McDowell JE, *et al.*: **The impact of acute stress on the neural processing of food cues in bulimia nervosa: Replication in two samples.** *J Abnorm Psychol.* 2017; **126**(5): 540–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
69. Rosenberg N, Bloch M, Ben Avi I, *et al.*: **Cortisol response and desire to binge following psychological stress: comparison between obese subjects with and without binge eating disorder.** *Psychiatry Res.* 2013; **208**(2): 156–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
70. Gluck ME, Geliebter A, Lorence M: **Cortisol stress response is positively correlated with central obesity in obese women with binge eating disorder (BED) before and after cognitive-behavioral treatment.** *Ann NY Acad Sci.* 2004; **1032**: 202–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
71. Schulz S, Laessle RG: **Stress-induced laboratory eating behavior in obese women with binge eating disorder.** *Appetite.* 2012; **58**(2): 457–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
72. **F** Klatzkin RR, Baldassaro A, Hayden E: **The impact of chronic stress on the predictors of acute stress-induced eating in women.** *Appetite.* 2018; **123**: 343–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
73. Mason AE, Lustig RH, Brown RR, *et al.*: **Acute responses to opioidergic blockade as a biomarker of hedonic eating among obese women enrolled in a mindfulness-based weight loss intervention trial.** *Appetite.* 2015; **91**: 311–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
74. Tryon MS, Carter CS, Decant R, *et al.*: **Chronic stress exposure may affect the brain's response to high calorie food cues and predispose to obesogenic eating habits.** *Physiol Behav.* 2013; **120**: 233–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
75. **F** Klatzkin RR, Gaffney S, Cyrus K, *et al.*: **Stress-induced eating in women with binge-eating disorder and obesity.** *Biol Psychol.* 2018; **131**: 96–106.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
76. **F** Naish KR, Laliberte M, MacKillop J, *et al.*: **Systematic review of the effects of acute stress in binge eating disorder.** *Eur J Neurosci.* 2018.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
77. Becker DF, Grilo CM: **Childhood maltreatment in women with binge-eating disorder: Associations with psychiatric comorbidity, psychological functioning, and eating pathology.** *Eat Weight Disord.* 2011; **16**(2): e113–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
78. Fuemmeler BF, Dedert E, McClemon FJ, *et al.*: **Adverse childhood events are associated with obesity and disordered eating: Results from a U.S. population-based survey of young adults.** *J Trauma Stress.* 2009; **22**(4): 329–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
79. St-Hilaire A, Steiger H, Liu A, *et al.*: **A prospective study of effects of prenatal maternal stress on later eating-disorder manifestations in affected offspring: Preliminary indications based on the Project Ice Storm cohort.** *Int J Eat Disord.* 2015; **48**(5): 512–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
80. Su X, Xu B, Liang H, *et al.*: **Prenatal maternal bereavement and risk of eating disorders in infants and toddlers: A population-based cohort study.** *BMC Psychiatry.* 2015; **15**: 229.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
81. Gluck ME, Geliebter A, Hung J, *et al.*: **Cortisol, hunger, and desire to binge eat following a cold stress test in obese women with binge eating disorder.** *Psychosom Med.* 2004; **66**(6): 876–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
82. Het S, Vocks S, Wolf JM, *et al.*: **Blunted neuroendocrine stress reactivity in young women with eating disorders.** *J Psychosom Res.* 2015; **78**(3): 260–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
83. Pankevich DE, Bale TL: **Stress and sex influences on food-seeking behaviors.** *Obesity (Silver Spring).* 2008; **16**(7): 1539–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
84. Yau YH, Potenza MN: **Stress and eating behaviors.** *Minerva Endocrinol.* 2013; **38**(3): 255–67.
[PubMed Abstract](#) | [Free Full Text](#)
85. **F** Wonderlich JA, Breithaupt L, Thompson JC, *et al.*: **The impact of neural responses to food cues following stress on trajectories of negative and positive affect and binge eating in daily life.** *J Psychiatr Res.* 2018; **102**: 14–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
86. **F** Fischer S, Breithaupt L, Wonderlich J, *et al.*: **Impact of the neural correlates of stress and cue reactivity on stress related binge eating in the natural environment.** *J Psychiatr Res.* 2017; **92**: 15–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
87. Whirledge S, Cidlowski JA: **Estradiol antagonism of glucocorticoid-induced GILZ expression in human uterine epithelial cells and murine uterus.** *Endocrinology.* 2013; **154**(1): 499–510.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
88. **F** Ramos-Ortola DL, Doreste-Mendez RJ, Alvarado-Torres JK, *et al.*: **Ovarian hormones modify anxiety behavior and glucocorticoid receptors after chronic social isolation stress.** *Behav Brain Res.* 2017; **328**: 115–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
89. **F** Wiersliels KR, Wicks B, Simko H, *et al.*: **Sex differences in corticotropin releasing factor-evoked behavior and activated networks.** *Psychoneuroendocrinology.* 2016; **73**: 204–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
90. Carey MP, Detered CH, de Koning J, *et al.*: **The influence of ovarian steroids on hypothalamic-pituitary-adrenal regulation in the female rat.** *J Endocrinol.* 1995; **144**(2): 311–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
91. Viau V, Meaney MJ: **Variations in the hypothalamic-pituitary-adrenal response to stress during the estrous cycle in the rat.** *Endocrinology.* 1991; **129**(5): 2503–11.
[PubMed Abstract](#) | [Publisher Full Text](#)
92. **F** Freitas-de-Melo A, Damián JP, Hötzel MJ, *et al.*: **Progesterone pretreatment increases the stress response to social isolation in ewes.** *Hormones (Athens).* 2016; **15**(1): 81–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
93. Hollenstein K, Janett F, Bleul U, *et al.*: **Influence of estradiol on adrenal activity in ovariectomized cows during acute stress.** *Anim Reprod Sci.* 2006; **93**(3–4): 292–302.
[PubMed Abstract](#) | [Publisher Full Text](#)
94. Komesaroff PA, Esler M, Clarke IJ, *et al.*: **Effects of estrogen and estrous cycle on glucocorticoid and catecholamine responses to stress in sheep.** *Am J Physiol.* 1998; **275**(4 Pt 1): E671–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
95. Burgess LH, Handa RJ: **Chronic estrogen-induced alterations in adrenocorticotropin and corticosterone secretion, and glucocorticoid receptor-mediated functions in female rats.** *Endocrinology.* 1992; **131**(3): 1261–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
96. Wilson ME, Legendre A, Pazol K, *et al.*: **Gonadal steroid modulation of the limbic-hypothalamic-pituitary-adrenal (LHPA) axis is influenced by social status in female rhesus monkeys.** *Endocrine.* 2005; **26**(2): 89–97.
[PubMed Abstract](#) | [Publisher Full Text](#)
97. Dakin RS, Walker BR, Seckl JR, *et al.*: **Estrogens protect male mice from obesity**

- complications and influence glucocorticoid metabolism. *Int J Obes (Lond)*. 2015; **39**(10): 1539–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
98. Weiser MJ, Foradori CD, Handa RJ: **Estrogen receptor beta activation prevents glucocorticoid receptor-dependent effects of the central nucleus of the amygdala on behavior and neuroendocrine function.** *Brain Res*. 2010; **1336**: 78–88.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 99. Young EA, Altemus M, Parkison V, *et al.*: **Effects of estrogen antagonists and agonists on the ACTH response to restraint stress in female rats.** *Neuropsychopharmacology*. 2001; **25**(6): 881–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
 100. **F** Michopoulos V: **Stress-induced alterations in estradiol sensitivity increase risk for obesity in women.** *Physiol Behav*. 2016; **166**: 56–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 101. Miller WJS, Suzuki S, Miller LK, *et al.*: **Estrogen receptor (ER)beta isoforms rather than ERalpha regulate corticotropin-releasing hormone promoter activity through an alternate pathway.** *J Neurosci*. 2004; **24**(47): 10628–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
 102. Kato J, Hirata S, Nozawa A, *et al.*: **Gene expression of progesterone receptor isoforms in the rat brain.** *Horm Behav*. 1994; **28**(4): 454–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
 103. Shughrue PJ, Lane MV, Merchenthaler I: **Comparative distribution of estrogen receptor-alpha and -beta mRNA in the rat central nervous system.** *J Comp Neurol*. 1997; **388**(4): 507–25.
[PubMed Abstract](#) | [Publisher Full Text](#)
 104. **F** Albert K, Pruessner J, Newhouse P: **Estradiol levels modulate brain activity and negative responses to psychosocial stress across the menstrual cycle.** *Psychoneuroendocrinology*. 2015; **59**: 14–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 105. Friederich HC, Schild S, Schellberg D, *et al.*: **Cardiac parasympathetic regulation in obese women with binge eating disorder.** *Int J Obes (Lond)*. 2006; **30**(3): 534–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
 106. Young EA, Altemus M: **Puberty, ovarian steroids, and stress.** *Ann N Y Acad Sci*. 2004; **1021**: 124–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
 107. Andreano JM, Cahill L: **Menstrual cycle modulation of medial temporal activity evoked by negative emotion.** *Neuroimage*. 2010; **53**(4): 1286–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 108. Ferree NK, Kamat R, Cahill L: **Influences of menstrual cycle position and sex hormone levels on spontaneous intrusive recollections following emotional stimuli.** *Conscious Cogn*. 2011; **20**(4): 1154–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 109. Jacobs EG, Holsen LM, Lancaster K, *et al.*: **17 β -estradiol differentially regulates stress circuitry activity in healthy and depressed women.** *Neuropsychopharmacology*. 2015; **40**(3): 566–76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 110. Goldstein JM, Jerram M, Poldrack R, *et al.*: **Hormonal cycle modulates arousal circuitry in women using functional magnetic resonance imaging.** *J Neurosci*. 2005; **25**(40): 9309–16.
[PubMed Abstract](#) | [Publisher Full Text](#)
 111. Ossewaarde L, van Wingen GA, Rijpkema M, *et al.*: **Menstrual cycle-related changes in amygdala morphology are associated with changes in stress sensitivity.** *Hum Brain Mapp*. 2013; **34**(5): 1187–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
 112. Ossewaarde L, Hermans EJ, van Wingen GA, *et al.*: **Neural mechanisms underlying changes in stress-sensitivity across the menstrual cycle.** *Psychoneuroendocrinology*. 2010; **35**(1): 47–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
 113. **F** Chung KC, Peisen F, Kogler L, *et al.*: **The Influence of Menstrual Cycle and Androstadienone on Female Stress Reactions: An fMRI Study.** *Front Hum Neurosci*. 2016; **10**: 44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 114. Roca CA, Schmidt PJ, Altemus M, *et al.*: **Differential menstrual cycle regulation of hypothalamic-pituitary-adrenal axis in women with premenstrual syndrome and controls.** *J Clin Endocrinol Metab*. 2003; **88**(7): 3057–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
 115. **F** Herrera AY, Nielsen SE, Mather M: **Stress-induced increases in progesterone and cortisol in naturally cycling women.** *Neurobiol Stress*. 2016; **3**: 96–104.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 116. da Silva CC, Lazzaretti C, Fontanive T, *et al.*: **Estrogen-dependent effects on behavior, lipid-profile, and glycemic index of ovariectomized rats subjected to chronic restraint stress.** *Behav Processes*. 2014; **103**: 327–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
 117. Gamaro GD, Prediger ME, Lopes JB, *et al.*: **Interaction between estradiol replacement and chronic stress on feeding behavior and on serum leptin.** *Pharmacol Biochem Behav*. 2003; **76**(2): 327–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
 118. Johnson ZP, Lowe J, Michopoulos V, *et al.*: **Oestradiol differentially influences feeding behaviour depending on diet composition in female rhesus monkeys.** *J Neuroendocrinol*. 2013; **25**(8): 729–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 119. van Lier E, Carriquiry M, Meikle A: **Sex steroid modulation of cortisol secretion in sheep.** *Animal*. 2014; **8**(6): 960–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 120. **F** Baker JH, Runfola CD: **Eating disorders in midlife women: A perimenopausal eating disorder?** *Maturitas*. 2016; **85**: 112–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 121. **F** Baker JH, Peterson CM, Thornton LM, *et al.*: **Reproductive and Appetite Hormones and Bulimic Symptoms during Midlife.** *Eur Eat Disord Rev*. 2017; **25**(3): 188–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 122. **F** Mangweth-Matzek B, Hoek HW, Rupp CI, *et al.*: **The menopausal transition—a possible window of vulnerability for eating pathology.** *Int J Eat Disord*. 2013; **46**(6): 609–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 123. Mangweth-Matzek B, Hoek HW, Pope HG: **Pathological eating and body dissatisfaction in middle-aged and older women.** *Curr Opin Psychiatry*. 2014; **27**(6): 431–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
 124. Evuarherhe O, Leggett J, Waite E, *et al.*: **Reversal of the hypothalamo-pituitary-adrenal response to oestrogens around puberty.** *J Endocrinol*. 2009; **202**(2): 279–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
 125. Holder MK, Blaustein JD: **Puberty and adolescence as a time of vulnerability to stressors that alter neurobehavioral processes.** *Front Neuroendocrinol*. 2014; **35**(1): 89–110.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 126. Patacchioli FR, Simeoni S, Monnazzi P, *et al.*: **Menopause, mild psychological stress and salivary cortisol: Influence of long-term hormone replacement therapy (HRT).** *Maturitas*. 2006; **55**(2): 150–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
 127. **F** Villada C, Espin L, Hidalgo V, *et al.*: **The influence of coping strategies and behavior on the physiological response to social stress in women: The role of age and menstrual cycle phase.** *Physiol Behav*. 2017; **170**: 37–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 128. Kajantie E, Phillips DI: **The effects of sex and hormonal status on the physiological response to acute psychosocial stress.** *Psychoneuroendocrinology*. 2006; **31**(2): 151–78.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Yong Xu

Children's Nutrition Research Center, Department of Pediatrics, Baylor College of Medicine, Houston, TX, 77030, USA

Competing Interests: No competing interests were disclosed.

2. Jessica Baker

UNC Center of Excellence for Eating Disorders, Department of Psychiatry, University of North Carolina, Chapel Hill, NC, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research