RESEARCH ARTICLE

Burden of health behaviours and socioeconomic position on health care expenditure in Ontario [version 1; peer review: 1 approved, 1 approved with reservations]

Douglas G. Manuel ¹⁻⁶, Carol Bennett^{1,2}, Richard Perez^{1,2}, Andrew S. Wilton², Adrian Rohit Dass⁷, Audrey Laporte^{2,7}, David A. Henry^{2,7,8}

¹Ottawa Hospital Research Institute, Ottawa, Canada

²ICES, Toronto and Ottawa, Canada

³Statistics Canada, Ottawa, Canada

⁴Department of Family Medicine, University of Ottawa, Ottawa, Canada

⁵School of Epidemiology and Public Health, University of Ottawa, Ottawa, Canada

⁶Bruyère Research Institute, Ottawa, Canada

⁷Institute of Health Policy, Management and Evaluation, University of Toronto, Toronto, Canada

⁸Centre for Research in Evidence-Based Practice, Bond University, Gold Coast, Australia

V1 First published: 18 Mar 2019, 8:303
<https://doi.org/10.12688/f1000research.18205.1>

Latest published: 16 Oct 2019, 8:303
<https://doi.org/10.12688/f1000research.18205.2>

Abstract

Background: Smoking, unhealthy alcohol consumption, poor diet and physical inactivity are leading risk factors for morbidity and mortality, and contribute substantially to overall healthcare costs. The availability of health surveys linked to health care provides population-based estimates of direct healthcare costs. We estimated health behaviour and socioeconomic-attribute healthcare costs, and how these have changed during a period when government policies have aimed to reduce their burden.

Methods: The Ontario samples of the Canadian Community Health Surveys (conducted in 2003, 2005, and 2007-2008) were linked at the individual level to all records of health care use of publicly funded healthcare. Generalized linear models were estimated with a negative binomial distribution to ascertain the relationship of health behaviours and socioeconomic risk factors on health care costs. The multivariable cost model was then applied to unlinked, cross-sectional CCHS samples for each year from 2004 to 2013 to examine the evolution of health behaviour and socioeconomic-attributable direct health care expenditures over a 10-year period.

Results: We included 80,749 respondents, aged 25 years and older, and 312,952 person-years of follow-up. The cost model was applied to 200,324 respondents aged 25 years and older (CCHS 2004 to 2013). During the 10-year period from 2004 to 2013, smoking, unhealthy alcohol consumption, poor diet and physical inactivity attributed to 22% of Ontario's direct health care costs. Ontarians in the most

Open Peer Review

Approval Status

	1	2
version 2 (revision) 16 Oct 2019		 view
version 1 18 Mar 2019	 view	 view

1. **Suneela Mehta** , University of Auckland, Auckland, New Zealand

2. **Melody Ding** , The University of Sydney, Sydney, Australia

Any reports and responses or comments on the article can be found at the end of the article.

disadvantaged socioeconomic position contributed to 15% of the province's direct health care costs. Taken together, health behaviours and socioeconomic position were associated with 34% (\$134 billion) of direct health care costs (2004 to 2013). Over this time period, we estimated a 1.9% reduction in health care expenditure (\$5.0 billion) attributable to improvements in some health behaviours, most importantly reduced rates of smoking.

Conclusions: Health behaviours and socioeconomic position cause a large direct health care system cost burden.

Keywords

Burden, smoking, alcohol, diet, physical activity, healthcare cost

Corresponding author: Douglas G. Manuel (dmanuel@ohri.ca)

Author roles: **Manuel DG:** Conceptualization, Formal Analysis, Methodology, Project Administration, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Bennett C:** Conceptualization, Formal Analysis, Methodology, Project Administration, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Perez R:** Conceptualization, Formal Analysis, Methodology, Writing – Review & Editing; **Wilton AS:** Conceptualization, Formal Analysis, Investigation, Methodology, Writing – Review & Editing; **Rohit Dass A:** Formal Analysis, Investigation, Methodology, Writing – Review & Editing; **Laporte A:** Conceptualization, Investigation, Methodology, Supervision, Writing – Review & Editing; **Henry DA:** Conceptualization, Investigation, Methodology, Project Administration, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This study was supported by ICES, which is funded by an annual grant from the Ontario Ministry of Health and Long-Term Care (MOHLTC). The opinions, results and conclusions reported in this paper are those of the authors and are independent from the funding sources. No endorsement by ICES or the Ontario MOHLTC is intended or should be inferred.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 Manuel DG *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Manuel DG, Bennett C, Perez R *et al.* **Burden of health behaviours and socioeconomic position on health care expenditure in Ontario [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2019, 8:303 <https://doi.org/10.12688/f1000research.18205.1>

First published: 18 Mar 2019, 8:303 <https://doi.org/10.12688/f1000research.18205.1>

Introduction

Smoking, unhealthy alcohol consumption, poor diet and physical inactivity are leading risk factors for morbidity and mortality worldwide¹. Despite this knowledge, prevalence of these risk factors remains high and reduction efforts may be hindered by failure to understand the full human and cost burdens these risk factors impose on societies². In an era of increasing health care expenditure most political focus has been on payments for services and the growing impacts of diagnostic and therapeutic technologies. There is also a need to consider costs resulting from upstream health behaviours, and how these have changed over time, to help prioritize public health strategies and support public health decision makers. These relationships are likely to be complex because of some conflicting trends in the prevalence of health behaviours^{3,4}.

While health behaviours have a leading role in morbidity and mortality, it is also recognized that there is uneven distribution of health across socioeconomic position (i.e., social and economic factors that influence what position individuals hold within the structure of a society⁵). Canadian research indicates that individuals with lower socioeconomic position tend to be less healthy than those who enjoy greater educational, income and occupational advantages^{6–9}. As with health behaviour-attributable health care use, evidence of the economic cost of these health disparities helps us understand the health and financial benefits of reducing the gap⁶.

We sought to estimate the economic burden attributable to four health behaviour risk factors (smoking, unhealthy alcohol consumption, poor diet and physical inactivity), how these have changed over time, and how they interact with socioeconomic position. The study had three objectives: 1) to examine the direct healthcare costs associated with smoking, unhealthy alcohol consumption, poor diet and physical inactivity; 2) to examine the change in direct health care costs as a consequence of changes, over time, in health behaviours; and, 3) to examine the direct health care costs associated with socioeconomic position (i.e., education, family income, home ownership, and neighbourhood deprivation).

Past studies typically infer health care costs indirectly—where aggregate health care expenditure data is categorized by disease, for example. We used unique Canadian data that individually link respondents from large repeated population health surveys to comprehensive health care utilisation and cost data covering hospital and primary care sectors in Ontario. These data provide, to our knowledge, the largest and most complete population-based examination of the relationship between health behaviours and direct public healthcare costs. We believe this is the first study to measure directly how changes in health behaviours result in changes in health care use. The linked data also provide the means to assess the degree to which health costs are associated with socioeconomic inequalities.

Methods

Ethical approval

This study was approved by the Ottawa Health Science Network Research Ethics Board. Datasets were linked using unique encoded identifiers and analysed at ICES. ICES is an independent, non-profit research institute whose legal status under Ontario's health information privacy law allows it to collect and analyse health care and demographic data, without consent, for health system evaluation and improvement.

Study cohorts

We used the Ontario sample drawn from a national population health survey—the Canadian Community Health Survey (CCHS), conducted in 2003, 2005, 2007–2008, 2009–2010, 2011–2012, and 2013–2014 to develop linked and unlinked study cohorts. The CCHS is a cross-sectional survey conducted by Statistics Canada that collects data related to health determinants, health status and health care use—details of data collected and used in the analyses are provided below. The survey employs a complex multistage sampling strategy to randomly select households in each health region. Details of the survey methodology have previously been published¹⁰. A weight, which reflects the number of individuals represented in the target population, is assigned to each respondent; the target population includes individuals aged 12 years and older living in Canada's ten provinces and three territories. Individuals living on First Nation Reserves, institutionalized residents, full-time members of the Canadian Forces and residents of certain remote areas are excluded from the survey's sampling frame.

For the linked study cohort, the Ontario sample of the 2003, 2005, and 2007–2008 CCHS cycles provided 128,501 valid interviews. Of these respondents, a subset agreed to share and link their interview information and 101,506 were successfully linked to their provincial health card number using a deterministic and probabilistic algorithm. We included respondents aged 25 years and older if they were eligible for provincially funded health care and not pregnant at the time of survey administration. For individuals with multiple interviews, only the earliest interview was included. We excluded individuals who were lost to follow-up in the first year following their interview (i.e., they were not available at the beginning of the study). This resulted in a final cohort of 80,749 unique Ontario respondents (see Figure 1).

The Ontario sample of all six CCHS waves provided 200,324 valid respondents aged 25 years and older for the unlinked study cohort. This cohort was used to estimate direct health care costs by applying the multivariable model that was derived using the linked cohort CCHS data (see Model Development).

Behavioural and other risk factors for health care use

The CCHS waves were used to examine the following risk factors for their association with health care use: age, sex, four health behaviours (smoking, alcohol consumption, diet and physical

Figure 1. Creation of study cohort for multivariable cost model. CCHS, Canadian Community Health Survey; OHIP, Ontario Health Insurance Program.

activity; see Table 1), sociodemographic factors (immigrant status, education level, urban dwelling, neighbourhood deprivation, household income, home ownership, marital status), self-perceived stress, preventive health behaviour (flu vaccination), and health status indicators (body mass index, hypertension, diabetes, heart disease, cancer, history of stroke, dementia, and extent of difficulty in performing basic tasks or participating in activities).

Public health care spending data

We examined person-level health care costs across three sectors: 1) hospital care (inpatient hospitalizations, same day surgeries, emergency department visits, rehabilitation hospitals, and complex continuing care centres); 2) drugs (for Ontarians age 65 and older and Ontarians receiving social assistance, Ontario Drug Benefit costs were captured); and 3) community care (primary care billings, specialist billings, lab billings, capitation services, and home care services).

Costs to operate the health care system (e.g., health ministry administrative costs) and capital costs for large scale projects

(e.g., building new hospitals) are not reflected in the person-level costs. To account for these exclusions in the health care cost analysis, we obtained annual total health care expenditures from publicly available Ontario Ministry of Finance records (fiscal years 2003 to 2013, where fiscal year is April 1 to March 31), the Canadian Institute for Health Information's *National Health Expenditure Trends* publication, and the MOHLTC's *Report Card for the Ontario Drug Benefit Program* publications^{11–23}. The expenditures from each year were categorized into our three health care sectors and expenditures that did not correspond with any of the three sectors were assigned to an 'other' category.

All costs are expressed in 2014 Canadian dollars with past costs inflated using the annual general Consumer Price Index reported by Statistics Canada.

Model development

To develop multivariable cost models, Ontario respondents to the 2003, 2005 and 2007–2008 cycles of CCHS were linked, at the individual level, to all records of health care use that were paid for by the Ontario Ministry of Health and Long-Term Care

Table 1. Definitions of behavioural health risks.

Behaviour	Category*	Definition
Smoking	Heavy smoker	Current daily smoker (≥ 20 cigarettes/day)
	Light smoker	Current daily smoker (< 20 cigarettes/day) or current occasional smoker with ≥ 100 lifetime cigarettes
	Former heavy smoker	Former daily smoker (≥ 20 cigarettes/day)
	Former light smoker	Former daily smoker (< 20 cigarettes/day) or former occasional smoker with ≥ 100 lifetime cigarettes
	<i>Non-smoker</i>	<i>Never smoker or occasional smoker < 100 lifetime cigarettes</i>
Alcohol	Heavy drinker	Bingeing† or ≥ 21 (men) or ≥ 14 (women) drinks/week
	<i>Moderate drinker</i>	<i>≤ 21 (men) or ≤ 14 (women) drinks/week with no bingeing</i>
	Non-drinker	No alcohol consumption in the last 12 months
Diet	Poor diet	Index score† 0 to < 2.5
	Fair diet	Index score 2.5 to < 5
	<i>Adequate diet</i>	<i>Index score 5 to 10</i>
Physical activity	Inactive	0 to < 1.5 METs/day
	Moderately active	1.5 to < 3 METs/day
	<i>Active</i>	<i>≥ 3 METs/day</i>

*Highest risk levels are in bold and lowest risk levels (reference group) are in italics.

†Bingeing: five or more drinks on any day in the previous week or weekly bingeing behaviour in the previous month.

†Index score: the healthiness of a diet based on consumption of fruit and vegetables. Individuals start with 2 points and achieve up to 8 additional points for each average daily serving of fruits and vegetables (maximum score = 10). Points are deducted for daily fruit juice servings exceeding 1 (-2 points), no carrot consumption (-2 points), or daily potato consumption exceeding 1 serving for males and 0.7 servings for females (-2 points). Scores that result in negative values after deductions are recoded to zero, resulting in a final range of 0 to 10 for the index.

MET, metabolic equivalent of task (a measure of calories burned by type, duration and frequency of physical activity).

(MOHLTC). The cost associated with each record was estimated using costing methods developed for Ontario health administrative data²⁴. Briefly, a payer (the MOHLTC) cost perspective was taken, using person-level health care utilization and per-use fee information or budgetary data. Cost information for sectors (i.e., acute hospitalization, same day surgery, emergency department, inpatient rehabilitation, and complex continuing care) that are funded using global budgets (e.g., by institution) are determined using a top-down approach through case-mix methodology. Sectors that have fee payments associated with each use (e.g., prescription drugs, physician services, and home care) have costs estimated directly.

Beginning one year after survey administration, CCHS respondents were followed for a four-year period (between 2004 and 2013) to develop multivariable models estimating the effect of health behaviours on health care costs. We used a generalized linear model with a negative binomial distribution and an offset to account for variation in follow-up times to create separate, sex-specific models for each of the health care sectors: hospital care, drugs, and community care. To assess confounding and mediation, we use a pre-specified, stepwise modelling approach. We started with a *health behaviour model* followed by a basic *sociodemographic model*, a *primary attribution model* that adjusted for additional sociodemographic risk factors, a

distal mediator model that included health status indicators, and a *proximal mediator model* that included a measure of fragility (see Figure 2). This stepwise analysis resulted in 15 models for each sex.

The model building approach sought to address three issues in assessing the contribution of health behaviours to health care costs. First, we were interested in having appropriate adjustment for other risk factors for increased costs that are correlated with health behaviours (e.g., age and sociodemographic risk factors). Second, we were attentive to risk factors that may mediate the relationship between health behaviours and health care costs (e.g., body mass index or hypertension). Our concern was that their inclusion in the model could inappropriately attenuate the risk from health behaviours. Third, we considered pre-existing illness that may have led to health behaviour change. For example, as people become ill and frail they may become less physically active; in such a situation, physical inactivity may be associated with increased health care cost that is more appropriately identified as illness-associated inactivity.

Estimating population attributable fractions of health care costs

We calculated the proportion of the health care costs that can be attributed to health behaviours—the population attributable

Figure 2. Stepped approach to model building—consideration of risk factors leading to healthcare use and costs.

fraction—for fiscal years 2003 to 2013, for each health care sector. Using the unlinked CCHS cohort, we estimated annual population attributable fractions using each CCHS cycle and for years between CCHS cycles, by averaging the population attributable fractions from the preceding and succeeding year.

We used a factor-deleted approach to calculate population attributable fractions that involved three steps. In the first step, we estimated expected annual population health care costs for a specific sector by applying the corresponding sector-specific primary attribution models to the weighted CCHS cycle. In the second step, we repeated the calculation after recoding each respondent's health behaviour to the counterfactual reference or "no exposure" category. For example, taking the weighted cohort, we estimated hospital costs using all smoking exposures (i.e., current, former, and non-smokers) and re-estimated hospital costs assuming all current and former smokers were non-smokers. The difference between the two calculations was an estimate of the annual contribution of smoking to hospital costs. In the final step, we divided this difference by the original population estimate (from the first step) to produce a population attributable fraction. In our example, this would be the population attributable fraction of hospital costs associated with smoking. Health sector specific population attributable fractions were calculated for each health behaviour, and the combination of health behaviours. The same analysis was performed for different socioeconomic groups defined by education, family income, home ownership, and neighbourhood deprivation.

Estimating the health care cost burden of health behaviours and socioeconomic position in Ontario

We calculated annual estimates of costs attributable to health behaviours and socioeconomic position (fiscal years 2004 to 2013) by applying the sector-specific population attributable fractions to the annual public health care expenditures and summing the health care sector results together.

Estimating costs attributable to changes in health behaviours

The change in health care costs attributable to the change in health behaviours was estimated for fiscal years 2004 to 2013. A baseline population attributable fraction for total health care costs associated with all health behaviours was estimated for fiscal year 2003 using the previously described methods for population attributable fractions and attributable costs. The overall health care budget was estimated over the subsequent decade, assuming that health behaviours in 2003 remained constant (e.g., the baseline population attributable fraction did not change over time). The difference between the counterfactual health care budget and the actual health care budget provided an annual estimate of the change in health care costs attributable to changes in health behaviours.

Sensitivity analysis

We performed three sets of sensitivity analyses. First, we created five models representing a stepwise approach to assess confounding and mediation. The five models assessed health care costs and burden estimates with progressive adjustment for risk factors: age, sex and health behaviours (Model 1), basic sociodemographic characteristics (Model 2), additional sociodemographic characteristics (Model 3), health status mediators (Model 4), and fragility measures (Model 5) (Figure 2). The estimate derived from Model 3 was assumed to be our most accurate and appropriate estimate of the attributable burden due to health behaviours. The estimates derived from Model 1 (simply age and health behaviours) and Model 5 (the over-adjusted model) were used as upper and lower bounds of uncertainty.

For our second sensitivity analysis, we compared age-standardized cost ratios after excluding the top 5% of health care users to assess the possibility of overly influential respondents. The use of health care varies considerably between people, particularly for hospital care and other speciality services. Only a small proportion of people are hospitalized and, of those

hospitalized, a small proportion has multiple admissions and complicated long hospital stays. The skewed distribution of health care services has potential to distort the attributable health care expenditure analysis because a small proportion of CCHS respondents may have a strong influence on overall or total population estimates.

Third, we replicated analyses using an inverse propensity-weighted model to assess robustness of the health care cost ratios attributable to smoking. The inverse propensity-weighted model, a complimentary approach to the generalized linear model, is an alternative approach to adjust cost ratios for multi-variable risk factors. Our inverse propensity-weighted analyses

included several covariates in addition to all those used in the multivariable analyses (i.e., the primary attribution model): household type, highest level of household education, main source of household income, labour force participation, sense of belonging to the community, regional health authority, and survey cycle.

Results

The population attributable fractions for the four behavioural risks were calculated using responses from 80,749 Ontarians surveyed between 2003 and 2008. In total, there were 312,952 person-years of follow-up. Characteristics of the study cohort are presented in [Table 2](#).

Table 2. Baseline description of the study cohorts.

	Male cohort			Female cohort		
	Survey sample* %	Person-years	Represented population† %	Survey sample* %	Person-years	Represented population† %
	(N=36,807)		(N=3,962,088)	(N=43,942)		(N=4,131,570)
Age group (years)						
25 to 29	7.4	10,668	9.1	7.3	12,642	8.8
30 to 34	9.0	13,001	10.0	8.5	14,714	9.1
35 to 39	10.3	14,957	11.5	9.1	15,888	11.5
40 to 44	11.1	16,183	14.5	8.8	15,326	12.5
45 to 49	9.0	13,059	11.5	8.1	14,149	11.6
50 to 54	9.6	13,847	10.2	9.4	16,256	10.2
55 to 59	10.0	14,420	9.6	10.3	17,760	9.1
60 to 64	9.0	12,897	7.2	9.0	15,451	7.2
65 to 69	7.9	11,152	5.6	8.0	13,568	6.2
70 to 74	6.8	9,408	4.6	7.5	12,650	5.0
75 to 79	5.3	7,004	3.4	6.5	10,654	4.2
80 to 84	3.1	3,907	2.0	4.8	7,575	2.8
85 to 89	1.2	1,356	0.7	2.1	3,257	1.3
90+	0.3	320	0.2	0.7	883	0.4
Health Behaviours						
Smoking status						
Heavy smoker	10.7	15,116	9.2	5.8	9,828	4.8
Light smoker	14.5	20,642	15.4	14.8	25,234	13.8
Former heavy smoker	20.0	27,998	16.0	9.5	16,010	7.7
Former light smoker	17.4	24,531	16.8	18.0	30,678	16.3
Non-smoker	36.5	52,432	41.8	51.0	87,361	56.6
Missing	1.1	1,460	0.9	1.0	1,661	0.8
Alcohol consumption						
Heavy drinker	12.3	17,695	10.9	3.4	5,762	3.3
Moderate drinker	70.7	100,787	71.8	72.2	124,162	70.0
Non-drinker	15.1	20,940	15.3	23.3	38,974	25.6
Missing	2.0	2,758	2.0	1.1	1,876	1.1
Physical activity						
Inactive	46.5	65,794	47.8	52.8	89,455	54.3
Moderately active	25.2	36,032	24.3	25.6	44,103	24.5

	Male cohort			Female cohort		
	Survey sample* %	Person-years	Represented population† %	Survey sample* %	Person-years	Represented population† %
	(N=36,807)		(N=3,962,088)	(N=43,942)		(N=4,131,570)
Active	26.1	37,473	25.1	20.8	35,913	19.7
Missing	2.2	2,880	2.8	0.9	1,302	1.5
Diet						
Poor diet	15.0	21,400	14.2	8.6	14,632	8.3
Fair diet	41.7	59,340	40.3	30.0	50,933	29.2
Adequate diet	38.6	55,167	40.7	58.2	99,981	59.0
Missing	4.7	6,272	4.8	3.3	5,227	3.6
Sociodemographic Indicators						
Immigrant status						
Immigrant	21.8	30,837	33.5	21.5	36,779	33.9
Non-immigrant	78.1	111,177	66.2	78.3	133,765	65.8
Missing	0.1	165	0.3	0.1	228	0.4
Ethnicity						
White	88.9	126,274	78.9	89.7	152,990	79.3
Non-white	10.7	15,357	20.5	10.0	17,183	20.1
Missing	0.4	548	0.6	0.4	600	0.5
Education						
Less than high school	18.9	25,984	14.8	20.2	33,702	16.5
High school graduate	22.7	32,530	22.2	25.0	42,730	24.9
Post-secondary graduate	57.5	82,404	61.8	54.1	93,212	57.7
Missing	0.9	1,262	1.2	0.7	1,129	0.9
Marital status						
Married/common-law	67.8	96,810	76.8	57.7	99,508	68.8
Other	32.2	45,306	23.2	42.3	71,197	31.2
Missing	0.0	64	0.0	0.0	68	0.0
Residence ownership						
Yes	79.3	113,229	79.9	75.9	130,283	77.8
No	20.5	28,730	19.7	23.9	40,246	21.9
Missing	0.2	221	0.3	0.1	244	0.3
Household income (\$)						
0 to 29,999	16.6	22,903	11.0	25.9	43,299	16.4
30,000 to 79,999	45.5	64,687	40.1	41.6	71,561	39.3
80,000+	32.2	46,643	40.3	23.5	40,774	31.0
Missing	5.7	7,946	8.6	9.0	15,140	13.3
Preventive healthcare						
Flu shot						
Yes	61.6	87,077	58.0	68.6	116,714	63.4
No	36.0	51,907	38.9	30.5	52,701	35.0
Missing	2.5	3,195	3.1	0.9	1,358	1.5
Geography						
Urban						
No	22.3	31,709	14.9	21.0	36,089	14.3
Yes	77.7	110,471	85.1	79.0	134,684	85.7
Neighbourhood deprivation						
High	15.1	21,394	11.8	16.3	27,519	12.6
Moderate	62.3	88,446	60.8	62.3	106,413	61.2

	Male cohort			Female cohort		
	Survey sample* %	Person-years	Represented population† %	Survey sample* %	Person-years	Represented population† %
	(N=36,807)		(N=3,962,088)	(N=43,942)		(N=4,131,570)
Low	20.6	29,532	25.3	19.3	33,198	24.2
Missing	2.0	2,807	2.1	2.2	3,643	2.0
General health indicators						
Self-perceived stress						
Quite a bit or extremely stressful	20.2	28,881	23.0	21.5	36,892	24.4
At most, a bit stressful	79.5	112,917	76.7	78.2	133,339	75.4
Missing	0.3	382	0.3	0.3	542	0.3
Body mass index						
Underweight	0.7	926	0.8	2.7	4,428	3.2
Normal	43.5	62,221	43.3	29.9	51,187	28.3
Overweight	15.6	22,224	13.9	12.5	21,474	10.9
Obese	4.8	6,884	4.0	6.3	10,740	5.3
Morbidly Obese	34.2	48,428	36.5	45.5	77,566	48.5
Missing	1.2	1,497	1.5	3.2	5,378	3.9
Indicators of illness						
Hypertension						
Yes	22.7	31,629	18.9	25.7	43,146	20.5
No	77.0	110,133	80.8	74.2	127,408	79.4
Missing	0.3	417	0.3	0.1	218	0.1
Diabetes						
Yes	8.7	11,757	7.2	7.5	12,319	6.2
No	91.3	130,305	92.7	92.5	158,364	93.8
Missing	0.1	118	0.1	0.1	90	0.0
Heart disease						
Yes	9.4	12,559	6.8	7.9	12,819	5.5
No	90.4	129,348	93.0	91.9	157,654	94.3
Missing	0.2	272	0.1	0.2	299	0.2
Cancer						
Yes	2.9	3,754	2.1	2.6	4,199	2.0
No	97.0	138,284	97.9	97.3	166,374	97.9
Missing	0.1	141	0.1	0.1	200	0.1
Stroke						
Yes	2.0	2,552	1.4	1.8	2,796	1.4
No	98.0	139,540	98.6	98.1	167,826	98.6
Missing	0.1	88	0.1	0.1	151	0.0
Dementia						
Yes	0.6	645	0.5	0.4	629	0.5
No	99.4	141,440	99.5	99.5	170,034	99.4
Missing	0.1	94	0.1	0.1	109	0.1
Fragility						
Help with basic tasks	6.6	8,570	5.4	13.6	22,096	11.4
Limitation due to health	21.7	30,749	18.8	20.3	34,852	17.8
No limitations	71.4	102,527	75.5	65.8	113,449	70.6
Missing	0.3	334	0.3	0.2	376	0.2

*Data source: Canadian Community Health Survey (CCHS) 2.1, 3.1 and 4.1 (2003, 2005 and 2007/08).

†Population estimated using the CCHS sampling weights.

Health behaviour attributable healthcare use

From fiscal years 2004 to 2013, 22% of Ontario's health care costs could be attributed to the four health behaviour risk factors (Figure 3). Physical activity had the largest attribution (13%), followed by smoking (10%). However, uncertainty for the burden estimates indicates potential overestimation for physical activity and underestimation for diet. Alcohol-attributable health care costs were also likely underestimated (see limitations section).

Population health impact of behavioural risks and socioeconomic position

During the 10-year period (2004 to 2013), \$89.3 billion in health care costs were attributable to health behaviours. In that same period, the costs attributable to health behaviours improved by 1.9% (23.3% of total healthcare costs in 2004 to 21.4% in 2013). If the proportion of health care expenditure that can be attributed to health behaviours (the population attributable fraction) had remained at 23.3%, use of health care would have been \$5.0 billion greater (what we term 'avoided cost').

Table 3 presents the burden of health behaviours related to health care costs (\$89.3 billion) and the costs avoided by the adoption of healthy behaviours (\$5.0 billion). Physical inactivity and smoking contributed the largest proportion of the burden (53% and 41%, respectively). However, a decline in smoking between 2004 and 2013 was responsible for 84% of the avoided costs.

The scenarios from our sensitivity analysis demonstrate results that were similar to our main analysis. Not unexpectedly, the attribution of health care costs to health behaviours decreased as we increased the number of risk factors adjusted for in the model. Excluding high-cost health care users demonstrated slightly attenuated age-standardised cost ratios for men and women for almost all health behaviour risks. The inverse propensity-weighted model, which adjusted for additional variables, had similar cost ratios to the main analysis from the multivariable model.

Between 2004 and 2013, \$60.7 billion dollars in health care costs (15% of all health care costs for Ontarians aged 25 years

Figure 3. Proportion of health care costs attributed to selected health behaviour risk factors for Ontarians aged 25 and older, 2004 to 2013. Error bars represent high and low boundaries on burden estimates.

Table 3. Health care attributable and avoided costs by health behaviour risk factor for Ontarians aged 25 and older, 2004 to 2013.

Risk	Attributable costs (\$89.3 billion)	Avoided costs (\$5.0 billion)
Smoking	41%	84%
Alcohol	1%	0%
Diet	5%	5%
Physical activity	53%	11%

and older) were attributable to low socioeconomic position (Figure 4). When health behaviours and socioeconomic position are considered jointly, the health care cost burden was \$134 billion (34% of all health care costs for those aged 25 years and older). The break down by health care sector was similar for health behaviour and socioeconomic position. The largest portion of the burden is due to hospital care costs (46% of health behaviour attribution and 54% of socioeconomic position attribution to health care costs); followed by community care costs (22% and 19%), 'other' health care costs (21% and 19%), and drug costs (10% and 9%).

Discussion

We estimated that smoking, unhealthy alcohol consumption, poor diet and physical inactivity attributed to 22% of Ontario's direct health care costs during the ten-year period from 2004 to 2013. During this same period, improving health behaviours equated to a nearly 2% reduction in direct health care expenditure. Ontarians in the most disadvantaged socioeconomic group contributed to 15% of the province's direct health care costs. Taken together, health behaviours and socioeconomic position contributed to a burden of \$134 billion in direct health care costs (Ontario, 2004 to 2013).

Estimates of the cost of modifiable health behaviours and socioeconomic risk factors provides evidence to allow policy-makers to prioritize interventions aimed at reducing health care costs. Our estimate of a 1.9% reduction in health care expenditure (i.e., \$5 billion) through improved health behaviours, suggests that investments in promotion of healthy living have potential for substantial savings in health care costs in Canada. In our study, reduced smoking was the main contributor to avoided health behaviour attributable health care costs (accounting for 84% of the 1.9% cost reduction). This large cost reduction reflects

prominent smoking prevention strategies that were introduced during the study period—including 100% smoke-free public places including restaurants²⁵. From 2004 to 2013 there was an 11% reduction in the attributable fraction of smoking, which was solely a consequence of a reduction in the prevalence of current smoking. The large remaining burden from health behaviours and social inequalities suggests that there are significant opportunities to further reduce health care costs through population health strategies.

It is difficult to compare our findings with previous studies for two main reasons. First, various studies have estimated the economic burden in terms of costs for treatment and management of chronic diseases related to smoking, alcohol, diet or physical activity, but very few have evaluated the simultaneous impact of multiple risk factors in a population. These latter studies have used traditional population aggregated-data attributable fraction methods to estimate economic burden^{26–31}. Second, the significant methodological differences between our study and previous literature limits direct comparison of findings. Briefly, traditional population attributable fraction methods identify diseases where health behaviours are risk factors, estimate the health care costs of these diseases, calculate the proportion of the disease that can be attributed to the risk factors (based on relative risk of disease from an external source and prevalence of exposure in the population of interest), and apply these population attributable fractions to the cost data. There are limitations associated with these methods that stem from combining ecological summary measures of exposure, outcome, and hazards across different sources of data³². Our use of multivariable algorithms and the direct attribution of health behaviours to health care costs offers advantages over analyses that have been performed to date: controlling for confounders, accounting for complexities in the relationship between multiple exposures and covariates, using

Figure 4. Health behaviour risk factors and socioeconomic position attribution to health care costs for Ontarians aged 25 and older, 2004 to 2013.

consistent definitions of exposure, and using specific measures of risk derived internally from the study population.

Our study has several limitations that we believe underestimate the actual burden of health care costs attributable to the four health behaviours. First, we excluded individuals younger than 25 years of age. In general, health care costs for this age group are small; however, alcohol burden for younger people is a notable omission. Alcohol has an important attribution for injury, suicide and other social burdens that occur disproportionately among young people^{33–35}. Second, our reliance on self-reported health risk exposures will generally result in an underestimation of risk burden, especially for food and alcohol^{36–39}. In what is referred to as “social desirability bias”, survey respondents tend to over-report what they perceive as healthy behaviour and underreport unhealthy behaviour⁴⁰. As an example, self-reported alcohol consumption in surveys accounts about half the volume of alcohol sold^{41–42}. While reporting accuracy affects all risks in this study, burden estimates are mostly affected when people report they are in the healthiest category and they are actually in an unhealthy category. Third, the survey’s brief questions about risks may not capture the full spectrum of behaviour. For example, our measure of physical activity (leisure-time physical activity) did not include active transportation (such as walking and bicycling to work), work activity, or sedentary time and our measure of diet (fruit and vegetable consumption) did not specifically ascertain intake of sodium, trans fats, calories or other aspects of healthy and unhealthy eating. Fourth, we used respondents’ answers to health behaviours and other risks that correspond to their health behaviour at the time of the survey. In general, studies that consider lifetime changes in risks generate higher burden estimates. Our physical activity burden estimates may be an exception, where reverse-causality (i.e., ill health is the cause of reduced physical activity) results in overestimation. Fifth, “other” health care costs—including health care system operating costs and capital costs—are not included our health care cost data and were estimated indirectly. Sixth, we did not include indirect costs (such as lost productivity, wages and income related to illness associated with unhealthy living, or costs borne by individuals to care for their illness), nor did we include costs for health care beyond those paid by the provincial government (e.g., employee health plans).

Despite these limitations, our study demonstrates the importance of integrating public health and prevention within the health care system. A greater investment in disease prevention and population health could help increase the sustainability of publicly funded health care by reducing spending on illness, particularly with respect to hospital care. Combining this effort with strategies that address social determinants of health could secure further benefits. Indeed, our study suggests that interventions outside the health care system, such as improving levels of income and education, and other risk factors that influence socioeconomic position, will reduce health care costs. The health system is one determinant of population health and attention must be paid to the needs of disadvantaged individuals, populations, and communities in order to avoid increasing

health disparities⁴³. While health inequities play a significant role in health system costs, estimating the cost of the gap in health care as a result of socioeconomic position can be difficult because of the complexity of the problem. Our study is one of the few that has been able to translate this gap into a dollar figure.

Conclusions

Our study shows that health behaviours and socioeconomic position contribute to a large health care cost burden. Better health and well-being is the primary goal of improving health behaviours and reducing social inequity. However, it is important to recognize that existing investments in public health also results in a large reduction in expenditure in the acute care sector, particularly hospital care. The health premium of prevention and social equity is an overlooked opportunity for a sustainable health care system.

Data availability

Underlying data

The dataset from this study is held securely in coded form at ICES. While data sharing agreements prohibit ICES from making the dataset publicly available, access may be granted to those who meet pre-specified criteria for confidential access, available at www.ices.on.ca/DAS. The full dataset creation plan and underlying analytic code are available from the authors upon request, understanding that the computer programs may rely upon coding templates or macros that are unique to ICES and are therefore either inaccessible or may require modification.

Extended data

Open Science Framework: Burden of health behaviours and socioeconomic position on health care expenditure in Ontario. <https://doi.org/10.17605/OSF.IO/KW4C5>⁴⁴.

This project contains the sensitivity analysis supplementary file.

Extended data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

Grant information

This study was supported by ICES, which is funded by an annual grant from the Ontario Ministry of Health and Long-Term Care (MOHLTC). The opinions, results and conclusions reported in this paper are those of the authors and are independent from the funding sources. No endorsement by ICES or the Ontario MOHLTC is intended or should be inferred.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We are grateful for the contributions of Sima Gandhi and Erika Yates as well as our Scientific and Policy Advisory Committees—namely, Peter Austin, Bernard Choi, Prabhat Jha, Isra Levy, Heather Manson, Claudia Sanmartin, Joanne Thanos, and Pegeen Walsh.

References

- Lim SS, Vos T, Flaxman AD, *et al.*: **A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010.** *Lancet*. 2012; **380**(9859): 2224–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ding D, Lawson KD, Kolbe-Alexander TL, *et al.*: **The economic burden of physical inactivity: a global analysis of major non-communicable diseases.** *Lancet*. 2016; **388**(10051): 1311–24.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Simmons G, Jackson R, Swinburn B, *et al.*: **The increasing prevalence of obesity in New Zealand: is it related to recent trends in smoking and physical activity?** *N Z Med J*. 1996; **109**(1018): 90–92.
[PubMed Abstract](#)
- Lin AM, Lin MP, Markovic D, *et al.*: **Less Than Ideal: Trends in Cardiovascular Health Among US Stroke Survivors.** *Stroke*. 2019; **50**(1): 5–12.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Krieger N, Williams DR, Moss NE: **Measuring social class in US public health research: concepts, methodologies, and guidelines.** *Annu Rev Public Health*. 1997; **18**: 341–78.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Social Determinants and Science Integration Directorate, Public Health Agency of Canada: **Report summary - The Direct Economic Burden of Socioeconomic Health Inequalities in Canada: An Analysis of Health Care Costs by Income Level.** *Health Promot Chronic Dis Prev Can*. 2016; **36**(6): 118–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Canadian Population Health Initiative: **Reducing gaps in health: A focus on socio-economic status in urban Canada.** Ottawa, Ontario: Canadian Institute for Health Information (CIHI). Institut canadien d'information sur la santé. 2008.
[Reference Source](#)
- McIntosh CN, Fines P, Wilkins R, *et al.*: **Income disparities in health-adjusted life expectancy for Canadian adults, 1991 to 2001.** *Health Rep*. 2009; **20**(4): 55–64.
[PubMed Abstract](#)
- Chief Public Health Officer of Canada: **The Chief Public Health Officer's report on the state of public health in Canada: addressing health inequalities.** Ottawa (ON); 2008.
[Reference Source](#)
- Beland Y: **Canadian community health survey--methodological overview.** *Health Rep*. 2002; **13**(2): 9–14.
[PubMed Abstract](#)
- Ontario Ministry of Finance, Public accounts of Ontario: **Annual Report and Consolidated Financial Statements, 2003-2004.** Toronto, ON: Ministry of Finance; 2004; Accessed January 7, 2019.
[Reference Source](#)
- Ontario Ministry of Finance, Public accounts of Ontario: **Annual Report and Consolidated Financial Statements, 2004-2005.** Toronto, ON: Ministry of Finance; 2005; Accessed January 7, 2019.
[Reference Source](#)
- Ontario Ministry of Finance, Public accounts of Ontario: **Annual Report and Consolidated Financial Statements, 2005-2006.** Toronto, ON: Ministry of Finance; 2006; Accessed January 7, 2019.
[Reference Source](#)
- Ontario Ministry of Finance, Public accounts of Ontario: **Annual Report and Consolidated Financial Statements, 2006-2007.** Toronto, ON: Ministry of Finance; 2007; Accessed January 7, 2019.
[Reference Source](#)
- Ontario Ministry of Finance, Public accounts of Ontario: **Annual Report and Consolidated Financial Statements, 2007-2008.** Toronto, ON: Ministry of Finance; 2008; Accessed January 7, 2019.
[Reference Source](#)
- Ontario Ministry of Finance, Public accounts of Ontario: **Annual Report and Consolidated Financial Statements, 2008-2009.** Toronto, ON: Ministry of Finance; 2009; Accessed January 7, 2019.
[Reference Source](#)
- Ontario Ministry of Finance, Public accounts of Ontario: **Annual Report and Consolidated Financial Statements, 2009-2010.** Toronto, ON: Ministry of Finance; 2010; Accessed January 7, 2019.
[Reference Source](#)
- Ontario Ministry of Finance, Public accounts of Ontario: **Annual Report and Consolidated Financial Statements, 2010-2011.** Toronto, ON: Ministry of Finance; 2011; Accessed January 7, 2019.
[Reference Source](#)
- Ontario Ministry of Finance, Public accounts of Ontario: **Annual Report and Consolidated Financial Statements, 2011-2012.** Toronto, ON: Ministry of Finance; 2012; Accessed January 7, 2019.
[Reference Source](#)
- Ontario Ministry of Finance, Public accounts of Ontario: **Annual Report and Consolidated Financial Statements, 2012-2013.** Toronto, ON: Ministry of Finance; 2013; Accessed January 7, 2019.
[Reference Source](#)
- Ontario Ministry of Finance, Public accounts of Ontario: **Annual Report and Consolidated Financial Statements, 2013-2014.** Toronto, ON: Ministry of Finance; 2014; Accessed January 7, 2019.
[Reference Source](#)
- Canadian Institute for Health Information: **National Health Expenditure Trends, 1975 to 2015.** Ottawa, ON; 2015; Accessed February 16, 2016.
[Reference Source](#)
- Ontario Ministry of Health and Long-Term Care, **Annual ODB Report Cards.** Accessed January 7, 2018.
[Reference Source](#)
- Wodchis WP, Bushmeneva K, Nikitovic M, *et al.*: **Guidelines on Person-Level Costing Using Administrative Databases in Ontario.** Toronto; 2013; **1**.
[Reference Source](#)
- Ontario Ministry of Health and Long Term Care: **Smoke-Free Ontario: The Next Chapter for a Healthier Ontario.** Queen's Printer for Ontario; 2018; Accessed January 7, 2019.
[Reference Source](#)
- Scarborough P, Bhatnagar P, Wickramasinghe KK, *et al.*: **The economic burden of ill health due to diet, physical inactivity, smoking, alcohol and obesity in the UK: an update to 2006-07 NHS costs.** *J Public Health (Oxf)*. 2011; **33**(4): 527–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Krueger H, Krueger J, Koot J: **Variation across Canada in the economic burden attributable to excess weight, tobacco smoking and physical inactivity.** *Can J Public Health*. 2015; **106**(4): e171–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cadilhac DA, Magnus A, Sheppard L, *et al.*: **The societal benefits of reducing six behavioural risk factors: an economic modelling study from Australia.** *BMC Public Health*. 2011; **11**: 483.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Krueger H, Koot J, Andres E: **The economic benefits of fruit and vegetable consumption in Canada.** *Can J Public Health*. 2017; **108**(2): e152–e61.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Manuel DG, Perez R, Sanmartin C, *et al.*: **Measuring Burden of Unhealthy Behaviours Using a Multivariable Predictive Approach: Life Expectancy Lost in Canada Attributable to Smoking, Alcohol, Physical Inactivity, and Diet.** *PLoS Med*. 2016; **13**(8): e1002082.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Levin ML, Bertell R: **RE: "simple estimation of population attributable risk from case-control studies.** *Am J Epidemiol*. 1978; **108**(1): 78–79.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tanuseputro P, Perez R, Rosella L, *et al.*: **Improving the estimation of the burden of risk factors: an illustrative comparison of methods to measure smoking-attributable mortality.** *Popul Health Metr*. 2015; **13**(1): 5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- MacKinnon N, Colman I: **Factors Associated with Suicidal Thought and Help-Seeking Behaviour in Transition-Aged Youth versus Adults.** *Can J Psychiatry*. 2016; **61**(12): 789–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Public Health Agency of Canada: **The Chief Public Health Officer's Report on the State of Public Health in Canada, 2015: Alcohol Consumption in Canada.** Ottawa, Canada; 2016.
[Reference Source](#)
- World Health Organization: **Alcohol and injury in emergency departments: summary of the report from the WHO Collaborative Study on Alcohol and Injuries.** 2007.
[Reference Source](#)
- Whitford JL, Widner SC, Mellick D, *et al.*: **Self-report of drinking compared to objective markers of alcohol consumption.** *Am J Drug Alcohol Abuse*. 2009; **35**(2): 55–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Shields M, Connor Gorber S, Tremblay MS: **Estimates of obesity based on self-report versus direct measures.** *Health Rep*. 2008; **19**(2): 61–76.
[PubMed Abstract](#)
- Wong SL, Shields M, Leatherdale S, *et al.*: **Assessment of validity of self-reported smoking status.** *Health Rep*. 2012; **23**(1): 47–53.
[PubMed Abstract](#)
- Muggah E, Graves E, Bennett C, *et al.*: **Ascertainment of chronic diseases using population health data: a comparison of health administrative data and patient self-report.** *BMC Public Health*. 2013; **13**: 16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kreuter F, Presser S, Tourangeau R: **Social desirability bias in CATI, IVR, and Web surveys the effects of mode and question sensitivity.** *Public Opin Q*. 2008; **72**(5): 847–65.
[Publisher Full Text](#)
- Gmel G, Rehm J: **Measuring alcohol consumption.** *Contemp Drug Probl*. 2004; **31**(3): 467–540.
[Reference Source](#)
- Rehm J, Patra J, Popova S: **Alcohol-attributable mortality and potential years of life lost in Canada 2001: implications for prevention and policy.** *Addiction*. 2006; **101**(3): 373–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wodchis WP, Austin PC, Henry DA: **A 3-year study of high-cost users of health care.** *CMAJ*. 2016; **188**(3): 182–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Manuel DG, Bennett C: **Burden of health behaviours and socioeconomic position on health care expenditure in Ontario.** 2019.
<http://www.doi.org/10.17605/OSF.IO/KW4C5>

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 04 July 2019

<https://doi.org/10.5256/f1000research.19913.r49969>

© 2019 Ding M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

? **Melody Ding**

Prevention Research Collaboration/the Charles Perkins Centre, Sydney School of Public Health, The University of Sydney, Sydney, NSW, Australia

Thank you for the opportunity to review this paper. In this study, the authors presented some novel and interesting analyses to explore health care cost attributable to unhealthy lifestyle patterns and socioeconomic position (SEP; or, perhaps more meaningful, socioeconomic inequalities?). Compared with existing literature on this topic, this study has a few unique contributions: 1) The authors modelled several lifestyle risk factors simultaneously, both in terms of contribution of individual risk factors and combined lifestyle profiles; 2) the authors used data across multiple years and considered the change in attributable and avoided costs over time; 3) the authors considered both lifestyle behaviours and low SEP as risk factors, both individually and combined. However, I have several recommendations for the authors to consider.

1. The information regarding the measurement of lifestyle risk factors and SEP is critical and should be presented with details. What measurement instruments were used? What are the reliability and validity of these measures, what is the rationale for the categorisation? Particularly, the small proportion of healthcare cost attributable to poor diet may be a result of the limitations of the measure (I consider the dietary measure to be a fruit and vegetable intake measure only).
2. Although it is interesting and informative to examine healthcare cost associated with low SEP, I think this idea may require a little bit more consideration. SEP is a relative concept, so what is the practical interpretation of PAF associated with SEP? Were the authors considering the healthcare cost that could have been averted if everyone was of high socioeconomic status? Particularly, how do we correctly interpret PAF associated with ethnicity? What are the counterfactuals?

To me, it may be more meaningful to estimate the PAF of socioeconomic inequality, rather than SEP.

3. It seems that the authors have a-priori decided to model women and men separately? What

is the rationale? However, later on in the paper, when describing the economic cost, all results were presented for both sexes combined. Is there an explanation for this?

4. It looks like the authors have conducted thorough and comprehensive statistical analysis. However, most results are not presented. For example, results from the regression analysis (at least for Model 3) could be of interest to readers.
5. The change analysis between 2004 and 2013: is this based on modelling using all data points or the two time points only?
6. Could you please provide more information and a reference for inverse propensity-weighted analyses?
7. Sorry if I have missed this point from the paper. The modelling of SEP is based on adjusted analysis considering lifestyle risk factors? As we know SEP is a strong predictor of lifestyle behaviour, so parts of the association between SEP and healthcare cost would be explained by lifestyle behaviours. It would be useful if the authors more explicitly define the part of PAF attributable SEP independent of lifestyle risk factors.
8. Given that the data spans over 10 years, have the authors considered inflation? Particularly healthcare specific inflation? It may be worth discussing that the change in PAF over time could be a result of a change in population health behaviour of that in the average costs of certain health care procedures/items.
9. Furthermore, the authors may consider the checklist we developed for reporting cost of illness analysis.¹

Minor comments/questions:

1. Abstract: Taken together, health behaviours and socioeconomic position were associated with 34% (\$134 billion)... suggest add "these" before health behaviours, because this is only referring to the 4 selected behaviours, not health behaviour in general.
2. Methods: could the authors provide some justification regarding why these lifestyle behaviours were selected?
3. Results: "uncertainty for the burden estimates indicates potential overestimation for physical activity and underestimation for diet." Could the authors provide some justification for this statement?
4. The units for the physical activity measure seem incorrect. Could it be MET hour, instead of MET?

References

1. Ding D, Kolbe-Alexander T, Nguyen B, Katzmarzyk PT, et al.: The economic burden of physical inactivity: a systematic review and critical appraisal. *Br J Sports Med*. 2017; **51** (19): 1392-1409
[PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Lifestyle epidemiology, physical activity and health, cost of illness analysis.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 23 April 2019

<https://doi.org/10.5256/f1000research.19913.r45912>

© 2019 Mehta S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Suneela Mehta

School of Population Health, Section of Epidemiology and Biostatistics, University of Auckland, Auckland, New Zealand

This elegant analysis aims to estimate direct healthcare costs in Ontario associated with four adverse health behaviours, the change in direct health care costs associated with these health behaviours over time and the direct healthcare costs associated with low socioeconomic position. This is an extremely valuable endeavour; clarifying the economic cost associated with health behaviour risk factors and lower socioeconomic status assists decision makers in prioritising public health strategies focussed on these domains. The introduction provides a thoughtful background to the study, the definition of the behavioural risk factors (Table 1) and representativeness of the cohort (Table 2) are clearly outlined and the discussion of the study

limitations is excellent.

However, some additional details and minor amendments would be useful:

1. The analysis relates to publicly-funded health care costs in Ontario but as an international reader, it would be helpful to get a brief background of the structure of health system in Canada, and any aspects that are unique to Ontario. In particular, what proportion of healthcare is delivered in the private health sector?
2. Related to point 1), how generalisable are the findings to Canada as a whole and other countries?
3. The descriptions of the public health care spending data considered (on page 4) and the linkage to person-level health care costs (outlined in the Model development section beginning on page 4) are very clear, but the incorporation of the costs to operate the health system into the modelling process needs some further explanation. Are these non-person level costs considered in the MOHLTC costing perspective developed for Ontario health administrative data that is mentioned at the top of page 5? Also, fifteen models have been developed (five per sex for each of the three identified health care sectors) so how have the expenditures associated with the 'Other' category mentioned on page 4 been considered in the analysis?
4. In the first paragraph of the section entitled "Sensitivity analysis", is the explanation of the stepwise model approach describing something different to the description of the model building process detailed on page 5? If not, then I would delete the latter and just leave the last sentence that mentions that models 1 and 5 were used as upper and lower bounds. Furthermore, I would move the information that the primary attribution models were assumed to provide the most accurate estimate of attributable burden of adverse health behaviours to the description on page 5.
5. It would be good to briefly outline the rationale for using a four year period of follow-up on page 5 where this is first mentioned. Also, the titles of tables A-5 to A-8 in the supplementary data state that a five year period between 2004 and 2013 was analysed so these should either be corrected if they are typos or further detail provided in the sensitivity analysis section of the methods.
6. Results are presented in Figure 4 for health care costs attributable to low socioeconomic position but there is no corresponding definition of low socioeconomic position in the methods or results.

Minor amendments:

1. In the methods section of the abstract, it would be helpful to add that the unlinked, cross-sectional CCHS samples for each year from 2004 to 2013 were Ontario samples only.
2. The sentence in the results section of the abstract beginning "Taken together, health behaviours..." would be more accurate if stated as "Taken together, these four adverse health behaviours and low socioeconomic position were associated with 34%". Similarly, the conclusion of the abstract and other sentences similarly worded throughout the paper as well as the labels assigned to Figure 4 would benefit from being reworded to something like "adverse health behaviours" and "low socioeconomic position".
3. In the results section describing the sensitivity analyses, it would be helpful to the reader to include the corresponding table numbers in the supplementary data.
4. The latter part of the sentence in the discussion on page 12 beginning "Fifth, "other" health care costs..." should read 'are not included in our health care cost data...'

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Epidemiology of chronic diseases, cardiovascular risk prediction modelling, linked administrative health data

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 02 May 2019

Doug Manuel, Ottawa Hospital Research Institute, Ottawa, Canada

We thank you for the constructive review and agree with your suggested edits to improve the comprehensiveness of the paper. Our plan is to submit a revised version, addressing the proportion of health care costs covered by Ontario's public health care system and generalizability of results; clarifying the methods for modelling person-level health care costs, the stepped approach to model building, and follow-up period; and, defining measurement of SEP. However, given the current version is under review by another reviewer, we will wait to submit a revised version.

Competing Interests: No competing interests were disclosed.

Author Response 04 Oct 2019

Doug Manuel, Ottawa Hospital Research Institute, Ottawa, Canada

Thank you for your constructive review. We provide a point-by-point summary of response to your comments below.

1. The analysis relates to publicly-funded health care costs in Ontario but as an international reader, it would be helpful to get a brief background of the structure of

health system in Canada, and any aspects that are unique to Ontario. In particular, what proportion of healthcare is delivered in the private health sector?

We have added a description of Canada's health care system to the methods section *Public health care spending data*.

2. Related to point 1), how generalisable are the findings to Canada as a whole and other countries?

We have added a new paragraph in the Discussion (paragraph 3) that focuses solely on generalizability.

3. The descriptions of the public health care spending data considered (on page 4) and the linkage to person-level health care costs (outlined in the Model development section beginning on page 4) are very clear, but the incorporation of the costs to operate the health system into the modelling process needs some further explanation. Are these non-person level costs considered in the MOHLTC costing perspective developed for Ontario health administrative data that is mentioned at the top of page 5? Also, fifteen models have been developed (five per sex for each of the three identified health care sectors) so how have the expenditures associated with the 'Other' category mentioned on page 4 been considered in the analysis?

Failure to indicate our method to incorporate 'other' health care costs (i.e., total provincial health care expenditures minus expenditures in hospital care, community care and drugs) was an oversight. We applied the population attributable fraction from community care to the indirectly estimated other health care costs. We have added this detail to the methods section *Estimating the health care cost burden of health behaviours and socioeconomic position in Ontario*.

4. In the first paragraph of the section entitled "Sensitivity analysis", is the explanation of the stepwise model approach describing something different to the description of the model building process detailed on page 5? If not, then I would delete the latter and just leave the last sentence that mentions that models 1 and 5 were used as upper and lower bounds. Furthermore, I would move the information that the primary attribution models were assumed to provide the most accurate estimate of attributable burden of adverse health behaviours to the description on page 5.

We have revised the text as suggested to address the confusion regarding the model building approach.

5. It would be good to briefly outline the rationale for using a four year period of follow-up on page 5 where this is first mentioned. Also, the titles of tables A-5 to A-8 in the supplementary data state that a five year period between 2004 and 2013 was analysed so these should either be corrected if they are typos or further detail provided in the sensitivity analysis section of the methods.

We have added details to the *Model development* section to clarify the choice of a four-year follow-up time frame.

The supplementary appendices were erroneously labelled. We have revised accordingly.

6. Results are presented in Figure 4 for health care costs attributable to low socioeconomic position but there is no corresponding definition of low socioeconomic position in the methods or results.

We have added details to the methods section *Estimating population attributable fractions of health care costs* to define the comparator group the the SEP analysis.

Minor amendments:

1. In the methods section of the abstract, it would be helpful to add that the unlinked, cross-sectional CCHS samples for each year from 2004 to 2013 were Ontario samples only.

Done

2. The sentence in the results section of the abstract beginning “Taken together, health behaviours...” would be more accurate if stated as “Taken together, these four adverse health behaviours and low socioeconomic position were associated with 34%”. Similarly, the conclusion of the abstract and other sentences similarly worded throughout the paper as well as the labels assigned to Figure 4 would benefit from being reworded to something like “adverse health behaviours” and “low socioeconomic position”.

Done

3. In the results section describing the sensitivity analyses, it would be helpful to the reader to include the corresponding table numbers in the supplementary data.

Done

4. The latter part of the sentence in the discussion on page 12 beginning “Fifth, “other” health care costs...” should read ‘are not included in our health care cost data...’

Done

Competing Interests: No competing interests were disclosed.

Author Response 04 Oct 2019

Doug Manuel, Ottawa Hospital Research Institute, Ottawa, Canada

Thank you for your helpful review. We have added a number of details to respond to your points. See our point-by-point response below.

1. The information regarding the measurement of lifestyle risk factors and SEP is critical and should be presented with details. What measurement instruments were used? What are the reliability and validity of these measures, what is the rationale for the categorisation? Particularly, the small proportion of healthcare cost attributable to poor diet may be a result of the limitations of the measure (I consider the dietary measure to be a fruit and vegetable intake measure only).

We have added details to the methods section *Behavioural and other risk factors for health care use* to more fully describe measurement of health behaviour and sociodemographic risk factors. We have also expanded the discussion *Limitations* section.

2. Although it is interesting and informative to examine healthcare cost associated with low SEP, I think this idea may require a little bit more consideration. SEP is a relative concept, so what is the practical interpretation of PAF associated with SEP? Were the authors considering the healthcare cost that could have been averted if everyone was of high socioeconomic status? Particularly, how do we correctly interpret PAF associated with ethnicity? What are the counterfactuals?

We agree, SEP is a relative concept. We have framed the analysis as an equity gap and added details to the *Estimating population attributable fractions of health care costs* to describe the exposures used for the analysis.

3. To me, it may be more meaningful to estimate the PAF of socioeconomic inequality, rather than SEP.

See above.

4. It seems that the authors have a-priori decided to model women and men separately? What is the rationale? However, later on in the paper, when describing the economic cost, all results were presented for both sexes combined. Is there an explanation for this?

Yes, we did a priori model men and women separately. There were several considerations such as differential misclassification bias of exposure (e.g., men and women have different pattern of misclassification error for weight and height). Many/most studies of the underlying leading causes of diseases sex-stratify based, for example, on the observed effect modification for cardiovascular disease and most leading causes of cancer. The results were then combined to clearly present our main focus on health behaviours and SEP.

5. It looks like the authors have conducted thorough and comprehensive statistical analysis. However, most results are not presented. For example, results from the regression analysis (at least for Model 3) could be of interest to readers.

We agree with your suggestion. As you can appreciate, there was a wide range of components for the study. However, on reflection we agree that these models are of

interest, particularly for replication or for use by other jurisdictions who may use the estimated to indirectly model burden. These models will be added to the on-line figure repository. (The analyst who created those models is currently on leave).

6. The change analysis between 2004 and 2013: is this based on modelling using all data points or the two time points only?

We have revised the methods section *Estimating costs attributable to changes in health behaviours* to more clearly indicate that this analysis is based on annual estimates.

7. Could you please provide more information and a reference for inverse propensity-weighted analyses?

We have added additional details and references to the description of the inverse propensity-weighted sensitivity analysis.

8. Sorry if I have missed this point from the paper. The modelling of SEP is based on adjusted analysis considering lifestyle risk factors? As we know SEP is a strong predictor of lifestyle behaviour, so parts of the association between SEP and healthcare cost would be explained by lifestyle behaviours. It would be useful if the authors more explicitly define the part of PAF attributable SEP independent of lifestyle risk factors.

The findings for both health behaviours and SEP were estimated using Model 3 simultaneously for health behaviours and SEP. We have clarified this in the manuscript. We agree with the reviewer's perspective that SEP and health behaviours have strong relationships with complex pathways.

As such, Model 3 likely represents a somewhat conservative estimate of the independent effects of either health behaviours or SEP. The sensitivity analyses were performed with less and more specified models to gauge the potential effect of adding or removing mediators. Model 1 was a simple model of health behaviours without SEP measures. Of note, we didn't create to complementary model—i.e., a simple model of SEP without health behaviour measures.

Given the complex relationships, we recognize that there are alternative methods approaches that we hope will be further explored in future studies.

9. Given that the data spans over 10 years, have the authors considered inflation? Particularly healthcare specific inflation? It may be worth discussing that the change in PAF over time could be a result of a change in population health behaviour of that in the average costs of certain health care procedures/items.

For this study, all costs are expressed in 2014 Canadian dollars with past costs inflated using the annual general Consumer Price Index from Statistics Canada. It is unclear what is meant by "...could be a result of a change in population health behaviour of that in the average costs of certain health care procedures/items."

10. Furthermore, the authors may consider the checklist we developed for reporting

cost of illness analysis.

We've completed the checklist and added it to the on-line repository. We had been holding off using a reporting guideline until those for population modelling studies (<http://www.mrc-epid.cam.ac.uk/ph-modelling-guidelines/>) became available. We have also performed studies that have examined how modelling/burden studies can be reported and found the CHEERS guidelines are relevant.

Minor comments/questions:

1. Abstract: Taken together, health behaviours and socioeconomic position were associated with 34% (\$134 billion)... suggest add "these" before health behaviours, because this is only referring to the 4 selected behaviours, not health behaviour in general.

We have modified the language in the abstract to more clearly indicate the attributable costs are related to select health behaviours and socioeconomic measures.

2. Methods: could the authors provide some justification regarding why these lifestyle behaviours were selected?

In the Introduction we highlight our focus on smoking, unhealthy alcohol consumption, poor diet and physical inactivity as the leading risk factors for morbidity and mortality worldwide.

3. Results: "uncertainty for the burden estimates indicates potential overestimation for physical activity and underestimation for diet." Could the authors provide some justification for this statement?

We have added details to the results section *Health behaviour attributable healthcare use* to indicate we are referring to the sensitivity analyses represented by the error bars in Figure 3.

4. The units for the physical activity measure seem incorrect. Could it be MET hour, instead of MET?

We have clarified in the table that our measure was MET-hours/day.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research