Check for updates

RESEARCH ARTICLE

Adaptation, fitness landscape learning and fast evolution

[version 1; peer review: 2 approved with reservations]John Reinitz¹, Sergey Vakulenko²⁻⁴, Dmitri Grigoriev⁵, Andreas Weber⁶ ¹Departments of Statistics, Ecology and Evolution, Molecular Genetics and Cell Biology, University of Chicago, Chicago, IL, USA²Saint Petersburg University of Industrial Technologies and Design, Saint Petersburg, Russian Federation³Institute for Mechanical Engineering Problems, Russian Academy of Sciences, Saint Petersburg, Russian Federation⁴Saint Petersburg National Research University of Information Technologies, Mechanics and Optics, Saint Petersburg, Russian Federation⁵CNRS, Mathématiques, Université de Lille, Villeneuve d'Ascq, France⁶Department of Computer Science, University of Bonn, Bonn, Germany**v1** First published: 01 Apr 2019, 8:358
<https://doi.org/10.12688/f1000research.18575.1>Latest published: 13 Sep 2019, 8:358
<https://doi.org/10.12688/f1000research.18575.2>**Abstract**

We consider evolution of a large population, where fitness of each organism is defined by many phenotypical traits. These traits result from expression of many genes. Under some assumptions on fitness we prove that such model organisms are capable, to some extent, to recognize the fitness landscape. That fitness landscape learning sharply reduces the number of mutations needed for adaptation. Moreover, this learning increases phenotype robustness with respect to mutations, i.e., canalizes the phenotype. We show that learning and canalization work only when evolution is gradual. Organisms can be adapted to many constraints associated with a hard environment, if that environment becomes harder step by step. Our results explain why evolution can involve genetic changes of a relatively large effect and why the total number of changes are surprisingly small.

Keywords

evolution, gene networks, fitness landscape learning

Open Peer Review**Approval Status**

	1	2
version 2		
(revision)
13 Sep 2019
version 1
01 Apr 2019

1. **Eors Szathmáry** , MTA Centre for Ecological Research, Tihany, Hungary
 Parmenides Center for the Conceptual Foundations of Science, Pullach, Germany
 Eötvös University, Budapest, Hungary

2. **Aviv Bergman**, Albert Einstein College of Medicine, New York, USA
Yehonatan Sella, Albert Einstein College of Medicine, New York, USA

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding authors: John Reinitz (reinitz@galton.uchicago.edu), Sergey Vakulenko (vakulenfr@mail.ru)

Author roles: **Reinitz J:** Conceptualization, Methodology, Writing – Review & Editing; **Vakulenko S:** Conceptualization, Formal Analysis, Investigation, Methodology, Software, Writing – Original Draft Preparation, Writing – Review & Editing; **Grigoriev D:** Formal Analysis, Methodology, Validation, Writing – Review & Editing; **Weber A:** Software, Visualization, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The second author was supported by the grant of Russian Ministry of Education, 2012-1.2.1-12-000-1013-016. Additionally, the second author was financially supported by Government of Russian Federation, Grant Grant 08-08. D. Grigoriev is grateful to the grant RSF 16-11-10075 and to both MCCME and MPI f"ur Mathematik for wonderful working conditions and inspiring atmosphere. J. Reinitz and S. Vakulenko were supported by US NIH grant RO1 OD010936 (formerly RO1 RR07801).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 Reinitz J *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Reinitz J, Vakulenko S, Grigoriev D and Weber A. **Adaptation, fitness landscape learning and fast evolution [version 1; peer review: 2 approved with reservations]** F1000Research 2019, 8:358 <https://doi.org/10.12688/f1000research.18575.1>

First published: 01 Apr 2019, 8:358 <https://doi.org/10.12688/f1000research.18575.1>

1. Introduction

A central idea of modern biology is that evolution proceeds by mutation and selection. This process may be represented as a walk in a fitness landscape leading to fitness increase and slow adaptation¹. According to classical ideas this walk can be considered a sequence of small random steps with small phenotypic effects. Nevertheless, there is a limited amount of experimental support for this idea² and some experimental evidence that evolution can involve genetic changes of a relatively large effect and that the total number of changes are surprisingly small³. Another intriguing fact is that organisms are capable of making adaptive predictions of environmental changes⁴.

To explain those facts new evolutionary concepts have been suggested (see the review by 5 and references therein). The main idea is that a population can “learn” (recognize) fitness landscapes^{5–7}. The approach developed in these works is a generalization of ideas from machine learning in which learning (regression to data) is viewed as selection and generalization (interpolation or extrapolation) is viewed as adaptation.

A mathematical basis for investigation of evolution learning problems has been developed by 8. However, this work uses a simplified model, where organisms are represented as Boolean circuits seeking an “ideal answer” to environmental challenges. These circuits involve N_g Boolean variables that can be interpreted as genes, and the ideal circuit answer maximizes the fitness. A similar model was studied numerically by 7 to confirm the theory of “facilitated variation” explaining the appearance of genetic variations which can lead to large phenotypic ones. In the work by 9 a theory of the evolution of these Boolean circuits was advanced. It was shown that, under some conditions—weak selection, see 10—a polynomially large population over polynomially many generations (polynomial in N_g) will end up almost surely consisting exclusively of assignments, which satisfy all constraints. This theorem can shed light on the problem of the evolution of complex adaptations since that satisfiability problem can be considered as a rough mathematical model of adaptation to many constraints.

In 6 it is shown that, in the regime of weak selection, population evolution can be described by the multiplicative weight update algorithm (MWUA), which is a powerful tool, well known in theoretical computer science and a generalization of such famous algorithms as Adaboost and others¹¹. Note that in 6 infinitely large populations are investigated whereas the results of 9 hold only for finite populations and take into account genetic drift.

Evolution Computation(EC) problems are considered recently by many papers^{12–16} mainly for artificial test fitness functions like OneMax or LeadingOnes (for an overview of EC problems, see 17).

In this paper, we investigate adaptation and the fitness landscape learning problem for more realistic fitness function. This fitness can model adaptation for insects and connected with a fundamental hard combinatorial problem: K -SAT.

The main results can be outlined as follows. We show that, in a fixed environment, genes can serve as learners in the machine learning sense. Indeed, if an organism has survived for a long period, this fact alone constitutes important information, which can be used. The biological interpretation of this fact is simple: if a population is large enough and mutations are sufficiently rare, deleterious mutations are eliminated by purifying selection. Hence, those non-neutral mutant alleles which have become fixed in natural populations will, with probability close to 1, be adaptive and cause a positive increment of fitness (see Theorem 3.1 and Theorem 3.2 in Subsection 3.1 and Subsection 3.2). We obtain mathematical results, which allows us to estimate the reduction of mutation number due to that learning landscape procedure. Learning can sharply reduce the number of mutations needed to form a phenotypic trait useful for adaptation that is consistent with experimental data mentioned above (see 3).

Another important result is as follows. We estimate the accuracy of fundamental Nagylaki equations^{6,10} for a realistic population model, where the population size is bounded and a non-zero mutation rate is taken into account (in the case of asexual reproduction). Those accuracy estimates are fulfilled for all possible values of mutation rates and population sizes.

2. Model

In this section, we describe our model and mathematical approach.

2.1 Genome

We assume that the genotype can be described by Boolean strings of length N_g , where N_g is the number of genes. Then

$$s = (s_1, s_2, \dots, s_{N_g}), \quad s_i \in S = \{0, 1\}, \quad s \in S^{N_g}, \quad (2.1)$$

where $s_i = 1$ means that gene i is activated (switched on) and $s_i = 0$ means that it is repressed (switched off). Correspondence between Boolean hypercube and genotypes is considered for example in 12.

2.2 Phenotypic traits

Although phenotype is controlled by genes, it is also influenced by environmental conditions and various epigenetic processes. In this paper, we suppose that phenotypic traits are controlled by genotype only. We consider levels f_j of expressions of those traits as real variables in the interval $(0, 1)$. Then the vector $f = (f_1, \dots, f_{N_b})$ can be considered to represent the organismal phenotype. We suppose that

$$f_j = f_j(s), \quad j = 1, \dots, N_b, \quad (2.2)$$

where $f_j \in (0, 1)$ is a real valued function of the Boolean string s , the genotype.

Only a part of s_i is involved in f_j . Namely, for each j we have a set of indices $K_j = \{i_1, i_2, \dots, i_{n_j}\}$ such that f_j depends on s_i with $i \in K_j$, so that

$$f_j(s) = f_j(s_{i_1}, s_{i_2}, \dots, s_{i_{n_j}}),$$

where $i_l \in K_j$ and n_j is the number of genes involved in the control of the trait expression. The representation of phenotype by the quantities f_j is suggestive of quantitative traits because the f_j are real valued. The limiting values of 0 or 1 suggest another interpretation, however, in terms of cell type. Multicellular organisms consist of cells of different types. One can suppose that the organismal phenotype is defined completely by the corresponding cell pattern. The cell type j is determined by morphogenes, which can be identified as gene products or signaling molecules that can change cell type (or genes that code for signaling molecules that can determine cell types or cell-cell interactions and then finally the cell pattern). The morphogene activity is defined by (2.2).

We further suppose:

Assumption M. Assume activities f_j have the following properties.

The sets K_j are independent uniformly random subsets of $S_g = \{1, \dots, N_g\}$

$$K_j = \{i_1, \dots, i_{n_j}\}, \quad i_l \in S_g, \quad l = 1, \dots, n_j. \quad (2.3)$$

We denote the total number of genes involved in regulation of all f_j by N_r , where $N_r = \sum n_j \leq KN_b$.

Assumption M implies that the genetic control of the phenotype is organized, in a sense, randomly, and that only a portion of the full set of genes controls phenotypic traits. That modularity of gene control is well known from experimental data (see 18, 19) and for evolution computation problems it was studied, for example, in 16.

Consider an example, where the assumption M holds, where we have a saturated expression, inspired by earlier work^{20,21}. Let

$$f_j = s \left(\sum_{i=1}^{N_g} w_{ji} s_i - h_j \right), \quad (2.4)$$

where $j = 1, \dots, N_b$. Here $\sigma(z)$ is a sigmoidal function of real z such that

$$\sigma(+\infty) = 1, \quad \sigma(-\infty) = 0, \quad \sigma'(z) > 0 \quad \forall z \quad (2.5)$$

and w_{ji}, h_j are some coefficients (their meaning will be explained below). As an example, we can take $\sigma(S) = (1 + \exp(-bS))^{-1}$, where $b > 0$ is a sharpness parameter. Note that for large b this sigmoidal function tends to the step function and for $b = +\infty$ our model becomes a Boolean one. The parameters h_j defines thresholds for trait expression²⁰. The relation (2.4) can be interpreted as a simple mathematical model for quantitative trait locus (QTL) action.

To understand the role of h_j consider a trait f_j and suppose that for a well adapted organism $f_j \approx 1$. Let, for simplicity, w_{ji} take the values 1, 0, or -1. Then the parameter h_j defines how many genes involved in the control of the f_j expression should be activators and how many should be repressors. Let the numbers of activator and repressor genes be n_j^+ respectively.

Then $f_j \approx 1$ if $n_j^+ - n_j^- \gg h_j$.

One can suppose that h_j describes a direct influence of environment on phenotype, such as stress, that can exert epigenetic effects. In Section 2.5 using data from 3 we will show that the model defined by (2.4) are capable to describe main topological characteristics of really observed fitness functions in the case of mimicry, camouflage and thermoregulation for insects.

Let us introduce the matrix $\mathbf{W}$ of size $N_b \times N_g$ with the entries w_{ji} . The coefficients w_{ji} determine the effects of terminal differentiation genes (see 23), and hence encodes the genotype-phenotype map. We assume that the coefficients w_{ji} are random, with the probability that $w_{ji} > 0$ or that $w_{ji} < 0$ is $\beta/2N$, where $\beta > 0$ is a parameter. This quantity $\beta \ll N$ defines a genetic redundancy, i.e., averaged numbers of genes that control a trait. Note that then large $\beta \gg 1$ one has $n_j < 4\beta$ with the probability $\Pr_\beta$, which is exponentially close to 1: $\Pr_\beta > 1 - \exp(-0.1\beta)$, thus, the number n_j are bounded.

2.3 Fitness

We know little about the details of how fitness relates to the phenotype of multicellular organisms, and for that reason classic neo-Darwinian theory takes fitness to be a function of genotype. Some models which take account of epistasis have been proposed²⁴. The random field models assign fitness values to genotypes independently from a fixed probability distribution. They are close to mutation selection models introduced by 25, and can be named House of Cards (HoC) model. The best known model of this kind is the NK model introduced by Kauffman and Weinberger²⁶, where each locus interacts with K other loci. Rough Mount Fuji (RMF) models are obtained by combining a random HoC landscape with an additive landscape models²⁷. In evolution computations (EC) some artificial fitness models were used, for example OneMax and Leading Ones to test evolution algorithms, see for example¹⁵.

In this work, we use the classical approach of R. Fisher by introducing an explicit representation of phenotype, f , and allow it to determine fitness through interaction with an environment b . That is, we assume that the phenotype is completely determined by the phenotype trait expression, and thus the fitness depends on the genotype s via f_j .

We express the relative fitness F and its dependence on environment b via an auxiliary function W via the relation

$$F(s, b) = K_F \exp(W(s, b)), \quad (2.6)$$

where K_F is a positive constant and $b = (b_1, \dots, b_{N_b})$ is a vector consisting of coefficients b_j , respectively. Below we refer to W as a fitness potential, and we assume that

$$W(s, b) = \sum_{j=1}^{N_b} b_j f_j(s). \quad (2.7)$$

Sometimes, if the parameter b is fixed, we shall omit the corresponding argument in notation for W and F .

We consider fitness as a numerical measure of interactions between the phenotype and an environment. For a fixed environment, this idea gives us the fitness of classical population genetics. A part of the fitness, however, depends on the organism developing properly and for now we represent it as independent of the environment, although we are aware that this is not always the case. Note that some coefficients b_j may be negative and others may be positive, and that the model (2.7) can describe gene epistatic effects via dependence of f_j on s if f_j are nonlinear in s .

The expression (2.7) can serve as a rough approximation of the fitness function in the case of insects such as grasshoppers or fruit flies. In fact, important factors, which determine insect survival, are thermoregulation, mimicry and camouflage levels^{18,22,28}. All those factors depend on colour pigmentation pattern. Blackwhite pigmentation patterns can be roughly described by vectors $f = (f_1, f_2, \dots, f_{N_b})$, where $f_j \approx 1$ and $f_j \approx 0$ mean that the cell j is black, or white, respectively. Then thermoregulation depends on $\sum_j f_j$. The mimicry level can be approximately defined by expression $\sum_j |f_j - f_j^*|$, where f_j^* is a target pattern corresponding to an insect to mimic. Colour patterns can be also described by classical RGB formalism.

The representation of the fitness as a sum of terms (2.7) is of course a rough approximation; however if assumption **M** holds that representation is consistent with important observed facts. First, mutations have been identified that alter one part of the pigment pattern without affecting any other. This independence of different pattern parts can be explained by the modular organization of the genetic regulation that controls pigmentation. In the course of evolution, different aspects of the pigment pattern have clearly evolved independently of each other¹⁸. Second, the topology of the fitness landscapes was studied in ²² by field experiments in the case of insect mimicry. Main conclusions are as follows. A number of studies of fitness landscapes in natural populations have demonstrated low fitness of intermediate phenotypes, i.e., existence of valley in the fitness landscape. It is found²² that natural selection promotes genetic architecture preventing the expression of intermediate phenotypes. Close fitness peaks are separated by ridges, favouring colour pattern switches and allowing drift from local peaks.

In Section 2.5 we will show that the fitness model defined by (2.4) and (2.7) have those topological properties.

2.4 Population dynamics model

For simplicity, we consider populations with asexual reproduction. (Although a part of the results remain valid for sexual reproduction, as we discuss at the end of this subsection). We choose initial genotypes randomly from a gene pool and assign them to organisms. This choice is invariant with respect to the population member, i.e., the probability to assign a given genotype s to a member of the population does not depend on that member.

In each generation, there are $N_{\text{pop}}(t)$ individuals, the genome of each of which is denoted by $s(t)$, where $t = 0, 1, 2, \dots$ stands for the evolution step number). Following the classical Wright-Fisher

ideas, we suppose that generations do not overlap. In each generation (i.e., for each t), the following three steps are performed:

1. Each individual s at each evolution step can mutate with probability p_{mut} per gene;
2. At evolution step t each individual with a genotype s produces k progeny, where k is a random non-negative integer, distributed according to the Poisson law

$$P_k = \frac{q^k}{k!} \exp(-q), \quad (2.8)$$

where $q = F(s)$ is the fitness of that individual;

3. To take into account ecological restrictions on the population size, we introduce the maximal population size N_{popmax} . If $N'(t) > N_{\text{popmax}}$, where $N'(t)$ is the number of progeny produced by the population at step t , we kill randomly selected individuals in a population-dependent manner. The probability of the death of an individual is given by $p_{\text{kill}}(N') = 1 - (N_{\text{popmax}}/N'(t))$. If $N'(t) \leq N_{\text{popmax}}$, we do nothing. We refer to this as the “massacre procedure.”

Conditions 1 and 2 imply that mutations in the genotypes create a new genetic pool and then a new round of selection starts. Condition 3 expresses the fundamental ecological limitation that all environments can only support populations of a limited size. If $N_{\text{popmax}} \gg 1$ then by (2.8) and the Central Limit Theorem one can show, under some additional conditions, (see Section 4) that fluctuations of the population size are small, and thus the population is ecologically stable and $N_{\text{pop}}(t) \approx N_{\text{popmax}}$.

In the limit case of infinitely large populations we will write the discrete dynamical equation for the time evolution of the frequency $X(s, t)$ of the genotype s in the population as

$$X(s, t+1) = \bar{F}(t)^{-1} X(s, t) F(s), \quad X(s, 0) = X_0(s), \quad (2.9)$$

where $\bar{F}(t)$ is the average fitness of the population at the moment t defined by

$$\bar{F}(t) = \sum_{s \in S(t)} X(s, t) F(s), \quad (2.10)$$

where $S(t)$ is the set of genotypes existing in the population at time t (the genetic pool) and $X(s, t) = N(s, t)/N_{\text{pop}}(t)$ is the frequency of the genotype s . Here $N(s, t)$ denotes the number of the population members with the genotype s at the step t .

The equations (2.9) do not take mutations into account. They describe only changes in the genotype frequencies because of selection at the t -th time step. The same equations govern evolution in the case of sexual reproduction in the limit of weak selection^{6,10}. Note that for an evolution defined by (2.9), the average fitness $\bar{F}(t)$ defined by (2.10) satisfies Fisher's theorem, so that this function increases at each time step t : $\bar{F}(t+1) \geq \bar{F}(t)$.

2.5 Adaptation as a hard combinatorial problem

Adaptation (i.e., maximization of fitness in a changing environment) is a very hard problem since over evolutionary history we

observe the coevolution of many traits accompanied by changes in many genes. In its general context, this is a problem in the theory of macroevolution, which in general requires the integration of population genetics and developmental biology for its full understanding. There are two key components of this problem. First, development is itself a dynamical process operating over time. Second, there is a combinatorial component of development wherein different combinations of gene must be expressed in different cell types. This combinatorial aspect of the problem means that straightforward theoretical methods of considering the relationship between gene expression and a changing environment that have been very successful in single celled organisms²⁹ cannot be applied to metazoa. In this work, for the sake of tractability, we focus on the combinatorial aspect of the problem and neglect developmental dynamics. Even at the highly simplified level of our model, adaptation is a hard computational problem, as we now demonstrate.

Consider the case, where f_j are defined by relations (2.4) and assume that

- i) σ is the step function;
- ii) $b_j > 0$.

As a consequence of the second assumption, F attains its maximum for $f_1 = 1, f_2 = 1, \dots, f_{N_b} = 1$. Let us show that, even in this particular case, the problem of the fitness maximization with respect to s is very complex. In fact, for a choice of h_j it reduces to the famous NP-complete problem, so-called K -SAT, which has received a great deal of attention in the last few decades (see 30–35). The K -SAT can be formulated as follows.

K -SAT problem. *Let us consider the set $V_n = \{s_1, \dots, s_n\}$ of Boolean variables $s_i \in \{0, 1\}$ and a set $\mathcal{C}_m$ of m clauses. The clauses C_j are disjunctions (logical ORs) involving K literals $z_{j1}, z_{j2}, \dots, z_{jK}$, where each z_{ji} is either s_i or the negation $\bar{s}_i$ of s_i . The problem is to test whether one can satisfy all of the clauses by an assignment of Boolean variables.*

Cook and Levin^{30,31} have shown that the K -SAT problem is NP-complete and therefore in general it is not feasible in a reasonable running time. In subsequent studies—for instance, by 32—it was shown that K -SAT of a random structure is feasible under the condition that $N_b < \alpha_c(K)N_g$, where $\alpha_c(K) \approx 2^K \log 2$ for large K .

The set $\mathcal{C}_K$ of solutions of random K -SAT has a nontrivial structure depending on parameter $\alpha = N_b/N_g$ ^{33,35}. For sufficiently small $\alpha < \alpha_g(K)$, where $\alpha_g(K) \approx 2^K \log(K)/K$ is some critical parameter, the set $\mathcal{C}_K$ forms a giant cluster, where nearest solutions are connected by a single flip and one can go from a solution to another by a sequence of single flips (pointed mutations)³⁵. For $\alpha \in (\alpha_g, \alpha_d)$, where $\alpha_d(k) < \alpha_c$ is another critical value, solutions form a set of disconnected clusters. The local search algorithms do not work in the domain $\alpha > \alpha_g$.

Probably, for evolution context K -SAT was applied first in 36, where it was used for an investigation of speciation problem.

To see the connection of our model with K -SAT, consider equation (2.4) supposing that $w_{ji} \in \{1, 0, -1\}$ and $h_j = -C_j + 0.5$, where C_j is the number of negative w_{ji} in the sum $s_j = \sum_{i=1}^{N_g} w_{ji}s_i$. We set $m = N_b$ and $n = N_g$. Under this choice of h_j , the terms $\sigma(s_j)$ can be represented as disjunctions of literals z_j . Each literal z_j equals either s_j or $\bar{s}_j$, where $\bar{s}_j$ denotes negation of s_j . To maximize the fitness, we must assign s_j such that all disjunctions will be satisfied. If we fix the number n_j of the literals participating in each disjunction (clause) and set $n_j = K$, this assignment problem is precisely the K -SAT problem formulated above.

Reduction to the K -SAT problem is a transparent way of representing the idea that multiple constraints need to be satisfied. The number K defines the gene redundancy and the probability of gene pleiotropy. Remind that pleiotropy occurs when one gene influences two or more seemingly unrelated phenotypic traits. The threshold h_j and K define the number of genes which need be flipped in order to attain a high expression of the trait f_j . Note that gene pleiotropy is a fundamental characteristics³⁷, which is studied for real organisms only recently (see 19). We can compare experimental observations and consequence of model (2.4), which is a generalisation of K -SAT (compare plot Figure 1 and plots on Figure 1 in 19). So, we can fit our model parameters using real data. Moreover, we can check validity of our model by the following arguments.

We note that, in the case of giant cluster formation, the topological properties of the solution set $\mathcal{C}_K$ mentioned above, outline the properties of really observed fitness landscapes²²: existence of many peaks, valleys and ridges connecting peaks. Namely, existence of many solutions of K -SAT, when a giant cluster exists, means that the landscape has a number of peaks separated by valleys. On the other hand, connectance of solutions within the giant cluster can be interpreted that there exist ridges that connect peaks.

Note that there are important differences between K -SAT in Theoretical Computer Science and fitness maximization problems. First, the signs of b_j are unknown for real biological situations since the fitness landscape is unknown. Second, our adaptation problem involves the threshold parameters h_j (see (2.4)). In contrast to K -SAT, in our case the Boolean circuit is plastic, because the h_j are not fixed.

If the b_j are unknown, the adaptation (fitness maximization) problem becomes even harder because we do not know the function to optimize. Therefore, many algorithms for K -SAT are useless for biological adaptation problems. Below we will nonetheless obtain some analytical results based on the assumption that b_j are random.

3. Main theorems

The subsequent material is organized as follows. First we formulate a result on regulation mechanism power. Furthermore, we prove two fitness landscape learning theorems.

3.1 Fitness landscape learning theorems

For simplicity, we consider asexual reproduction. To obtain similar results for sexual reproduction, one can consider a weak

Figure 1. Frequency distributions of degree of gene pleiotropy for model (2.4) with the parameters $N_g = 4000$, $\beta = 4$, $h = 0$, $N_b = 3000$.

selection regime and use the results of 10, where eq. (2.9) are derived.

Let us introduce two sets of indices I_+ and I_- , such that $I_+ \cup I_- = \{1, \dots, N_b\}$. We refer to these sets in the sequel as positive and negative sets, respectively. We have

$$I_+ = \{j \in \{1, \dots, N_b\} | b_j > 0\}, \quad (3.1)$$

$$I_- = \{j \in \{1, \dots, N_b\} | b_j < 0\}. \quad (3.2)$$

The biological interpretation of that definition is transparent: the expression of the traits f_j with $j \in I_+$ increases the fitness and for $j \in I_-$ expression of the trait decreases the fitness.

Let s and $\bar{s}$ be two genotypes. Then we denote by $\text{Diff}(s, \bar{s})$ the set of positions i such that $s_i \neq \bar{s}_i$:

$$\text{Diff}(s, \bar{s}) = \{i \in \{1, \dots, N_g\} | s_i \neq \bar{s}_i\}.$$

The set $\text{Diff}(s, \bar{s})$ indicates which genes in s should be flipped in order to obtain $\bar{s}$.

We formulate two theorems on fitness landscape learning. First we consider the case of infinitely large populations.

Theorem 3.1. Suppose that the evolution of the genotype frequencies $X(s, t)$ is determined by equations (2.9) and (2.10). Moreover, assume that

I for all $t \in [T_1, T_1 + T_c]$, where $T_1, T_c > 0$ are integers, the population contains two genotypes s and $\bar{s}$ such that the frequencies $X(s, t)$ and $X(\bar{s}, t)$ satisfy

$$X(s, T_1) = p_0 > 0, \quad X(\bar{s}, T_1 + T_c) = p_1 > 0, \quad (3.3)$$

II we have

$$\text{Diff}(s, \bar{s}) \subset K_j, \quad (3.4)$$

for some j . In other words, the genes s_i such that $s_i \neq \bar{s}_i$ are involved in a single regulation set K_j ; and finally,

III Let

$$\delta_j = |f_j(s) - f_j(\bar{s})| > 0, \quad |b_j| > 0, \quad (3.5)$$

and

$$T_c > \frac{-\log(p_0 p_1)}{|b_j| \delta_j}. \quad (3.6)$$

Then, if

$$f_j(s) < f_j(\bar{s}), \quad (3.7)$$

we have $j \in I_+$. If $f_j(s) > f_j(\bar{s})$, then $j \in I_-$.

Before proving this, let us make some comments. The biological meaning of the theorem is simple: for simple fitness models, where unknown parameters b_j are involved in a linear way, in the limit of infinitely large populations fitness landscape learning is possible.

Moreover, note that we do not make any specific assumptions about the nature of mutation, but only that all genetic variation between s and $\bar{s}$ are contained in a single regulatory set K_j .

The assertion of Theorem 3.1 is not valid if the set $\text{Diff}(s, \bar{s})$ belongs to a union of different regulation sets K_j , $j = j_1, \dots, j_p$.

with $p > 1$. This effect of belonging to different sets K_j is pleiotropy in gene regulation. Note that if $N_b \ll N_g$ then the pleiotropy probability is small for large genome lengths N_g . On the contrary, if $N_b \gg N_g$ then assumption **II** is invalid.

Assumption **II** looks natural if when we deal with point mutations. In fact, if $\bar{s}$ is obtained from s by a single point mutation then condition (3.4) always holds for some j . For small mutation rates the probability of two point mutations is essentially below than the probability of a single mutation.

To conclude let us note that Theorem gives a rough estimate for the learning time T_c :

$$T_c = O\left(-\frac{\log(p_0 p_1)}{|b_j| \delta_j}\right).$$

Proof. The main idea is simple. Negative mutations lead to elimination of mutant genotypes from the population, and the corresponding frequencies become, for large times, exponentially small.

Assume that (3.7) holds. Let $j \in I_-$, and thus $b_j < 0$. Consider the quantity

$$Q(t) = \frac{X(s, t)}{X(\bar{s}, t)} = \frac{N(s, t)}{N(\bar{s}, t)}. \quad (3.8)$$

According to assumption **II**

$$\Delta W = W(s) - W(\bar{s}) = b_j (f_j(s) - f_j(\bar{s})). \quad (3.9)$$

Assumption **III** entails that

$$\Delta W \geq |b_j| \delta_j. \quad (3.10)$$

Relations (2.6) and (3.10) imply

$$\frac{F(s)}{F(\bar{s})} = \exp(\Delta W) \geq \exp(|b_j| \delta_j).$$

By (2.9) and the last inequality we find that for $T > T_1$

$$Q(T) \geq Q(T_1) \exp(|b_j| \delta_j (T - T_1)). \quad (3.11)$$

Consider inequality (3.11) for $T = T_1 + T_c$. Let us note that in the relation $Q(T_1) = X(s, T_1)/X(\bar{s}, T_1)$ the numerator is p_0 whereas the denominator ≤ 1 . Thus, $Q(T_1) \geq p_0$. The same arguments show that $Q(T_1 + T_c) \leq 1/p_1$. Therefore, by (3.11) one obtains that

$$\frac{1}{p_1 p_0} \geq \exp(|b_j| \delta_j T_c). \quad (3.12)$$

This inequality leads to a contradiction for T_c satisfying (3.6), thus completing the proof.

3.2 The case of finite populations

Theorem 3.1 can be extended to the case of finite populations and non-zero mutation rates. is small. To formulate this generalization, we need an additional assumption about the fitness function. Suppose that

$$\begin{aligned} 1 < c_F &< \min_{s \in S(t)} F(s), \\ \max_{s \in S(t)} F(s) &< C_F, \quad \forall t \in [T_1, T_1 + T_c] \end{aligned} \quad (3.13)$$

where $c_F, C_F > 0$ are constants independent of t . For example, if

$$\sum_{j=1}^{N_b} |b_j| < \gamma,$$

then $c_F = K_F \exp(-\gamma)$ and $C_F = K_F \exp(\gamma)$ and (3.13) holds if $K_F > \exp(\gamma)$.

Condition (3.13) means that each individual gives birth to at least c_F and at most C_F descendants, where those bounds do not depend on the population size and evolution step.

Let

$$N_{\text{pop}}(T_1) = N_{\text{popmax}}. \quad (3.14)$$

Note that for simplicity in the next Theorem 3.2 we consider point mutations (bit flipping) only. The model used here cannot represent mutations of arbitrarily small effect, but it can include insertions or deletions. In contrast to Theorem 3.2, Theorem 3.1 is valid for all kinds of mutations.

Then we have

Theorem 3.2. Consider the population dynamics defined by model 1-3 in subsection 2.4. Assume conditions (3.14) and **M** hold, and assumptions (3.3), (3.4), (3.5), (3.7) of Theorem 3.1 are satisfied. Suppose

$$X(s, t) \geq p_0 \quad \forall t \in [T_1, T_1 + T_c], \quad (3.15)$$

$$X(\bar{s}, t) \geq p_1 \quad \forall t \in [T_1, T_1 + T_c]. \quad (3.16)$$

Then if $j \in I_-$ the inequality

$$p_1 < p_0^{-1} \exp(-0.5 |b_j| \delta_j T_c) \quad (3.17)$$

is fulfilled with the probability $\Pr_v$ such that

$$\Pr_v > (1 - \rho(p_0) - \rho(p_1))^{T_c}, \quad (3.18)$$

where for large N_{popmax} and $p_{\text{mut}} \rightarrow 0$

$$\rho(p) = \exp\left(-(\ln 2 - 1/2) p_{\text{mut}} c_F p k N_{\text{popmax}}\right).$$

Interpretation of Theorem 3.2

It is interesting to compare Theorem 3.1 and Theorem 3.2. The previous one asserts that for infinite populations the probability of the event $j \in I_-$ is zero whereas the second one claims that this probability becomes exponentially small as the population size increases.

This theorem also shows that evolution can make a statistical test checking the hypothesis H_- that $j \in I_-$ against the hypothesis H_+

that $j \in I_+$. Suppose that H_- is true. Let V be the event that the frequency $X(\bar{s}, t)$ of the genotype $\bar{s}$ in the population is larger than p_1 within a sufficiently large time T_c . According to estimate (3.18), the probability of the event V is so small that it is almost unbelievable. Therefore, the hypothesis H_- should be rejected. We will refer T_c as the *checking time*.

Ideas for the proof. The main idea is the same as that for the previous theorem: we compare the frequencies of the organisms with the genotype $\bar{s}$ and the organisms with the genotype s . However, the proof includes a number of technical details connected with estimates of mutation effects and fluctuations. The formal proof can be found in Section 4. It is based on estimates of the accuracy of the Nagylaki equations (in the case of asexual reproduction).

4 Proof of theorems

Let us prove Theorem 3.2.

4.1 Main tools and auxiliary Lemmas

Let us introduce notation and make some preliminary remarks. Remind that we denote by $N(s, t)$ the number of the population members with the genotype s at the moment t . Let $X(t)$ be the set of all population members at the moment t . For each $x \in X(t)$ let us denote by $N'(x, t)$ the number of progeny born by the individual x at the moment t before the massacre (see point 3 of model from Subsection 2.4). Let $s_g(x)$ be the genotype of x . Then, according to (2.8), the mean of $N'(x, t)$ is

$$EN'(x, t) = F(s_g(x)), \quad (4.1)$$

where EX denotes the expected value of X . By $\bar{N}(s, t)$ we denote the number of all progeny born by individuals with the genotype s at the moment t before the massacre. Since all progeny are produced independently and randomly, the previous relation gives

$$E\bar{N}(s, t) = N(s, t)F(s). \quad (4.2)$$

Our main analytical tools are the Chernoff bounds and the Hoeffding inequalities. We also use the Markov inequality: for a positive random quantity X and $a > 0$ one has

$$\Pr\{X > a\} \leq \frac{EX}{a}. \quad (4.3)$$

Moreover, we use two elementary estimates. Let $\mathcal{A}$ be an event in stochastic population dynamics. We denote by $\text{Not } \mathcal{A}$ the negation (complement) of $\mathcal{A}$ and by $\Pr(\mathcal{A}|\mathcal{B})$ the conditional probability of $\mathcal{A}$ under the condition $\mathcal{B}$. For events $\mathcal{A}, \mathcal{B}_1, \dots, \mathcal{B}_n$ we have

$$\begin{aligned} \Pr(\mathcal{A}) &= \Pr(\mathcal{A}|\mathcal{B}_1 \dots \mathcal{B}_n) + \\ &\quad \Pr(\mathcal{A}|\text{Not}(\mathcal{B}_1 \dots \mathcal{B}_n)) \\ &\leq \Pr(\mathcal{A}|\mathcal{B}_1 \dots \mathcal{B}_n) + \sum_{j=1}^n \Pr(\text{Not } \mathcal{B}_j) \end{aligned} \quad (4.4)$$

For two events $\mathcal{A}, \mathcal{B}$ one has

$$\Pr(\mathcal{A}|\mathcal{B}) \geq 1 - \Pr(\text{Not } \mathcal{A}) - \Pr(\text{Not } \mathcal{B}). \quad (4.5)$$

Lemma 4.1. Let X_i be independent random quantities, where $i = 1, \dots, n$. Let each X_i be distributed according to the Poisson law with the average $EX_i = \mu_i$. Let us denote

$$X = \sum_{j=1}^n X_j, \quad \bar{\mu} = \frac{\sum_{j=1}^n \mu_j}{n}.$$

Then for all $\delta > 0$

$$\Pr\{X > (1 + \delta)\bar{\mu}n\} \leq \exp(-\bar{\mu}d(\delta)n), \quad (4.6)$$

where

$$d(\delta) = (1 + \delta)\ln(1 + \delta) - \delta.$$

Similarly,

$$\Pr\{X < (1 - \delta)\bar{\mu}n\} \leq \exp(-\bar{\mu}d(-\delta)n). \quad (4.7)$$

Proof. Note that for any $\lambda > 0$

$$\begin{aligned} \Pr\{X > (1 + \delta)\bar{\mu}n\} &= \\ \Pr\{\exp(\lambda X) > \exp(\lambda(1 + \delta)\bar{\mu}n)\} \end{aligned} \quad (4.8)$$

Since X_j are independent quantities, we have

$$E \exp(\lambda X) = \prod_{j=1}^n E \exp(\lambda X_j).$$

The straight forward computation shows that

$$E \exp(\lambda X_j) = \exp((e^\lambda - 1)\mu_j).$$

Therefore, due to the Markov inequality (4.3) and estimate (4.8) one has

$$\Pr\{X > (1 + \delta)\bar{\mu}n\} \leq \exp(n\bar{\mu}f(\lambda)),$$

where

$$f(\lambda) = \exp(\lambda) - 1 - \lambda(1 + \delta).$$

We minimize f with respect to λ and obtain (4.6). To derive (4.7), we use

$$\begin{aligned} \Pr\{X < (1 - \delta)\bar{\mu}n\} &= \\ \Pr\{\exp(-\lambda X) > \\ \exp(-\lambda(1 - \delta)\bar{\mu}n)\} \end{aligned} \quad (4.9)$$

and repeat the same arguments. The Lemma is proved.

Lemma 4.2. Let X_i be independent random quantities, where $i = 1, \dots, n$ such that $X_i \in \{0, 1\}$ and $EX_i = p$. Then

$$\Pr\{2X < pn\} \leq \exp(-g(p)n), \quad (4.10)$$

where

$$g(p) = -\frac{p \ln 2}{2} - \ln(1 - \frac{p}{2}). \quad (4.11)$$

Proof. Note that for any $\lambda > 0$

$$\Pr\{X < pn/2\} = \Pr\{\exp(-\lambda X) > \exp(-\lambda pn/2)\}. \quad (4.12)$$

Since X_j are independent quantities, we have

$$E \exp(-\lambda X) = \prod_{j=1}^n E \exp(-\lambda X_j).$$

Note that $E \exp(-\lambda X_j) = p \exp(-\lambda) + 1 - p$. Let

$$G(\lambda, p) = \lambda p/2 + \ln(p \exp(-\lambda) + 1 - p).$$

We take $\lambda = \ln 2$ and find that $G(\ln 2, p) = -g(p)$. Now by using the Markov inequality (4.3) and estimate (4.12) one obtains (4.10). The Lemma is proved.

We also use the following Chernoff-Hoeffding theorem. Let X_i be i.i.d. quantities such that $X_i \in \{0, 1\}$ and $EX_i = p$, where $i = 1, \dots, n$. Then for $X = \sum_{j=1}^n X_j$ one has

$$\Pr\{X > (p + \varepsilon)n\} \leq \exp(-D(p + \varepsilon \| p)n), \quad (4.13)$$

where $D(x \| y)$ is the Kullback-Leibler divergence

$$D(x \| y) = x \ln(x/y) + (1-x) \ln((1-x)/(1-y)). \quad (4.14)$$

Moreover, we will use the Hoeffding Theorem: if i.i.d. quantities $X_i \in [0, 1]$ with the probability 1 then

$$\Pr\{|X - EX| > a\} \leq 2 \exp(-2a^2/n). \quad (4.15)$$

4.2 Main lemmas

First we estimate the population size fluctuations.

Lemma 4.3. Let $\bar{N}(t)$ be the number of all progeny, born in the population at the moment t before the massacre, and $\varepsilon_1 > 0$ be a small number. Then

$$\bar{N}(t) \in J_{\varepsilon_1}(t) = \left[(1 - \varepsilon_1) \bar{F}(t) N_{\text{pop}}(t), (1 + \varepsilon_1) \bar{F}(t) N_{\text{pop}}(t) \right] \quad (4.16)$$

with probability

$$\Pr_{\bar{N}} > 1 - \eta_0(\varepsilon_1), \quad (4.17)$$

where

$$\eta_0(\varepsilon_1) = \exp(-d(\varepsilon_1) c_F N_{\text{pop}}(t)) + \exp(-d(-\varepsilon_1) c_F N_{\text{pop}}(t)). \quad (4.18)$$

Proof. Let $n'(x, t)$ denote the number of progeny produced by the individual x before the massacre at the t -th evolution step. The number $\bar{N}(t)$ is the sum

$$\bar{N}(t) = \sum_{x \in X(t)} n'(x, t)$$

of the mutually independent random quantities. According to (4.2), the average $EN'(x, t)$ is $F(s_g(x))$. Therefore,

$$E\bar{N}(t) = \sum_{x \in X(t)} EN'(x, t) \quad (4.19)$$

$$= \sum_{x \in X(t)} F(s_g(x)) \quad (4.20)$$

$$= N_{\text{pop}}(t) \bar{F}(t). \quad (4.21)$$

We set

$$n = N_{\text{pop}}(t), \quad \mu_x = F(s_g(x)), \quad \bar{\mu} = \bar{F}(t)$$

and use the Lemma 4.1 that gives us (4.17).

Lemma 4.4. Let $\varepsilon_2 \in (0, 1)$ be fixed and condition (3.13) be fulfilled. Assume, moreover, that

$$2N_{\text{popmax}} \geq N_{\text{pop}}(t) \geq \kappa N_{\text{popmax}}, \quad (4.22)$$

where

$$\kappa \in (C_F^{-1}, 1) \quad (4.23)$$

and $c_F > 1$ is defined by (3.13). Let us define the event $\mathcal{D}_{\varepsilon_2}(t)$ by

$$\mathcal{D}_{\varepsilon_2}(t) = \left\{ |N_{\text{pop}}(t+1) - N_{\text{popmax}}| < \varepsilon_2 N_{\text{popmax}} \right\} \quad (4.24)$$

Then one has

$$\Pr(\mathcal{D}_{\varepsilon_2}(t)) > 1 - \eta(\varepsilon_2), \quad (4.25)$$

where

$$\begin{aligned} \eta(\varepsilon_2) = & \exp(-d(\tilde{\varepsilon}) \kappa N_{\text{popmax}}) + \\ & \exp(-d(-\tilde{\varepsilon}) \kappa N_{\text{popmax}}) + \\ & 2 \exp\left(-\frac{2\varepsilon_2^2 N_{\text{popmax}}}{2(1+\tilde{\varepsilon})C_F}\right) \end{aligned} \quad (4.26)$$

and

$$\tilde{\varepsilon} = 1 - (\kappa c_F)^{-1}. \quad (4.27)$$

Proof. Let $\xi(x)$ be random quantities defined as follows: $\xi(x) = 1$ if the individual x is survived as a result of massacre (see point 3 of our model from Subsection 2.4), and $\xi(x) = 0$ otherwise. Let $X'(t)$ be the set of progeny produced by all individuals from the population. Then the number $N_{\text{sur}}(t) = N_{\text{pop}}(t+1)$ of finally survived progeny can be computed as follows:

$$N_{\text{sur}}(t) = \sum_{x \in X'(t)} \xi(x).$$

Note that $|X'(t)| = \bar{N}(t)$. Moreover, $E\xi(x) = N_{\text{popmax}} / \bar{N}(t)$ for $\bar{N}(t) \geq N_{\text{popmax}}$. Therefore, if $\bar{N}(t) \geq N_{\text{popmax}}$ then

$$EN_{\text{sur}}(t) = N_{\text{popmax}} \cdot \quad (4.28)$$

Let us define the event

$$\mathcal{B}(t) = \{\bar{N}(t) \in J_{\tilde{\varepsilon}}(t)\}, \quad (4.29)$$

where the interval $J_{\tilde{\varepsilon}}(t)$ is defined by (4.16) and $\tilde{\varepsilon}$ is defined by (4.27). By (4.4) we have

$$\Pr(\text{Not } \mathcal{D}_{\varepsilon_2}(t)) \leq \Pr(\text{Not } \mathcal{D}_{\varepsilon_2}(t) | \mathcal{B}(t)) + \Pr(\text{Not } \mathcal{B}(t)). \quad (4.30)$$

Now we apply the Hoeffding inequality (4.15). For each $\varepsilon_2 > 0$ we obtain

$$\Pr(\text{Not } \mathcal{D}_{\varepsilon_2}(t)) < 2 \exp\left(-\frac{2\varepsilon_2^2 EN_{\text{sur}}(t)^2}{\bar{N}(t)}\right).$$

If $\mathcal{B}(t)$ takes place, then $\bar{N}(t) \geq N_{\text{popmax}}$ and consequently

$$\frac{2\varepsilon_2^2 EN_{\text{sur}}(t)^2}{\bar{N}(t)} > \frac{2\varepsilon_2^2 N_{\text{popmax}}}{2(1+\tilde{\varepsilon})C_F}. \quad (4.31)$$

Therefore,

$$\Pr(\text{Not } \mathcal{D}_{\varepsilon_2}(t) | \mathcal{B}(t)) < 2 \exp\left(-\frac{2\varepsilon_2^2 N_{\text{popmax}}}{2(1+\tilde{\varepsilon})C_F}\right). \quad (4.32)$$

Moreover, by Lemma 4.3

$$\Pr(\text{Not } \mathcal{B}(t)) < \exp\left(-d(\tilde{\varepsilon})\kappa N_{\text{popmax}}\right) + \exp\left(-d(-\tilde{\varepsilon})\kappa N_{\text{popmax}}\right). \quad (4.33)$$

Inequalities (4.30), (4.32) and (4.33) prove (4.25).

The following lemma, in particular, allows us to obtain equations (2.9) and (2.10) in the limit of infinite populations and for small mutation probabilities.

Recall that $\bar{N}(s, t)$ denotes the number of non-mutated progeny generated by the individuals with the genotype s before the massacre. Let $N_{\text{sur}}(s, t)$ be the number of those progeny that survived after that massacre.

Lemma 4.5. Let ε_0 be a positive number satisfying (4.75) and

$$\kappa N_{\text{popmax}} < N_{\text{pop}}(t) < 2N_{\text{popmax}}. \quad (4.34)$$

Then one has

$$N(s, t+1) > (1-\varepsilon_0)F(s)\bar{F}(t)^{-1}N(s, t) \quad (4.35)$$

with the probability $\Pr_{s,t,+}$ such that

$$\Pr_{s,t,+} > 1 - \sum_{i=1}^5 R_i(s, t), \quad (4.36)$$

where

$$R_1(s, t) = \exp\left(-d(1)c_F N_{\text{pop}}(t)\right) + \exp\left(-d(-1)c_F N_{\text{pop}}(t)\right), \quad (4.37)$$

$$R_2(s, t) = \exp(-d(0.5)c_F N(s, t)) + \exp(-d(-0.5)c_F N(s, t)), \quad (4.38)$$

$$R_3(s, t) = 2 \exp\left(\frac{-\varepsilon_0^2}{16C_F^2} c_F N(s, t)\right), \quad (4.39)$$

$$R_4(s, t) = \exp\left(-0.5(2 \ln 2 p_{\text{mut}} + (1-p_{\text{mut}}) \ln\left(\frac{1-2p_{\text{mut}}}{1-p_{\text{mut}}}\right)) \cdot c_F N(s, t)\right), \quad (4.40)$$

$$R_5(s, t) = \exp\left(-d(1)c_F N(s, t)\right) + \exp\left(-d(-1)c_F N(s, t)\right), \quad (4.41)$$

Similarly,

$$N(s, t+1) < (1+\varepsilon_0)F(s)\bar{F}(t)^{-1}N(s, t) \quad (4.42)$$

with the probability $\Pr_{s,t,-}$ such that

$$\Pr_{s,t,-} > 1 - \sum_{i=1}^5 R_i(s, t). \quad (4.43)$$

Proof. Step 1, estimates of fluctuations. First let us estimate the fluctuations of the number $\bar{N}(s, t)$. For each $\varepsilon_2 > 0$ let us define the event

$$\mathcal{A}_{s,\varepsilon_2}(t) = \{|\bar{N}(s, t) - E\bar{N}(s, t)| > \varepsilon_2 E\bar{N}(s, t)\}. \quad (4.44)$$

By Lemma 4.1 one has

$$\Pr(\mathcal{A}_{s,\varepsilon_2}(t)) < \exp\left(-d(\varepsilon_2)E\bar{N}(s, t)\right) + \exp\left(-d(-\varepsilon_2)E\bar{N}(s, t)\right) \quad (4.45)$$

Note that

$$E\bar{N}(s, t) = F(s)N(s, t) > c_F N(s, t). \quad (4.46)$$

As a result, by (4.46) we obtain

$$\Pr(\mathcal{A}_{s,\varepsilon_2}(t)) < \exp(-d(\varepsilon_2)c_F N(s, t)) + \exp(-d(-\varepsilon_2)c_F N(s, t)) \quad (4.47)$$

Step 2. Here we estimate the number of progeny that survived as a result of the massacre procedure (point 3 of the population dynamics model, see subsection 2.9). Let $X'(s, t)$ be the set of progeny produced by individuals with the genotype s . Then the number $N_{\text{sur}}(s, t)$ of survived progeny x for individuals x belonging to the set $Z'(s, t)$ is

$$N_{\text{sur}}(s, t) = \sum_{x \in X'(s, t)} \xi(x),$$

where $\xi(x)$ are defined in the proof of the previous Lemma. For $\varepsilon_3 > 0$ we consider the event

$$\mathcal{A}_{\text{sur},s,\varepsilon_3}(t) = \{|N_{\text{sur}}(s, t) - EN_{\text{sur}}(s, t)| > \varepsilon_3 E[N_{\text{sur}}(s, t)]\}, \quad (4.48)$$

Let us estimate the probability $\Pr(\mathcal{A}_{\text{sur},s}(t))$. According to the Hoeffding Theorem (4.15)

$$\Pr(\mathcal{A}_{\text{sur},s,\varepsilon_3}(t)) < 2 \exp\left(-2\varepsilon_3^2 E[N_{\text{sur}}(s, t)]^2 \bar{N}(s, t)^{-1}\right). \quad (4.49)$$

Note that $\xi(x)$ and $\xi(y)$ are independent quantities for different x and y , thus under the condition $\bar{N}(t) > N_{\text{popmax}}$

$$EN_{\text{sur}}(s, t) = \sum_{x \in X'(s, t)} E\xi(x) = \bar{N}(s, t) \frac{N_{\text{popmax}}}{\bar{N}(t)},$$

therefore,

$$\Pr(\mathcal{A}_{\text{sur},s,\varepsilon_3}(t)) < 2 \exp\left(-2\varepsilon_3^2 \bar{N}(s, t) \left(\frac{N_{\text{popmax}}}{\bar{N}(t)}\right)^2\right). \quad (4.50)$$

Let us define the events $\mathcal{B}_s(t)$ and $\mathcal{B}(t)$ by

$$\begin{aligned} \mathcal{B}_s(t) &= \{\bar{N}(s, t) > (1 - \varepsilon_2)E\bar{N}(s, t)\}, \end{aligned} \quad (4.51)$$

$$\begin{aligned} \mathcal{B}(t) &= \{\bar{N}(t) < (1 + \varepsilon_1)E\bar{N}(t)\}. \end{aligned} \quad (4.52)$$

Then using (4.4) one has

$$\begin{aligned} \Pr(\mathcal{A}_{\text{sur},s,\varepsilon_3}(t)) &\leq \Pr(\mathcal{A}_{\text{sur},s}(t) | \mathcal{B}_s(t) \mathcal{B}(t)) + \\ &\quad \Pr(\text{Not } \mathcal{B}_s(t)) + \\ &\quad \Pr(\text{Not } \mathcal{B}(t)). \end{aligned} \quad (4.53)$$

We observe that under conditions $\mathcal{B}_s(t)$ and $\mathcal{B}(t)$

$$\begin{aligned} \bar{N}(s, t) \left(\frac{N_{\text{popmax}}}{\bar{N}(t)}\right)^2 &< (1 - \varepsilon_2)(1 + \varepsilon_1)^{-2} \cdot \\ &\quad E\bar{N}(s, t) \cdot \\ &\quad \left(\frac{N_{\text{popmax}}}{E\bar{N}(t)}\right)^2. \end{aligned} \quad (4.54)$$

In that estimate let us set $\varepsilon_2 = 0.5$ and $\varepsilon_1 = 1$. Taking into account that $E\bar{N}(t) = \bar{F}(t)N_{\text{pop}}(t) < 2C_F N_{\text{popmax}}$, we have that

$$\Pr(\mathcal{A}_{\text{sur},s,\varepsilon_3}(t) | \mathcal{B}_s(t) \mathcal{B}(t)) < \bar{R}_3(\varepsilon_3, s, t), \quad (4.55)$$

where

$$\bar{R}_3(\varepsilon_3, s, t) = 2 \exp\left(-0.25\varepsilon_3^2 C_F^2 c_F N(s, t)\right). \quad (4.56)$$

Moreover, according to (4.47)

$$\Pr(\text{Not } \mathcal{B}_s(t)) < R_2(s, t), \quad (4.57)$$

and due to (4.17)

$$\Pr(\text{Not } \mathcal{B}(t)) < R_1(s, t), \quad (4.58)$$

where R_1, R_2 are defined by (4.37) and (4.38). Finally,

$$\Pr(\mathcal{A}_{\text{sur},s,\varepsilon_3}(t)) < R_1(s, t) + R_2(s, t) + \bar{R}_3(\varepsilon_3, s, t). \quad (4.59)$$

Step 3, estimate of the number of mutants.

Let us estimate how many individuals with genotypes s can mutate. The probability of mutation is p_{mut} . Let $N_{\text{mut}}(s, t)$ be the number of such mutants. Let us define the event $\mathcal{A}_{\text{mut},s}(t)$ by

$$\mathcal{A}_{\text{mut},s}(t) = \{N_{\text{mut}}(s, t) > 2p_{\text{mut}}\bar{N}(s, t)\}, \quad (4.60)$$

Since the random quantity $N_{\text{mut}}(s, t)$ is subject to the Bernoulli law, we can apply the Chernoff-Hoeffding inequality (4.13). Then we obtain that

$$\Pr(\mathcal{A}_{\text{mut},s}(t)) < \exp\left(-D(2p_{\text{mut}} \| p_{\text{mut}}) \bar{N}(s, t)\right), \quad (4.61)$$

where, according to definition (4.14) of $D(x \| y)$, one has

$$D(2p_{\text{mut}} \| p_{\text{mut}}) = g(p_{\text{mut}})$$

and g is defined by (4.11).

Using (4.4) one has

$$\Pr(\mathcal{A}_{\text{mut},s}(t)) \leq \Pr(\mathcal{A}_{\text{mut},s}(t) | \mathcal{B}_s(t)) + \Pr(\text{Not } \mathcal{B}_s(t)). \quad (4.62)$$

As a result, by Lemma 4.3 one finds

$$\Pr(\mathcal{A}_{\text{mut},s}(t)) \leq R_4 + R_5, \quad (4.63)$$

where R_4, R_5 are defined by (4.40) and (4.41).

To prove (4.35), we set $\varepsilon_3 = \varepsilon_0/2$. Taking into account condition (4.75) for ε_0 we see that if the both events $\text{Not } \mathcal{A}_{\text{mut},s}(t)$ and $\mathcal{A}_{\text{sur},s,\varepsilon_0/2}(t)$ take place, then inequality (4.35) is fulfilled. Thus

$$\begin{aligned} \Pr(\text{Not } \mathcal{A}_{\text{mut},s}(t) \text{Not } \mathcal{A}_{\text{sur},s,\varepsilon_0/2}(t)) &\geq \\ 1 - \Pr(\mathcal{A}_{\text{mut},s}(t)) - \Pr(\mathcal{A}_{\text{sur},s,\varepsilon_0/2}(t)) &> \\ 1 - \sum_{i=1}^5 R_i, \end{aligned}$$

where R_i are defined by (4.37)–(4.41).

Finally, taking into account the results of steps 1, 2 and 3 we see that estimate (4.35) holds with the probability $\Pr_{t,+}$. It completes the proof of (4.35). The second inequality (4.42) can be obtained in the same way.

4.3 Remaining part of the proof of Theorem 3.2

We use the same idea that in the proof of Theorem 3.1 but first we establish uniform bounds for the population size and other quantities involved in the proof.

Step 1 Here we estimate the population size. Let us set

$$\varepsilon_2 = 1 - \kappa > 0$$

in Lemma 4.4. Let us consider the events $\mathcal{D}_{\varepsilon_2}(t)$ defined by (4.24) in Lemma 4.4. If the events $\mathcal{D}_{\varepsilon_2}(t)$ take place for all $t \in [T_1, T_1 + T_c]$ and $N_{\text{pop}}(0) = N_{\text{popmax}}$, we have that

$$2N_{\text{popmax}} > N_{\text{pop}}(t) \quad (4.64)$$

$$> \kappa N_{\text{popmax}} \quad \forall t \in [T_1, T_1 + T_c] \quad (4.65)$$

Then conditions (3.15), (3.16) of Theorem 3.2 imply

$$N(s, t) > \kappa p_0 N_{\text{popmax}}, \quad N(\bar{s}, t) > \kappa p_1 N_{\text{popmax}}. \quad (4.66)$$

Those inequalities imply the following estimates for the quantities R_i defined by (4.37)–(4.41):

$$R_i(s, t) > q_i(p_0), \quad R_i(\bar{s}, t) > q_i(p_1), \quad (4.67)$$

where q_i are defined by

$$q_1 = \exp\left(- (2 \ln 2 - 1) c_F \kappa N_{\text{popmax}}\right) + \exp\left(- c_F \kappa N_{\text{popmax}}\right), \quad (4.68)$$

$$q_2(p) = 2 \exp\left(- (3/2 \ln(3/2) - 1/2) c_F p \kappa N_{\text{popmax}}\right) + \exp\left(- 1/2 (1 - \ln 2) c_F p \kappa N_{\text{popmax}}\right), \quad (4.69)$$

$$q_3(p) = 2 \exp\left(- \frac{\varepsilon_0^2}{16 C_F^2} \kappa c_F p N_{\text{popmax}}\right), \quad (4.70)$$

$$q_4(p) = \exp\left(- 0.5 U(p_{\text{mut}}) c_F p \kappa N_{\text{popmax}}\right), \quad (4.71)$$

$$q_5(p) = \exp\left(- (2 \ln 2 - 1) c_F \kappa p N_{\text{popmax}}\right) + \exp\left(- c_F p \kappa N_{\text{popmax}}\right), \quad (4.72)$$

where

$$U(p) = 2 \ln 2 p + (1 - p) \ln((1 - 2p)/(1 - p)), \quad (4.73)$$

and

$$\begin{aligned} \tilde{q} = & \exp\left(- d(\tilde{\varepsilon}) \kappa N_{\text{popmax}}\right) + \\ & \exp\left(- d(-\tilde{\varepsilon}) \kappa N_{\text{popmax}}\right) + \\ & 2 \exp\left(- \frac{2(1 - \kappa)^2 N_{\text{popmax}}}{2(1 + \tilde{\varepsilon}) C_F}\right). \end{aligned} \quad (4.74)$$

where $\tilde{\varepsilon} = 1 - (\kappa c_F)^{-1}$, and

$$\begin{aligned} \varepsilon_0 = & \frac{1 - \exp(-b_j \delta_j T_c / 2)}{1 + \exp(-b_j \delta_j T_c / 2)} \\ & > 4 p_{\text{mut}} C_F. \end{aligned} \quad (4.75)$$

For each $p \in (0, 1)$ let us define an auxiliary function

$$\rho(p) = q_1 + q_2(p) + q_3(p) + q_4(p) + q_5(p) + \tilde{q}, \quad (4.76)$$

where $q_i, \tilde{q}$ are defined by relations (4.68)–(4.72). We can find asymptotics of ρ under natural assumptions that $p_{\text{mut}} \rightarrow 0$ and $N_{\text{popmax}} \rightarrow \infty$ while all the rest parameters are fixed. Then the leading term in the right hand side of (4.36) is q_4 and $U(p_{\text{mut}}) = (2 \ln 2 - 1) p_{\text{mut}} + O(p_{\text{mut}}^2)$. As a result, we have

$$\begin{aligned} \rho(p) = & \exp\left(- (\ln 2 - 1/2) p_{\text{mut}} c_F p \kappa N_{\text{popmax}}\right) \\ & (1 + o(1)), \quad p_{\text{mut}} \rightarrow 0. \end{aligned} \quad (4.77)$$

Step 2. Let $Q(t)$ is defined by (3.8) and, moreover, let $j \in \underline{I}$. We use Lemma 4.5 inductively for genotypes s and $\bar{s}$. Let us set

$$\theta = \frac{1 - \varepsilon_0}{1 + \varepsilon_0},$$

where ε is defined by (4.75). We remark that the inequality

$$Q(T_c + t + 1) \geq Q(T_c + t) \theta \exp(b_j \delta_j T_c) \quad (4.78)$$

holds with a probability $\Pr_{Q,t} > 0$. Let us obtain a uniform estimate of that probability. Let $\mathcal{E}(t)$ be the event that (4.78) holds at the step t . Using (4.4) we have

$$\begin{aligned} \Pr(\text{Not } \mathcal{E}(t)) \leq & \Pr(\text{Not } \mathcal{E}(t) | \mathcal{D}_{\varepsilon_2}(t)) + \\ & \Pr(\text{Not } \mathcal{D}_{\varepsilon_2}(t)), \end{aligned} \quad (4.79)$$

where, according to Lemma 4.4, the probability of the event $\text{Not } \mathcal{D}_{\varepsilon_2}(t)$ is less than η , where η is defined by (4.25), and

$$\Pr(\text{Not } \mathcal{E}(t) | \mathcal{D}_{\varepsilon_2}(t)) < \tilde{q} + \rho(p_0) + \rho(p_1).$$

We conclude by (4.5) that

$$\Pr_{Q,t} > Z, \quad Z = 1 - \tilde{q} - \rho(p_0) - \rho(p_1), \quad (4.80)$$

where $\tilde{q}$ is defined by (4.74). This estimate is uniform in $t \in [1, \dots, T_c]$. By (4.80) we obtain then that the inequality

$$Q(T_c + T_1) \geq Q(T_1) \bar{\theta}, \quad \bar{\theta} = \frac{1 - \varepsilon_0}{1 + \varepsilon_0} \exp(b_j \delta_j T_c) \quad (4.81)$$

is satisfied with the probability $\Pr_v$ such that

$$\Pr_v > Z^{T_c}. \quad (4.82)$$

For ε_0 defined by (4.75), one has

$$Q(T_c + T_1) \geq Q(T_1) \exp(b_j \delta_j T_c / 2).$$

Now repeating the same arguments that in the end of the proof of Theorem 3.1, and taking into account asymptotics (4.77), we obtain the conclusion of Theorem 3.2.

5 Discussion

In this paper, we proposed a model for fitness landscape learning, which extends earlier work by 7–9 in two ways. First, we use hybrid circuits involving two kinds of variables. The first class of variables are real valued in the interval $(0, 1)$ and can be interpreted as relative levels of phenotypic traits, other variables are Boolean and can be interpreted as genes. Second, we use a threshold scheme of regulation, which is inspired by ideas of the paper by 20. All variables are involved in gene regulation via thresholds.

The work presented here is a major extension of a long term effort to explicitly model the effects of phenotypic buffering in evolution by considering a class of Boolean and mixed Boolean-continuous models in which the phenotype is represented explicitly and the degree of phenotypic buffering can be controlled in various ways. For example, we have demonstrated that the idea of an “evolutionary capacitor”^{38,39} can be implemented by explicit control of phenotypic buffering in a hub-and-spokes architecture²³ and that in a more general class of genetic architecture numerical simulations show that an intermediate level of buffering is optimal for evolution in a changing environment⁴⁰.

The results reported here are very promising, since they are consistent with the results of recent experiments by 41 and 42 on heat shock stress. The essential mechanism is that the exploration of the fitness landscape by the genetic network in such a way that future mutations are more likely to be adaptive. We have shown that, at least for some fitness landscapes, rapid evolutionary changes—perhaps instances of the “hopeful monsters” of Goldschmidt⁴³—can be created by a combination of random small mutations and epigenetic effects. The main idea is that small mutations pave the way for large epigenetic or genetic changes. The hypothetical mechanism, which we propose, can be outlined as follows (see Figure 2, Figure 3).

The expression of genes involved in the expression of phenotypic traits depends on threshold parameters h_j , which take three values: a large negative one, a neutral value close to zero and a large positive one. First the threshold parameter h_j is small and thus the phenotypic trait is sensitive with respect to even small mutations. Those mutations play a fundamental role working as scouts exploring environments (see Figure 3). If a mutation occurred and the corresponding mutant has survived within $T_c \gg 1$ generations then according to Theorem 3.1 and Theorem 3.2 these events mean that that mutation increases the fitness that allows the network to estimate the correct direction of evolution. Then gene regulation detects that increase to change the threshold according to simple rules. Namely, if the trait is less expressed in that mutant with respect to wild type parent, the gene regulation system decreases the threshold up to the large negative value. On the contrary, if the trait is strongly expressed in the mutant, the gene regulation system increases the threshold up to the large positive value. This simple

Figure 2. This graph illustrates Goldschmidt's leaps. At initial moment the trait expressions take the values $x = 0.5$, $y = 0.5$. According to Fisher's ideas, random large mutations decrease the fitness $F = K_F \exp(W)$. (Changes of $x = F_1$, $y = f_2$, which are induced by mutations, are shown by red vectors.) Thus such mutations produce non-viable organisms.

Figure 3. This plot illustrates the main ideas of evolution based on the fitness landscape learning. At the initial time the trait expressions take the values $x = 0.5$, $y = 0.5$. Evolutionary changes go in two stages. First we make small random mutations (shown by red vectors), which explore the fitness landscape. If such a mutation is not eliminated from the population, this means that a correct evolution direction is found, and gene regulation system makes a big leap (shown by the green vector) in the direction of that small mutation. Such a two step model can be called clever Goldschmidt leaps. Note that evolution is gradual, and the existence of clusters of almost identical genes involved in the same QTL increases the chances to create a clever Goldschmidt hopeful monster.

regulation control not only sharply reduces the number of mutations needed for adaptation, but also canalizes the phenotype since for large thresholds the trait expression level becomes insensitive with respect to mutations. We suppose that these threshold modifications can be inherited.

So, we propose the mechanism: small mutations serve as scouts finding the way for large epigenetic or genetic changes, which can be performed by gene regulatory system.

The mechanism may also explain the results of 4 on prediction of environmental changes. In fact, let us suppose that environment varies in time. The first, perhaps relatively small, variations can trigger the threshold mechanism described above. As a result, the population will be adapted to the subsequent changes in advance.

Our results show that evolution can proceed rapidly because it reduces the number of mutations required for adaptive change.

The primary limitation of our results is that the representation of the evolving genetic network is limited to the network of gene controlling phenotype, represented here by the Boolean strings s . Other model variables represent the coarse-grained activities of genes. One class is the terminal differentiation genes represented by w_{ij} , and another are the genes or epigenetic factors controlling the thresholds h and their associated learning rules. A more careful consideration of the relationship

of these moieties to observable molecular entities is an important objective of future work. At the mathematical level, the key analytical results were obtained in a simplified context that falls short of a realistic level of pleiotropy and thus of the level of NP-hard complexity exhibited by fully pleiotropic forms of our model. We believe that our analytical results can be generalized, which we plan to address in future work.

Data availability

All data underlying the results are available as part of the article and no additional source data are required.

Grant information

The second author was supported by the grant of Russian Ministry of Education, 2012-1.2.1-12-000-1013-016. Additionally, the second author was financially supported by Government of Russian Federation, Grant Grant 08-08.

D. Grigoriev is grateful to the grant RSF 16-11-10075 and to both MCCME and MPI für Mathematik for wonderful working conditions and inspiring atmosphere.

J. Reinitz and S. Vakulenko were supported by US NIH grant RO1 OD010936 (formerly RO1 RR07801).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Orr HA: **The genetic theory of adaptation: a brief history.** *Nat Rev Genet.* 2005; **6**(2): 119–127.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Orr HA, Coyne JA: **The genetics of adaptation: a reassessment.** *Am Nat.* 1992; **140**(5): 725–742.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Zeyl C: **The number of mutations selected during adaptation in a laboratory population of *Saccharomyces cerevisiae*.** *Genetics.* 2005; **169**(4): 1825–1831.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Mitchell A, Romano GH, Groisman B, *et al.*: **Adaptive prediction of environmental changes by microorganisms.** *Nature.* 2009; **460**(7252): 220–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Watson RA, Szathmáry E: **How Can Evolution Learn?** *Trends Ecol Evol.* 2016; **31**(2): 147–157.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Chastain E, Livnat A, Papadimitriou C, *et al.*: **Algorithms, games, and evolution.** *Proc Natl Acad Sci U S A.* 2014; **111**(29): 10620–10623.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. Parter M, Kashtan N, Alon U: **Facilitated variation: how evolution learns from past environments to generalize to new environments.** *PLoS Comput Biol.* 2008; **4**(11): e1000206.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. Valiant LG: **Evolvability.** *Journal of the ACM.* 2009; **56**(1): 1–21.
[Publisher Full Text](#)
9. Livnat A, Papadimitriou C, Rubinfeld A, *et al.*: **Satisfiability and evolution.** In *Annu IEEE Symp Found Comput Sci FOCS.* 2014; 524–530.
[Publisher Full Text](#)
10. Nagylaki T: **The evolution of multilocus systems under weak selection.** *Genetics.* 1993; **134**(2): 627–647.
[PubMed Abstract](#) | [Free Full Text](#)
11. Arora S, Hazan E, Kale S: **The Multiplicative Weights Update Method: A Meta-Algorithm and Applications.** *Theory Comput.* 2012; **8**: 121–164.
[Publisher Full Text](#)
12. Paixão T, Badkobeh G, Barton N, *et al.*: **Toward a unifying framework for evolutionary processes.** *J Theor Biol.* 2015; **383**: 28–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Mühlenbein H, Schlierkamp-Voosen D: **Predictive Models for the Breeder Genetic Algorithm I. Continuous Parameter Optimization.** *Evol Comput.* 1993; **1**(1): 25–49.
[Publisher Full Text](#)
14. Chatterjee K, Pavlogiannis A, Adlam B, *et al.*: **The time scale of evolutionary innovation.** *PLoS Comput Biol.* 2014; **10**(9): e1003818.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Heredia JP, Trubenová B, Sudholt D, *et al.*: **Selection Limits to Adaptive Walks on Correlated Landscapes.** *Genetics.* 2017; **205**(2): 803–825.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Doerr B, Doerr C, Kötzing T: **Solving Problems with Unknown Solution Length at (Almost) No Extra Cost.** In *Proceedings of the 2015 on Genetic and Evolutionary Computation Conference GECCO - '15.* Madrid, Spain, ACM Press, 2015; 831–838.
[Publisher Full Text](#)
17. Rudolph G: **Finite Markov Chain Results in Evolutionary Computation: A Tour d'Horizon.** *Fundamenta Informaticae.* 1998; **35**(1–4): 67–89.
[Publisher Full Text](#)
18. Witcopp PJ, Carroll SB, Kopp A: **Evolution in black and white: genetic control of pigment patterns in *Drosophila*.** *Trends Genet.* 2003; **19**(9): 495–504.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Wang Z, Liao BY, Zhang J: **Genomic patterns of pleiotropy and the evolution of complexity.** *Proc Natl Acad Sci U S A.* 2010; **107**(42): 18034–18039.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Stern C: **Selection for subthreshold differences and the origin of pseudoexogenous adaptations.** *Am Nat.* 1958; **92**(866): 313–316.
[Publisher Full Text](#)
21. Mjolsness E, Sharp DH, Reinitz J: **A connectionist model of development.** *J Theor Biol.* 1991; **152**(4): 429–453.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Arias M, le Poul Y, Chouteau M, *et al.*: **Crossing fitness valleys: empirical estimation of a fitness landscape associated with polymorphic mimicry.** *Proc Biol Sci.* 2016; **283**(1829): pii: 20160391.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Grigoriev D, Reinitz J, Vakulenko S, *et al.*: **Punctuated evolution and robustness in morphogenesis.** *Biosystems.* 2014; **123**: 106–113.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Franke J, Klözer A, de Visser JA, *et al.*: **Evolutionary accessibility of mutational pathways.** *PLoS Comput Biol.* 2011; **7**(8): e1002134.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Kingman JFC: **A simple model for the balance between selection and mutation.** *J Appl Probab.* 1978; **15**(1): 1–12.
[Publisher Full Text](#)
26. Kauffman SA, Weinberger ED: **The NK model of rugged fitness landscapes and its application to maturation of the immune response.** *J Theor Biol.* 1989; **141**(2): 211–245.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Alta T, Iwakura M, Hsui Y: **A cross-section of the fitness landscape of dihydrofolate reductase.** *Protein Eng.* 2001; **14**(9): 633–638.
[Publisher Full Text](#)
28. Leimar O, Tullberg B, James M: **Mimicry, saltational evolution, and the crossing of fitness valleys.** Chapter 16 in: E.I. Svensson and R. Calsbeek, eds. *The Adaptive Landscape in Evolutionary Biology*, 2012; 257–270.
[Publisher Full Text](#)
29. Savageau MA: **Demand theory of gene regulation. II. Quantitative application to the lactose and maltose operons of *Escherichia coli*.** *Genetics.* 1998; **149**(4): 1677–1691.
[PubMed Abstract](#) | [Free Full Text](#)
30. Cook SA: **The complexity of theorem-proving procedures.** In *Proc third Annu ACM Symp Theory Comput. – STOC '71.* New York, USA, ACM Press. 1971; 151–158.
[Publisher Full Text](#)
31. Levin LA: **Universal enumeration problems (Russian).** *Probl Peredai Inf.* 1973; **9**(3): 115–116.
[Reference Source](#)
32. Friedgut E: **Sharp thresholds of graph properties, and the *k*-sat problem.** *J Am Math Soc.* 1999; **12**(04): 1017–1055.
[Publisher Full Text](#)
33. Moore C, Mertens S: **The Nature of Computation.** Oxford University Press, 2011.
[Publisher Full Text](#)
34. Mertens S, Mézard M, Zecchina R: **Threshold values of random *k*-sat from the cavity method.** *Random Struct Algor.* 2006; **28**(3): 340–373.
[Publisher Full Text](#)
35. Achlioptas D, Coja-Oghlan A: **Algorithmic barriers from phase transitions.** In *Proceedings 49th FOCS.* 2008; 793–802.
[Publisher Full Text](#)
36. Gravner J, Pitman D, Gavrilits S: **Percolation on fitness landscapes: effects of correlation, phenotype, and incompatibilities.** *J Theor Biol.* 2007; **248**(4): 627–645.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Wagner GP, Zhang J: **The pleiotropic structure of the genotype-phenotype map: the evolvability of complex organisms.** *Nat Rev Genet.* 2011; **12**(3): 204–213.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Rutherford SL, Lindquist S: **Hsp90 as a capacitor for morphological evolution.** *Nature.* 1998; **396**(6709): 336–342.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Masel J, Siegal ML: **Robustness: mechanisms and consequences.** *Trends Genet.* 2009; **25**(9): 395–403.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Jiang P, Kreitman M, Reinitz J: **The relationship between robustness and evolution.** *bioRxiv.* 2018; 268862.
[Publisher Full Text](#)
41. Fanti L, Piacentini L, Cappucci U, *et al.*: **Canalization by Selection of *de Novo* Induced Mutations.** *Genetics.* 2017; **206**(4): 1995–2006.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Klosin A, Casas E, Hidalgo-Carcedo C, *et al.*: **Transgenerational transmission of environmental information in *C. elegans*.** *Science.* 2017; **356**(6335): 320–323.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Goldschmidt R: **The Material Basis of Evolution.** Yale Univ. Press, New Haven CT, 1940.
[Reference Source](#)

Open Peer Review

Current Peer Review Status: ? ?

Version 1

Reviewer Report 13 June 2019

<https://doi.org/10.5256/f1000research.20332.r48858>

© 2019 Bergman A et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Yehonatan Sella

Systems & Computational Biology Department, Albert Einstein College of Medicine, New York, NY, USA

Aviv Bergman

Systems & Computational Biology Department, Albert Einstein College of Medicine, New York, NY, USA

In the current work the authors study a population genetics model in which fitness is a linear function of a set of phenotypic traits, and where the genotype-to-phenotype map is given by a linear transformation composed with sigmoidal functions. Despite the seeming simplicity of the fitness function, the authors make the case that optimizing fitness is a hard, NP-complete, problem. Under this model, they study the extent to which the fitness landscape (that is, the question of which phenotypic traits contribute positively to fitness, and which contribute negatively) can be inferred from the distribution of these traits in the population after being subject to evolution for a moderate span of time. They then connect this question with that of whether learning algorithms (potentially epigenetic in nature) can help optimize and speed up evolution by learning the fitness landscape.

Toward the goal of inferring the fitness landscape, the authors prove two theorems. Theorem 3.1 concerns a simplified model of an infinite population with no mutations, while Theorem 3.2 concerns a more complex model of a finite population with mutation, stochastic number of descendants and a culling process. In either case, the result is that under certain assumptions, if a mutant genotype is present in the population with high enough frequency after a long enough period of elapsed time, then we can confidently infer that any phenotypic trait differential between the wildtype and the mutant is associated with a higher fitness.

While Theorem 3.1 ignores mutation, even Theorem 3.2 seems at odds with mutation-selection balance. Even a mutant with lower fitness will be present in the population at a frequency that is on the order of the rate of mutation, while the theorem seems to claim that the frequency of such a mutant will be exponentially small with high probability. The authors should resolve this apparent discrepancy. In addition, the main ideas of the proof would be more clearly

communicated if the authors would include a treatment of the intermediate model of an infinite population with mutation.

As for the discussion of the way learning can speed up the evolutionary process, this part of the paper remains unclear and underdeveloped. The authors discuss a two-step evolutionary process in which the first step consists of small mutations in order to explore the fitness landscape, and the next step involves changing the thresholds involved in the genotype-to-phenotype map in a way that promotes phenotypic traits associated with higher fitness. While this idea is interesting and worth exploring, a few issues arise. A conceptual issue remaining to be addressed is whether the threshold h_i is part of the genotype and what mechanism are needed to alter its value. According to the authors, the h_i 's can be modified genetically or epigenetically. If epigenetically, it is not apparent what are the environmental cues that will lead to such learning let alone the actual mechanism of modifying them. If genetically, it is similarly unclear in what way the learning of the fitness landscape is being stored, if at all, in the genotype, and what is the connection to the Theorems of chapter 3. The theorems in chapter 3 rely on observing the frequency of a genotype in the population, but such information is not stored in individual genotypes. Additionally, if the thresholds are understood to be variable and subject to selection, then the fitness-maximization problem in fact becomes easy (in contrast to the prior analysis of it as NP-complete), unless we impose restrictions on the range of the threshold. Indeed, one can set the thresholds at positive or negative infinity depending on whether the corresponding trait is positive or negative in order to effectively keep the trait on or off regardless of genotype.

Finally, as this manuscript addresses the relation between learning and the rate of evolution, it would benefit from including a reference to one of the most relevant and intuitive articles written in the late 80's by Geoffrey E. Hinton & Steven J. Nowlan, "How Learning Can Guide Evolution" *Complex Systems*, 1, 495-502¹. In it, Hinton and Nowlan showed how learning alters the shape of the fitness landscape and thereby provides easier evolutionary paths towards sets of co-adapted alleles. Hinton and Nowlan demonstrated that this effect allows learning individuals to evolve faster than non-learners. Though the learning model presented by Hinton and Nowlan operates at "somatic" timescale, the analogy to mutations at the evolutionary timescale can be drawn.

To conclude, though it can be improved by the above suggestions, this article touches upon very interesting and important issues in the field of evolutionary biology which are still only lightly investigated, and highlight what might be a fruitful path towards better understanding.

References

1. Hinton GE, Nowlan SJ: How Learning Can Guide Evolution. *Complex Systems*. 1987; **1**.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Not applicable

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Theoretical/Mathematical Systems and Evolutionary Biology.

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however we have significant reservations, as outlined above.

Reviewer Report 23 April 2019

<https://doi.org/10.5256/f1000research.20332.r46602>

© 2019 Szathmáry E. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Eors Szathmáry

¹ Evolutionary Systems Research Group, MTA Centre for Ecological Research, Tihany, Hungary

² Parmenides Center for the Conceptual Foundations of Science, Pullach, Germany

³ Department of Plant Systematics, Ecology and Theoretical Biology, Eötvös University, Budapest, Hungary

This is a new paper in a series of exciting papers about phenotypic evolution, based on models of genes genetic regulatory networks, and phenotypes.

There are many technical details in the paper that most biologists will find difficult to follow – I would put at least some of these items into an Appendix. Also, I am wondering how many (even theoretical) biologists would readily grasp the meaning of the K-SAT problem as defined. A more accessible formulation (in addition to the strictly technical one) could do a lot for easier understanding (one of the papers by the authors contains a didactic figure, for example).

I consider this paper a serious attempt at broadening our views of the role of learning dynamics in evolution, but I think the message could be made a lot clearer. The punch line in a way is in the Discussion that I find both intriguing and baffling.

First, as it is portrayed in this paper, the effect of genes is filtered through a genetic regulatory network. It seems that, just like in Ref 40¹, that this filter itself does not evolve, or at least the dynamics is not given. In other words, how do the weights in the network evolve? Note that in the Watson-Wagner paper in *Evolution*² it was the evolution of the weights that was related to Hebbian dynamics. So, how do we stand on this?

Second, in this paper the genetic control of the thresholds is also kept implicit, but the evolution

of threshold values does play an important role in the Goldschmidtian argument. Is the change in thresholds genetic or epigenetic? If the latter, then it ought to be part of the regulatory network. I guess this is what the statement “gene regulation detects that increase to change the threshold according to simple rules”. How these rules would be implemented in mechanistic terms remains obscure. And how would they arise in evolution? The simple rule implies feedback from expression levels to threshold values.

Third, do I understand it correctly that here, in contrast to Ref 40¹, thresholds can be negative? But a large negative threshold value would mean that expression levels will increase, since negative times negative is positive (Eq. 2.4). If the statement “take three values: a large negative one, a neutral value close to zero and a large positive one” instead refers to gene expression levels rather than threshold values (the latter would in this case be nonnegative) then I do not understand the thought example. Increasing the threshold would reduce the expression level that was selected to go up in the first place!

These points seem to me crucial, so their clarification is badly needed. I would appreciate a few numerical examples along with at least hints to answers to the more conceptual questions on the dynamics.

Also, according to the cited Nagylaki dynamics³ the population evolves almost as if it were in linkage equilibrium – which cannot hold for the asexual population considered by the authors. There are also some minor issues in the paper: in Eq. 2.4 “s” should be replaced by a sigma, and there are also examples where plurals and singulars in the same sentence do not match.

References

1. Jiang P, Kreitman M, Reinitz J: The relationship between robustness and evolution. *bioRxiv*. 2018. [Publisher Full Text](#)
2. Watson RA, Wagner GP, Pavlicev M, Weinreich DM, et al.: The evolution of phenotypic correlations and. *Evolution*. 2014; **68** (4): 1124-38 [PubMed Abstract](#) | [Publisher Full Text](#)
3. Nagylaki T: The evolution of multilocus systems under weak selection. *Genetics*. 1993; **134** (2): 627-47 [PubMed Abstract](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Not applicable

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Theoretical evolutionary biology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research