Check for updates

RESEARCH ARTICLE

Anti-infective potential of hydroalcoholic extract of *Punica granatum* peel against gram-negative bacterial pathogens

[version 1; peer review: 1 approved, 2 approved with reservations]

Chinmayi Joshi , Pooja Patel , Vijay Kothari

Institute of Science, Nirma University, Ahmedabad, Gujarat, 382481, India

V1 First published: 18 Jan 2019, 8:70
<https://doi.org/10.12688/f1000research.17430.1>
 Latest published: 08 Apr 2019, 8:70
<https://doi.org/10.12688/f1000research.17430.2>

Abstract

Background: *Punica granatum* extracts have been prescribed in traditional medicine for management of a variety of disease conditions including microbial infections. Generation of scientific evidence for validation of *P. granatum* peel extract's anti-pathogenic efficacy is required.

Methods: Hydroalcoholic extract of *P. granatum* peel (PGPE), prepared by microwave assisted extraction method was evaluated for its quorum-modulatory potential against two different human-pathogenic bacteria viz. *Chromobacterium violaceum* and *Pseudomonas aeruginosa*.

Results: This extract was able to modulate *in vitro* production of quorum sensing-regulated pigments in both these test bacteria at ≥ 5 $\mu\text{g/ml}$. Virulence traits of *P. aeruginosa* like haemolytic activity, and biofilm formation were negatively affected by the test extract, and it also made *P. aeruginosa* more susceptible to lysis by human serum. Antibiotic susceptibility of both test bacteria was modulated owing to pre-treatment with PGPE. Exposure of these test pathogens to PGPE (≥ 0.5 $\mu\text{g/ml}$) effectively reduced their virulence towards the nematode *Caenorhabditis elegans*. Repeated subculturing of *P. aeruginosa* on PGPE-supplemented growth medium did not induce resistance to PGPE in this notorious pathogen, and this extract was also found to exert a post-extract effect on *P. aeruginosa*. Individual constituent phytocompounds of PGPE were found to be less efficacious than the whole extract. PGPE seemed to interfere with the *signal-response* machinery of *P. aeruginosa* and *C. violaceum*. PGPE also exhibited notable prebiotic potential by promoting growth of probiotic strains-*Bifidobacterium bifidum* and *Lactobacillus plantarum* at ≤ 50 $\mu\text{g/ml}$.

Conclusions: This study indicates PGPE to be an effective antipathogenic and prebiotic preparation, and validates its therapeutic use mentioned in traditional medicine. This study also

Open Peer Review

Approval Status

	1	2	3
version 2 (revision) 08 Apr 2019			
version 1 18 Jan 2019	 view	 view	 view

1. **Saif Hameed**, Amity University Haryana, Gurugram, India
2. **Sisir Nandi** , Global Institute of Pharmaceutical Education and Research, Affiliated to Uttarakhand Technical University, Kashipur, India
Uttarakhand Technical University, Kashipur, India
3. **Shilpa Deshpande Kaistha** , Chhatrapati Shahu Ji Maharaj University (CSJMU), Kanpur, India

Any reports and responses or comments on the article can be found at the end of the article.

emphasizes the need for testing any bioactive extract at broadest possible concentration range, particularly *in vivo*, so that an accurate picture of dose-response relationship can emerge.

Keywords

Punica granatum peel, Quorum Sensing, Post Extract Effect, Prebiotic, Microwave Assisted Extraction

This article is included in the **Pathogens** gateway.

This article is included in the **Antimicrobial Resistance** collection.

Corresponding author: Vijay Kothari (vijay.kothari@nirmauni.ac.in)

Author roles: **Joshi C:** Data Curation, Investigation, Writing – Original Draft Preparation; **Patel P:** Data Curation, Investigation, Writing – Original Draft Preparation; **Kothari V:** Conceptualization, Formal Analysis, Methodology, Resources, Supervision, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work received no extramural funding, and was fully supported by our departmental research budget (Institute of Science, Nirma University).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 Joshi C *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Joshi C, Patel P and Kothari V. **Anti-infective potential of hydroalcoholic extract of Punica granatum peel against gram-negative bacterial pathogens [version 1; peer review: 1 approved, 2 approved with reservations]** F1000Research 2019, 8:70 <https://doi.org/10.12688/f1000research.17430.1>

First published: 18 Jan 2019, 8:70 <https://doi.org/10.12688/f1000research.17430.1>

Introduction

Antimicrobial resistance (AMR) has been recognized as a global problem. Globalization and international travel has increased the vulnerability of any country to infections prevalent in other countries. Rapid spread of resistant factors across different species of pathogenic bacteria, particularly through horizontal gene transfer permitting promiscuous exchange of genetic material, presents a global threat to public health. It is estimated that every 5 minutes one child on the earth dies because of bacterial resistance to antibiotics (ESRC Report, 2014). One of the most problematic features of AMR is that resistance to a new antimicrobial can begin simultaneous to its development, meaning that simply developing new bactericidal antibiotics is not the panacea.

Hitherto, antimicrobial therapy has largely remained dependent on conventional bactericidal antibiotics. However, owing to their killing effect, such antibiotics pose considerably strong selection pressure on the pathogen populations to evolve resistant phenotypes. This necessitates thinking of alternative approaches for handling microbial infections, which asks for identifying novel targets, other than those (e.g. cell wall synthesis, nucleic acid/protein synthesis, cell membrane) targeted by current antibiotics (Beceiro *et al.*, 2013; Fair & Tor, 2014; Kohanski *et al.*, 2010). Among such possible novel targets, quorum sensing (QS) has attracted considerable attention from the researchers working in the area of AMR. QS is a mechanism of chemical signalling (via acyl homoserine lactones (AHL) in gram-negative bacteria, and autoinducing peptides (AIP) in gram-positive bacteria) based process, which allows the bacterial population to modulate its behaviour in relation to its cell density (Moore *et al.*, 2015). Though a large part of the bacterial genome (including the genes coding for virulence) is regulated by QS, it is not that essential for their survival as an individual cell, hence quorum-modulatory agents can be expected to exert their anti-infective/antipathogenic effect on susceptible bacterial populations, without necessarily putting a strong selection pressure on them.

Though the modern medicine has almost always focused on search of purified active molecules as therapeutic leads, knowledge of Indian/Chinese/Arabian traditional medicine makes us aware of a variety of plant extracts and inorganic formulations prescribed for management of infections. One such widely mentioned item in these ancient systems of complementary and alternative medicine, including the Indian system *Ayurved*, is *Punica granatum* L. extracts. This plant belongs to the family Lythraceae (Punicaceae), and the common English name is pomegranate. Different parts of this plant have been applied in traditional medicine to treat a variety of health problems. A peep into literature with particular focus on its peel reveals that in Egyptian culture, common ailments such as inflammation, diarrhea, intestinal worms, cough, and infertility were treated by exploiting pomegranate peel extract (Ismail *et al.*, 2012). *P. granatum* has been indicated in traditional Indian and Iranian medicine for its antimicrobial potential, for treatment of throat infections (e.g. sore throat caused by bacterial infection), diarrhea, wound healing, etc. (Hosein Farzaei *et al.*, 2014). In

Charak Samhita (Sutrsthana) pomegranate has been mentioned as an important ingredient of a *Yavagu* formulation for treatment of dysentery. In India, Tunisia, and Guatemala, decoction of dried peels of *P. granatum* is employed externally as well as internally as an astringent and germicide, and utilized for treating aphthae and diarrhoea. In *Ayurvedic* medicine, this plant described as ‘*dadim*’ (its Sanskrit name), is considered as a ‘blood purifier’ and suggested to cure parasitic infections (Colombo *et al.*, 2013). A search in the IMPPAT database (<https://cb.imsc.res.in/imppat/basicsearch/therapeutics>) using the search term “*Punica granatum*” yields 75 different therapeutic uses, of which multiple conditions (vaginal trichomoniasis, pharyngitis, periodontitis, paracoccidioidomycosis, malaria, leishmaniasis, Klebsiella Pneumonia, *Helicobacter pylori* infection, gonorrhea, gingivitis, genital herpes, dysentery, diarrhea, conjunctivitis, cholera, etc.) for which *P. granatum* is indicated can involve microbial infections. Ethnobotanical use of *P. granatum* for treatment of diarrhoea has been documented from South Africa too (Mathabe *et al.*, 2006). Mexican traditional medicine also mentions utility of *P. granatum* in treatment of gastrointestinal disorders (Alanis *et al.*, 2005). Based on Iranian traditional medicinal practice, Hajimahmoodi *et al.* 2011 demonstrated peel extracts of eight cultivars of pomegranate to be effective against *Helicobacter pylori*. One ethnomedical practice reported from the Paliyar tribe from Tamilnadu in India involves taking internally, dried fruit coat of *P. granatum* after grinding and mixing with water, to treat stomach ache and diarrhoea (Duraipandiyan *et al.*, 2006). Ayurvedic pharmacopeia of India mentions formulations containing rind of *P. granatum* to be useful in conditions such as fever, dysentery, bacteremia, and infections of oral cavity (Khare, 2004). Decoction containing *P. granatum* peel is used for gastrointestinal benefit in Algeria too (Gharzouli *et al.*, 1999). Despite their wide use in medicinal texts of various cultures, efficacy of pomegranate extracts needs to be validated through appropriate experiments for their wider acceptance in the modern world.

This study aimed at investigating the possible anti-infective potential of pomegranate peel extract against two gram-negative bacteria, *Pseudomonas aeruginosa* and *Chromobacterium violaceum*. *P. aeruginosa* is amongst the most notorious bacterial pathogens, associated with chronic and acute urinary and respiratory tract infections, and its carbapenem-resistant phenotype has been listed as a pathogen of ‘critical’ priority against which new antimicrobials are urgently warranted (WHO report, 2017). *C. violaceum* is widely used in QS studies and is also being viewed as an emerging pathogen (Kothari *et al.*, 2017). WHO advises for future research to focus on the development of new antibiotics specifically against drug-resistant gram-negative bacteria (WHO report, 2017). In both gram-negative bacteria selected in this study, pigment production is a trait, which is reported to be under control (largely but not fully) of their QS machinery, and this was used as a marker trait by us while assaying effect of our extract on these bacteria. QS-regulated pigments produced by these bacteria are: pyoverdine and pyocyanin in *P. aeruginosa* (Adonizio *et al.*, 2007; Lee & Zhang, 2015), and violacein in *C. violaceum* (Vasavi *et al.*, 2013).

Methods

Materials

All the growth media/ media ingredients/ assay reagents, and antibiotics were procured from Himedia, unless specified otherwise. TLC plates, and all organic solvents used in this study were procured from Merck. Whatman paper, and bacteriological filters were from Axiva (Haryana). Catechin was from Sigma Aldrich (USA). Triton X-100 was from CDH (New Delhi). All experiments were performed with the same set of reagents.

Test organisms

C. violaceum (MTCC 2656), and *Lactobacillus plantarum* (MTCC 2621), were procured from MTCC (Microbial Type Culture Collection, Chandigarh), whereas *Bifidobacterium bifidum* (NCDC 255) was procured from NCDC (National Collection of Dairy Cultures), NDRI (National Dairy Research Institute, Karnal). *C. violaceum* was grown in nutrient broth (HiMedia, Mumbai), *L. plantarum* was grown in Lactobacillus MRS medium (HiMedia, Mumbai), and *B. bifidum* was grown on MRS with 0.05% cysteine. Incubation temperature and time for *C. violaceum*, *L. plantarum*, and *B. bifidum* was 37°C, and 22–24 h. Antibiotic susceptibility profile of the bacterial strains used in this study was generated using the antibiotic discs-Dodeca Universal – I, Dodeca G - III – Plus, and Icosa Universal -2 (HiMedia, Mumbai). *C. violaceum* was found to be resistant to cefadroxil (30 µg), ampicillin (10 µg), cloxacillin (1 µg), and penicillin (10 µg). Culture of *P. aeruginosa* was obtained from Microbiology Department, M.G. Science Institute, Ahmedabad. *Pseudomonas* agar (HiMedia, Mumbai) was used for the maintenance of the culture. Strain of *P. aeruginosa* was found to be resistant to amoxicillin (30 µg), cefadroxil (30 µg), ampicillin (10 µg), cloxacillin (1 µg), penicillin (10 µg), chloramphenicol (30 µg), cefixime (5 µg), clindamycin (2 µg), and nitrofurantoin (300 µg).

Plant material

Peels of *P. granatum* were procured from the fruits purchased from local market in the city of Ahmedabad. Plant material (ref no: GU/Bot/29/4/2015) was authenticated for its unambiguous identity by Dr Archana Mankad (Department of Botany, Gujarat University, Ahmedabad). The plant name was checked against <http://www.theplantlist.org> on April 20, 2018. Collected peels were shade-dried, before being used for extract preparation. We also analysed another pomegranate fruit extract as 'Pomella' by Pharmanza Herbal Pvt. Ltd., which contained not less than 30% punicalagin

Extract preparation

A hydroalcoholic extract of the plant material was prepared in 50% ethanol using the microwave-assisted extraction (MAE) method (Kothari *et al.*, 2009). Dry peel powder (total 5 g; 1 g in each vessel) was soaked into the solvent in a ratio of 1:50, and subjected to microwave heating (Electrolux EM30EC90SS) at 720 W. Total heating time was 120 seconds, with intermittent cooling. This was followed by centrifugation (at 10,000 rpm for 15 min.), and filtration with Whatman paper #1 (Axiva, Haryana). Solvent was evaporated from the filtered extract and then the

dried extract was reconstituted in dimethyl sulfoxide [DMSO (absolute); Merck] for broth dilution assay. Reconstituted extract was stored under refrigeration for further use. Extraction efficiency was calculated as percentage weight of the starting dried plant material, and its value was 42% w/w.

Broth dilution assay

Assessment of QS-regulated pigment production by test pathogens in presence or absence of the test formulation, was done using broth dilution assay (Patel *et al.*, 2017). Organisms were challenged with different concentrations (0.5-500 µg/ml) of *P. granatum* peel extract (PGPE). Nutrient broth or *Pseudomonas* broth (peptic digest of animal tissue 20 g/l, potassium sulphate 10 g/l, magnesium chloride 1.4 g/l, pH 7.0 ± 0.2) was used as a growth medium. Inoculum standardized to 0.5 McFarland turbidity standard was added at 10% v/v, to the media supplemented with required concentration of PGPE, followed by incubation at appropriate temp for each organism. Appropriate vehicle control containing DMSO was also included in the experiment, along with abiotic control (containing extract and growth medium, but no inoculum). Catechin (50 µg/ml; Sigma Aldrich, USA) was used as positive control, since it is already known QS inhibitor (Nazzaro *et al.*, 2013; Vandeputte *et al.*, 2010).

Measurement of bacterial growth and pigment production

At the end of the incubation, bacterial growth was quantified at 764 nm (Joshi *et al.*, 2016). This was followed by pigment extraction and quantification, as per the method described below for each of the pigments. Purity of each of the extracted pigment was confirmed by running a UV-vis scan (Agilent Cary 60 UV-visible spectrophotometer). Appearance of single major peak (at the λ_{max} reported in literature) was taken as indication of purity.

Violacein extraction

Extraction of violacein from *C. violaceum* culture was executed as described by Choo *et al.* (2006). A total of 1 ml of the culture broth was centrifuged (Eppendorf 5417 R) at 15,300g for 10 min at room temperature, and the resulting supernatant was discarded. The remaining cell pellet was resuspended into 1 ml of DMSO, and vortexed, followed by centrifugation at 15,300g for 10 min. The purple coloured violacein was extracted in the supernatant; OD was measured at 585 nm. Violacein unit was calculated as OD_{585}/OD_{764} . This parameter was calculated to nullify the effect of any change in cell density on pigment production.

Pyoverdine and pyocyanin extraction

Extraction of pyoverdine and pyocyanin from *P. aeruginosa* culture was achieved as described in El-Fouly *et al.* (2015) and Unni *et al.* (2014), respectively. A total of 1 ml of the culture broth was mixed with chloroform (Merck, Mumbai) in 2:1 proportion followed by centrifugation at 12,000 rpm (13,520g; REMI CPR-24 Plus) for 10 min. This resulted in formation of two immiscible layers. OD of the upper water-soluble phase containing yellow-green fluorescent pigment pyoverdine was measured at 405 nm. Pyoverdine Unit was calculated as OD_{405}/OD_{764} .

The lower chloroform layer containing pyocyanin was mixed with 0.1 N HCl (Merck; at the rate of 20%v/v), resulting in a colour change from blue to pink. Absorbance of this pyocyanin in acidic form was measured at 520 nm. Pyocyanin Unit was calculated as OD_{520}/OD_{764} .

N- acyl-homoserine lactone (AHL) augmentation assay

AHL extraction: AHL was extracted from the bacterial culture broth as described by [Chang et al. \(2014\)](#). OD of overnight grown bacterial culture was standardized to 1.00 at 764 nm. It was centrifuged at 5000g for 5 min. Cell free supernatant was filter sterilized using 0.45 µm filter (Axiva, Haryana), and was mixed with equal volume of acidified ethyl acetate [0.1% formic acid (Merck) in Ethyl acetate (Merck)]. Ethyl acetate layer was collected, and evaporated at 55°C, followed by reconstitution of the dried crystals in 100 µl phosphate buffer saline (pH 6.8). Identity of thus extracted AHL was confirmed by thin-layer chromatography (TLC). R_f value of purified AHL from *C. violaceum* culture, while performing TLC [Methanol (60): Water (40); TLC Silica gel 60 F₂₅₄ plates; Merck] [McClean et al., \(1997\)](#) was found to be 0.70, near to that (0.68) reported for N-hexonyl homoserine lactone (C6-HSL) ([Shaw et al., 1997](#)). TLC of AHL extracted from *P. aeruginosa* resulted in three spots corresponding to R_f values of 0.43, 0.68, and 0.92 near to those (0.41, 0.68, 0.84) for C8-HSL, C6-HSL, and C4-HSL respectively.

Bacterial culture growing in presence of test formulation was supplemented with 2%v/v AHL after 6 hours of incubation, and at the end of a total 24-hour incubation pigments were extracted from AHL-supplemented experimental tubes, as well as AHL-non-supplemented control tubes. If the QS-regulated pigment production is found inhibited in both these tubes in comparison to bacteria growing in absence of extract as well as AHL, then the effect of test extract was interpreted as signal-response inhibitor, because if the test formulation would have acted as a signal-supply inhibitor, then exogenous supply of AHL should restore pigment formation by the bacteria.

Hemolysis assay

This assay was done as described by [Neun et al. \(2015\)](#). OD_{764} of overnight grown culture was standardized to 1.00. Cell free supernatant was prepared by centrifugation at 15,300 g for 10 min. A total of 10 µl human blood (collected in heparinized vial) was incubated with this cell free supernatant for 2 h at 37°C, followed by centrifugation at 800 g for 15 min; 1% Triton X-100 (CDH, New Delhi) was used as a positive control. Phosphate buffer saline was used as a negative control. OD of the supernatant was read at 540 nm, to quantify the amount of haemoglobin released.

Assay of bacterial susceptibility to lysis in presence of human serum

This assay was performed as described by [Ferro et al. \(2016\)](#). Serum was separated by centrifuging blood at 1,500 rpm (800g) for 10 min. Bacterial culture grown in media with and without PGPE was centrifuged, and the cell pellet was reconstituted in PBS, so that the resulting suspension attains $OD_{764}=1$. A total of 200 µl of this bacterial suspension from control or

experimental tubes was mixed with 740 µl of PBS and 60 µl of serum. After incubation for 24 h at 37°C, absorbance was read at 764 nm. DMSO -treated cells (0.5%v/v) suspended in PBS served as control, against which OD of the PGPE-treated cells (serum-exposed) was compared. Tubes containing bacterial cells exposed neither to DMSO nor serum were also included in the experimental set-up, to nullify any interference from autolysis.

Human blood was obtained from the authors, who each gave their written informed consent. The use of this blood was approved by the Institutional Ethics Committee of the Institute of Science, Nirma University (approval no: EC/NU/18/IS/4).

Catalase assay

The OD_{764} of the culture was adjusted to 1.00. Next, 400 µl of phosphate buffer was added into a 2 ml vial followed by 400 µl H_2O_2 . To this 200 µl of the bacterial culture was added, and the mixture was incubated for 10 min. Then 10 µM of sodium azide was added to stop the reaction ([Iwase et al., 2013](#)), followed by centrifugation at 12,000 rpm for 10 min. OD of the supernatant was measured at 240 nm to quantify remaining H_2O_2 ([Weydert & Cullen, 2010](#)), with phosphate buffer as blank. Disappearance of the substrate H_2O_2 was taken as the measure of the catalase activity.

Assay for biofilm formation, eradication and viability

In this assay, the control and experimental groups contained nine test tubes. In each group, three subgroups were made. First subgroup of three test tubes in the experimental group contained *Pseudomonas* broth supplemented with PGPE, whereas remaining six tubes contained *Pseudomonas* broth with no PGPE on first day of experiment. All these tubes were inoculated with inoculum (10%v/v) standardized to 0.5 McFarland turbidity standard (making the total volume in the tube 1 ml), followed by incubation at 37°C for 24 h under static condition, which resulted in formation of biofilm as a ring on walls of the glass tubes. This biofilm was quantified by crystal violet assay ([Patel et al., 2013](#)), preceded by quantification of bacterial cell density and pigment. Content from the remaining six test tubes from rest of the two subgroups were discarded following cell density and pigment estimation, and then the biofilms remaining on inner surface of these tubes were washed with phosphate buffer saline (PBS; pH 7) to remove loosely attached cells. Now, 2 ml of minimal media (Sucrose 15 g/l, K_2HPO_4 5.0 g/l, NH_4Cl 2 g/l, NaCl 1 g/l, $MgSO_4$ 0.1 g/l, yeast extract 0.1 g/l, pH 7.4±0.2) containing PGPE, was added into each of these tubes, so as to cover the biofilm completely, and tubes incubated for 24 h at 37°C. At the end of incubation, one subgroup of 3 tubes was subjected to crystal violet assay to know whether any eradication of the pre-formed biofilm has occurred under the influence of PGPE, and the last subgroup of 3 tubes was subjected to viability assessment through MTT assay. For the crystal violet assay, the biofilm- containing tubes (after discarding the inside liquid) were washed with PBS in order to remove all non-adherent (planktonic) bacteria, and air-dried for 15 min. Then, each of the washed tubes was stained with 1.5 ml of 0.4% aqueous crystal violet solution for 30 min. Afterwards, each tube was washed twice with 2 ml sterile distilled water and immediately de-stained

with 1500 µl of 95% ethanol. After 45 min of de-staining, 1 ml of de-staining solution was transferred into separate tubes, and read at 580 nm. For the MTT assay (Trafny *et al.*, 2013), the biofilm-containing tubes (after discarding the inside liquid) were washed with PBS in order to remove all non-adherent (planktonic) bacteria, and air-dried for 15 min. Then 900 µl of minimal media was added into each tube, followed by addition of 100 µl of 0.3% MTT [3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide, HiMedia]. After 2 h incubation at 37°C, resulting purple formazan derivatives were dissolved in DMSO and measured at 560 nm.

Cell surface hydrophobicity (CSH) assay

Bacterial surface hydrophobicity was measured using the bacterial adhesion to hydrocarbon (BATH) assay as described in Hui & Dykes (2012). *P. aeruginosa* culture was collected at stationary phase and pelleted by centrifugation (NF800R, NUVE, Belgium) at 7,000 rpm for 10 min. This pellet was washed twice in phosphate buffer saline (PBS; pH 7.4), and then resuspended in PBS with PGPE (25 and 50 µg/ml) to OD₇₆₄ = 1.00. Same procedure was repeated with *P. aeruginosa* culture without PGPE, as a control. Each bacterial suspension was then incubated for 1 h at room temperature. A 2-ml sample of each suspension was collected and absorbance (A) at 764 nm was measured, using PBS as the blank. Next, 1 ml xylene (HiMedia, Mumbai) was added to the 2-ml cell suspension, and this mixture was vortexed for 2 min. The phases were then allowed to separate for 1 h. The absorbance of the aqueous phase (A₀) was again determined. Results were expressed as % attachment to xylene = $(1 - A/A_0) \times 100$.

Determination of the effect of PGPE on antibiotic susceptibility of the test organism

After *in vitro* assessment of QS modulatory (QSM) property of the test formulation, effect of this PGPE on antibiotic susceptibility of the test pathogen was investigated. The bacterial cells pre-treated with PGPE were subsequently challenged with sub-MIC concentration of antibiotic. All the antibiotics were procured from HiMedia, Mumbai. Names and concentrations of the antibiotics used can be seen in Figure 1c, Figure 2c, Figure 5c and Figure 7e.

In vivo assay

In vivo efficacy of the PGPE was evaluated using the nematode worm *Caenorhabditis elegans* as the model host, using the method described by Eng & Nathan (2015) with some modification. This worm was maintained on Nematode Growing Medium (NGM; 3 g/l NaCl, 2.5 g/l peptone, 1 M CaCl₂, 1 M MgSO₄, 5 mg/ml cholesterol, 1 M phosphate buffer of pH 6, 17 g/l agar-agar) with *Escherichia coli* OP50 (procured from LabTIE B.V., JR Rosmalen, the Netherlands) as the feed. Worm population to be used for the *in vivo* assay was kept on NGM plates not seeded with *E. coli* OP50 for three days, before being challenged with the test pathogen.

Test bacterium was incubated with the PGPE for 22–24h. Following incubation, OD₇₆₄ of the culture suspension was equalized to that of the DMSO control. 100 µl of this bacterial suspension

was mixed with 900 µl of the M9 buffer containing 10 worms (L3–L4 stage). This experiment was performed in 24-well (sterile, non-treated) polystyrene plates (HiMedia), and incubation was carried out at 22°C. Number of live vs. dead worms was counted everyday till five days by putting the plate (with lid) under light microscope (4X). Standard antibiotic- (ampicillin 500 µg/ml; gentamicin 0.1 µg/ml) and catechin- (50 µg/ml) treated bacterial suspension were used as positive control. Straight worms were considered to be dead. On last day of the experiment, when plates could be opened, their death was also confirmed by touching them with a straight wire, wherein no movement was taken as confirmation of death.

Investigating whether the bacterium becomes resistant to PGPE upon repeated exposure

P. aeruginosa was subcultured on PGPE (500 µg/ml)-containing agar medium 10 times, and this PGPE-habituated culture was then challenged with PGPE (500 µg/ml) in liquid media for broth dilution assay, wherein bacterial cell density and pigment production were quantified as detailed in preceding sections. Virulence of this PGPE-habituated culture towards *C. elegans* was also assessed using the same methodology as described above under the heading ‘*In vivo* assay’.

Chromatographic analysis

Chromatographic fingerprint of PGPE was generated through HPLC (Shimadzu Lab Solutions) by running the extract through a C18 column (Shimadzu) for 75 min. Mobile phase consisted of a mixture of trichloroacetic acid (0.05%, w/v) in water, and trichloroacetic acid (0.05%, w/v) in methanol, wherein proportion of latter was varied over a gradient of 10–80%. Flow rate was 1 ml/min. Detection wavelength of the UV detector was set at 258 nm. Punicalagin A and punicalagin B (both at 100 ppm; Sigma Aldrich) were used as standard compounds.

Molecular docking

Structures of the pure compounds were downloaded from PubChem: catechin (PubChem CID: 73160), gallic acid (GA) (PubChem CID: 370), quercetin (PubChem CID: 5280343), ellagic acid (PubChem CID: 5281855), cinnamic acid (PubChem CID: 444539), and chlorogenic acid (PubChem CID: 1794427). Structure of the bacterial target proteins was downloaded from the Protein Data Bank (PDB): CviR of *C. violaceum* (PDB ID 3QP4), LasR (PDB ID 2UV0), and PqsR (PDB ID 4JVC) of *P. aeruginosa*. AutoDock Vina (Trott & Olson, 2010) was used for docking studies, Discovery Studio Visualizer 4.0.1 was used to study protein-ligand interactions, Pymol viewer (version v1.3r1) was used to convert protein-ligand docked structures from .pdbqt to .pdb format, and MglTools was used to prepare protein and ligand files for docking.

Statistical analysis

All the *in vitro* experiments were performed in triplicate, and measurements are reported as mean ± standard deviation (SD) of three independent experiments. *In vivo* experiments were run in four replicates. Statistical significance of the data was evaluated by applying *t*-test using Microsoft Excel. *p* values ≤0.05 were considered to be statistically significant.

Results

C. violaceum

C. violaceum was challenged with the PGPE at 0.5–500 µg/ml (Figure 1A). PGPE had a negative effect on bacterial growth only at concentrations ≥ 25 µg/ml; however, an increase in concentration could not inhibit growth to any greater extent. In fact, statistically all the concentrations in the range 50–500 µg/ml had equivalent effect on *C. violaceum* growth. On the other hand, production of the QS-regulated pigment violacein was sensitive to PGPE at lesser concentrations (i.e. at ≥ 5 µg/ml), and the inhibitory effect increased with increase in concentration of the extract. PGPE can be said to exert a purely QSM effect at 5–10 µg/ml, as at these concentrations, it could inhibit pigment production significantly without inhibiting *C. violaceum* growth.

Once the QS-inhibitory potential of PGPE against *C. violaceum* was confirmed, we experimented to know whether it acts as a signal-supply inhibitor or signal-response inhibitor. To know this, we added AHL (2% v/v; 6 h post-inoculation) into quorum

inhibited culture of *C. violaceum* growing in the presence of PGPE (10–100 µg/ml). Following AHL (signal) augmentation, the quorum inhibitory effect of PGPE on violacein production was not reversed (Figure 1B), indicating that PGPE acts as a signal-response inhibitor.

When PGPE-treated *C. violaceum* culture was subsequently challenged with sub-MIC concentrations of four antibiotics belonging to four different classes, extract pre-treatment was found to enhance susceptibility of this bacterium to all these antibiotics. However, pre-treatment with higher (25 µg/ml) PGPE concentration was found to be no better than the lesser (10 µg/ml) concentration, with respect to making bacteria more susceptible to given antibiotic, except in the case of ampicillin (Figure 1C).

PGPE was found to curb haemolytic potential of *C. violaceum* in a dose-dependent fashion, wherein this extract at 100 µg/ml could reduce haemolysis by $\sim 36\%$ (Table 1). This extract could also reduce the catalase activity of *C. violaceum* marginally;

Figure 1. Effect of PGPE on *C. violaceum*. 'Control' in this figure is the vehicle control (0.5%v/v DMSO), which did not exert any effect on growth and violacein production of *C. violaceum*. (A) Effect of PGPE on growth and QS-regulated violacein production in *C. violaceum*: Bacterial growth was measured as OD₇₆₄; OD of violacein was measured at 585 nm, and Violacein Unit was calculated as the ratio OD₅₈₅/OD₇₆₄ (an indication of violacein production per unit of growth); Catechin (50 µg/ml) did not exert any effect on growth of *C. violaceum*, but inhibited violacein production by 47.69±0.03%. (B) PGPE acts as a signal-response inhibitor against *C. violaceum*. (C) PGPE-pre-treatment enhances susceptibility of *C. violaceum* to different antibiotics. (D) PGPE-treatment attenuates virulence of *C. violaceum* towards *C. elegans*: Catechin (50 µg/ml) and ampicillin (500 µg/ml) employed as positive controls conferred 100% protection on worm population, and pre-treatment of bacteria with PGPE at concentrations (10, 25, 50 and 100 µg/ml) other than those shown in figure allowed 75%, 77.5%, 80%, and 75% worm survival, respectively; DMSO present in the 'vehicle control' at 0.5%v/v did not affect virulence of the bacterium towards *C. elegans*; DMSO (0.5%v/v) and PGPE at tested concentrations showed no toxicity towards the worm. **p*<0.05, ***p*<0.01, ****p*<0.001; AS, antibiotic susceptibility; QS, quorum sensing; PGPE, *Punica granatum* peel extract.

Table 1. Effect of PGPE on catalase and haemolytic activity of test bacteria.

Organism	Concentration of extract, µg/ml	% Change in catalase activity (mean ± SD)	% Inhibition in Hemolysis (Mean ± SD)
<i>C. violaceum</i>	10	-4.65* ± 2.46	8.83* ± 3.46
	25	-5.81* ± 3.28	22.09** ± 2.40
	50	-8.13* ± 1.64	28.72** ± 3.79
	100	-11.62* ± 6.57	35.91*** ± 3.02
<i>P. aeruginosa</i>	10	0.79 ± 0.62	-9.19 ± 6.27
	25	0.78 ± 0.44	15.47* ± 3.33
	50	-1.77 ± 0.47	22.61* ± 4.89
	100	-2.56* ± 0.15	33.33** ± 3.83

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; Catalase assay was done by monitoring disappearance of H_2O_2 at 240 nm; Hemoglobin concentration was measured at OD_{540} . DMSO in 'vehicle control' tube had no effect on catalase and haemolytic activity of any of the three bacteria; Chloramphenicol (0.5 µg/ml) enhanced catalase activity of *C. violaceum* by $11.23\% \pm 0.01$; Tetracycline (0.5 µg/ml) inhibited catalase activity of *P. aeruginosa* by $21.51\% \pm 0.02$.

however, the effect on this oxidative stress response enzyme activity did not increase with increase in PGPE concentration (Table 1).

In vitro demonstration of the QSM potential of PGPE was followed by assessment of its anti-infective potential *in vivo*, employing *C. elegans* as the model host, wherein PGPE was found to confer survival benefit on *C. elegans* upon challenge with *C. violaceum* at all the concentrations (0.5–100 µg/ml) tested (Figure 1D). Notably, even the concentrations (0.5–2.5 µg/ml) having no significant *in vitro* effect on *C. violaceum* growth and pigment production, were found to be effective *in vivo*. Maximum *in vivo* benefit was obtained at 5 µg/ml PGPE concentration, and worm survival percentage at any of the higher concentrations were statistically no better than that obtained at 5 µg/ml. Onset of death in the worm population was also delayed by 2 days (i.e. death started on first day-post infection in control wells, as against on third day in experimental wells) when the challenger bacteria were pre-treated with PGPE. Movement of the surviving worm in 'vehicle control' wells was slowed down, whereas worms surviving in the 'experimental' wells showed normal movement. Further, worms in the experimental well corresponding to 100 µg/ml PGPE could also generate progeny worms, which were not observed in any other well.

P. aeruginosa

P. aeruginosa was challenged with PGPE at 0.5–500 µg/ml, wherein growth of this bacterium got notably inhibited 50 µg/ml onwards. Pyocyanin production was affected negatively from 5 µg/ml onwards in a concentration-dependent manner, whereas pyoverdine production was enhanced from 2.5 µg/ml onwards in a largely concentration-dependent fashion except at 500 µg/ml (Figure 2A). When AHL was added exogenously to the quorum-modulated *P. aeruginosa* culture, the inhibitory effect

of PGPE on pyocyanin and its stimulatory effect on pyoverdine production was not reversed (Figure 2B) indicating that PGPE acts as a signal-response inhibitor against this pathogen.

Effect of PGPE pre-treatment on antibiotic susceptibility of *P. aeruginosa* was also investigated. However, no major changes were observed in bacterial susceptibility to four different test antibiotics, when challenged with these antibiotics following PGPE exposure (Figure 2C). This extract was not found to alter catalase activity much; however, haemolytic potential of *P. aeruginosa* was notably affected 25 µg/ml onwards (Table 1). PGPE at 50 µg/ml could enhance (by 13.55%; $p = 0.03$) susceptibility of this bacterial pathogen to lysis by human serum (Figure 2D).

PGPE was able to reduce *P. aeruginosa* biofilm formation up to 17–30%, when tested at 10–50 µg/ml (Figure 2E). When this extract was applied on pre-formed biofilm, it could eradicate the biofilm by ~15–19%; and biofilm viability (metabolic activity as measured by MTT assay) was reduced by ~13–22%. However, whether this observed reduction in viability is due to lesser number of cells, or some other reason, remains to be investigated. Reduced biofilm formation by *P. aeruginosa* in presence of PGPE may partly be due to this extract's ability to reduce CSH marginally, which is an important determinant of biofilm forming ability in bacteria (Hui and Dykes, 2012). Xylene adherence of *P. aeruginosa* treated with PGPE at 25 and 50 µg/ml was found to be reduced by 14.19% ($p < 0.05$; % adherence = 44.74) and 7.82% ($p < 0.05$; % adherence = 38.37) respectively, against 30.55% adherence of control.

In vivo assay did confirm the anti-infective potential of PGPE against *P. aeruginosa*. At concentrations as low as 0.5–1 µg/ml, it could support overall survival of the worm population up to

Figure 2. Effect of PGPE on various traits of *P. aeruginosa*, in vitro. 'Control' in this figure is the vehicle control (0.5%v/v DMSO), which did not exert any effect on growth and pigment production of *P. aeruginosa*. **(A)** Effect of PGPE on growth and QS-regulated pigment production in *P. aeruginosa*: Bacterial growth was measured as OD₇₆₄; OD of pyoverdine was measured at 405 nm, and Pyoverdine Unit was calculated as the ratio OD₄₀₅/OD₇₆₄ (an indication of pyoverdine production per unit of growth), Pyocyanin Unit was calculated as the ratio OD₅₂₀/OD₇₆₄ (an indication of pyocyanin production per unit of growth); Catechin (50 µg/ml) inhibited pyoverdine and pyocyanin production by 17.13±0.06% and 23.65±0.04% respectively, without affecting the bacterial growth. **(B)** PGPE acts as a *signal-response inhibitor* against *P. aeruginosa*. **(C)** *P. aeruginosa* challenged with antibiotics following pre-treatment with PGPE. **(D)** PGPE modulates susceptibility of *P. aeruginosa* to lysis by human serum: 'Control' in serum-dependent lysis assay was PGPE-unexposed cells of *P. aeruginosa* incubated with human serum. **(E)** Effect of PGPE on *P. aeruginosa* biofilm formation, eradication, and viability. Crystal violet assay was performed to measure biofilm formation, and biofilm eradication, followed by the measurement of OD at 580 nm; Cell viability in biofilm was estimated through MTT assay, wherein OD was measured at 560 nm. **(F)** PGPE-treatment reduces the virulence of *P. aeruginosa* towards *C. elegans*: Catechin (50 µg/ml) and gentamicin (0.1 µg/ml) employed as a positive controls conferred 100% and 80% protection on worm population respectively. Pre-treatment of bacteria with PGPE at concentrations (2.5, 5, 10, 25, and 100 µg/ml) other than shown in figure allowed 73.75, 75, 77.5, 77.5, and 75% worm survival respectively; DMSO present in the 'vehicle control' at 0.5%v/v did not affect virulence of the bacterium towards *C. elegans*; DMSO (0.5%v/v) and PGPE at tested concentrations showed no toxicity towards the worm. **(G)** Effect of PGPE on *P. aeruginosa* growth, Pyoverdine Unit, and Pyocyanin Unit remained unaltered after repeated exposure to PGPE (500 µg/ml). **p*<0.05, ***p*<0.01, ****p*<0.001. AS, antibiotic susceptibility; QS, quorum sensing; PGPE, *Punica granatum* peel extract.

56–71% in face of *P. aeruginosa* challenge, as against only 12.5% worm survival in control population. Higher PGPE concentrations did not offer any better protection to *C. elegans* against *P. aeruginosa* (Figure 2F). Notably, *in vitro* experiments showed PGPE to have no effect on *P. aeruginosa* growth below 5 µg/ml, and no effect on its QS-regulated pigments below 2.5 µg/ml; whereas PGPE at 0.5–1 µg/ml was able to confer significant survival benefit on *C. elegans* against *P. aeruginosa* challenge. Onset of death in worm population was also delayed by 1–3 days, when *P. aeruginosa* was exposed to PGPE before being allowed to infect *C. elegans*.

After confirming the anti-infective potential of PGPE against *P. aeruginosa*, we investigated whether this bacterium can develop resistance to PGPE upon repeated exposure to this extract. We sub-cultured *P. aeruginosa* on PGPE (500 µg/ml) containing agar medium 10 times, and this PGPE-habituated culture was then challenged with PGPE (500 µg/ml) in liquid media for broth dilution assay, wherein effect of PGPE on growth and pigment production in *P. aeruginosa* was not found to be much different than that on *P. aeruginosa* with no previous PGPE exposure (Figure 2G). Similarly, there was no difference in the virulence of *P. aeruginosa* towards *C. elegans*, irrespective of whether it was previously habituated to PGPE. (Figure 2F).

With *P. aeruginosa*, we did an additional experiment to investigate whether the daughter cells of PGPE-exposed *P. aeruginosa* bear any attenuation of their virulence owing to their parent cells being exposed to PGPE. When the PGPE-exposed *P. aeruginosa* was subsequently subcultured on PGPE-free media, pigment production in cells obtained after first such subculturing was still altered in a pattern similar to the PGPE-exposed parent culture (Figure 3A). Though QS-regulated

pigment production remained deviated from ‘control’ upon second subculturing too, it did not follow the same pattern, as observed with cells obtained after first subculturing on PGPE-free media. With each subculturing, virulence of *P. aeruginosa* towards *C. elegans* approached nearer to that of ‘control’, but it still did not get equivalent to ‘control’ (Figure 3B), clearly suggesting that effect of PGPE-treatment was not completely absent even from the progeny cells (who never were directly exposed to PGPE), whose parent cells received single PGPE-exposure. This phenomenon of long-lasting effect after single-time exposure to the antimicrobial agent is described as ‘post-antimicrobial effect’. Here, in case of a plant extract, we prefer to call it post-extract effect (PEE), which refers to the persistent suppression of one or more bacterial traits (e.g. growth/virulence, etc.) after one-time-exposure to antimicrobial agents, and may last for many hours depending on the concentration of test agent and the susceptibility of the target bacterium (Pfaller *et al.*, 2004; Ramadan *et al.*, 1995; Ramanuj *et al.*, 2012). During actual animal/human infections, such phenomenon can be of high significance, as though the infectious bacteria may multiply inside the host system, their progenies may not have the virulence at par with the parent cells.

Characterization of PGPE

Once *in vitro* and *in vivo* anti-pathogenic potential of PGPE was confirmed, we proceeded to chromatographic fingerprinting of this extract. The resultant chromatogram is shown in Figure 4. Since punicalagin is one of the most widely reported bioactive metabolite from *P. granatum*, we quantified its amount in our extract, which was found to be present at 6.6%. Earlier we had analyzed another pomegranate fruit extract marketed as ‘Pomella’ by Pharmanza Herbal Pvt. Ltd., which contained not less than 30% punicalagin for its QS-modulatory potential (extended data,

Figure 3. Post extract effect of PGPE (10 µg/ml) on *P. aeruginosa*. ‘Control’ in this figure is the vehicle control (0.5%v/v DMSO), which did not exert any effect on growth and pigment production of *P. aeruginosa*. (A) Effect of PGPE on growth and QS-regulated pigment production in *P. aeruginosa* after subculturing of cells in PGPE-free media. Bacterial growth was measured as OD₇₆₄; OD of pyoverdine was measured at 405 nm, and Pyoverdine Unit was calculated as the ratio OD₄₀₅/OD₇₆₄ (an indication of pyoverdine production per unit of growth). Pyocyanin Unit was calculated as the ratio OD₅₂₀/OD₇₆₄ (an indication of pyocyanin production per unit of growth). (B) PGPE-treatment reduces the virulence of *P. aeruginosa* towards *C. elegans* even after subculturing of cells in PGPE-free media. Catechin (50 µg/ml) and gentamicin (0.1 µg/ml) employed as a positive controls conferred 100% and 80% protection on worm population respectively; DMSO present in the ‘vehicle control’ at 0.5%v/v did not affect virulence of the bacterium towards *C. elegans*; DMSO (0.5%v/v) and PGPE at tested concentrations showed no toxicity towards the worm. **p*<0.05, ***p*<0.01, ****p*<0.001; AS, antibiotic susceptibility; QS, quorum sensing; PGPE, *Punica granatum* peel extract.

Figure 4. HPLC profile of PGPE. Peak 11 and 14 occupied 24.61% and 27.48% of total area, which corresponds to Punicalagin A and B, respectively. PGPE, *Punica granatum* peel extract.

Figure S1A) (Joshi *et al.*, 2018). Though there were no major differences with respect to the *in vitro* effect on growth and pyocyanin production by *P. aeruginosa*, pyoverdine production was affected much more by Pomella than PGPE. However, *in vivo* efficacy of PGPE was bit better than that of Pomella; PGPE at 0.5–50 µg/ml allowed 56–80% worm survival, as against 55–65% worm survival supported by ‘Pomella’ at

identical concentrations (extended data, Figure S1B) (Joshi *et al.*, 2018). From this, we may speculate punicalagin not to be the major constituent responsible for *P. granatum* extract’s anti-pathogenic activity. Against *C. violaceum* too, except at 250 µg/ml, effect of PGPE on violacein production was statistically not different from that of Pomella, with latter exerting a bit more inhibitory effect on growth of this bacterium (extended

data, [Figure S1C](#)) ([Joshi et al., 2018](#)). However, role of punicalagins cannot be completely ruled out, as they are known to be metabolized inside human system to urolithins, and latter has been reported to be capable of exerting anti-QS effect ([Giménez-Bastida et al., 2012](#)). [Larrosa et al. \(2010\)](#) suggested the possibility of urolithin-A being the most active anti-inflammatory compound derived from pomegranate ingestion in healthy subjects.

QSM potential of *P. granatum* peel phytochemicals in pure form

Other than punicalagin, pomegranate peel extracts have been reported to contain pyrogallol, catechin, rutin, cinnamic acid, benzoic acid, chlorogenic acid, acacetin, ferulic acid, kampferol, genistein ([Ali et al., 2014](#)), coumaric acid ([Khan et al., 2017](#)) tannins (punicalin, pedunculagin, GA and ellagic acid) ([Ismail et al., 2012](#); [Pagliarulo et al., 2016](#)), and quercetin ([Saxena et al., 2017](#)). To gain more insight, we docked all these reported compounds against LuxR analogues of *C. violaceum* and *P. aeruginosa*, against which our extract was found to act as signal-response inhibitor. Results of this molecular docking exercise are presented in [Table 3](#). The amino acid residues involved in this on-screen ligand-receptor binding, for compounds

used in wet-lab study, are shown in extended data, [Table S1](#) ([Joshi et al., 2018](#)).

Pure compounds against *C. violaceum*. Against the LuxR analogue of *C. violaceum*, docking scores higher than that of its natural ligand (C6-HSL) were obtained for acacetin, catechin, and quercetin. However, for further *in vitro* experiments, we did not set any cut-off value with respect to docking score, instead we experimented with all those pure compounds from those listed in [Table 2](#), which were available in our lab viz. catechin, quercetin, chlorogenic acid, cinnamic acid, and GA. Cinnamic acid had no effect on *C. violaceum* growth and pigment production in the concentration range 5–100 µg/ml, except 26% reduction in violacein production at 100 µg/ml ([Figure 5A](#)). Hence all subsequent experiments with cinnamic acid were performed at 100 µg/ml. Cinnamic acid was able to enhance *C. violaceum* susceptibility to streptomycin by 11.94%, but had no effect on its susceptibility to cephalixin and tetracycline ([Figure 5C](#)). Haemolytic potential and catalase activity of *C. violaceum* were curbed to a smaller (8.47% and 2.68%, respectively), albeit statistically significant extent ([Table 3](#)). This phenylpropanoid metabolite was also able to confer some survival benefit (24% lesser worm death) on *C. elegans* ([Figure 5D](#)). Cinnamic acid

Table 2. Docking score of compounds with various targets.

Compound	PubChem CID	Binding affinity (kcal/mol)		
		<i>C. violaceum</i> (CviR)	<i>P. aeruginosa</i> (LasR)	<i>P. aeruginosa</i> (PqsR)
(Natural ligand) Autoinducer ^a	C6-HSL: 3462373 C4-HSL: 11513882 PQS: 2763159	-7.6	-7.2	-6.2
Acacetin	5280442	-9.0	-5.9	-6.9
Benzoic acid	243	-7.1	-6.1	-5.0
Catechin	73160	-8.5	-9.8	-6.8
Chlorogenic acid	1794427	-7.3	-5.7	-7.5
Cinnamic acid	444539	-7.1	-6.9	-4.7
Coumaric acid	637542	-6.4	-7.1	-5.3
Ellagic acid	5281855	-6.8	-8.8	-6.6
Ferulic acid	445858	-6.4	-7.5	-5.7
Gallic acid	370	-5.9	-6.3	-5.8
Genistein	5280961	-6.3	-9.4	-6.5
Kaempferol	5280863	-6.8	-7.2	-6.9
Pedunculagin	442688	-7.6	-6.4	-8.1
Punicalagin	44584733	-7.0	-7.4	-9.3
Punicalin	5388496	-7.4	-6.9	-8.3
Pyrogallol	1057	-5.2	-6.1	-5.1
Quercetin	5280343	-8.5	-9.9	-7.2
Rutin	5280805	-7.3	-6.7	-8.2

^aC6-HSL autoinducer of *C. violaceum*, C4-HSL autoinducer of Las QS system of *P. aeruginosa*, and PQS quinolone signal of PQS system of *P. aeruginosa* were used for docking as control.

Figure 5. Effect of cinnamic acid and chlorogenic acid on *C. violaceum*. 'Control' in this figure is the vehicle control (0.5%v/v DMSO), which did not exert any effect on growth and violacein production of *C. violaceum*. (A) Effect of cinnamic acid on growth and violacein production in *C. violaceum*. Bacterial growth was measured as OD₇₆₄; OD of violacein was measured at 585 nm, and Violacein Unit was calculated as the ratio OD₅₈₅/OD₇₆₄ (an indication of violacein production per unit of growth). (B) Effect of chlorogenic acid on growth and violacein production in *C. violaceum*. Bacterial growth was measured as OD₇₆₄; OD of violacein was measured at 585 nm, and violacein unit was calculated as the ratio OD₅₈₅/OD₇₆₄ (an indication of violacein production per unit of growth). (C) Pre-treatment of *C. violaceum* with chlorogenic acid and cinnamic acid enhances its susceptibility to streptomycin and cephalaxin. (D) Cinnamic acid, chlorogenic acid, and catechin reduced virulence of *C. violaceum* towards *C. elegans*. Catechin (50 μg/ml) and ampicillin (500 μg/ml) employed as positive controls conferred 100% protection on worm population. DMSO present in the 'vehicle control' at 0.5%v/v did not affect virulence of the bacterium towards *C. elegans*; DMSO (0.5%v/v) and compounds at tested concentrations showed no toxicity towards the worm. **p*<0.05, ***p*<0.01, ****p*<0.001. AS, antibiotic susceptibility; QS, quorum sensing.

induced violacein inhibition was not reversed upon exogenous supply of AHL (extended data, Figure S2) (Joshi *et al.*, 2018), indicating its mode of QS-inhibition to be signal-response inhibition.

Chlorogenic acid could not affect *C. violaceum* growth and pigment production at any of the test concentrations (5–100 μg/ml), except 16.21% inhibition of violacein production at 75 μg/ml (Figure 5B). Notably, violacein production was not affected at higher concentration (i.e. 100 μg/ml). Hence, all subsequent experiments with chlorogenic acid were performed at 75 μg/ml. At this concentration chlorogenic acid could make *C. violaceum* 8.91% more susceptible to streptomycin; however, it did not alter this bacterium's susceptibility to cephalaxin and tetracycline (Figure 5C). Chlorogenic acid was also able to inhibit catalase activity and haemolytic activity of *C. violaceum* by 3.84% and

10.16% respectively (Table 3). It could also offer a sizable (49%) survival benefit to *C. elegans*, when challenged with *C. violaceum* (Figure 5D). AHL augmentation of the *C. violaceum* culture experiencing chlorogenic acid induced QS inhibition, did not result in reversal of this inhibition (extended data, Figure S2) (Joshi *et al.*, 2018), indicating this plant metabolite to act as a signal-response inhibitor.

GA was tested at 10–200 μg/ml for its potential quorum modulatory action against *C. violaceum*, and it was found to inhibit the growth of this bacterium in a dose-dependent fashion (Figure 6A). Violacein production was negatively affected from 100 μg/ml onwards, and this inhibition was reversed upon external AHL supply (Figure 6B). Since GA was indicated to act as a signal-supply inhibitor, we docked it against the LuxI analogue of *C. violaceum* i.e. CviI, the enzyme responsible

AH
 -6.9
 uce

en
–20
a pr
n w
duct
in
cati
erva
que
the
cata
erve

echi
erea
ward
bitor
y eff
men

in
ha
sta
Ir
all
Q
in
(F

Pr
in,
qu
re
ag
ba
ac
ef
fe
fo
Pr
ch
P.
ho

Figure 6. Effect of GA, quercetin, and catechin on *C. violaceum*. Bacterial growth was measured as OD₇₆₄; OD of violacein was measured at 585 nm, and Violacein Unit was calculated as the ratio OD₅₈₅/OD₇₆₄ (an indication of violacein production per unit of growth). 'Control' in this figure is the vehicle control (0.5% v/v DMSO), which did not exert any effect on growth and violacein production of *C. violaceum*. (A) Effect of gallic acid on growth and violacein production in *C. violaceum*. (B) GA acts as a signal-supply inhibitor. (C) Effect of quercetin on growth and violacein production in *C. violaceum*. (D) Quercetin acts as a signal-response inhibitor. (E) Effect of catechin on growth and violacein production in *C. violaceum*. (F) Catechin acts as a signal-response inhibitor. *p<0.05, **p<0.01, ***p<0.001. QS, quorum sensing; GA, gallic acid.

to ofloxacin and gentamicin. With respect to antibiotic susceptibility result obtained with cinnamic acid (Figure 7E) were identical to those with chlorogenic acid. AHL augmentation was able to reverse effect of chlorogenic as well as cinnamic acid on production of both the QS-regulated pigments of

P. aeruginosa (Figure 7D), suggesting these compounds to disturb the QS machinery of this pathogen by acting as signal-supply inhibitors. Both these compounds were able to negate haemolytic ability of *P. aeruginosa* notably (19.11%; $p=0.04$ and 35.29%; $p=8.31363E-05$ reduction respectively by cinnamic acid

and chlorogenic acid), whereas catalase activity was enhanced to a marginal (1.47%) extent only by chlorogenic acid. *In vivo* efficacy of chlorogenic acid was found to be better than that of cinnamic acid (Figure 7F).

Catechin could reduce production of pyocyanin and pyoverdine both, in *P. aeruginosa*, without affecting its growth (Figure 7C). However, there was not much difference in the effects exerted by different concentrations of catechin. For

Figure 7. Effect of pure compounds on QS-regulated traits of *P. aeruginosa*. Bacterial growth was measured as OD₇₆₄; OD of pyoverdine was measured at 405 nm, and Pyoverdine Unit was calculated as the ratio OD₄₀₅/OD₇₆₄ (an indication of pyoverdine production per unit of growth), Pyocyanin Unit was calculated as the ratio OD₅₂₀/OD₇₆₄ (an indication of pyocyanin production per unit of growth). 'Control' in this figure is the vehicle control (0.5%v/v DMSO), which did not exert any effect on growth and pigment production of *P. aeruginosa*. (A) Effect of cinnamic acid on growth and QS-regulated pigment production in *P. aeruginosa*. (B) Effect of chlorogenic acid on growth and QS-regulated pigment production in *P. aeruginosa*. (C) Effect of catechin on growth and QS-regulated pigment production in *P. aeruginosa*. (D) Cinnamic acid and chlorogenic acid act as signal-supply inhibitors against *P. aeruginosa*, whereas catechin acts as a signal-response inhibitor. (E) *P. aeruginosa* receiving pre-incubation with cinnamic acid or chlorogenic acid becomes more susceptible to cephalaxin. (F) Pre-treatment of *P. aeruginosa* with cinnamic acid or chlorogenic acid reduces its virulence towards *C. elegans*: Catechin (50 μg/ml) and gentamicin (0.1 μg/ml) employed as positive controls conferred 100% and 80% protection on worm population respectively; DMSO present in the 'vehicle control' at 0.5%v/v did not affect virulence of the bacterium towards *C. elegans*; DMSO (0.5%v/v) and compounds at tested concentrations showed no toxicity towards the worm. **p*<0.05, ***p*<0.01, ****p*<0.001; AS, Antibiotic susceptibility; QS, Quorum sensing.

example, the effect on pyoverdine production exerted by all concentrations in the range 10-100 µg/ml was statistically same; and so was the case for the effect exerted by catechin on pyocyanin production in the concentration range of 100-200 µg/ml. Catalase and haemolytic activity of *P. aeruginosa* were both negatively affected by catechin (Table 3). Catechin seemed to act as a signal-response inhibitor against *P. aeruginosa* (Figure 7D).

GA and quercetin were tested against *P. aeruginosa* at 10-200 µg/ml. Both these compounds were able to inhibit the bacterial growth, and this growth inhibitory effect was more profound in case of quercetin. Production of both the QS-regulated pigments in *P. aeruginosa* was enhanced by GA. Quercetin had a negative effect on pyocyanin production, whereas pyoverdine production was enhanced by it from

10 µg/ml onwards. Catalase activity remained unaffected in GA-treated *P. aeruginosa* culture, whereas it was affected to a minor extent in quercetin-treated culture. Both of these plant compounds were able to reduce biofilm-forming capacity of *P. aeruginosa* notably (Figure 8A, B).

Growth promoting effect of PGPE on probiotic strains. PGPE was able to enhance growth of *L. plantarum* and *B. bifidum* at 10-50 µg/ml. Higher concentration (100 µg/ml) did not seem to promote their growth further (Figure 9).

Discussion

Our *in vitro* and *in vivo* experiments indicated hydroalcoholic extract of *P. granatum* peel to possess appreciable antipathogenic activity against both the test bacteria. Antimicrobial activity of pomegranate extracts against bacteria, fungi, and

Figure 8. Effect of GA and quercetin on various traits of *P. aeruginosa*, in vitro. 'Control' in this figure is the vehicle control (0.5%v/v DMSO), which did not exert any effect on growth and pigment production of *P. aeruginosa*; Bacterial growth was measured as OD₇₆₄; OD of pyoverdine was measured at 405 nm, and Pyoverdine Unit was calculated as the ratio OD₄₀₅/OD₇₆₄ (an indication of pyoverdine production per unit of growth), Pyocyanin Unit was calculated as the ratio OD₅₂₀/OD₇₆₄ (an indication of pyocyanin production per unit of growth); Crystal violet assay was performed to measure biofilm formation, and biofilm eradication, followed by the measurement of OD at 580 nm. (A) Effect of GA on growth and QS-regulated pigment production in *P. aeruginosa*. (B) Effect of quercetin on growth and QS-regulated pigment production in *P. aeruginosa*. *p<0.05, **p<0.01, ***p<0.001; AS, antibiotic susceptibility; QS, quorum sensing; GA, gallic acid.

Figure 9. Growth promoting effect of *Punica granatum* peel extract on probiotic strains. Bacterial growth was measured as OD₆₆₀. *p<0.05, **p<0.01, ***p<0.001. PGPE: *Punica granatum* peel extract.

plasmodium has previously been reported by many researchers (Rahmani *et al.*, 2017). Other researchers have reported *P. granatum* extracts to possess antimicrobial and/or anti-QS effect. However almost all of them have shown the effective concentrations to be much higher than what we report for PGPE in the present study. Ethanolic extracts of pomegranate were shown to inhibit methicillin-resistant *S. aureus* isolates at 200–400 µg/ml (Voravuthikunchai & Kitpipit, 2005). A high tannin pomegranate extract was reported to be effective against MRSA at 1–5 mg/ml (Su *et al.*, 2012). *Helicobacter pylori* isolates were shown to be inhibited by pomegranate extracts at 50–100 µg disc⁻¹. Pomegranate peel extract was reported to inhibit *Salmonella typhimurium* and *Salmonella aureus* on meat surfaces, at 250 µg/ml (Tayel *et al.*, 2012). Effective concentrations reported in our study are also lower than the quorum-modulatory concentrations reported by Yang *et al.*, (2016). They reported tannin-rich fraction from pomegranate rind (TFPR) caused a reduction (~50-70%) in *C. violaceum* pigmentation at 0.312–0.625 mg/ml, whereas comparable pigment inhibition was achieved by our extract at 50-150 µg/ml (Figure 1A). Oh *et al.* (2015) reported pomegranate peel extract to inhibit bacterial growth at 3–12 mg/ml. Our extract could inhibit violacein production by 78.26% at 10 times lesser concentration (200 µg/ml; Figure 1A) than the concentration (2 mg/ml) they reported for their extract to achieve a comparable (78.5%) violacein inhibition in *C. violaceum*. Inhibition of biofilm formation of human pathogens by *P. granatum* extract at <40 µg/ml reported by Bakkiyaraj *et al.* (2013) matches with our results of the present study (Figure 2E). Nuamsetti *et al.* (2012) reported pomegranate extracts to exert antibacterial activity against four different bacteria with MIC values reaching 103.6-207 mg/ml.

Much evidence for antimicrobial action of pomegranate extracts has come from *in vitro* assays, whose confirmation through *in vivo* assays is necessary (Howell & D'Souza, 2013). Dazal *et al.* (2015) reported ethanolic extract of *P. granatum* pericarp to inhibit QS in *P. aeruginosa* at 219.78–2197.80 µg/ml; whereas our extract could modulate production of both QS-regulated pigments from 5 µg/ml onwards *in vitro*; notable *in vivo* efficacy was observed at 0.5 µg/ml onwards. An ethanolic extract of pomegranate exhibited MICs of 0.49–1.95 mg/ml and MBCs of 1.95–3.91 mg/ml against *E. coli* O157:H7. Compared to these relatively higher effective concentrations, PGPE reported in the present study exhibited notable (>50% worm survival, as against 12.5-35% in control wells) *in vivo* efficacy at lower (0.5-5 µg/ml) concentrations (Figure 1D and Figure 2F).

Higher anti-infective efficacy shown by our extract reported in this study may be attributed to the extraction method employed by us i.e. MAE. This method is known to be capable of fast extraction of phenolics Proestos & Komaitis, (2008), and also suitable for heat-labile phytoconstituents (Gupta *et al.*, 2012). The fact that same extract when prepared using different extraction methods, can vary with respect to its efficacy, has previously also been emphasized by us (Kothari *et al.*, 2012).

This study has found PGPE to be an effective quorum modulator against two gram-negative bacterial pathogens. An

ideal quorum-modulator is expected to have no or minimal effect on bacterial growth, and thus to exert lesser selection pressure on the target pathogenic population. PGPE reported in this study was observed to exert a purely quorum-modulatory effect against *C. violaceum* till 10 µg/ml, and *P. aeruginosa* till 2.5 µg/ml. It is generally expected that owing to lesser effect on bacterial growth, bacterial populations are likely to develop resistance to quorum-targeting antimicrobials at a slower pace. Further, in the case of plant extracts, there is always a possibility of multiple phytochemicals being simultaneously responsible for the observed antipathogenic effect, and they may be targeting multiple cellular components in the target bacteria, hence the susceptible pathogen population may find it difficult to develop resistance against them. Precisely this was observed in our study with *P. aeruginosa*, wherein even repeated exposure of this bacterium to PGPE did not induce resistance, and even the PGPE-habituated *P. aeruginosa* remained susceptible to this extract, *in vitro* (Figure 2G) as well as *in vivo* (Figure 2F).

One of the frequently observed problem with conventional antibiotic treatments is that besides killing the target pathogen, they simultaneously disturb the normal human microbiota, which may lead to the condition of dysbiosis. Hence, it is desirable from an anti-infective preparation that it should not have any negative effect on the normal microbiota organisms. This study has found PGPE to promote growth of two bacteria (*L. plantarum* and *B. bifidum*), which are part of normal human gut biota (Figure 9). From these results, PGPE can be said to have appreciable prebiotic potential. Li *et al.* (2015) had also indicated pomegranate whole fruit extract to work as a potential prebiotic, with beneficial effect on *Bifidobacterium* and *Lactobacillus*. Pomegranate extracts can be expected to exert meaningful prebiotic and anti-infective effect *in vivo* in human body, as they are rich in ellagitannins, which are hydrolysed in the gut to ellagic acid, which further is metabolized by the colon microbiota to form urolithin A and urolithin B. These urolithins at micromolar concentrations were indicated to be potent *in vivo* QS-inhibitors by Giménez-Bastida *et al.* (2012). Prebiotic preparations with bifidogenic property, besides improving gastrointestinal function, can also find applications in management of conditions like depression and psychological distress (Messaoudi *et al.*, 2011). Overall GI tract benefits expected from *P. granatum* extracts, as indicated in ethnomedicinal records (Colombo *et al.*, 2013; Hollebeek *et al.*, 2012) can be in part due to its prebiotic potential, which makes use of pomegranate in management of inflammatory bowel disease appear relevant.

Our results regarding effect of PGPE on haemolytic activity of both the test bacteria, and its effect on *P. aeruginosa* susceptibility to human serum also indicate the high probability of this extract to be of therapeutic relevance. Haemolysis has been considered as an important virulence factor, and infection by haemolytic bacteria may lead to severe anaemia (Orf & Cunningham, 2015). By reducing the haemolytic activity of the pathogens, PGPE-like extracts can significantly reduce iron-availability for the pathogens, as lysis of red blood cells is one of the effective strategies for many pathogens to ensure sufficient supply of iron, necessary for their survival inside

host. Enhanced susceptibility of *P. aeruginosa*, under influence of PGPE, to lysis by human serum is also of clinical significance, as this can aid the clearance of pathogenic bacteria by human body, while facing bacteremia. This may represent an additional possible mechanism by which PGPE may confer *in vivo* protection against bacterial infection.

Any quorum-inhibitory substance may act either by targeting the signal-synthesis machinery, sequestering the already synthesized signals, or by interfering with the signal-reception (Zhang & Li, 2016). Though different phytochemicals of a given crude extract may have different mode of action, as a whole extract PGPE was found to act as a signal-response inhibitor against *C. violaceum* (Figure 1B) and *P. aeruginosa* (Figure 2B). Among the pure compounds tested in this study, except GA, remaining four compounds seemed to act as a signal-response inhibitor against *C. violaceum*. Notably, cinnamic acid and chlorogenic acid were found to behave as signal-supply inhibitors against *P. aeruginosa*, and as signal-response inhibitor against *C. violaceum*. This suggests that same plant compound may act differently against different bacteria. Here this fact becomes even more interesting, given both *C. violaceum* and *P. aeruginosa* are gram-negative bacteria, and a heavy overlap is suggested to be there among the QS machinery of different members of gram-negative bacterial group (Papenfort & Bassler, 2016). Since PGPE contains a mixture of signal-supply and -response inhibitors, it can be difficult for the susceptible bacteria to develop simultaneous resistance to both mechanisms (i.e. against the whole extract). Compared to this, it may be easier for the bacteria to develop resistance against single-molecule antibiotics with bactericidal action.

At 0.5-1 µg/ml PGPE showed *in vitro* effect neither on growth, nor on pigment production in any of the test bacteria. In case of *C. violaceum*, any *in vitro* effect was not observed till 2.5 µg/ml. However, all these concentrations that were 'not effective' *in vitro* were able to significantly reduce virulence of these bacterial pathogens towards *C. elegans* (Figure 1D and Figure 2F). This fact highlights the necessity of assaying any test extract at broadest possible concentration range *in vivo*, including at least one level below the lowest effective *in vitro* concentration. Usually *in vitro* screens are performed at a broader concentration range to identify the most effective concentration(s), and *in vivo* assays are conducted at this narrowed concentration range; though logical, this approach carries an inherent risk of missing out some concentrations meaningful *in vivo*. A simple explanation which can be proposed for this apparent mismatch between *in vitro* and *in vivo* results is that, *in vitro* we look for effect of test extract on a limited number of parameters (e.g. only two parameters, cell density and pigment production, in current study), whereas *in vivo* assays provide the test extract an opportunity to interact with multiple targets in host as well as pathogen. We may not be able to study all these targets, but the overall effect can be detected in terms of survival benefit of the model host. Even the magnitude of difference in *in vitro* effects of different concentrations, may not always match with their *in vivo* effect. For example, *in vitro* pyoverdine by *P. aeruginosa* was overproduced 2.51-fold more at 50 µg/ml than that at 25 µg/ml PGPE; but there was no significant difference in the *in vivo* efficacy of these two

concentrations. On the same line, *in vitro* effect of PGPE at 25-100 µg/ml on *C. violaceum* pigment production was 42-56%, whereas *in vivo* effect of these concentrations fell within a narrower range of 75-80% (i.e. a 5% *in vivo* difference for 14% *in vitro* difference in efficacy values).

Results of this study also suggest that while screening natural/synthetic compounds for their possible effect on bacterial QS, what we should observe for is 'quorum modulation' and not just 'quorum inhibition'. This is particularly evident from the effect of PGPE on pyoverdine production in *P. aeruginosa*, wherein this virulence factor production is not inhibited by PGPE, but promoted to quite notable extent, and this pyoverdine-enhancing (modulatory) effect of PGPE does not prevent it from being effective *in vivo*. Similar enhancement in pigment production by *Serratia marcescens* challenged with an anti-infective quorum-modulatory polyherbal *Panchvalkal* formulation was reported by us earlier (Patel *et al.*, 2017), and the formulation was effective *in vivo*.

While investigating the effect of PGPE and/or its constituent phytochemicals against the two gram-negative bacteria in this study, the dose-response relationship was not always found to be linear. Since dose-response relationships can be described by various models (Calabrese, 2004), we made an effort to see, which model fits best to results obtained in this study. For example, effect of PGPE on *C. violaceum* growth did not follow a linear pattern. Concentrations till 2.5 µg/ml affected neither cell density nor pigment production (Figure 1A); this phenomenon can be said to fit in the threshold model, wherein the biological effect is not observed until the threshold concentration is crossed. This concept of threshold model also seems applicable while describing effect of PGPE on *P. aeruginosa* cell density and pigment production, as none of the parameters seem to be affected until the threshold concentration of 1 µg/ml was not crossed (Figure 2A). At higher concentrations, the effect on pyoverdine production seemed to somewhat follow an inverted U-shaped hormetic model. Whether such a pattern would have continued at still higher concentrations, cannot be predicted without actually performing experiment at those concentrations. Similar analysis with respect to pure compounds revealed that effect of GA on *C. violaceum* cell density and pigment production (Figure 6A) can be said to follow the linear non-threshold model, within the tested concentration range. The term 'hormesis' is used in general to describe a biphasic dose response (Mattson, 2008), and this concept can be relevant while describing 'adaptive stress response' in bacteria challenged with some antimicrobial/quorum modulatory agent. Hormetic response can be said to occur (or not to occur) within the concentration range actually tested during a particular experiment, but it need not to be extrapolated into the realm of uncertain i.e. concentrations not really tried, on either side of the test concentration range.

Overall, the whole extract seemed to be more efficacious than any of the pure compounds tested. For example, at 10 µg/ml, PGPE affected pyoverdine production in *P. aeruginosa* much more than either GA or quercetin. Such synergistic action of the pomegranate constituents, apparently being superior to that of its individual constituents, has been indicated in literature

(Jurenka, 2008). Among the many constituents of PGPE, GA in this study was found to act as a signal-supply inhibitor against *C. violaceum*, whereas all other compounds as well as the whole extract were found to act as signal-response inhibitor against this bacterium. There may be few other unknown compounds present in this extract, which may either inhibit AHL synthesis, sequester synthesized AHLs, or bind to the LuxR component. It is likely to be difficult for the bacteria to develop complete resistance against any such poly-component preparation, as different compounds will target different components of the bacterial QS machinery. Anand *et al.* (2013) has also indicated the combination of LuxR non-competitive inhibitors and LuxI inhibitors, as a robust therapeutic strategy to act multiplicatively across a broad parameter range.

In silico docking exercise indicated both, quercetin and, catechin to have affinity for CviR more than its natural ligand C6-HSL (Table 2), i.e. a high probability of them being capable of acting as signal-response inhibitor. *In vitro* experiments validated this possibility (Figure 6D, F). Though the docking scores of both these phytochemicals against CviR were identical, *in vitro* violacein production was inhibited bit more by quercetin. Gallic acid, which showed second lowest affinity (-5.9 kcal/mol) for CviR among all docked phytochemicals, was also not able to act as signal-response inhibitor, instead it acted as signal-supply inhibitor against *C. violaceum*, and its docking score against the signal synthesizing enzyme AHL synthase was also better i.e. (-7.9 kcal/mol). Cinnamic acid displayed lowest *in silico* binding affinity against *P. aeruginosa* receptor protein PqsR (Table 2), and this corroborates well with the fact revealed from *in vitro* experiment that it acts as a signal-supply inhibitor (Figure 7D). Though the binding affinity of catechin and quercetin both, against *P. aeruginosa* protein LasR was identical as per docking score, indicating a high probability of them acting as a signal-response inhibitor with respect to pyoverdine, whose production is controlled by the las system, and *in vitro* experiments did find catechin to act as a signal-response inhibitor (Figure 7D); pyoverdine production was affected much more by quercetin than by catechin. If we focus on docking scores of only those five compounds, with which *in vitro* experiments were also conducted in this study, against both the *P. aeruginosa* receptors, quercetin gives an impression of being more effective than GA, and *in vitro* experiments also show quercetin to have more effect on pigment production by this bacterium, than GA (Figure 8A and 8B). Similarly docking scores of catechin against both *P. aeruginosa* receptor proteins were better than that of cinnamic acid (Table 2), and *in vivo* assay also found catechin to be more efficacious than cinnamic acid (Figure 7F). Overall there seems to be an empirical match between the *in silico* and wet-lab experiments.

This study has found PGPE to be an effective quorum-modulator against two different antibiotic resistant strains of gram-negative bacteria. In general, it is more challenging to find antimicrobials effective against gram-negative bacteria, than against gram-positive bacteria, mainly for the reason that the former possess an additional barrier in form of outer membrane, posing extra-challenge to the intracellular entry of any antimicrobial. Besides affecting QS-regulated pigment production in the test bacteria, this extract could also effectively modulate other traits, including

catalase activity, haemolytic activity, biofilm formation, susceptibility of bacteria to antibiotics and to lysis by human-serum factors. Though researchers have reported the antimicrobial potential of *P. granatum* extracts (Al-Zoreky *et al.*, 2009; Choi *et al.*, 2011; Prashanth *et al.*, 2001) or nanoparticles derived from it (Devanesan *et al.*, 2018), reports describing *in vivo* anti-infective efficacy of PGPE with focus on its quorum modulatory potential are still warranted. This extract was not found to inhibit growth of test bacteria too heavily, at effective quorum-modulatory concentrations, and this is what is expected from a 'good' quorum-modulator (i.e. having no or minimal effect on growth of the test bacterium, and exerting no great selection pressure on the pathogen population).

Further studies regarding elucidation of the molecular mechanism associated with such appreciable anti-infective activity of PGPE is warranted, as that will give us a better insight into its mode of action. This study also highlights the importance of using different phytochemicals in combination (e.g. as a whole extract; concept of synergy), rather than always focusing on the single-active-molecule approach. Results reported here can be said to be supportive of the concept of polyherbalism (Parasuraman *et al.*, 2014) widely employed by the practitioners of complementary and alternative medicine.

Data availability

Underlying data

All raw data containing the underlying data behind each figure and table are available on figshare. DOI: <https://doi.org/10.6084/m9.figshare.7471979> (Joshi *et al.*, 2018).

Extended data

Extended data are available on figshare. DOI: <https://doi.org/10.6084/m9.figshare.7471979> (Joshi *et al.*, 2018).

Figure S1. Antipathogenic effect of Pomella on *P. aeruginosa*.

Figure S2. Quorum-inhibitory effect of cinnamic and chlorogenic acid on violacein production in *C. violaceum* is not reversed following augmentation of the quorum-inhibited culture by AHL

Table S1. Binding affinity of different phytochemicals when docked against LuxR homologue of test pathogens.

Data are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

Grant information

This work received no extramural funding, and was fully supported by our departmental research budget (Institute of Science, Nirma University).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

Authors thank NERF (Nirma Education and Research Foundation), Ahmedabad for financial and infrastructural support; Viridis

Biopharma Pvt. Ltd., Mumbai for HPLC analysis of PGPE; Pharmanza Herbal Pvt. Ltd. for consenting to mention data on their product 'Pomella'; Dr. Archana Mankad, Department of

Botany, Gujarat University for authentication of plant material; and Microbiology Department, M.G. Science Institute, Ahmedabad for providing culture of *P. aeruginosa*.

References

- Adonizio A, Kong KF, Mathee K: **Inhibition of quorum sensing-controlled virulence factor production in *Pseudomonas aeruginosa* by South Florida plant extracts.** *Antimicrob Agents Chemother.* 2007; **52**(1): 198–203.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Alanis AD, Calzada F, Cervantes JA, *et al.*: **Antibacterial properties of some plants used in Mexican traditional medicine for the treatment of gastrointestinal disorders.** *J Ethnopharmacol.* 2005; **100**(1–2): 153–157.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ali SI, El-Baz FK, El-Emary GAE, *et al.*: **HPLC-Analysis of polyphenolic compounds and free radical scavenging activity of pomegranate fruit (*Punica granatum* L.).** *Int J Pharm Clin Res.* 2014; **6**: 348–355.
[Reference Source](#)
- Al-Zoreky NS: **Antimicrobial activity of pomegranate (*Punica granatum* L.) fruit peels.** *Int J Food Microbiol.* 2009; **134**(3): 244–248.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Anand R, Rai N, Thattai M: **Interactions among quorum sensing inhibitors.** *PLoS One.* 2013; **8**(4): e62254.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bakkiyaraj D, Nandhini JR, Malathy B, *et al.*: **The anti-biofilm potential of pomegranate (*Punica granatum* L.) extract against human bacterial and fungal pathogens.** *Bioloouling.* 2013; **29**(8): 929–937.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Beceiro A, Tomás M, Bou G: **Antimicrobial resistance and virulence: a successful or deleterious association in the bacterial world?** *Clin Microbiol Rev.* 2013; **26**(2): 185–230.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Calabrese EJ: **Hormesis: a revolution in toxicology, risk assessment and medicine.** *EMBO Rep.* 2004; **5** Spec No: S37–S40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chang CY, Krishnan T, Wang H, *et al.*: **Non-antibiotic quorum sensing inhibitors acting against *N*-acyl homoserine lactone synthase as druggable target.** *Sci Rep.* 2014; **4**: 7245.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Choi JG, Kang OH, Lee YS, *et al.*: **In Vitro and In Vivo Antibacterial Activity of *Punica granatum* Peel Ethanol Extract against *Salmonella*.** *Evid Based Complement Alternat Med.* 2011; **2011**: 690518.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Choo JH, Rukayadi Y, Hwang JK: **Inhibition of bacterial quorum sensing by vanilla extract.** *Lett Appl Microbiol.* 2006; **42**(6): 637–641.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Colombo E, Sangiovanni E, Dell'Agli M: **A review on the anti-inflammatory activity of pomegranate in the gastrointestinal tract.** *Evid Based Complement Alternat Med.* 2013; **2013**: 247145.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dazal MV, Gonzales E, Sabino AK, *et al.*: **Controlling biofilm formation by inhibiting the quorum-sensing activity of *Pseudomonas aeruginosa* using the ethanolic extracts of *Piper nigrum* (Piperaceae) fruit, *Punica granatum* (Lythraceae) Pericarp, and *Pisum sativum* (Fabaceae) seed.** *Int J Pharm Sci Drug Res.* 2015; **7**(4): 349–353.
[Reference Source](#)
- Devanesan S, AlSalhi MS, Balaji RV, *et al.*: **Antimicrobial and Cytotoxicity Effects of Synthesized Silver Nanoparticles from *Punica granatum* Peel Extract.** *Nanoscale Res Lett.* 2018; **13**(1): 315.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Duraipandian V, Ayyanar M, Ignacimuthu S: **Antimicrobial activity of some ethnomedicinal plants used by Palayir tribe from Tamil Nadu, India.** *BMC Complement Altern Med.* 2006; **6**: 35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- El-Fouly MZ, Sharaf AM, Shahin AM, *et al.*: **Biosynthesis of pyocyanin pigment by *Pseudomonas aeruginosa*.** *J Radiat Res Appl Sci.* 2015; **8**(1): 36–48.
[Publisher Full Text](#)
- Eng SA, Nathan S: **Curcumin rescues *Caenorhabditis elegans* from a *Burkholderia pseudomallei* infection.** *Front Microbiol.* 2015; **6**: 290.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fair RJ, Tor Y: **Antibiotics and bacterial resistance in the 21st century.** *Perspect Medicin Chem.* 2014; **6**: 25–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ferro TA, Araújo JM, Dos Santos Pinto BL, *et al.*: **Cinnamaldehyde Inhibits *Staphylococcus aureus* Virulence Factors and Protects against Infection in a *Galleria mellonella* Model.** *Front Microbiol.* 2016; **7**: 2052.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gharzouli K, Khennouf S, Amira S, *et al.*: **Effects of aqueous extracts from *Quercus ilex* L. root bark, *Punica granatum* L. fruit peel and *Artemisia herba-alba* Asso leaves on ethanol-induced gastric damage in rats.** *Phytother Res.* 1999; **13**(1): 42–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Giménez-Bastida JA, González-Sarriás A, Larrosa M, *et al.*: **Ellagitannin metabolites, urolithin A glucuronide and its aglycone urolithin A, ameliorate TNF- α -induced inflammation and associated molecular markers in human aortic endothelial cells.** *Mol Nut Food Res.* 2012; **56**(5): 784–796.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gupta A, Naranimal M, Kothari V: **Modern extraction methods for preparation of bioactive plant extracts.** *Int J Appl Nat Sci.* 2012; **1**(1): 8–26.
[Reference Source](#)
- Hajimahmoodi M, Shams-Ardakani M, Saniee P, *et al.*: **In vitro antibacterial activity of some Iranian medicinal plant extracts against *Helicobacter pylori*.** *Nat Prod Res.* 2011; **25**(11): 1059–1066.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hollebeeck S, Larondelle Y, Schneider YJ, *et al.*: **The use of pomegranate (*Punica granatum* L.) phenolic compounds as potential natural prevention against IBDs.** *In Inflammatory bowel disease-advances in pathogenesis and management.* InTech. 2012.
[Publisher Full Text](#)
- Hoseini Farzaei M, Abbasabadi Z, Reza Shams-Ardakani M, *et al.*: **A comprehensive review of plants and their active constituents with wound healing activity in traditional Iranian medicine.** *Wounds.* 2014; **26**(7): 197–206.
[PubMed Abstract](#)
- Howell AB, D'Souza DH: **The pomegranate: effects on bacteria and viruses that influence human health.** *Evid Based Complement Alternat Med.* 2013; **2013**: 606212.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hui YW, Dykes GA: **Modulation of cell surface hydrophobicity and attachment of bacteria to abiotic surfaces and shrimp by Malaysian herb extracts.** *J Food Prot.* 2012; **75**(8): 1507–1511.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ismail T, Sestili P, Akhtar S: **Pomegranate peel and fruit extracts: a review of potential anti-inflammatory and anti-infective effects.** *J Ethnopharmacol.* 2012; **143**(2): 397–405.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Iwase T, Tajima A, Sugimoto S, *et al.*: **A simple assay for measuring catalase activity: a visual approach.** *Sci Rep.* 2013; **3**: 3081.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Joshi C, Kothari V, Patel P: **Importance of Selecting Appropriate Wavelength, While Quantifying Growth and Production of Quorum Sensing Regulated Pigments in Bacteria.** *Recent Pat Biotechnol.* 2016; **10**(2): 145–152.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Joshi C, Patel P, Kothari V: **Anti-infective potential of hydroalcoholic extract of *Punica granatum* peel against gram-negative bacterial pathogens.** *Figshare.* Fileset. 2018.
<http://www.doi.org/10.6084/m9.figshare.7471979.v2>
- Jurenka JS: **Therapeutic applications of pomegranate (*Punica granatum* L.): a review.** *Altern Med Rev.* 2008; **13**(2): 128–144.
[PubMed Abstract](#)
- Khan I, Rahman H, Abd El-Salam NM, *et al.*: ***Punica granatum* peel extracts: HPLC fractionation and LC MS analysis to quest compounds having activity against multidrug resistant bacteria.** *BMC Complement Altern Med.* 2017; **17**(1): 247.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Khare CP: **Indian herbal remedies: rational Western therapy, ayurvedic, and other traditional usage.** Botany. Springer science & business media. 2004.
[Publisher Full Text](#)
- Kohanski MA, Dwyer DJ, Collins JJ: **How antibiotics kill bacteria: from targets to networks.** *Nat Rev Microbiol.* 2010; **8**(6): 423–435.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kothari V, Gupta A, Naranimal M: **Comparative study of various methods for extraction of antioxidant and antibacterial compounds from plant seeds.** *J Nat Rem.* 2012; **12**(2): 162–173.
[Reference Source](#)
- Kothari V, Punjabi A, Gupta S: **Optimization of microwave-assisted extraction of**

Annona squamosa seeds. *Icfai University J Life Sci.* 2009; 3(1): 55–60.

Reference Source

- Kothari V, Sharma S, Padia D: **Recent research advances on *Chromobacterium violaceum*.** *Asian Pac J Trop Med.* 2017; 10(8): 744–752.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Larrosa M, González-Sarriá A, Yáñez-Gascón MJ, *et al.*: **Anti-inflammatory properties of a pomegranate extract and its metabolite urolithin-A in a colitis rat model and the effect of colon inflammation on phenolic metabolism.** *J Nutr Biochem.* 2010; 21(8): 717–725.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lee J, Zhang L: **The hierarchy quorum sensing network in *Pseudomonas aeruginosa*.** *Protein Cell.* 2015; 6(1): 26–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Li Z, Summanen PH, Komoriya T, *et al.*: **Pomegranate ellagitannins stimulate growth of gut bacteria *in vitro*: Implications for prebiotic and metabolic effects.** *Anaerobe.* 2015; 34: 164–168.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mathabe MC, Nikolova RV, Lall N, *et al.*: **Antibacterial activities of medicinal plants used for the treatment of diarrhoea in Limpopo Province, South Africa.** *J Ethnopharmacol.* 2006; 105(1–2): 286–293.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mattson MP: **Hormesis defined.** *Ageing Res Rev.* 2008; 7(1): 1–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- McClean KH, Winson MK, Fish L, *et al.*: **Quorum sensing and *Chromobacterium violaceum*: exploitation of violacein production and inhibition for the detection of N-acylhomoserine lactones.** *Microbiology.* 1997; 143(Pt 12): 3703–3711.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Messaoudi M, Lalonde R, Violle N, *et al.*: **Assessment of psychotropic-like properties of a probiotic formulation (*Lactobacillus helveticus* R0052 and *Bifidobacterium longum* R0175) in rats and human subjects.** *Br J Nutr.* 2011; 105(5): 755–764.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Moore JD, Rossi FM, Welsh MA, *et al.*: **A Comparative Analysis of Synthetic Quorum Sensing Modulators in *Pseudomonas aeruginosa*: New Insights into Mechanism, Active Efflux Susceptibility, Phenotypic Response, and Next-Generation Ligand Design.** *J Am Chem Soc.* 2015; 137(46): 14626–14639.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nazzaro F, Fratianni F, Coppola R: **Quorum sensing and phytochemicals.** *Int J Mol Sci.* 2013; 14(6): 12607–12619.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Neun BW, Ilinskaya AN, Dobrovolskaia MA: **Analysis of hemolytic properties of nanoparticles.** NCL method ITA-1 Version 1.2, Nanotechnology Characterization Laboratory, Frederick, MD. 2015.
[Reference Source](#)
- Nuamsetti T, Dechayuenyong P, Tantipabulvut S: **Antibacterial activity of pomegranate fruit peels and arils.** *Sci Asia.* 2012; 38(3): 319–322.
[Publisher Full Text](#)
- Oh SK, Chang HJ, Chun HS, *et al.*: **Pomegranate (*Punica granatum* L.) peel extract inhibits quorum sensing and biofilm formation potential in *Yersinia enterocolitica*.** *Microbiol Biotechnol Lett.* 2015; 43–4: 357–366.
[Publisher Full Text](#)
- Orf K, Cunningham AJ: **Infection-related hemolysis and susceptibility to Gram-negative bacterial co-infection.** *Front Microbiol.* 2015; 6: 666.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Pagliarulo C, De Vito V, Picariello G, *et al.*: **Inhibitory effect of pomegranate (*Punica granatum* L.) polyphenol extracts on the bacterial growth and survival of clinical isolates of pathogenic *Staphylococcus aureus* and *Escherichia coli*.** *Food Chem.* 2016; 190: 824–831.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Papenfort K, Bassler BL: **Quorum sensing signal-response systems in Gram-negative bacteria.** *Nat Rev Microbiol.* 2016; 17(9): 576–588.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Parasuraman S, Thing GS, Dhanaraj SA: **Polyherbal formulation: Concept of ayurveda.** *Pharmacogn Rev.* 2014; 8(16): 73–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Patel I, Patel V, Thakkar A, *et al.*: ***Tamarindus indica* (Cesalpiniaceae), and *Syzygium cumini* (Myrtaceae) seed extracts can kill multidrug resistant *Streptococcus mutans* in Biofilm.** *J Nat Remed.* 2013; 13(2): 81–94.
[Reference Source](#)
- Patel P, Joshi C, Palep H, *et al.*: **Anti-infective potential of a quorum modulatory polyherbal extract (*Panchvalka*) against certain pathogenic bacteria.** *J Ayurveda Integr Med.* 2017.
[Publisher Full Text](#)
- Pfaller MA, Sheehan DJ, Rex JH: **Determination of fungicidal activities against yeasts and molds: lessons learned from bactericidal testing and the need for standardization.** *Clin Microbiol Rev.* 2004; 17(2): 268–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Prashanth D, Asha MK, Amit A: **Antibacterial activity of *Punica granatum*.**

Fitoterapia. 2001; 72(2): 171–173.

PubMed Abstract | Publisher Full Text

- Proestos C, Komaltis M: **Application of microwave-assisted extraction to the fast extraction of plant phenolic compounds.** *LWT Food Sci Technol.* 2008; 41(4): 652–659.
[Publisher Full Text](#)
- Rahmani AH, Alsahli MA, Almatroodi SA: **Active constituents of Pomegranates (*Punica granatum*) as potential candidates in the management of health through modulation of biological activities.** *Pharmacogn J.* 2017; 9(5): 689–695.
[Publisher Full Text](#)
- Ramadan MA, Tawfik AF, Shibl AM, *et al.*: **Post-antibiotic effect of azithromycin and erythromycin on streptococcal susceptibility to phagocytosis.** *J Med Microbiol.* 1995; 42(5): 362–366.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ramanuj K, Bachani P, Kothari V: ***In vitro* antimicrobial activity of certain plant products / seed extracts against multidrug resistant *Propionibacterium acnes*, *Malassezia furfur*, and aflatoxin producing *Aspergillus flavus*.** *Res Pharm.* 2012; 2(3): 22–31.
[Reference Source](#)
- [Report]: **Anti-microbial resistance: setting the social science agenda.** Report of an ESRC Working Group. 2014; 1.
[Reference Source](#)
- [Report]: **Global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics.** World Health Organization Report. 2017.
[Reference Source](#)
- Saxena R, Sharma R, Nandy BC: **Chromatographic determination of phenolic profile from *Punica granatum* fruit peels.** *Int Res J Pharm.* 2017; 8(1): 61–65.
[Publisher Full Text](#)
- Shaw PD, Ping G, Daly SL, *et al.*: **Detecting and characterizing N-acyl-homoserine lactone signal molecules by thin-layer chromatography.** *Proc Natl Acad Sci U S A.* 1997; 94(12): 6036–6041.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Su X, Howell AB, D'Souza DH: **Antibacterial effects of plant-derived extracts on methicillin-resistant *Staphylococcus aureus*.** *Foodborne Pathog Dis.* 2012; 9(6): 573–578.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tayel AA, El-Tras WF, Moussa SH, *et al.*: **Surface decontamination and quality enhancement in meat steaks using plant extracts as natural biopreservatives.** *Foodborne Pathog Dis.* 2012; 9(8): 755–761.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Trafny EA, Lewandowski R, Zawistowska-Marciniak I, *et al.*: **Use of MTT assay for determination of the biofilm formation capacity of microorganisms in metalworking fluids.** *World J Microbiol Biotechnol.* 2013; 29(9): 1635–1643.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Trott O, Olson AJ: **AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading.** *J Comput Chem.* 2010; 31(2): 455–461.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Unni K, Priji P, Geoffroy V, *et al.*: ***Pseudomonas aeruginosa* BUP2—A novel strain isolated from malabari goat produces Type 2 pyoverdine.** *Adv Biosci Biotechnol.* 2014; 5(11): 874–885.
[Publisher Full Text](#)
- Vandeputte OM, Kiendrebeogo M, Rajaonson S, *et al.*: **Identification of Catechin as one of the flavonoids from *Combretum albidiflorum* bark extract that reduces the production of quorum-sensing-controlled virulence factors in *Pseudomonas aeruginosa* PAO1.** *Appl Environ Microbiol.* 2010; 76(1): 243–253.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Vasavi HS, Arun AB, Rekha PD: **Inhibition of quorum sensing in *Chromobacterium violaceum* by *Syzygium cumini* L. and *Pimenta dioica* L.** *Asian Pac J Trop Biomed.* 2013; 3(12): 954–959.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Voravuthikunchai SP, Kitpipit L: **Activity of medicinal plant extracts against hospital isolates of methicillin-resistant *Staphylococcus aureus*.** *Clinic Microbiol Infect.* 2005; 11(6): 510–512.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Weydert CJ, Cullen JJ: **Measurement of superoxide dismutase, catalase and glutathione peroxidase in cultured cells and tissue.** *Nat Protoc.* 2010; 5(1): 51–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Yang Q, Wang L, Gao J, *et al.*: **Tannin-Rich Fraction from Pomegranate Rind Inhibits Quorum Sensing in *Chromobacterium violaceum* and Biofilm Formation in *Escherichia coli*.** *Foodborne Pathog Dis.* 2016; 13(1): 28–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zhang W, Li C: **Exploiting Quorum Sensing Interfering Strategies in Gram-Negative Bacteria for the Enhancement of Environmental Applications.** *Front Microbiol.* 2016; 6: 1535.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 25 February 2019

<https://doi.org/10.5256/f1000research.19060.r43898>

© 2019 Kaistha S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Shilpa Deshpande Kaistha

Department of Microbiology, Chhatrapati Shahu Ji Maharaj University (CSJMU), Kanpur, India

This study provides comprehensive experimental evidence for the activity of Punica granatum peel extract in modulating pathogenicity of *Pseudomonas aeruginosa* and quorum sensing pigment producing *Chromobacterium violaceum*. The effect of PGPE on various aspect of pathogenicity related to antibiotic sensitivity, biofilm formation, surface hydrophobicity, quorum sensing, hemolysin & catalase production and *in vivo* *C. elegans* infection model together make a comprehensive and elegant study of the herbal extract activity. The effect of concentration dependent PGPE treatment on various parameters and subsequent generations of the pathogens make for an interesting read. It is recommended that the article be indexed with some minor improvements as suggested below:

Introduction:

At several places, relevant references may be cited.

Material and Methods:

1. Pigment production can be separate heading and subheadings for each pigment, to make clear. Growth measurement can be included in the write up.
2. Details of Chloramphenicol enhanced catalase maybe provided.

Results:

1. The quality of all the figures requires significant improvement. The details of significant values maybe kept in the figures to avoid crowding and enhance understanding. Font size of Figures, for example Figure 2C, needs to be clarified. The standard deviations are not visible in most of the figures.
2. A brief summary of comprehensive results at the end of each section will help the reader grasp the crux of the experiment without having to go through the written text regarding working concentrations of the extracts etc.

Discussion:

1. Requires a section wherein the findings are succinctly concluded regarding the findings for *Ps. aeruginosa* and *C. violaceum*.

2. Please include in Discussion why does the PGPE enhance probiotic activity while showing anti-infective activity against the pathogens studied.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Antimicrobial resistance and biocontrol strategies

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 21 February 2019

<https://doi.org/10.5256/f1000research.19060.r43897>

© 2019 Nandi S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Sisir Nandi

¹ Department of Pharmaceutical Chemistry, Global Institute of Pharmaceutical Education and Research, Affiliated to Uttarakhand Technical University, Kashipur, India

² Uttarakhand Technical University, Kashipur, India

Chinmayi and co-workers studied hydroalcoholic extract of *P. granatum* peel (PGPE), prepared by microwave assisted extraction method for its quorum-modulatory potential against two different human-pathogenic bacteria viz. *Chromobacterium violaceum* and *Pseudomonas aeruginosa*. This is a very significant work to produce antimicrobials. The following minor comments should be

considered before acceptance of this article.

1. "Though the modern medicine has almost always focused on search of purified active molecules as therapeutic leads, knowledge of Indian/Chinese/Arabian traditional medicine makes us aware of a variety of plant extracts and inorganic formulations prescribed for management of infections". Cite the references.
2. "One such widely mentioned item in these ancient systems of complementary and alternative medicine, including the Indian system *Ayurved*, is *Punica granatum* L. extracts". Cite the references.
3. "In India, Tunisia, and Guatemala, decoction of dried peels of *P. granatum* is employed externally as well as internally as an astringent and germicide, and utilized for treating aphthae and diarrhoea". Cite the references.
4. "Klebseiella Pneumonia" Spelling mistake is there. It should be corrected as *klebsiella pneumonia*.
5. "Incubation temperature and time for *C. violaceum*, *L. plantarum*, and *B. bifidum* was 37°C, and 22–24 h". Cite the references.
6. "Antibiotic susceptibility profile of the bacterial strains used in this study was generated using the antibiotic discs- Dodeca Universal – I, Dodeca G - III – Plus, and Icosa Universal -2 (HiMedia, Mumbai)". Cite the references.
7. "The plant name was checked against <http://www.theplantlist.org> on April 20, 2018. Collected peels were shade-dried, before being used for extract preparation. Biological activity of the plant fruit even depends on the time and season of productions". This fruit is generally produced in season in the Northern Hemisphere from September to February and in the Southern Hemisphere from March to May. Please comment on this. Authors are being also suggested to carry out the same experiment using the Peels of *P. granatum* were procured from different times. (optional)
8. "Purity of each of the extracted pigment was confirmed by running a UV-vis scan (Agilent Cary 60 UV-visible spectrophotometer). Appearance of single major peak (at the $\lambda_{\max}$ reported in literature) was taken as indication of purity". Cite reference.
9. Fig 4 is not clear and acceptable in the present form. Please prepare a high resolution figure as per the authors instruction.
10. Why did the author do molecular docking study? It should be clearly mentioned in the methods section along with references^{1,2}. Docking score is not the ultimatum measurement of potency or binding affinity of the ligand. I suggest giving the figures of mode of ligand-receptor interactions and discussing the crucial interacting amino acids of the protein.
11. "One ethnomedical practice reported from the Paliyar tribe from Tamilnadu in India involves taking internally, dried fruit coat of *P. granatum* after grinding and mixing with water, to treat stomach ache and diarrhoea (Duraipandiyan *et al.*, 2006)". Correct the typographical error of Tamilnadu.
12. The present version contains many grammatical and typographical errors. Authors are suggested to do these corrections.
13. "Inoculum standardized to 0.5 McFarland turbidity standard was added at 10% v/v, to the media supplemented with required concentration of PGPE, followed by incubation at appropriate temp for each organism". Correct the spelling error of temp.
14. Separate conclusion should be given based on experimental and in-silico results.

References

1. Nandi S: Recent Advances in Ligand and Structure Based Screening of Potent Quorum Sensing Inhibitors Against Antibiotic Resistance Induced Bacterial Virulence. *Recent Patents on Biotechnology*. 2016; **10** (2): 195-216 [Publisher Full Text](#)
2. Nandi S, Bagchi MC: 3D-QSAR and molecular docking studies of 4-anilinoquinazoline derivatives: a rational approach to anticancer drug design. *Mol Divers*. 2010; **14** (1): 27-38 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

No

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Synthesis of bioactive molecules, drug design, discovery of lead and molecular modeling.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 19 February 2019

<https://doi.org/10.5256/f1000research.19060.r43899>

© 2019 Hameed S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Saif Hameed

Amity University Haryana, Gurugram, India

In this study, the authors analyze an "Anti-infective potential of hydroalcoholic extract of Punica

granatum peel against gram-negative bacterial pathogens" with various experiments QS, biofilm hemolytic activity, catalase activity, surface hydrophobicity. Authors also performed *in-vivo* activity with *C. elegans* model organism and also analyzed with docking tool. While this study is comprehensive and important for better understanding of gram negative bacteria (*C. violaceum* and *P. aeruginosa*) physiology and pathogenesis, experiments are enough and paper sounds good but requires minor corrections with respect to the overall presentation of the manuscript.

Comment 1: In the introduction, the ESRC reference should be changed as it is from 2014.

Comment 2: Brief description about AMR and how bacteria develop these mechanism would be better.

Comment 3: Broth dilution assay: Give the proper reference of MIC determination method and how they defined MIC for example (MIC50, MIC80 or MIC90) or complete at the end point of experiment. Better to put the reference of MIC according to CLSI guidelines.

Comment 4: Measurement of bacterial growth and pigment production: better to merge this sentence with above and keep the pigment production in separate heading.

Comment 5: *In vivo* assay: last day of the experiment, when plates could be opened, their death was also confirmed by touching them with a straight wire, wherein no movement was taken as confirmation of death. Please mention what metal was used.

Comment 6: In Results section:

- Figure 1A: is not clear and difficult to understand. The values shown in boxes or brackets - what are these values defining? Are they necessary to show? If yes, better to show in table form.
- Similarly for Fig 1C.
- Figures 1D: Better to show *C. elegans* survival graph along with their pictures (Fig. 1D 2F, 3B).
- Figure 2: very overcrowded and messy. Is it necessary to show 5 figures A,B,C,D,E,F in a single panel? If there is any figure limitation then show only relevant ones and put rest as supplementary figures.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Comments on this article

Version 1

Reader Comment 19 Jan 2019

Muhammad Shahzad Aslam, Xiamen University Malaysia, Malaysia

I understand the Violacein extraction method and the Pyoverdine and pyocyanin extraction method but I could not find Preparation of media and its composition according to paper the authors have cited "Inhibition of bacterial quorum sensing by vanilla extract"

The quality of picture in Figures 7 and 8 is unclear.

In Figure 6 what is meant by Violacein unit?

Did you characterize the compounds you extracted? How you validate Violacein extraction method and Pyoverdine and pyocyanin extraction method?

Competing Interests: Nil

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research