REVIEW

Recent advances in marburgvirus research [version 1; peer review: 3 approved]

Judith Olejnik^{1,2}, Elke Mühlberger^{1,2}, Adam J. Hume ^{1,2}

¹Department of Microbiology, Boston University School of Medicine, Boston, Massachusetts, 02118, USA

²National Emerging Infectious Diseases Laboratories, Boston University, Boston, Massachusetts, 02118, USA

V1 First published: 21 May 2019, 8(F1000 Faculty Rev):704
<https://doi.org/10.12688/f1000research.17573.1>

Latest published: 21 May 2019, 8(F1000 Faculty Rev):704
<https://doi.org/10.12688/f1000research.17573.1>

Abstract

Marburgviruses are closely related to ebolaviruses and cause a devastating disease in humans. In 2012, we published a comprehensive review of the first 45 years of research on marburgviruses and the disease they cause, ranging from molecular biology to ecology. Spurred in part by the deadly Ebola virus outbreak in West Africa in 2013–2016, research on all filoviruses has intensified. Not meant as an introduction to marburgviruses, this article instead provides a synopsis of recent progress in marburgvirus research with a particular focus on molecular biology, advances in animal modeling, and the use of Egyptian fruit bats in infection experiments.

Keywords

Marburg virus, marburgviruses, filovirus, filoviruses, Egyptian roussette, viral proteins

Open Peer Review

Approval Status

	1	2	3
version 1 21 May 2019			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Ilhem Messaoudi**, University of California, Irvine, USA
2. **Chad E Mire**, University of Texas Medical Branch, Galveston, USA
3. **M. Javad Aman**, Integrated Biotherapeutics, Inc., Rockville, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Adam J. Hume (hume@bu.edu)

Author roles: Olejnik J: Writing – Original Draft Preparation, Writing – Review & Editing; Mühlberger E: Writing – Original Draft Preparation, Writing – Review & Editing; Hume AJ: Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under award numbers R01-AI133486 and R21-AI135912.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 Olejnik J *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Olejnik J, Mühlberger E and Hume AJ. **Recent advances in marburgvirus research [version 1; peer review: 3 approved]** F1000Research 2019, 8(F1000 Faculty Rev):704 <https://doi.org/10.12688/f1000research.17573.1>

First published: 21 May 2019, 8(F1000 Faculty Rev):704 <https://doi.org/10.12688/f1000research.17573.1>

Introduction

Marburg virus (MARV) is a member of the *Marburgvirus* genus that contains two different viruses: MARV and Ravn virus (RAVV). Both viruses are represented by numerous isolates¹. Filovirus taxonomy is confusing, and for those who do not know the difference between Marburg virus (the virus MARV), *Marburgvirus* (the genus), and marburgvirus (MARV and RAVV), we recommend browsing through the *Guide to the Correct Use of Filoviral Nomenclature* by Kuhn².

MARV is closely related to the better-known Ebola virus (EBOV) and causes a similarly severe disease in humans. Some of the unique characteristics of filovirus outbreaks were reported for MARV disease (MVD) long before they were noticed in EBOV disease (EVD). This includes persistent infection, sexual transmission, and long-term sequelae³. There are also heart-breaking reports of social stigmatization and severe chronic health issues as recalled by survivors of the 1967 MVD outbreak in Marburg, Germany⁴. Notably, one of the patients from that outbreak temporarily lost the ability to write and calculate and never completely recovered from concentration disorders, reflecting the severe neurological consequences in survivors of filovirus disease⁴.

MVD remains a global health threat with outbreaks continuing to occur in Central Africa, including two outbreaks in Uganda in 2012 and 2014^{5,6}. Jointly with EVD, MVD is listed on the World Health Organization 2018 Priority Diseases List (<https://www.who.int/blueprint/priority-diseases/en/>). This list is used as a tool to determine which diseases and pathogens should be prioritized for research, development of countermeasures, and emergency response preparedness.

In 2012, we published a comprehensive overview on marburgviruses and the disease they cause, ranging from ecology to molecular biology and treatment options⁷. A number of areas of significant progress in the marburgvirus field since 2012—including the development of tools to study MARV replication and transcription and to rescue MARV clones^{8,9}, advances in filovirus countermeasures^{10,11}, and vaccine development¹²—have recently been summarized. Progress in the development of MARV vaccines and antiviral treatment options has led to phase 1 clinical trials to evaluate their safe use in humans^{13–15}. Rather than repeating what has been covered in our previous review and these other excellent reviews, this article instead will focus on (1) recent progress in marburgvirus molecular biology, (2) novel developments in the study of marburgviruses in Egyptian fruit bats, and (3) advances in the use of animal models to study marburgvirus infection, including their use in resolving isolate-specific differences in virulence and pathogenicity.

1. Molecular biology

During the last 6 years, much research has focused on a more detailed understanding of the different steps of the marburgvirus replication cycle as well as virus–host interactions using crystal structure, biochemistry, and bioinformatics approaches. Live-cell imaging studies have been instrumental in developing a deeper understanding of marburgvirus assembly and particle release.

a. Marburg virus genome

Marburgviruses belong to the group of non-segmented negative-strand RNA viruses. A detailed overview of viral genome organization, cis-acting elements, and genome replication and transcription strategies is provided in 7,16. High-throughput sequencing of MARV RNA combined with bioinformatics and statistical analysis has provided new insights into viral genome plasticity and MARV evolution. This includes codon usage analysis¹⁷, phylogenetics^{18,19}, and the identification of hot spots of U–C substitutions^{20,21}. Although the function of these U–C substitutions is unknown, it is suggestive of adenosine deaminase (ADAR) editing^{20,21}. Deep sequencing of MARV Angola RNA obtained from infected cells and infected non-human primates (NHPs) determined novel editing sites in the nucleoprotein (NP) and L open reading frames, increasing the potential coding capacity of these viral genes with as-yet-unknown functions²⁰.

b. Marburg virus assembly

The MARV RNA genome is in close association with the viral nucleocapsid proteins, including NP (enwraps viral RNA), the RNA-dependent RNA polymerase complex consisting of L (enzymatic moiety of the polymerase complex), and viral protein 35 (VP35) (polymerase cofactor), VP30 (function not clear; possibly a transcription regulator), and VP24 (involved in nucleocapsid formation and maturation). Viral genome replication and transcription occur in the cytoplasm of the infected cells in viral inclusions which are highly ordered aggregations of nucleocapsids²². Live-cell imaging has shed some light on the trafficking of mature MARV nucleocapsids from viral inclusions to the sites of budding and helped to identify viral and cellular proteins involved in this process. Mature MARV nucleocapsids are transported along actin filaments from the viral inclusions to the plasma membrane, where they recruit the viral matrix protein VP40. Only nucleocapsids that are associated with VP40 are transported into filopodia²³. These long cellular protrusions are the main budding sites of MARV particles. MARV NP contains a late domain motif (PSAP) that recruits tumor susceptibility gene 101 (Tsg101), a component of the vesicular transport system ESCRT-I (endosomal sorting complexes required for transport I), to the viral inclusions. NP–Tsg101 interaction is required for the actin-dependent transport of MARV nucleocapsids into the filopodia²⁴. The MARV glycoprotein (GP) is also recruited to VP40-enriched membranes by a tubulin-dependent process²⁵.

c. Structure–function of Marburg virus proteins

Structural analyses have provided a deeper insight into the interactions and functions of almost all seven MARV proteins²⁶. Comparison of the MARV proteins with their ebolavirus homologs has shown a varying degree of structural resemblance that often correlates with functional similarity^{27–36}. Intriguingly, despite considerable structural conservation, some of the MARV proteins are functionally different from their ebolavirus homologs, with VP24 being a prime example of this.

Viral protein 24. VP24 is required for nucleocapsid formation and assembly for both viruses, but only ebolavirus VP24 blocks type I interferon (IFN) signaling (reviewed in 37,38). Crystal structure analysis revealed that the ebolavirus and MARV VP24

cores are structurally similar, supporting their common function in nucleocapsid formation and assembly. One difference between MARV and ebolavirus VP24 proteins is an extended β -sheet that is formed by MARV VP24 amino acid residues 201 to 217³⁵. Although this structural difference does not explain why MARV VP24 is not immunosuppressive³⁵, it is connected to a novel function of MARV VP24. The extended β -shelf contains the binding domain for the cellular adaptor protein Kelch-like ECH-associated protein 1 (Keap1). Keap1 is a central player in the oxidative stress response pathway and a suppressor of the antioxidant response transcription factor nuclear factor (erythroid-derived 2)-like 2 (Nrf2). MARV VP24 binds to Keap1 via the “K loop” that comprises amino acids 202 to 209^{39,40}. This leads to the nuclear accumulation of Nrf2 and consequently to the activation of an antioxidative and cytoprotective response in MARV-infected cells^{39,41}. In contrast, ebolavirus VP24 proteins, which do not have the K loop, are not able to induce a cytoprotective response^{39,41}. Keap1 is also a regulator of the nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) pathway. It targets I κ B kinase β (IKK β) for degradation which disables the cell to respond to NF- κ B-activating stimuli. MARV VP24 prevents Keap1 from binding to IKK β , allowing the cell to respond to NF- κ B-activating stimuli⁴².

Viral protein 30. Another example for functional difference despite structural conservation is VP30. Although EBOV VP30 has been shown to act as a transcription activator and regulator of viral RNA synthesis^{16,32}, the role of MARV VP30 is less well understood. Despite the high degree of structural homology between MARV and EBOV VP30, EBOV VP30 enhances transcriptional activity in a minigenome system at least 50-fold and is essential for detectable reporter gene activity, whereas MARV VP30 increases reporter gene activity only moderately (about twofold) in the MARV minigenome system^{43–45}. However, MARV VP30 is essential for viral replication^{46,47}. Crystal structure analysis confirmed the conformational similarity of EBOV and MARV VP30 proteins. Both EBOV and MARV VP30 C-terminal domains bind a peptide in the respective NP C-terminal region. The EBOV and MARV NP–VP30 complexes are remarkably similar, although slight differences might account for the lower binding affinity observed for the MARV NP–VP30 complex (K_D of 14 μ M compared with K_D of 5.7 μ M for EBOV complex). NP–VP30 complex formation was shown to be essential for EBOV minigenome activity³². Whether the observed differences in VP30–NP binding affinity play a role in the lower efficiency of MARV VP30 to enhance transcription remains to be determined.

Both EBOV and MARV VP30 proteins are phosphorylated, and phosphorylation of EBOV VP30 blocks its transcriptional activity^{48–52}. A recent study showed that phosphorylation of MARV VP30, like that of EBOV, interferes with its enhancing function in the minigenome system⁴⁵.

Viral protein 35. MARV VP35 plays an important role in various steps of the viral replication cycle. It is a component of the viral RNA-dependent RNA polymerase, required for nucleocapsid formation, and suppresses antiviral host responses, including type I IFN production, protein kinase R (PKR) activation,

and dendritic cell maturation^{53–56}. Crystal structure analysis provided deeper insights into the various functions of MARV VP35. Similar to EBOV VP35^{57,58}, MARV VP35 chaperones NP by inhibiting NP oligomerization and RNA binding. This facilitates the release of NP-bound viral RNA and gives the viral polymerase access to the RNA template. It also prevents unspecific NP-RNA aggregation^{33,34}. In regard to its function as a suppressor of type I IFN production, MARV VP35 seems to be less efficient than its EBOV homolog^{59–61}. This difference might be due to differences in the binding modes and affinities of EBOV and MARV VP35 proteins to double-stranded RNA, potentially impacting the inhibition of downstream antiviral pathways^{59,60,62}. Interestingly, compared with MARV VP35, RAVV VP35 seems to be more efficient in suppressing type I IFN production⁶¹.

Viral protein 40. MARV VP40 is a multifunctional protein that mediates virus egress, regulates viral transcription and replication, and counteracts the innate immune response by blocking signal transducer and activator of transcription (STAT)1 and STAT2 signaling⁶³. Recent research on VP40 has focused on the mechanisms underlying VP40–membrane interaction and budding. Crystal structure analysis has shown that MARV VP40 in solution forms a dimer with an N-terminal dimer interface. The C-terminal domains of the VP40 dimer, which form a flat cationic surface that interacts with anionic phospholipids at the inner leaflet of the plasma membrane^{29,36,64}, are required for budding but not for the immunosuppressive function of VP40²⁹. Hydrogen–deuterium exchange mass spectrometry has suggested that this interaction facilitates VP40 assembly and oligomerization⁶⁵.

VP40 hijacks cellular vesicular transport systems to bud from the infected cells, including the coat protein complex II (COPII) vesicular transport system and the ESCRT system³⁷. A screen to find additional cellular proteins that bind to the late domain motif (PPxY) within EBOV VP40 identified the co-chaperone protein BCL2-associated athanogene 3 (BAG3). In contrast to other late domain binding proteins that promote filovirus budding, BAG3 inhibits EBOV and MARV VP40-mediated budding and therefore could be part of a cellular antiviral defense system⁶⁶.

It has been shown previously that VP40 is a species-specific virulence factor and that changes in VP40 are required for marburgvirus adaptation to rodent hosts (reviewed in 67; an overview of the various MARV and RAVV isolates is provided in 7). Mouse adaptation of two marburgvirus isolates—RAVV and MARV Ci67—led to several mutations in the VP40 gene^{68,69}. Although the wild-type versions of RAVV and MARV Ci67 VP40 antagonized the type I IFN response inefficiently in mouse cells, the mouse-adapted VP40 mutants retained this function in both human and mouse cells^{70,71}. Intriguingly, the underlying amino acid changes required to facilitate type I IFN antagonism in mouse cells differed in the mouse-adapted RAVV and MARV Ci67 VP40 proteins^{70,71}. Other functional changes that correlated with mouse adaption affected the ability of VP40 to mediate budding. Whereas wild-type RAVV VP40 efficiently mediated budding from both human and mouse cells, mouse-adapted RAVV VP40 was restricted in this function by tetherin in human

cells⁷². Mutations in VP40 induced by guinea pig adaptation of MARV Musoke resulted in increased budding and a gain of viral fitness in guinea pig cells while not altering the type I IFN antagonizing function of VP40 in human or guinea pig cells⁷³. This shows that VP40 mutations which occur during rodent adaptation not only enhance type I IFN antagonism in these hosts but also promote VP40-mediated budding. MARV adaptation to guinea pigs resulted in mutations in both VP40 and the viral polymerase (L). Minigenome data suggest that there is a synergistic effect of these mutations in guinea pig cells, leading to enhanced replication activity⁷⁴. Because of the role of VP40 as a virulence factor, the inhibition of type I IFN signaling by VP40 represents an intriguing potential target for the development of anti-MARV therapeutics. Unfortunately, this is currently limited by our poor understanding of the exact molecular mechanisms underlying this inhibition.

Glycoprotein. Given its exposure on the surface of viral particles, which makes it the perfect target for antibody recognition, and its crucial role in attachment and fusion, MARV GP has been extensively studied in the last 6 years. This includes the identification and characterization of host factors involved in attachment and entry^{75–77}, the role of GP₂ in membrane fusion and tetherin antagonism^{30,31,78,79}, and the effects of steric shielding of host surface proteins, including major histocompatibility complex I, integrin β 1, and Fas, by MARV GP^{80,81}. GP shielding of surface-expressed Fas interfered with the induction of Fas-mediated apoptosis and could help to protect MARV-infected cells from premature cell death⁸¹. Interestingly, steric shielding of host proteins was more pronounced for MARV Angola GP than for MARV Musoke GP, indicating that it might play a role in virus-specific pathogenicity⁸⁰ (see section 3b). Much progress has also been achieved in the structural analysis of GP and its interaction with protective antibodies, which are promising candidates for post-exposure treatment^{82,83}. An excellent overview by King *et al.* on the structural features of protective epitopes on filovirus GPs highlights the striking differences between the neutralizing epitopes on EBOV and MARV GPs⁸⁴. To date, only two MARV GP epitopes that are targeted by protective antibodies have been identified: the receptor binding site and the wing domain that is located within the MARV GP₂ subunit and is not present in ebolavirus GPs. The MARV receptor binding site seems to be easily accessible to neutralizing antibodies whether GP is cleaved or not, whereas the respective region on ebolavirus GPs is shielded by a glycan cap and is exposed only after GP cleavage^{85,86}. The receptor binding site antibodies have shown promising potential for use as therapeutics, as they are protective when administered up to 5 days post-exposure in NHPs⁸³. Testing of antibodies directed against the wing domain have been less encouraging: protection of mice is observed when given 1 hour after infection⁸⁷, highlighting a large disparity in therapeutic potential between antibodies directed against these two regions of GP. Owing to differences in structure and identified protective epitopes, the search for a pan-filovirus antibody has remained unfruitful so far, although pan-ebolavirus protective antibodies have been identified^{88–95}.

2. Host response to Marburg virus infection

Recent studies on the host response to MARV infection have focused on identifying signatures of innate and adaptive immunity

as well as correlates of disease severity and outcome in both human MVD survivors and MARV-infected NHPs. Although both EBOV and MARV infections are lethal in cynomolgus macaques, transcriptional analysis revealed unique immune signatures associated with each virus and suggested a more pronounced immune dysregulation in MARV infection⁹⁶. The MARV-specific gene expression profile included the upregulation of complement system genes, genes involved in neutrophil and monocyte recruitment, and innate immune signaling genes⁹⁶. Distinct immune responses, which are predictive of clinical outcomes, have been detected in both human survivors and macaques infected with MARV: although lethal infection of rhesus macaques with MARV Angola was linked to T-helper cell type 2 (Th2)-skewed responses⁹⁷, human survivors of MARV infection exhibited Th1-skewed CD4⁺ T-cell responses⁹⁸. The ability to identify these predictive responses in patients could result in more effective patient triage and administration of targeted therapies.

Another major focus of MVD research has been the pursuit of earlier diagnostics, and particular attention has been on the early identification of unique signatures of marburgvirus infection. MARV infection of NHPs led to the activation of peripheral blood mononuclear cells, as reflected by the induction of the type I IFN response in rhesus macaques⁹⁹ and increased expression of heat shock proteins in cynomolgus macaques^{97,100}. Upregulation of type I IFN-stimulated genes was detected as early as 1 day post-infection in MARV-infected rhesus macaques, prior to the onset of symptoms⁹⁷. These data indicate that new diagnostic tools could be developed to detect these unique signatures of marburgvirus infections prior to the onset of viremia or symptoms, allowing earlier detection and treatment of MVD.

A major unanswered question in the filovirus field is how long human survivors remain protected against subsequent exposure to autologous virus. Although long-lived antibody responses have been found in human survivors of Sudan and Bundibugyo viruses (both ebolaviruses) as well as MARV¹⁰¹, MARV neutralizing antibody responses were lower and showed a faster decline in contrast to the prolonged presence of neutralizing antibodies after Sudan virus infection^{98,102}. Low titers of neutralizing antibodies were also observed in cynomolgus macaques after vaccination against MARV. Nevertheless, animals were protected against lethal MARV challenge for over 1 year¹⁰³. These findings emphasize the need for a better understanding of the differences in host response to diverse filoviruses and to determine long-term effects of vaccine candidates.

3. Marburg virus prevalence and pathology in bats

Another area of progress in the marburgvirus field is focused on the natural reservoir of these viruses. Prior work showed that the common Egyptian fruit bat (*Rousettus aegyptiacus*) is a reservoir for marburgviruses, having demonstrated positive serology, detection of viral RNA, and the ability to culture infectious virus from the bats, a feat still unparalleled in the ebolavirus field. Since our previous review of the marburgvirus literature⁷, major advances have been made in our understanding of the ecology of marburgviruses in Egyptian fruit bats, including the outcome of experimental infection of these bats with MARV.

a. Ecology of marburgviruses in bats

Expanding upon the handful of reports prior to 2012, more recent studies have investigated the prevalence of marburgviruses in various bat species. These reports primarily investigated the prevalence of Egyptian fruit bat exposure to MARV and RAVV as determined by serology and, to a lesser extent, quantitative real-time polymerase chain reaction (RT-PCR), finding varying degrees of prevalence in different geographic populations. The finding of higher rates of Egyptian fruit bats with detectable marburgvirus RNA that repopulated Kitaka mine¹⁰⁴ versus the population that was present prior to extermination¹⁰⁵ led the authors to propose that the new founding population may have been susceptible to MARV infection and, following multiple introductions of diverse marburgviruses, led to the high rates of RT-PCR-positive bats. This finding should serve as a warning for possible future attempts at controlling MARV outbreaks by trying to eradicate local bat colonies; such an approach may backfire.

Three separate studies analyzing MARV seropositivity and RNA in Egyptian fruit bats in Uganda, South Africa, and Zambia all detected cyclical temporal patterns: the lowest prevalence was observed during the birthing season and prevalence increased thereafter, particularly among juvenile bats^{106–108}. Intriguingly, 83% of previous marburgvirus outbreaks in the human population coincided with these peaks in viral abundance in Egyptian fruit bats¹⁰⁶, further implying the importance of Egyptian fruit bats as the prime source for MARV spillover to humans. The fact that similar data were obtained in diverse bat populations that differ drastically in their geographic distribution and mating patterns (once versus twice per year) strengthens the possibility that marburgvirus prevalence, at some level, may be linked to behavioral patterns of these bats.

Because MARV infection appears to be quite mild in Egyptian fruit bats (see section 2b) and prevalence is fairly low, some have speculated either that *R. aegyptiacus* is not the reservoir species or that it is unlikely to be the only species required for the enzootic maintenance of marburgviruses. To date, however, investigations of marburgvirus seroprevalence in other bat species^{105,109} and marburgvirus presence in *R. aegyptiacus*-associated ticks (*Ornithodoros faini*¹¹⁰) have been unfruitful in finding other likely reservoir species.

b. Experimental infection of Egyptian fruit bats

Critical advances in our understanding of MARV infection of Egyptian fruit bats have been facilitated by the establishment of captive, breeding colonies of Egyptian fruit bats at the Center for Emerging and Zoonotic Diseases in Sandringham, South Africa, and at the Centers for Disease Control and Prevention in Atlanta, Georgia, USA. Studies reporting primary MARV infection of Egyptian fruit bats from these two colonies have yielded strikingly similar results; although MARV is able to replicate in bats, viremia remains low, there is no resulting sickness or pathology in the bats, and the bats are able to quickly clear the infection^{111–114}. Whereas both groups observed MARV RNA in oral and rectal swabs of infected Egyptian fruit bats^{112,114}, only one of the groups was able to culture live virus from oral and rectal swabs from the infected bats¹¹².

An important aspect in considering the ecology of MARV in *R. aegyptiacus* populations is whether prior infection with MARV prevents subsequent reinfection. Two separate challenge studies clearly showed that Egyptian fruit bats previously infected with MARV were resistant to challenge by a homologous MARV isolate^{115,116}.

c. Transmission of Marburg virus by Egyptian fruit bats

Recent studies investigating the ability of MARV-infected Egyptian fruit bats to spread the virus to uninfected, co-housed animals have resulted in more divergent results. Whereas Paweska *et al.* found no evidence of transmission despite robust infection of the injected bats¹¹⁴, Schuh *et al.* did¹¹⁷. MARV was detected in blood and oral samples from the bats that were infected by horizontal transmission, providing a possible mechanism of viral transmission (saliva transmission). The different outcomes could be due to the different virus isolates used in these studies or differences in the genetic background of the bats themselves.

d. Molecular biology of Marburg virus infection in bat cells

In addition to advances in studying infections in bats, much recent study has focused on cellular responses in bat cells to MARV infection. Multiple transcriptomic analyses of MARV infection of *R. aegyptiacus* cell lines have come to similar conclusions. Infection of the cell lines R06E (derived from fetal body cells)¹¹⁸ and R0Ni/7.1 (adult kidney)¹¹⁹ appears to follow patterns similar to those of transcriptional responses in human cells, although viral replication was not as robust in the bat cells as in Huh7 and HepG2 human cells^{120–122}. Whereas two of these reports found either no induction or minimal induction of type I, II, or III IFNs or IFN-stimulated genes in bat cells infected with MARV^{120,121}, one of these reports found low to modest upregulation of these genes¹²². This may be due to differences in experimental design or differences in the MARV isolates used in the studies. Interestingly, one of these reports found that MARV upregulated some unannotated antiviral paralogs¹²¹.

Despite a relative dearth of bat-specific antibodies, some non-transcriptomic analyses have begun analyzing molecular mechanisms of MARV infection in bat cell lines. Although EBOV VP35 had previously been shown to inhibit PKR in human cells, a recent report describes the ability of MARV VP35 to inhibit PKR activation in a cell type-specific manner, with VP35 being unable to inhibit bat PKR in the RoNi/7.1 cell line⁵⁵. An analysis of fibroblast cell lines from four different bat species revealed that species-specific differences in Niemann–Pick C1 (NPC1) accounted for differences in EBOV GP-mediated, but not MARV GP-mediated, infectivity¹²³.

4. Advances in animal modeling of marburgvirus infections

Historically, individual studies analyzing marburgviruses have been performed using single isolates of MARV or RAVV, and comparative studies have been sparse¹²⁴. However, fueled by discussions about the impact of viral sequence variations on pathogenic potential that arose during the West African EBOV outbreak^{125–128} and by the advent of high-throughput sequencing technologies, marburgvirus isolate variations are now

more acknowledged, and marburgvirus diversity has become a research topic. Well-established and newly developed animal models are useful tools to conduct these comparative studies. A list of marburgvirus variants commonly used in animal experiments during the last 6 years is provided in [Table 1](#).

a. Newly developed animal models to study Marburg virus and Ravn virus infection

Since our previous review in 2012, a number of new animal models of MVD have been developed. Detailed comparisons of these and previously established MVD animal models have recently been published^{129–131}. Instead of repeating the content of these comprehensive reviews, we will briefly introduce these animal models and then focus on their use in resolving

isolate-specific differences in virulence and pathogenicity as well as their use in modeling post-recovery sequelae.

Recently developed animal models that allow the study of lethal MVD without virus adaptation include STAT2 knockout hamsters¹³², humanized mice¹³³, and marmosets¹³⁴. Infection of humanized mice with MARV, compared with infection with EBOV, was associated with lower overall weight loss, whereas the viral titers were similar¹³³. Surprisingly, ferrets infected with MARV or RAVV, in contrast to those infected with EBOV, did not develop symptoms of disease, regardless of dose, route of infection, and virus isolate^{135–138}. Immunocompetent small-animal infection models for MARV and RAVV, including mice, hamsters, and guinea pigs, require virus adaptation. Recent

Table 1. Experimentally used marburgviruses.

Year	Country of origin	Country of isolation	Isolate	Reference
1967	Presumably Uganda	Germany	MARV Ci67 (Marburg virus/H.sapiens-tc/GER/1967/Hesse-Cieplik)	Lofts <i>et al.</i> (2011) ⁶⁸ , Coffin <i>et al.</i> (2018) ¹³⁹
			MARV Voegel (Marburg virus/H.sapiens-tc/GER/1967/Hesse-Voegel)	Atkins <i>et al.</i> (2018) ¹³²
			MARV Pop (Marburg virus/H.sapiens-tc/GER/1967/Hesse-Popp)	Smither <i>et al.</i> (2013) ¹³⁴
1975	Rhodesia (now Zimbabwe)	South Africa	MARV Ozolins (Marburg virus/H.sapiens-tc/ZAF/1975/Sinoia-Ozolins)	Nicholas <i>et al.</i> (2018) ¹⁴⁰
			MARV Hogan (Marburg virus/H.sapiens-tc/ZAF/1975/Sinoia-Hogan)	Paweska <i>et al.</i> (2012) ¹¹¹
1980	Kenya	Kenya	MARV Musoke (Marburg virus/H.sapiens-tc/KEN/1980/Mt. Elgon-Musoke)	Atkins <i>et al.</i> (2018) ¹³² , Blair <i>et al.</i> (2018) ¹⁴¹ , Nicholas <i>et al.</i> (2018) ¹⁴⁰ , Daddario-DiCaprio <i>et al.</i> (2006) ¹⁴² , Coffin <i>et al.</i> (2018) ¹³⁹ , Daddario-DiCaprio <i>et al.</i> (2006) ¹⁴³ , Mire <i>et al.</i> (2014) ¹⁰³
1987	Kenya	Kenya	RAVV (Ravn virus/H.sapiens-tc/KEN/1987/Kitum Cave-810040)	Atkins <i>et al.</i> (2018) ¹³² , Lofts <i>et al.</i> (2011) ⁶⁸ , Cross <i>et al.</i> (2015) ¹⁴⁴ , Nicholas <i>et al.</i> (2018) ¹⁴⁰ , Daddario-DiCaprio <i>et al.</i> (2006) ¹⁴² , Thi <i>et al.</i> (2017) ¹⁴⁵ , Mire <i>et al.</i> (2017) ⁸³
1998–2000	COD	COD	MARV SPU 148/99/1	Paweska <i>et al.</i> (2015) ¹¹⁴ , Storm <i>et al.</i> (2018) ¹¹⁶
2004–2005	Angola	Angola	MARV-Angola-AGO-2005	Qiu <i>et al.</i> (2014) ¹⁴⁶ , Wong <i>et al.</i> (2018) ¹⁴⁷
			MARV-Angola-AGO-2005-368	Lavender <i>et al.</i> (2018) ¹³³ , Wong <i>et al.</i> (2018) ¹³⁶ , Marzi <i>et al.</i> (2016) ¹⁴⁸ , Fernando <i>et al.</i> (2015) ¹⁴⁹ , Nicholas <i>et al.</i> (2018) ¹⁴⁰ , Marzi <i>et al.</i> (2018) ⁹⁶
			MARV Angola (Marburg virus/H.sapiens-tc/AGO/2005/Angola-1379c)	Atkins <i>et al.</i> (2018) ¹³² , Johnston <i>et al.</i> (2015) ¹⁵⁰ , Lin <i>et al.</i> (2015) ¹⁵¹ , Alfson <i>et al.</i> (2018) ¹⁵² , Cross <i>et al.</i> (2015) ¹⁴⁴ , Blair <i>et al.</i> (2018) ¹⁴¹ , Cooper <i>et al.</i> (2018) ¹⁵³ , Woolsey <i>et al.</i> (2018) ¹⁵⁴ , Thi <i>et al.</i> (2017) ¹⁴⁵ , Mire <i>et al.</i> (2017) ⁸³
			MARV-Angola-1381	Ewers <i>et al.</i> (2016) ¹⁵⁵
2007	Uganda	Uganda	MARV-371-Bat2007 (Marburg virus/R.aegyptiacus-tc/UGA/2007/371Bat-811277)	Amman <i>et al.</i> (2015) ¹¹² , Jones <i>et al.</i> (2015) ¹¹³ , Schuh <i>et al.</i> (2017) ^{115,117}

This table shows Marburg virus (MARV) and Ravn virus (RAVV) variants that have been used in experimental animal infections highlighted in this review. "tc" in the virus name indicates tissue culture passaging. This information is not available for all listed viruses, but it is likely that all isolates were passaged more than once. COD, Democratic Republic of the Congo.

virus-adapted systems include a hamster model for MARV Angola that recapitulates the disease observed in humans and NHPs¹⁴⁸ and MARV Angola mouse models^{146,156}. Similar to what was observed before for mouse-adapted MARV isolates Ci67 and RAVV⁶⁷, deep sequencing of the MARV Angola genome during mouse adaptation revealed, among other mutations, adaptive changes in the VP40 open reading frame^{68,156}, emphasizing the importance of VP40 as a species-specific virulence factor (see “Viral protein 40” section above).

b. Differential virulence of Marburg virus variants

MARV Musoke was isolated in 1980 in Kenya from a physician who contracted the disease while attending a patient with MVD. Whereas the index case succumbed to the infection, the secondary case survived. There is only a single virus isolate from this MVD episode, and it was isolated from the surviving patient¹⁵⁷. The MARV Angola variant comes from the 2005 outbreak of MVD in Angola with at least 252 cases, including 227 fatalities. Multiple virus isolates exist from this outbreak¹⁵⁷. It is important to note that even individual isolates may vary from lab to lab, as they have typically been passaged in cell culture multiple times, disseminated to other laboratories, and possibly passaged further, leading to a diversity of virus stocks with different histories of propagation and thus potentially divergent genetic and phenotypic identities¹⁵². Passaging of viruses in cell culture is known to lead to the accumulation of defective interfering viral particles and higher particle/plaque-forming unit ratio⁹⁹. This was also shown for MARV Angola and EBOV; virus passaged multiple times in cell culture was found to be associated with increased survival and delayed disease progression in NHPs^{152,158}. Given the much longer history of the stocks derived from the MARV Musoke isolate, it is conceivable that MARV Musoke stocks have accumulated defective particles over time and that the compositions of the MARV Musoke and MARV Angola virus stocks differ, which should be taken into account when interpreting differences in pathogenicity.

Comparative studies in small-animal models that require virus adaptation revealed a range in pathogenicity, with MARV Angola being more pathogenic than other marburgvirus variants, including RAVV^{144,146,147}. Different results were obtained using the STAT2-deficient hamster infection model that does not require virus adaptation. In this infection model, fast disease progression and death were observed with MARV Musoke and Voegelé, a delayed but equally lethal infection was observed with MARV Angola, and RAVV resulted in a symptomatic but non-lethal infection¹³².

The pathogenicity of different marburgvirus variants was also studied in NHP models which recapitulate the disease in humans most faithfully. Due to advances in telemetry tracking of clinical data, disease symptoms now can be more easily monitored in this infection model¹⁵⁵. Fatal MARV infection is associated with increased lymphocyte and hepatocyte apoptosis in both rhesus and cynomolgus macaques¹⁴⁰. In the rhesus and cynomolgus macaque models, as in the small-animal models using adapted marburgviruses, MARV Angola showed higher virulence

compared with other marburgvirus variants, including MARV Musoke, Ci67, Ozolin, and RAVV^{140–143,149,154,159}. The increased virulence of MARV Angola could be explained, at least partially, by an enhanced replication efficiency of this virus, as shown in cell culture experiments¹⁶⁰. Conflicting results have been reported for RAVV infections of NHPs. Although RAVV infection of rhesus macaques was 100% lethal in two studies^{83,145}, complete survival was noted in a separate study¹⁴⁰. Overall, these studies highlight the need for further comparative analysis of different marburgvirus variants in different animal models.

c. Persistent Marburg virus infection in non-human primate models

Although persistent MARV infection and long-lasting health issues after recovery have been noted for a number of survivors of MVD, including those from the initial outbreak in 1967³, this issue became a research topic only after the devastating EVD outbreak in West Africa. Recent studies using rhesus macaques have shown that immune-privileged sites, including the eyes, female and male genital tracts, and mammary glands, are infected during acute MARV infection¹⁵³. Similar to human survivors of EBOV and MVD^{161–163}, some cynomolgus macaque survivors (including those that had been treated in antiviral drug testing) developed persistent MARV infection in the eyes and testes¹³⁹. These results indicate that NHP survivors of experimental MARV infection could serve as useful models to study sequelae of MVD. It remains to be determined whether persistent infection also occurs in other animal models of filovirus infection.

Abbreviations

BAG3, BCL2-associated athanogene 3; EBOV, Ebola virus; ESCRT, endosomal sorting complexes required for transport; EVD, Ebola virus disease; GP, glycoprotein; IFN, interferon; IKK β , I κ B kinase β ; Keap1, kelch-like ECH-associated protein 1; MARV, Marburg virus; MVD, Marburg virus disease; NF- κ B, nuclear factor kappa-light-chain-enhancer of activated B cells; NHP, non-human primate; NP, nucleoprotein; Nrf2, nuclear factor (erythroid-derived 2)-like 2; PKR, protein kinase R; RAVV, Ravn virus; RT-PCR, real-time polymerase chain reaction; STAT, signal transducer and activator of transcription; Tsg101, tumor susceptibility gene 101; VP, viral protein

Grant information

This work was supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under award numbers R01-AI133486 and R21-AI135912.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

The authors are grateful to Jens Kuhn (NIH/NIAID/DCR/Integrated Research Facility at Fort Detrick, Frederick, MD, USA), for help with Table 1.

References

1. Kuhn JH, Bao Y, Bavari S, *et al.*: **Virus nomenclature below the species level: a standardized nomenclature for natural variants of viruses assigned to the family *Filoviridae*.** *Arch Virol.* 2013; **158**(1): 301–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
2. Kuhn JH: **Guide to the Correct Use of Filoviral Nomenclature.** *Curr Top Microbiol Immunol.* 2017; **411**: 447–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Slenczka W: **Filovirus Research: How it Began.** *Curr Top Microbiol Immunol.* 2017; **411**: 3–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Moos F: **In uns und um uns. Meine Begegnung mit dem Marburg-Virus.** 1. edition ed. Frankfurt am Main: Mabuse Verlag GmbH; 2015.
[Reference Source](#)
5. Nyakarahuka L, Ojwang J, Tumusiime A, *et al.*: **Isolated Case of Marburg Virus Disease, Kampala, Uganda, 2014.** *Emerging Infect Dis.* 2017; **23**(6): 1001–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
6. **Outbreak news. Marburg haemorrhagic fever, Uganda.** *Wkly Epidemiol Rec.* 2012; **87**(45): 437–8.
[PubMed Abstract](#)
7. Brauburger K, Hume AJ, Mühlberger E, *et al.*: **Forty-five years of Marburg virus research.** *Viruses.* 2012; **4**(10): 1878–927.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. Schmidt KM, Mühlberger E: **Marburg Virus Reverse Genetics Systems.** *Viruses.* 2016; **8**(6): pii: E178.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. Hoenen T, Brandt J, Cai Y, *et al.*: **Reverse Genetics of Filoviruses.** *Curr Top Microbiol Immunol.* 2017; **411**: 421–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Connor J, Kobinger G, Olinger G: **Therapeutics Against Filovirus Infection.** *Curr Top Microbiol Immunol.* 2017; **411**: 263–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Cross RW, Mire CE, Feldmann H, *et al.*: **Post-exposure treatments for Ebola and Marburg virus infections.** *Nat Rev Drug Discov.* 2018; **17**(6): 413–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Reynolds P, Marzi A: **Ebola and Marburg virus vaccines.** *Virus Genes.* 2017; **53**(4): 501–15.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Heald AE, Charleston JS, Iversen PL, *et al.*: **AVI-7288 for Marburg Virus in Nonhuman Primates and Humans.** *N Engl J Med.* 2015; **373**(4): 339–48.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Sarwar UN, Costner P, Enama ME, *et al.*: **Safety and immunogenicity of DNA vaccines encoding Ebolavirus and Marburgvirus wild-type glycoproteins in a phase I clinical trial.** *J Infect Dis.* 2015; **211**(4): 549–57.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Kibuuka H, Berkowitz NM, Millard M, *et al.*: **Safety and immunogenicity of Ebola virus and Marburg virus glycoprotein DNA vaccines assessed separately and concomitantly in healthy Ugandan adults: a phase 1b, randomised, double-blind, placebo-controlled clinical trial.** *Lancet.* 2015; **385**(9977): 1545–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Brauburger K, DeFlübé LR, Mühlberger E: **Filovirus Transcription and Replication.** In: Pattnaik AK, Whitt MA, editors. *Biology and Pathogenesis of Rhabdo- and Filoviruses.* Singapore: World Scientific Publishing Co. Pte. Ltd.; 2015; 515–5555.
[Publisher Full Text](#)
17. Nasrullah I, Butt AM, Tahir S, *et al.*: **Genomic analysis of codon usage shows influence of mutation pressure, natural selection, and host features on Marburg virus evolution.** *BMC Evol Biol.* 2015; **15**: 174.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Carroll SA, Towner JS, Sealy TK, *et al.*: **Molecular evolution of viruses of the family *Filoviridae* based on 97 whole-genome sequences.** *J Virol.* 2013; **87**(5): 2608–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Peterson AT, Holder MT: **Phylogenetic assessment of filoviruses: how many lineages of Marburg virus?** *Ecol Evol.* 2012; **2**(8): 1826–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Shabman RS, Jabado OJ, Mire CE, *et al.*: **Deep sequencing identifies noncanonical editing of Ebola and Marburg virus RNAs in infected cells.** *mBio.* 2014; **5**(6): e02011.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Khrustalev VV, Barkovsky EV, Khrustaleva TA: **Local Mutational Pressures in Genomes of Zaire Ebolavirus and Marburg Virus.** *Adv Bioinformatics.* 2015; **2015**: 678587.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Dolnik O, Stevermann L, Kolesnikova L, *et al.*: **Marburg virus inclusions: A virus-induced microcompartment and interface to multivesicular bodies and the late endosomal compartment.** *Eur J Cell Biol.* 2015; **94**(7–9): 323–31.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Schudt G, Kolesnikova L, Dolnik O, *et al.*: **Live-cell imaging of Marburg virus-infected cells uncovers actin-dependent transport of nucleocapsids over long distances.** *Proc Natl Acad Sci U S A.* 2013; **110**(35): 14402–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Dolnik O, Kolesnikova L, Welsch S, *et al.*: **Interaction with Tsg101 is necessary for the efficient transport and release of nucleocapsids in marburg virus-infected cells.** *PLoS Pathog.* 2014; **10**(10): e1004463.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. **F** Mittler E, Schudt G, Halwe S, *et al.*: **A Fluorescently Labeled Marburg Virus Glycoprotein as a New Tool to Study Viral Transport and Assembly.** *J Infect Dis.* 2018; **218**(suppl_5): S318–S326.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
26. Kirchdoerfer RN, Wasserman H, Amarasinghe GK, *et al.*: **Filovirus Structural Biology: The Molecules in the Machine.** *Curr Top Microbiol Immunol.* 2017; **411**: 381–417.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Baker LE, Ellena JF, Handing KB, *et al.*: **Molecular architecture of the nucleoprotein C-terminal domain from the Ebola and Marburg viruses.** *Acta Crystallogr D Struct Biol.* 2016; **72**(Pt 1): 49–58.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Bruhn JF, Kirchdoerfer RN, Urata SM, *et al.*: **Crystal Structure of the Marburg Virus VP35 Oligomerization Domain.** *J Virol.* 2017; **91**(2): pii: e01085-16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Oda SI, Noda T, Wijesinghe KJ, *et al.*: **Crystal Structure of Marburg Virus VP40 Reveals a Broad, Basic Patch for Matrix Assembly and a Requirement of the N-Terminal Domain for Immunosuppression.** *J Virol.* 2016; **90**(4): 1839–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Koellhoffer JF, Malashkevich VN, Harrison JS, *et al.*: **Crystal structure of the Marburg virus GP2 core domain in its postfusion conformation.** *Biochemistry.* 2012; **51**(39): 7665–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Liu N, Tao Y, Brenowitz MD, *et al.*: **Structural and Functional Studies on the Marburg Virus GP2 Fusion Loop.** *J Infect Dis.* 2015; **212** Suppl 2: S146–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Kirchdoerfer RN, Moyer CL, Abelson DM, *et al.*: **The Ebola Virus VP30-NP Interaction Is a Regulator of Viral RNA Synthesis.** *PLoS Pathog.* 2016; **12**(10): e1005937.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Liu B, Dong S, Li G, *et al.*: **Structural Insight into Nucleoprotein Conformation Change Chaperoned by VP35 Peptide in Marburg Virus.** *J Virol.* 2017; **91**(16): pii: e00825-17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. **F** Zhu T, Song H, Peng R, *et al.*: **Crystal Structure of the Marburg Virus Nucleoprotein Core Domain Chaperoned by a VP35 Peptide Reveals a Conserved Drug Target for Filovirus.** *J Virol.* 2017; **91**(18): pii: e00996-17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
35. Zhang AP, Bornholdt ZA, Abelson DM, *et al.*: **Crystal structure of Marburg virus VP24.** *J Virol.* 2014; **88**(10): 5859–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. **F** Bhattarai N, Go JB, Gerstman BS, *et al.*: **Plasma membrane association facilitates conformational changes in the Marburg virus protein VP40 dimer.** *RSC Adv.* 2017; **7**(37): 22741–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
37. Kolesnikova L, Nanbo A, Becker S, *et al.*: **Inside the Cell: Assembly of Filoviruses.** *Curr Top Microbiol Immunol.* 2017; **411**: 353–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Olejnik J, Hume AJ, Leung DW, *et al.*: **Filovirus Strategies to Escape Antiviral Responses.** *Curr Top Microbiol Immunol.* 2017; **411**: 293–322.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Edwards MR, Johnson B, Mire CE, *et al.*: **The Marburg virus VP24 protein interacts with Keap1 to activate the cytoprotective antioxidant response pathway.** *Cell Rep.* 2014; **6**(6): 1017–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Johnson B, Li J, Adhikari J, *et al.*: **Dimerization Controls Marburg Virus VP24-dependent Modulation of Host Antioxidative Stress Responses.** *J Mol Biol.* 2016; **428**(17): 3483–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Page A, Volchkova VA, Reid SP, *et al.*: **Marburgvirus hijacks nrf2-dependent pathway by targeting nrf2-negative regulator keap1.** *Cell Rep.* 2014; **6**(6): 1026–36.
[PubMed Abstract](#) | [Publisher Full Text](#)
42. Edwards MR, Basler CF: **Marburg Virus VP24 Protein Relieves Suppression of the NF- κ B Pathway Through Interaction With Kelch-like ECH-Associated Protein 1.** *J Infect Dis.* 2015; **212** Suppl 2: S154–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. Wenigenath J, Kolesnikova L, Hoenen T, *et al.*: **Establishment and application of an infectious virus-like particle system for Marburg virus.** *J Gen Virol.* 2010; **91**(Pt 5): 1325–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Albariño CG, Uebelhoefer LS, Vincent JP, *et al.*: **Development of a reverse genetics system to generate recombinant Marburg virus derived from a bat isolate.** *Virology.* 2013; **446**(1–2): 230–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
45. **F** Tigabu B, Ramanathan P, Ivanov A, *et al.*: **Phosphorylated VP30 of Marburg**

- Virus Is a Repressor of Transcription.** *J Virol.* 2018; **92**(21): pii: e00426-18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
46. Enterlein S, Volchkov V, Weik M, *et al.*: **Rescue of recombinant Marburg virus from cDNA is dependent on nucleocapsid protein VP30.** *J Virol.* 2006; **80**(2): 1038–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 47. Fowler T, Bamberg S, Möller P, *et al.*: **Inhibition of Marburg virus protein expression and viral release by RNA interference.** *J Gen Virol.* 2005; **86**(Pt 4): 1181–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 48. Modrof J, Möritz C, Kolesnikova L, *et al.*: **Phosphorylation of Marburg virus VP30 at serines 40 and 42 is critical for its interaction with NP inclusions.** *Virology.* 2001; **287**(1): 171–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
 49. Modrof J, Mülhberger E, Klenk HD, *et al.*: **Phosphorylation of VP30 impairs ebola virus transcription.** *J Biol Chem.* 2002; **277**(36): 33099–104.
[PubMed Abstract](#) | [Publisher Full Text](#)
 50. Martínez MJ, Biedenkopf N, Volchkova V, *et al.*: **Role of Ebola virus VP30 in transcription reinitiation.** *J Virol.* 2008; **82**(24): 12569–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 51. Biedenkopf N, Hartlieb B, Hoenen T, *et al.*: **Phosphorylation of Ebola virus VP30 influences the composition of the viral nucleocapsid complex: impact on viral transcription and replication.** *J Biol Chem.* 2013; **288**(16): 11165–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 52. **F** Kruse T, Biedenkopf N, Hertz EPT, *et al.*: **The Ebola Virus Nucleoprotein Recruits the Host PP2A-B56 Phosphatase to Activate Transcriptional Support Activity of VP30.** *Mol Cell.* 2018; **69**(1): 136–145.e6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 53. Ramanan P, Edwards MR, Shabman RS, *et al.*: **Structural basis for Marburg virus VP35-mediated immune evasion mechanisms.** *Proc Natl Acad Sci U S A.* 2012; **109**(5): 20661–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 54. Albariño CG, Wiggleson Guerrero L, Spengler JR, *et al.*: **Recombinant Marburg viruses containing mutations in the IID region of VP35 prevent inhibition of Host immune responses.** *Virology.* 2015; **476**: 85–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 55. Hume A, Mülhberger E: **Marburg Virus Viral Protein 35 Inhibits Protein Kinase R Activation in a Cell Type-Specific Manner.** *J Infect Dis.* 2018; **218**(suppl_5): S403–S408.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 56. Yen BC, Basler CF: **Effects of Filovirus Interferon Antagonists on Responses of Human Monocyte-Derived Dendritic Cells to RNA Virus Infection.** *J Virol.* 2016; **90**(10): 5108–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 57. Leung DW, Borek D, Luthra P, *et al.*: **An Intrinsically Disordered Peptide from Ebola Virus VP35 Controls Viral RNA Synthesis by Modulating Nucleoprotein-RNA Interactions.** *Cell Rep.* 2015; **11**(3): 376–89.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 58. Kirchdoerfer RN, Abelson DM, Li S, *et al.*: **Assembly of the Ebola Virus Nucleoprotein from a Chaperoned VP35 Complex.** *Cell Rep.* 2015; **12**(1): 140–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 59. Bale S, Julien JP, Bornholdt ZA, *et al.*: **Marburg virus VP35 can both fully coat the backbone and cap the ends of dsRNA for interferon antagonism.** *PLoS Pathog.* 2012; **8**(9): e1002916.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 60. **F** Edwards MR, Liu G, Mire CE, *et al.*: **Differential Regulation of Interferon Responses by Ebola and Marburg Virus VP35 Proteins.** *Cell Rep.* 2016; **14**(7): 1632–40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 61. **F** Guito JC, Albariño CG, Chakrabarti AK, *et al.*: **Novel activities by ebolavirus and marburgvirus interferon antagonists revealed using a standardized in vitro reporter system.** *Virology.* 2017; **501**: 147–65.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 62. Xue Q, Zheng QC, Zhang JL, *et al.*: **Exploring the mechanism how Marburg virus VP35 recognizes and binds dsRNA by molecular dynamics simulations and free energy calculations.** *Biopolymers.* 2014; **101**(8): 849–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
 63. **F** Koehler A, Pfeiffer S, Kolesnikova L, *et al.*: **Analysis of the multifunctionality of Marburg virus VP40.** *J Gen Virol.* 2018; **99**(12): 1614–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 64. Wijesinghe KJ, Stahelin RV: **Investigation of the Lipid Binding Properties of the Marburg Virus Matrix Protein VP40.** *J Virol.* 2015; **90**(6): 3074–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 65. **F** Wijesinghe KJ, Urata S, Bhattarai N, *et al.*: **Detection of lipid-induced structural changes of the Marburg virus matrix protein VP40 using hydrogen/deuterium exchange-mass spectrometry.** *J Biol Chem.* 2017; **292**(15): 6108–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 66. **F** Liang J, Sagum CA, Bedford MT, *et al.*: **Chaperone-Mediated Autophagy Protein BAG3 Negatively Regulates Ebola and Marburg VP40-Mediated Egress.** *PLoS Pathog.* 2017; **13**(1): e1006132.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 67. Banadyga L, Dolan MA, Ebihara H: **Rodent-Adapted Filoviruses and the Molecular Basis of Pathogenesis.** *J Mol Biol.* 2016; **428**(17): 3449–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 68. Lofts LL, Wells JB, Bavari S, *et al.*: **Key genomic changes necessary for an in vivo lethal mouse marburgvirus variant selection process.** *J Virol.* 2011; **85**(8): 3905–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 69. Warfield KL, Bradfute SB, Wells J, *et al.*: **Development and characterization of a mouse model for Marburg hemorrhagic fever.** *J Virol.* 2009; **83**(13): 6404–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 70. Valmas C, Basler CF: **Marburg virus VP40 antagonizes interferon signaling in a species-specific manner.** *J Virol.* 2011; **85**(9): 4309–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 71. Feagins AR, Basler CF: **Amino Acid Residue at Position 79 of Marburg Virus VP40 Confers Interferon Antagonism in Mouse Cells.** *J Infect Dis.* 2015; **212** Suppl 2: S219–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 72. Feagins AR, Basler CF: **The VP40 protein of Marburg virus exhibits impaired budding and increased sensitivity to human tetherin following mouse adaptation.** *J Virol.* 2014; **88**(24): 14440–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 73. Koehler A, Kolesnikova L, Welzel U, *et al.*: **A Single Amino Acid Change in the Marburg Virus Matrix Protein VP40 Provides a Replicative Advantage in a Species-Specific Manner.** *J Virol.* 2016; **90**(3): 1444–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 74. Koehler A, Kolesnikova L, Becker S: **An active site mutation increases the polymerase activity of the guinea pig-lethal Marburg virus.** *J Gen Virol.* 2016; **97**(10): 2494–500.
[PubMed Abstract](#) | [Publisher Full Text](#)
 75. O'Hearn A, Wang M, Cheng H, *et al.*: **Role of EXT1 and Glycosaminoglycans in the Early Stage of Filovirus Entry.** *J Virol.* 2015; **89**(10): 5441–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 76. Cheng H, Koning K, O'Hearn A, *et al.*: **A parallel genome-wide RNAi screening strategy to identify host proteins important for entry of Marburg virus and H5N1 influenza virus.** *Virol J.* 2015; **12**: 194.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 77. **F** Kondoh T, Letko M, Munster VJ, *et al.*: **Single-Nucleotide Polymorphisms in Human NPC1 Influence Filovirus Entry Into Cells.** *J Infect Dis.* 2018; **218**(suppl_5): S397–S402.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 78. Olson MA, Lee MS, Yeh IC: **Membrane insertion of fusion peptides from Ebola and Marburg viruses studied by replica-exchange molecular dynamics simulations.** *J Comput Chem.* 2017; **38**(16): 1342–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
 79. Gniß K, Fiedler M, Krämer-Kühl A, *et al.*: **Analysis of determinants in filovirus glycoproteins required for tetherin antagonism.** *Viruses.* 2014; **6**(4): 1654–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 80. Noyori O, Matsuno K, Kajihara M, *et al.*: **Differential potential for envelope glycoprotein-mediated steric shielding of host cell surface proteins among filoviruses.** *Virology.* 2013; **446**(1–2): 152–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
 81. Noyori O, Nakayama E, Maruyama J, *et al.*: **Suppression of Fas-mediated apoptosis via steric shielding by filovirus glycoproteins.** *Biochem Biophys Res Commun.* 2013; **441**(4): 994–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 82. Dye JM, Herbert AS, Kuehne AI, *et al.*: **Postexposure antibody prophylaxis protects nonhuman primates from filovirus disease.** *Proc Natl Acad Sci U S A.* 2012; **109**(13): 5034–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 83. **F** Mire CE, Geisbert JB, Borisevich V, *et al.*: **Therapeutic treatment of Marburg and Ravn virus infection in nonhuman primates with a human monoclonal antibody.** *Sci Transl Med.* 2017; **9**(384): pii: eaai8711.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 84. **F** King LB, West BR, Schendel SL, *et al.*: **The structural basis for filovirus neutralization by monoclonal antibodies.** *Curr Opin Immunol.* 2018; **53**: 196–202.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 85. **F** King LB, Fusco ML, Flyak AI, *et al.*: **The Marburgvirus-Neutralizing Human Monoclonal Antibody MR191 Targets a Conserved Site to Block Virus Receptor Binding.** *Cell Host Microbe.* 2018; **23**(1): 101–109.e4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 86. **F** Sangha AK, Dong J, Williamson L, *et al.*: **Role of Non-local Interactions between CDR Loops in Binding Affinity of MR78 Antibody to Marburg Virus Glycoprotein.** *Structure.* 2017; **25**(12): 1820–1828.e2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 87. Fusco ML, Hashiguchi T, Cassan R, *et al.*: **Protective mAbs and Cross-Reactive mAbs Raised by Immunization with Engineered Marburg Virus GPs.** *PLoS Pathog.* 2015; **11**(6): e1005016.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 88. Saphire EO, Schendel SL, Fusco ML, *et al.*: **Systematic Analysis of Monoclonal Antibodies against Ebola Virus GP Defines Features that Contribute to Protection.** *Cell.* 2018; **174**(4): 938–952.e13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 89. Zhao X, Howell KA, He S, *et al.*: **Immunization-Elicited Broadly Protective**

- Antibody Reveals Ebola virus Fusion Loop as a Site of Vulnerability.** *Cell*. 2017; 169(5): 891–904.e15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
90. Bornholdt ZA, Herbert AS, Mire CE, *et al.*: **A Two-Antibody Pan-Ebola Virus Cocktail Confers Broad Therapeutic Protection in Ferrets and Nonhuman Primates.** *Cell Host Microbe*. 2019; 25(1): 49–58.e5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 91. Keck ZY, Enterlein SG, Howell KA, *et al.*: **Macaque Monoclonal Antibodies Targeting Novel Conserved Epitopes within Filovirus Glycoprotein.** *J Virol*. 2016; 90(1): 279–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 92. Holtsberg FW, Shulenin S, Vu H, *et al.*: **Pan-ebolavirus and Pan-filovirus Mouse Monoclonal Antibodies: Protection against Ebola and Sudan Viruses.** *J Virol*. 2015; 90(1): 266–78.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 93. Wec AZ, Herbert AS, Murin CD, *et al.*: **Antibodies from a Human Survivor Define Sites of Vulnerability for Broad Protection against Ebolaviruses.** *Cell*. 2017; 169(5): 878–890.e15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 94. **F** Howell KA, Qiu X, Brannan JM, *et al.*: **Antibody Treatment of Ebola and Sudan Virus Infection via a Uniquely Exposed Epitope within the Glycoprotein Receptor-Binding Site.** *Cell Rep*. 2016; 15(7): 1514–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 95. Brannan JM, He S, Howell KA, *et al.*: **Post-exposure immunotherapy for two ebolaviruses and Marburg virus in nonhuman primates.** *Nat Commun*. 2019; 10(1): 105.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 96. **F** Marzi A, Menicucci AR, Engelmann F, *et al.*: **Protection Against Marburg Virus Using a Recombinant VSV-Vaccine Depends on T and B Cell Activation.** *Front Immunol*. 2019; 9: 3071.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 97. Connor JH, Yen J, Caballero IS, *et al.*: **Transcriptional Profiling of the Immune Response to Marburg Virus Infection.** *J Virol*. 2015; 89(19): 9865–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 98. **F** Stonier SW, Herbert AS, Kuehne AI, *et al.*: **Marburg virus survivor immune responses are Th1 skewed with limited neutralizing antibody responses.** *J Exp Med*. 2017; 214(9): 2563–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 99. Thompson KA, Yin J: **Population dynamics of an RNA virus and its defective interfering particles in passage cultures.** *Virus J*. 2010; 7: 257.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 100. Caballero IS, Yen JY, Hensley LE, *et al.*: **Lassa and Marburg viruses elicit distinct host transcriptional responses early after infection.** *BMC Genomics*. 2014; 15: 960.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 101. **F** Natesan M, Jensen SM, Keasey SL, *et al.*: **Human Survivors of Disease Outbreaks Caused by Ebola or Marburg Virus Exhibit Cross-Reactive and Long-Lived Antibody Responses.** *Clin Vaccine Immunol*. 2016; 23(8): 717–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 102. Sobarzo A, Eskira Y, Herbert AS, *et al.*: **Immune memory to Sudan virus: comparison between two separate disease outbreaks.** *Viruses*. 2015; 7(1): 37–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 103. Mire CE, Geisbert JB, Agans KN, *et al.*: **Durability of a vesicular stomatitis virus-based marburg virus vaccine in nonhuman primates.** *PLoS One*. 2014; 9(4): e94355.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 104. Amman BR, Nyakarahuka L, McElroy AK, *et al.*: **Marburgvirus resurgence in Kitaka Mine bat population after extermination attempts, Uganda.** *Emerging Infect Dis*. 2014; 20(10): 1761–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 105. Towner JS, Amman BR, Sealy TK, *et al.*: **Isolation of genetically diverse Marburg viruses from Egyptian fruit bats.** *PLoS Pathog*. 2009; 5(7): e1000536.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 106. Amman BR, Carroll SA, Reed ZD, *et al.*: **Seasonal pulses of Marburg virus circulation in juvenile *Rousettus aegyptiacus* bats coincide with periods of increased risk of human infection.** *PLoS Pathog*. 2012; 8(10): e1002877.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 107. **F** Paweska JT, Jansen van Vuren P, Kemp A, *et al.*: **Marburg Virus Infection in Egyptian Rousette Bats, South Africa, 2013–2014.** *Emerging Infect Dis*. 2018; 24(6): 1134–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 108. **F** Changula K, Kajihara M, Mori-Kajihara A, *et al.*: **Seroprevalence of Filovirus Infection of *Rousettus aegyptiacus* Bats in Zambia.** *J Infect Dis*. 2018; 218(suppl_5): S312–S317.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 109. Ogawa H, Miyamoto H, Nakayama E, *et al.*: **Seroepidemiological Prevalence of Multiple Species of Filoviruses in Fruit Bats (*Eidolon helvum*) Migrating in Africa.** *J Infect Dis*. 2015; 212 Suppl 2: S101–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 110. Schuh AJ, Amman BR, Apanaskevich DA, *et al.*: **No evidence for the involvement of the argasid tick *Ornithodoros faini* in the enzootic maintenance of marburgvirus within Egyptian rousette bats *Rousettus aegyptiacus*.** *Parasit Vectors*. 2016; 9: 128.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 111. Paweska JT, Jansen van Vuren P, Masumu J, *et al.*: **Virological and serological findings in *Rousettus aegyptiacus* experimentally inoculated with vero cells-adapted hogan strain of Marburg virus.** *PLoS One*. 2012; 7(9): e45479.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 112. **F** Amman BR, Jones ME, Sealy TK, *et al.*: **Oral shedding of Marburg virus in experimentally infected Egyptian fruit bats (*Rousettus aegyptiacus*).** *J Wildl Dis*. 2015; 51(1): 113–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 113. Jones ME, Schuh AJ, Amman BR, *et al.*: **Experimental Inoculation of Egyptian Rousette Bats (*Rousettus aegyptiacus*) with Viruses of the *Ebolavirus* and *Marburgvirus* Genera.** *Viruses*. 2015; 7(7): 3420–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 114. Paweska JT, Jansen van Vuren P, Fenton KA, *et al.*: **Lack of Marburg Virus Transmission From Experimentally Infected to Susceptible In-Contact Egyptian Fruit Bats.** *J Infect Dis*. 2015; 212 Suppl 2: S109–18.
[PubMed Abstract](#) | [Publisher Full Text](#)
 115. **F** Schuh AJ, Amman BR, Sealy TK, *et al.*: **Egyptian rousette bats maintain long-term protective immunity against Marburg virus infection despite diminished antibody levels.** *Sci Rep*. 2017; 7(1): 8763.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 116. Storm N, Jansen Van Vuren P, Markotter W, *et al.*: **Antibody Responses to Marburg Virus in Egyptian Rousette Bats and Their Role in Protection against Infection.** *Viruses*. 2018; 10(2): pii: E73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 117. **F** Schuh AJ, Amman BR, Jones ME, *et al.*: **Modelling filovirus maintenance in nature by experimental transmission of Marburg virus between Egyptian rousette bats.** *Nat Commun*. 2017; 8: 14446.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 118. Jordan I, Horn D, Oehmke S, *et al.*: **Cell lines from the Egyptian fruit bat are permissive for modified vaccinia Ankara.** *Virus Res*. 2009; 145(1): 54–62.
[PubMed Abstract](#) | [Publisher Full Text](#)
 119. Kühl A, Hoffmann M, Müller MA, *et al.*: **Comparative analysis of Ebola virus glycoprotein interactions with human and bat cells.** *J Infect Dis*. 2011; 204 Suppl 3: S840–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 120. Hölzer M, Krähling V, Amman F, *et al.*: **Differential transcriptional responses to Ebola and Marburg virus infection in bat and human cells.** *Sci Rep*. 2016; 6: 34589.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 121. **F** Arnold CE, Guito JC, Altamura LA, *et al.*: **Transcriptomics Reveal Antiviral Gene Induction in the Egyptian Rousette Bat Is Antagonized *In Vitro* by Marburg Virus Infection.** *Viruses*. 2018; 10(11): pii: E607.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 122. Kuzmin IV, Schwarz TM, Ilinykh PA, *et al.*: **Innate Immune Responses of Bat and Human Cells to Filoviruses: Commonalities and Distinctions.** *J Virol*. 2017; 91(8): pii: e02471-16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 123. Ng M, Ndungo E, Kaczmarek ME, *et al.*: **Filovirus receptor NPC1 contributes to species-specific patterns of ebolavirus susceptibility in bats.** *eLife*. 2015; 4: pii: e11785.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 124. Burk R, Bollinger L, Johnson JC, *et al.*: **Neglected filoviruses.** *FEMS Microbiol Rev*. 2016; 40(4): 494–519.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 125. Marzi A, Chadinah S, Haddock E, *et al.*: **Recently Identified Mutations in the Ebola Virus-Makona Genome Do Not Alter Pathogenicity in Animal Models.** *Cell Rep*. 2018; 23(6): 1806–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 126. Urbanowicz RA, McClure CP, Sakuntabhai A, *et al.*: **Human Adaptation of Ebola Virus during the West African Outbreak.** *Cell*. 2016; 167(4): 1079–1087.e5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 127. Dietzel E, Schudt G, Krähling V, *et al.*: **Functional Characterization of Adaptive Mutations during the West African Ebola Virus Outbreak.** *J Virol*. 2017; 91(2): pii: e01913-16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 128. Wang MK, Lim SY, Lee SM, *et al.*: **Biochemical Basis for Increased Activity of Ebola Glycoprotein in the 2013–16 Epidemic.** *Cell Host Microbe*. 2017; 21(3): 367–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 129. Siragam V, Wong G, Qiu XG: **Animal models for filovirus infections.** *Zool Res*. 2018; 39(1): 15–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 130. Cross RW, Fenton KA, Geisbert TW: **Small animal models of filovirus disease: recent advances and future directions.** *Expert Opin Drug Discov*. 2018; 13(11): 1027–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
 131. Yamaoka S, Banadyga L, Bray M, *et al.*: **Small Animal Models for Studying Filovirus Pathogenesis.** *Curr Top Microbiol Immunol*. 2017; 411: 195–227.
[PubMed Abstract](#) | [Publisher Full Text](#)

132. Atkins C, Miao J, Kalveram B, *et al.*: **Natural History and Pathogenesis of Wild-Type Marburg Virus Infection in STAT2 Knockout Hamsters.** *J Infect Dis.* 2018; 218(suppl_5): S438–S447.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
133. Lavender KJ, Williamson BN, Saturday G, *et al.*: **Pathogenicity of Ebola and Marburg Viruses Is Associated With Differential Activation of the Myeloid Compartment in Humanized Triple Knockout-Bone Marrow, Liver, and Thymus Mice.** *J Infect Dis.* 2018; 218(suppl_5): S409–S417.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
134. Smither SJ, Nelson M, Eastaugh L, *et al.*: **Experimental respiratory Marburg virus haemorrhagic fever infection in the common marmoset (*Callithrix jacchus*).** *Int J Exp Pathol.* 2013; 94(2): 156–68.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
135. Cross RW, Mire CE, Borisevich V, *et al.*: **The Domestic Ferret (*Mustela putorius furo*) as a Lethal Infection Model for 3 Species of *Ebolavirus*.** *J Infect Dis.* 2016; 214(4): 565–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
136. **F** Wong G, Zhang Z, He S, *et al.*: **Marburg and Ravn Virus Infections Do Not Cause Observable Disease in Ferrets.** *J Infect Dis.* 2018; 218(suppl_5): S471–S474.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
137. Kozak R, He S, Kroeker A, *et al.*: **Ferrets Infected with Bundibugyo Virus or Ebola Virus Recapitulate Important Aspects of Human Filovirus Disease.** *J Virol.* 2016; 90(20): 9209–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
138. **F** Cross RW, Mire CE, Agans KN, *et al.*: **Marburg and Ravn Viruses Fail to Cause Disease in the Domestic Ferret (*Mustela putorius furo*).** *J Infect Dis.* 2018; 218(suppl_5): S448–S452.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
139. **F** Coffin KM, Liu J, Warren TK, *et al.*: **Persistent Marburg Virus Infection in the Testes of Nonhuman Primate Survivors.** *Cell Host Microbe.* 2018; 24(3): 405–416.e3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
140. **F** Nicholas VV, Rosenke R, Feldmann F, *et al.*: **Distinct Biological Phenotypes of Marburg and Ravn Virus Infection in Macaques.** *J Infect Dis.* 2018; 218(suppl_5): S458–S465.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
141. Blair PW, Keshthkar-Jahromi M, Psoter KJ, *et al.*: **Virulence of Marburg Virus Angola Compared to Mt. Elgon (Musoke) in Macaques: A Pooled Survival Analysis.** *Viruses.* 2018; 10(11): pii: E658.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
142. Daddario-DiCaprio KM, Geisbert TW, Geisbert JB, *et al.*: **Cross-protection against Marburg virus strains by using a live, attenuated recombinant vaccine.** *J Virol.* 2006; 80(19): 9659–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
143. **F** Daddario-DiCaprio KM, Geisbert TW, Ströher U, *et al.*: **Postexposure protection against Marburg haemorrhagic fever with recombinant vesicular stomatitis virus vectors in non-human primates: an efficacy assessment.** *Lancet.* 2006; 367(9520): 1399–404.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
144. Cross RW, Fenton KA, Geisbert JB, *et al.*: **Comparison of the Pathogenesis of the Angola and Ravn Strains of Marburg Virus in the Outbred Guinea Pig Model.** *J Infect Dis.* 2015; 212 Suppl 2: S258–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
145. Thi EP, Mire CE, Lee AC, *et al.*: **siRNA rescues nonhuman primates from advanced Marburg and Ravn virus disease.** *J Clin Invest.* 2017; 127(12): 4437–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
146. Qiu X, Wong G, Audet J, *et al.*: **Establishment and characterization of a lethal mouse model for the Angola strain of Marburg virus.** *J Virol.* 2014; 88(21): 12703–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
147. Wong G, Cao WG, He SH, *et al.*: **Development and characterization of a guinea pig model for Marburg virus.** *Zool Res.* 2018; 39(1): 32–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
148. Marzi A, Banadyga L, Haddock E, *et al.*: **A hamster model for Marburg virus infection accurately recapitulates Marburg hemorrhagic fever.** *Sci Rep.* 2016; 6: 39214.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
149. Fernando L, Qiu X, Melito PL, *et al.*: **Immune Response to Marburg Virus Angola Infection in Nonhuman Primates.** *J Infect Dis.* 2015; 212 Suppl 2: S234–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
150. Johnston SC, Lin KL, Twenhafel NA, *et al.*: **Dose Response of MARV/Angola Infection in Cynomolgus Macaques following IM or Aerosol Exposure.** *PLoS One.* 2015; 10(9): e0138843.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
151. Lin KL, Twenhafel NA, Connor JH, *et al.*: **Temporal Characterization of Marburg Virus Angola Infection following Aerosol Challenge in Rhesus Macaques.** *J Virol.* 2015; 89(19): 9875–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
152. Alfson KJ, Avena LE, Delgado J, *et al.*: **A Single Amino Acid Change in the Marburg Virus Glycoprotein Arises during Serial Cell Culture Passages and Attenuates the Virus in a Macaque Model of Disease.** *mSphere.* 2018; 3(1): pii: e00401-17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
153. **F** Cooper TK, Sword J, Johnson JC, *et al.*: **New Insights Into Marburg Virus Disease Pathogenesis in the Rhesus Macaque Model.** *J Infect Dis.* 2018; 218(suppl_5): S423–S433.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
154. Woolsey C, Geisbert JB, Matassov D, *et al.*: **Postexposure Efficacy of Recombinant Vesicular Stomatitis Virus Vectors Against High and Low Doses of Marburg Virus Variant Angola in Nonhuman Primates.** *J Infect Dis.* 2018; 218(suppl_5): S582–S587.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
155. Ewers EC, Pratt WD, Twenhafel NA, *et al.*: **Natural History of Aerosol Exposure with Marburg Virus in Rhesus Macaques.** *Viruses.* 2016; 8(4): 87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
156. **F** Wei H, Audet J, Wong G, *et al.*: **Deep-sequencing of Marburg virus genome during sequential mouse passaging and cell-culture adaptation reveals extensive changes over time.** *Sci Rep.* 2017; 7(1): 3390.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
157. Kuhn JH: **Filoviruses. A compendium of 40 years of epidemiological, clinical, and laboratory studies.** *Arch Virol Suppl.* editor: Springer Verlag; 2008; 20: 13–360.
[PubMed Abstract](#) | [Publisher Full Text](#)
158. Alfson KJ, Avena LE, Beadles MW, *et al.*: **Particle-to-PFU ratio of Ebola virus influences disease course and survival in cynomolgus macaques.** *J Virol.* 2015; 89(13): 6773–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
159. Geisbert TW, Daddario-DiCaprio KM, Geisbert JB, *et al.*: **Marburg virus Angola infection of rhesus macaques: pathogenesis and treatment with recombinant nematode anticoagulant protein c2.** *J Infect Dis.* 2007; 196 Suppl 2: S372–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
160. Alonso JA, Patterson JL: **Sequence variability in viral genome non-coding regions likely contribute to observed differences in viral replication amongst MARV strains.** *Virology.* 2013; 440(1): 51–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
161. Martini GA, Siegert R, editors: **Marburg virus disease.** New York: Springer; 1971.
[Publisher Full Text](#)
162. Vetter P, Kaiser L, Schibler M, *et al.*: **Sequelae of Ebola virus disease: the emergency within the emergency.** *Lancet Infect Dis.* 2016; 16(6): e82–e91.
[PubMed Abstract](#) | [Publisher Full Text](#)
163. Gear JS, Cassel GA, Gear AJ, *et al.*: **Outbreaks of Marburg virus disease in Johannesburg.** *Br Med J.* 1975; 4(5995): 489–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

- 1. M. Javad Aman**
Integrated Biotherapeutics, Inc., Rockville, MD, 20850, USA
Competing Interests: No competing interests were disclosed.
- 2. Chad E Mire**
Department of Microbiology and Immunology, University of Texas Medical Branch, Galveston, USA
Competing Interests: No competing interests were disclosed.
- 3. Ilhem Messaoudi**
Department of Molecular Biology and Biochemistry, University of California, Irvine, CA, USA
Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research