REVIEW

A new look at IgE beyond allergies [version 1; peer review: 2 approved]

Andrea J. Luker¹, Joseph C. Lownik^{1,2}, Daniel H. Conrad¹, Rebecca K. Martin¹

¹Department of Microbiology and Immunology, Virginia Commonwealth University, Richmond, VA, USA

²Center for Clinical and Translational Research, Virginia Commonwealth University, Richmond, VA, USA

V1 First published: 24 May 2019, 8(F1000 Faculty Rev):736
<https://doi.org/10.12688/f1000research.18186.1>

Latest published: 24 May 2019, 8(F1000 Faculty Rev):736
<https://doi.org/10.12688/f1000research.18186.1>

Abstract

Immunoglobulin E (IgE), though constitutively present at low levels, is most commonly studied in atopic disease where it plays a vital role in mast cell degranulation and in initiating a T helper 2 (Th2) response. With the advent of better detection assays, however, researchers are discovering the importance of IgE in actively contributing to many disease states and pathologies. This review will discuss the latest findings in IgE beyond its role in allergies and recently discovered roles for IgE in its cell-bound form on FcεRI-expressing effector cells like monocytes and dendritic cells. In terms of parasites, we will discuss helminth-induced IgE that appears to protect the worms from immune recognition and a tick-borne illness that elicits an IgE response against red meat. Next, we describe recent findings of how auto-reactive IgE can contribute to the progression of lupus and induce organ damage. Finally, we summarize the emerging roles of IgE in tumor surveillance and antibody-dependent cytotoxicity. We additionally discuss recent or ongoing clinical trials that either target harmful IgE or use the unique characteristics of the isotype.

Keywords

IgE, helminth, alpha-gal, auto-immunity, lupus, tumor, cancer

Open Peer Review

Approval Status

	1	2
version 1 24 May 2019

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Yui-Hsi Wang**, Sanofi, Cambridge, USA
2. **Edda Fiebiger**, Boston Children's Hospital, Boston,, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Rebecca K. Martin (rebecca.martin@vcuhealth.org)

Author roles: **Luker AJ:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Lownik JC:** Conceptualization, Writing – Original Draft Preparation; **Conrad DH:** Conceptualization, Supervision, Writing – Review & Editing; **Martin RK:** Conceptualization, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: RKM receives funding from the National Institute of Allergy and Infectious Diseases (NIH-NIAID 1R56AI139658-01A1 and NIH-NIAID R01AI18697A1-33-38).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 Luker AJ *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Luker AJ, Lownik JC, Conrad DH and Martin RK. **A new look at IgE beyond allergies [version 1; peer review: 2 approved]** F1000Research 2019, 8(F1000 Faculty Rev):736 <https://doi.org/10.12688/f1000research.18186.1>

First published: 24 May 2019, 8(F1000 Faculty Rev):736 <https://doi.org/10.12688/f1000research.18186.1>

Introduction

There are many hypotheses regarding the origin of the immunoglobulin E (IgE) response and its conservation throughout evolution. The more widely accepted theories include host protection against metazoan parasites, environmental toxins, and venoms. Although IgE-mediated degranulation is an effective means of initiating a protective immune response, the process can also cause significant damage to healthy tissue. IgE has long been viewed as the notorious Ig behind type I hypersensitivities and related diseases such as allergic rhinitis, anaphylaxis, and asthma. Classic IgE-mediated activation occurs through FcεRI on mast cells and basophils. Multivalent antigens cross-link the receptor-bound IgE to trigger degranulation and the release of pre-formed mediators. This pro-inflammatory milieu recruits immune cells and initiates a T helper 2 (Th2) immune response.

Today, millions of individuals suffer from allergies, asthma, and auto-immunity. This rapid rise in atopic disease may be explained by the “hygiene hypothesis”¹, which has been widely reviewed²⁻⁵. Briefly, this theory states that lower incidences of infection in developed countries lead to inappropriate immune activation by IgE against common environmental antigens and self-antigens. This hypothesis can be expanded to include the recent upsurge in food allergies, especially among young children^{6,7}. Recent research has suggested that a healthy and diverse gut microbiota can prevent the development of IgE-mediated food sensitivities^{8,9}.

Because asthma and allergies are a wide-scale problem across the globe, the majority of research into IgE revolves around these pathologies. Advances in analytical techniques as well as expanding scientific knowledge, however, have propelled a new wave of research into the role of IgE beyond the field of allergies. This review will span the latest findings on IgE in non-atopic pathological conditions. We will discuss a new take on “protective” IgE in helminth infection, revelations within a tick-borne disease, self-reactive IgE in auto-immunity, and a novel role for IgE in both preventing and treating cancer. Furthermore, we will discuss recent or ongoing clinical trials using IgE as a therapeutic tool in a variety of these scenarios.

IgE in helminth infection: Whom is it protecting?

It is traditionally understood that antibodies are produced by B2 cells. These cells undergo isotype switching and affinity maturation to produce antibodies with very high specificity toward an antigen. The resulting high-affinity, antigen-specific antibody response is important within the adaptive arm of the immune system¹⁰. In the mouse, however, there is a lesser-studied subset of B cell, the innate-like B1 cell, whose importance has begun to be revealed. Reports have described B1 cells as participants in resistance against intestinal bacteria, defense from viral pathogens, generating auto-immune responses, and responding to parasites¹¹⁻¹⁴. In each case, the role of the B1 cell is to immediately respond to infection with the production of low-affinity IgM or IgG antibodies¹⁵. Their B-cell receptor (BCR) repertoire is composed of vastly poly-specific receptors, meaning that B1 BCRs recognize a broad range of antigens¹⁵. The resulting antibodies, however, are a much lower affinity than those produced by B2 cells.

Gastrointestinal helminths are widely known to elicit a robust antigen-specific IgE response, although the reasons why are unclear because mast cells do not appear to play a major role in helminth clearance. Recently, our group showed that helminth infection also induces the production of low-affinity, poly-specific IgE¹⁶. Interestingly, this IgE is not derived from traditional B2 cells but instead is made by the innate-like B1 cells. This challenges the current understanding that B1 cells produce only IgM and IgG. Furthermore, Martin *et al.* revealed a competition between high-affinity B2 IgE and low-affinity B1 IgE to trigger or block mast cell degranulation, respectively¹⁶. Although mast cell degranulation does not clear the infection alone, it can enhance the anti-helminth response. We hypothesized that helminth may have evolved to induce this low-affinity B1 IgE within the host to prevent mast cell degranulation as a protective means of immune evasion.

The translational relevance of murine B1 cells in the human system, however, remains controversial because they are not well defined. The current markers used to identify human B1 cells are similarly found on activated cells and memory populations¹⁷. Human studies are also limited to circulating cells for collection and examination. For these reasons, it remains to be seen whether B1-derived IgE plays a clinical role in human disease.

IgE and ticks

Scientists have recently discovered a bizarre occurrence in certain tick-bitten individuals within the southern US and beyond. These patients develop a delayed anaphylactic reaction against red meat, and the average onset is 3 to 6 hours after ingestion¹⁸. The underlying cause is an IgE response against galactose-α-1,3-galactose (α-Gal), a carbohydrate moiety commonly found on glycoproteins. In particular, α-Gal is widely expressed within mammals but not within humans¹⁹. The unique nature of the intense hypersensitivity reaction to α-Gal has remained a curiosity. Some hypothesize that certain ticks produce α-Gal in order to hide from the immune system of mammals, which are tolerized to see α-Gal as “self”¹⁹. Humans, which are considered accidental hosts for ticks, however, will recognize the tick-expressed α-Gal as foreign and mount an immune response.

Additional questions remain surrounding the sensitization process following the tick bite as well as the delay in symptomology. The IgE responsible for red meat allergy is unique because of its specificity. Instead of recognizing a specific epitope on a specific protein, this IgE targets the α-Gal moiety that can be present on a variety of proteins and even lipids. This provides potential for “cross-reactivity” among all mammalian meats, including beef and pork.

The several-hour delay was originally thought to be caused by ineffective dendritic cell (DC) function. A recent study, however, showed that the presence of α-Gal on modified proteins actually increases the efficiency of antigen uptake²⁰. The authors further suggest that α-Gal may protect proteins from degradation, thereby slowing the process of major histocompatibility complex (MHC) presentation and antigen recognition²⁰. As mentioned before, α-Gal is also present on lipids. Fattier meats have been reported

to elicit more severe reactions in patients with red meat allergy²¹. One final theory attributes the late symptoms to the inherently slow metabolism of fatty lipids, resulting in the delayed release of α -Gal antigens.

A hidden role for IgE in auto-immune disease

At a basic level, auto-immunity is characterized by an immune response directed against “self”. Auto-reactive B or T cells recognize proteins or DNA fragments normally found in healthy tissue, causing severe and chronic damage. We have come to understand that some of these disorders are characterized by high levels of circulating auto-reactive IgM and IgG²². With the advent of better detection assays, however, it is now apparent that auto-reactive IgE is also present within these patients. In addition, researchers have recognized some self-antigens that also show cross-reactivity with exogenous or environmental allergens through molecular mimicry. Numerous reports have highlighted the elevated IgE seen in patients with atopic dermatitis²³, chronic urticaria²⁴, rheumatoid arthritis²⁵, or bullous pemphigoid²⁶. Indeed, there have been many reviews covering the decades of IgE studies in these diseases^{7,27–30}. The latest published research, however, has focused on systemic lupus erythematosus (SLE), the most common form of lupus.

IgE and basophils

SLE is characterized by chronic inflammation due to the overproduction of auto-antibodies. These antibodies, often against nuclear factors like DNA, form immune complexes (ICs) that deposit in tissues and cause organ damage. Several studies have revealed a role for IgE in SLE, especially in terms of IgE-DNA ICs^{31–34}. Work by Juan Rivera’s group has shown an important role for basophils in both murine models and SLE patients. The authors showed that auto-reactive IgE ICs activate basophils to produce interleukin-4 (IL-4) and contribute to an overall Th2 skewing³¹. This inflammatory environment resulted in increased auto-antibody production by plasma cells and the development of severe lupus nephritis³¹. The group went on to show that IgE deficiency delayed disease onset and prolonged survival³⁵. To authenticate the clinical relevance, they found increased basophil activation in SLE patients above healthy controls as well as the abnormal presence of basophils in the lymph node and spleen of two patients with SLE^{31,35}. Furthermore, the authors showed a clear association of high levels of anti-dsDNA IgE in SLE patients with lupus nephritis and a positive association between auto-reactive IgE and clinical disease activity^{31,36}. Overall, the authors hypothesize that IgE and basophils contribute to the disease pathology by *amplifying* the immune dysfunction.

IgE and dendritic cells

Significant strides toward understanding the role of IgE in SLE were made when another group revealed a role for plasmacytoid DCs (pDCs). Traditionally, pDCs are a specialized DC subset that produce type I interferons (IFNs) in response to bacterial or viral pathogen-associated molecular patterns (PAMPs)³⁷. Henault *et al.* designed a novel anti-dsDNA IgE molecule, which they used in conjunction with free DNA to form ICs (DNA-IgE IC)³⁴. These DNA-IgE ICs bound to Fc ϵ RI on pDCs to trigger robust IFN α

production through endosomal Toll-like receptor 9 (TLR9) activation. When co-cultured with B cells, pDCs activated by DNA-IgE IC caused a significant expansion of plasma cells. Furthermore, the authors showed close proximity between pDCs and B cells in kidney biopsies from patients with SLE. This suggests that the pDC-mediated B-cell proliferation seen *in vitro* may also be operant in SLE patients *in vivo*.

As a testament to the importance of these and many other findings, a small clinical trial (ClinicalTrials.gov identifier: NCT01716312) was conducted to test the effectiveness of treating SLE with the anti-IgE monoclonal antibody, omalizumab. The investigators hypothesized that neutralization of the circulating auto-reactive IgE would limit basophil and pDC activation, thereby dampening IFN α production. In a newly published report summarizing the results, the authors describe a maintained improvement in disease activity, as evaluated by using a patient scoring system (Systemic Lupus Erythematosus Disease Activity Index 2000, or SLEDAI 2K). Upon the conclusion of the 32-week omalizumab treatment period, patient scores began to worsen. The authors additionally analyzed type I IFN gene signatures and reported a trend toward improvement in patients receiving omalizumab. Although it did not reach significance within their small cohort, this attenuation effect was enhanced in those participants with high baseline scores. Additional studies will need to extend these results to include a larger cohort of participants. The results, however, are promising and lay an important framework for verifying the pathological relevance of IgE in SLE and other auto-immune diseases.

Another potential approach to reducing the effects of circulating IgE is through the use of soluble IgE receptors. The low-affinity IgE receptor, Fc ϵ RII or CD23, is well known to exist in both soluble and membrane-bound forms. CD23 is involved in a complex regulatory axis that controls the level of production of IgE^{38,39}. The high-affinity IgE receptor, Fc ϵ RI, exists mainly as a tetramer ($\alpha\beta\gamma_2$), although some human cells express a trimeric form lacking the amplifying beta-chain ($\alpha\gamma_2$)^{40,41}. In 1993, one group created a recombinant, soluble version of the human Fc ϵ RI α ectodomain (rsFc ϵ RI α)⁴². Using rsFc ϵ RI α both *in vitro* and *in vivo*, the group was able to completely block mast cell and basophil activation⁴². In 2011, another group reported their discovery of soluble Fc ϵ RI (sFc ϵ RI) within human sera⁴³. The roughly 40-kDa soluble form described by Dehlink *et al.* consists solely of the IgE-binding alpha-chain⁴³. sFc ϵ RI has been demonstrated to compete with membrane-bound Fc ϵ RI to bind IgE^{43,44}. Indeed, the authors hypothesize that this competitive mechanism could attenuate IgE-mediated DC activation⁴¹.

Recently, Moñino-Romero *et al.* sensitized mice with IgE in the presence of sFc ϵ RI and observed a reduction in the severity of experimental anaphylaxis⁴⁴. Interestingly, sFc ϵ RI was also effective in attenuating anaphylaxis when given post-sensitization⁴⁴. Within 24 hours of administration, sFc ϵ RI caused a reduction in the level of IgE bound to the surface of circulating basophils. Although it was found to be as effective as omalizumab in blocking IgE/Fc ϵ RI interaction⁴⁴, much more research is needed to fully understand the physiological role of sFc ϵ RI.

IgE and neoplastic disease

Cancer is a disease of excessive cell proliferation. The malignant cells become invasive, disrupting organ architecture and leaching nutrients from healthy tissue. Because uncontrolled growth can lead to organ failure and death, early detection is vital for patient survival. The body's defense against malignantly transformed cells relies mainly on the activity of CD8⁺ cytotoxic lymphocytes (CTLs). Tumors, however, can induce massive immune suppression to limit the activity of these effector T cells.

To prevent cancers from growing unchecked, there is great need to find new avenues to reverse the tolerance and induce an immune response within the tumor microenvironment. A popular antibody-based therapy consists of checkpoint inhibitors, which target T cell–deactivating molecules such as CTL-associated protein-4 (CTLA-4) and programmed death ligand protein-1 (PDL-1) in order to recover cytotoxic activity. Although many antibody drugs are on the market, they are all IgG-based. With five different antibody classes, it is possible that another isotype will become the next breakthrough in treating neoplastic disease.

Several epidemiological studies have compared IgE levels with incidences of cancer^{45–47}. The overall conclusion suggests that allergic disease may grant some protection against certain types of malignancies. This has led to the emergence of research into AllergoOncology, and there are several reviews of the subject^{48–50}. Elevated levels of IgE have been reported in patients with various types of cancer^{51–53}. Although there has not been any conclusive evidence for what purpose this IgE may serve or what antigens it may be specific to, several groups have investigated the unique role of IgE in tumor protection and divulged mechanisms through a variety of effector cells.

IgE as a memory response

In a direct investigation of the role of IgE in tumor development, one group explored adenocarcinoma growth in IgE-knockout (IgE-KO) mice⁵⁴. In un-immunized mice, the authors saw no difference in the tumor growth rate compared with that of wild-type (WT) mice. However, immunization with irradiated tumor cells 2 weeks prior to proper tumor challenge caused delayed tumor growth in the WT but not IgE-KO mice. Furthermore, mice with higher levels of IgE-secreting B cells (KN1) grew smaller tumors than WT mice and were completely protected in the immunization model. This IgE-protective effect was shown to rely on FcεRI and CD8⁺ cells⁵⁴.

Using IgE to detect tumor antigen

In addition to the classically IgE-activated mast cells and basophils, other cell types express FcεRI or FcεRII (CD23) or both. In particular, DCs are vital to both immune surveillance and priming immune responses. DCs take up free antigen through phagocytosis or capture IgG ICs carrying bound antigen, a process known as receptor-mediated endocytosis⁵⁵. These exogenous antigens then can be routed through a specialized pathway for presentation on MHC class I. This process, known as cross-presentation, permits antigen-specific activation of CD8⁺ CTLs. A 2015 report described a novel mechanism by which DCs can capture low levels of *free* antigen on FcεRI-bound IgE⁵⁶.

The authors were able to successfully show cross-presentation of soluble antigen to CD8⁺ cytotoxic T cells, resulting in proliferation and CTL activity. Furthermore, using antigen-specific IgE in a tumor vaccine model resulted in prolonged survival and generation of memory response upon re-challenge⁵⁶.

Auto-reactive IgE as a protective force

Crawford *et al.* showed a very different role for IgE in preventing epithelial carcinogenesis⁵⁷. Following topical application of a carcinogen, the group discovered an unexpected accumulation of IgE within the skin. DNA damage within the epithelial cells triggered a unique mechanism whereby γδ lymphocytes induced draining lymph node (dLN) B cells to produce a robust repertoire of auto-reactive IgE. Using mice that lacked either IgE (*IgH-7^{-/-}*) or FcεRI (*FcεRIα^{-/-}*), they specifically showed that IgE signaling was playing a protective role, as the genetic-deficient mice developed more aggressive tumors⁵⁷. This gives important insight into a previously unexplored immune mechanism that prevents the progression to epithelial malignancy.

Receptor-mediated cytotoxicity

In experiments using purified human peripheral blood mononuclear cells (PBMCs) transferred into nude mice, Karagiannis *et al.* showed that treatment with tumor antigen-specific IgE (MOv18) caused monocyte recruitment into the tumor site⁵⁸. The group also reported prolonged survival in treated mice, explained in part by an increase in antibody-dependent cell-mediated cytotoxicity (ADCC) that they observed *in vitro*. Interestingly, when cytotoxicity experiments were performed with non-specific IgE or PBMCs obtained from allergic patients, there was a significant decrease in activity⁵⁸. The same group later found that IgE can induce both monocyte-driven ADCC and antibody-dependent cell-mediated phagocytosis of ovarian tumor cells through FcεRI and CD23, respectively⁵².

An ongoing phase I clinical trial is using this chimeric MOv18 IgE, specific for the cancer-associated folate receptor-α⁵⁹. Another group is working to optimize the process of cloning and mass-producing recombinant IgE for use in pre-clinical and clinical studies⁶⁰. Together, these experiments suggest that tumor antigen-specific IgE can play a powerful role in eliciting and maintaining an immune response at the tumor site.

Conclusions and Future directions

Although IgE is normally the least abundant antibody found in circulation because of its short half-life, there are certain conditions that exhibit high levels of IgE. Typically, we associate this increased IgE titer with atopic diseases such as allergies and asthma. Furthermore, there is the additional nuance of discerning between antigen-specific and total IgE. This was particularly important in understanding the competition between antigen-specific B2 IgE and poly-clonal B1 IgE in helminth infection. This lesson can be applied to the many epidemiological studies of helminth infection in allergic patients. Whereas one study may find a positive correlation in IgE, another may not. Most likely, the difference in findings is due to the examination of total IgE rather than helminth-specific IgE. The additional translation between

animal studies with helminth infection and human studies has been difficult, as experimental outcomes in human studies are minimal or self-described observations, and longevity studies are hard to find.

Extremely high levels of other self-specific isotypes, such as IgM and IgG, are characteristic of certain auto-immune diseases. It was only recently that advances in basic research have allowed us to unmask the presence of auto-reactive IgE in a variety of pathologies. The expanded use of omalizumab in these conditions is seeing positive results with few side effects and thus will likely result in a powerful adjuvant therapy for diseases like SLE.

Multiple groups have uncovered unique roles for IgE in detecting malignant cells, slowing tumor progression, and even maintaining a memory response. An interesting feature of allergies is the lifelong sensitivity that can continue for years even after removing exposure to the antigen. Although it is unclear exactly

how this long-term hypersensitivity is maintained, the high-affinity memory response could be a beneficial avenue for surveying for tumor recurrence. Additionally, because IgE binds so tightly to its high-affinity receptor, FcεRI, cell-bound IgE could be retained within the tumor site for longer than other antibody classes. This could be a novel means of extending the activity of therapeutic drugs as well as a targeted means to elicit an immune response within the suppressive tumor microenvironment.

Grant information

RKM receives funding from the National Institute of Allergy and Infectious Diseases (NIH-NIAID 1R56AI139658-01A1 and NIH-NIAID R01AI18697A1-33-38).

The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Strachan DP: **Hay fever, hygiene, and household size.** *BMJ.* 1989; **299**(6710): 1259–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Versini M, Jeandel PY, Bashi T, *et al.*: **Unraveling the Hygiene Hypothesis of helminthes and autoimmunity: origins, pathophysiology, and clinical applications.** *BMC Med.* 2015; **13**: 81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Leong RW, Mitrev N, Ko Y: **Hygiene Hypothesis: Is the Evidence the Same All Over the World?** *Dig Dis.* 2016; **34**(1–2): 35–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Brooks C, Pearce N, Douwes J: **The hygiene hypothesis in allergy and asthma: an update.** *Curr Opin Allergy Clin Immunol.* 2013; **13**(1): 70–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Okada H, Kuhn C, Feillet H, *et al.*: **The ‘hygiene hypothesis’ for autoimmune and allergic diseases: an update.** *Clin Exp Immunol.* 2010; **160**(1): 1–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Anvari S, Miller J, Yeh CY, *et al.*: **IgE-Mediated Food Allergy.** *Clin Rev Allergy Immunol.* 2018; 1–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Pascal M, Perez-Gordo M, Caballero T, *et al.*: **Microbiome and Allergic Diseases.** *Front Immunol.* 2018; **9**: 1584.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Plunkett CH, Nagler CR: **The Influence of the Microbiome on Allergic Sensitization to Food.** *J Immunol.* 2017; **198**(2): 581–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Molloy J, Allen K, Collier F, *et al.*: **The potential link between gut microbiota and IgE-mediated food allergy in early life.** *Int J Environ Res Public Health.* 2013; **10**(12): 7235–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hansell CA, Schiering C, Kinstrie R, *et al.*: **Universal expression and dual function of the atypical chemokine receptor D6 on innate-like B cells in mice.** *Blood.* 2011; **117**(20): 5413–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Soriano FG, Barbeiro HV, Barbeiro DF: **Inflammatory response: role of B1 cells.** *Shock.* 2013; **39** Suppl 1: 5–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hayakawa K, Hardy RR: **Normal, autoimmune, and malignant CD5+ B cells: the Ly-1 B lineage?** *Annu Rev Immunol.* 1988; **6**: 197–218.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Förster I, Gu H, Rajewsky K: **Germline antibody V regions as determinants of clonal persistence and malignant growth in the B cell compartment.** *EMBO J.* 1988; **7**(12): 3693–703.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kocks C, Rajewsky K: **Stable expression and somatic hypermutation of antibody V regions in B-cell developmental pathways.** *Annu Rev Immunol.* 1989; **7**: 537–59.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Baumgarth N, Tung JW, Herzenberg LA: **Inherent specificities in natural antibodies: a key to immune defense against pathogen invasion.** *Springer Semin Immunopathol.* 2005; **26**(4): 347–62.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Martin RK, Damle SR, Valentine YA, *et al.*: **B1 Cell IgE Impedes Mast Cell-Mediated Enhancement of Parasite Expulsion through B2 IgE Blockade.** *Cell Rep.* 2018; **22**(7): 1824–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Griffin DO, Holodick NE, Rothstein TL: **Human B1 cells in umbilical cord and adult peripheral blood express the novel phenotype CD20+ CD27+ CD43+ CD70+.** *J Exp Med.* 2011; **208**(1): 67–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Carter MC, Ruiz-Esteves KN, Workman L, *et al.*: **Identification of alpha-gal sensitivity in patients with a diagnosis of idiopathic anaphylaxis.** *Allergy.* 2018; **73**(5): 1131–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Steinke JW, Platts-Mills TA, Commins SP: **The alpha-gal story: lessons learned from connecting the dots.** *J Allergy Clin Immunol.* 2015; **135**(3): 589–96; quiz 597.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Ristivojević MK, Grundström J, Tran TAT, *et al.*: **α-Gal on the protein surface affects uptake and degradation in immature monocyte derived dendritic cells.** *Sci Rep.* 2018; **8**(1): 12684.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Commins SP, Platts-Mills TA: **Delayed anaphylaxis to red meat in patients with IgE specific for galactose alpha-1,3-galactose (alpha-gal).** *Curr Allergy Asthma Rep.* 2013; **13**(1): 72–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Suurmond J, Diamond B: **Autoantibodies in systemic autoimmune diseases: specificity and pathogenicity.** *J Clin Invest.* 2015; **125**(6): 2194–202.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Altrichter S, Kriehuber E, Moser J, *et al.*: **Serum IgE autoantibodies target keratinocytes in patients with atopic dermatitis.** *J Invest Dermatol.* 2008; **128**(9): 2232–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kessel A, Helou W, Bamberger E, *et al.*: **Elevated serum total IgE—a potential marker for severe chronic urticaria.** *Int Arch Allergy Immunol.* 2010; **153**(3): 288–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Permin H, Wiik A: **The prevalence of IgE antinuclear antibodies in rheumatoid arthritis and systemic lupus erythematosus.** *Acta Pathol Microbiol Scand C.* 1978; **86C**(5): 245–9.
[PubMed Abstract](#) | [Publisher Full Text](#)

26. Arbesman CE, Wypych JI, Reisman RE, *et al.*: IgE levels in sera of patients with pemphigus or bullous pemphigoid. *Arch Dermatol.* 1974; 110(3): 378–81. [PubMed Abstract](#) | [Publisher Full Text](#)
27. Liu FT, Goodarzi H, Chen HY: IgE, mast cells, and eosinophils in atopic dermatitis. *Clin Rev Allergy Immunol.* 2011; 41(3): 298–310. [PubMed Abstract](#) | [Publisher Full Text](#)
28. **F** Maurer M, Altrichter S, Schmetzer O, *et al.*: Immunoglobulin E-Mediated Autoimmunity. *Front Immunol.* 2018; 9: 689. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
29. **F** Saniklidou AH, Tighe PJ, Fairclough LC, *et al.*: IgE autoantibodies and their association with the disease activity and phenotype in bullous pemphigoid: a systematic review. *Arch Dermatol Res.* 2018; 310(1): 11–28. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
30. **F** Sanjuan MA, Sagar D, Kolbeck R: Role of IgE in autoimmunity. *J Allergy Clin Immunol.* 2016; 137(6): 1651–61. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
31. **F** Charles N, Hardwick D, Daugas E, *et al.*: Basophils and the T helper 2 environment can promote the development of lupus nephritis. *Nat Med.* 2010; 16(6): 701–7. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
32. Rullo OJ, Tsao BP: Recent insights into the genetic basis of systemic lupus erythematosus. *Ann Rheum Dis.* 2013; 72 Suppl 2: ii56–61. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Sinico RA, Rimoldi L, Radice A, *et al.*: Anti-C1q autoantibodies in lupus nephritis. *Ann N Y Acad Sci.* 2009; 1173(1): 47–51. [PubMed Abstract](#) | [Publisher Full Text](#)
34. **F** Henault J, Riggs JM, Karnell JL, *et al.*: Self-reactive IgE exacerbates interferon responses associated with autoimmunity. *Nat Immunol.* 2016; 17(2): 196–203. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
35. **F** Dema B, Charles N, Pellefigues C, *et al.*: Immunoglobulin E plays an immunoregulatory role in lupus. *J Exp Med.* 2014; 211(11): 2159–68. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
36. Dema B, Pellefigues C, Hasni S, *et al.*: Autoreactive IgE is prevalent in systemic lupus erythematosus and is associated with increased disease activity and nephritis. *PLoS One.* 2014; 9(2): e90424. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Chistiakov DA, Orekhov AN, Sobenin IA, *et al.*: Plasmacytoid dendritic cells: development, functions, and role in atherosclerotic inflammation. *Front Physiol.* 2014; 5: 279. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Conrad DH, Ford JW, Sturgill JL, *et al.*: CD23: an overlooked regulator of allergic disease. *Curr Allergy Asthma Rep.* 2007; 7(5): 331–7. [PubMed Abstract](#) | [Publisher Full Text](#)
39. **F** Cooper AM, Hobson PS, Jutton MR, *et al.*: Soluble CD23 controls IgE synthesis and homeostasis in human B cells. *J Immunol.* 2012; 188(7): 3199–207. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
40. Kraft S, Kinet JP: New developments in FcεpsilonRI regulation, function and inhibition. *Nat Rev Immunol.* 2007; 7(5): 365–78. [PubMed Abstract](#) | [Publisher Full Text](#)
41. Platzter B, Ruiter F, van der Mee J, *et al.*: Soluble IgE receptors—elements of the IgE network. *Immunol Lett.* 2011; 141(1): 36–44. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Ra C, Kuromitsu S, Hirose T, *et al.*: Soluble human high-affinity receptor for IgE abrogates the IgE-mediated allergic reaction. *Int Immunol.* 1993; 5(1): 47–54. [PubMed Abstract](#) | [Publisher Full Text](#)
43. Dehlink E, Platzter B, Baker AH, *et al.*: A soluble form of the high affinity IgE receptor, Fc-epsilon-RI, circulates in human serum. *PLoS One.* 2011; 6(4): e19098. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
44. **F** Moñino-Romero S, Erkert L, Schmidthaler K, *et al.*: The soluble isoform of human FcεRI is an endogenous inhibitor of IgE-mediated mast cell responses. *Allergy.* 2019; 74(2): 236–45. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
45. van Hemelrijck M, Garmö H, Binda E, *et al.*: Immunoglobulin E and cancer: a meta-analysis and a large Swedish cohort study. *Cancer Causes Control.* 2010; 21(10): 1657–67. [PubMed Abstract](#) | [Publisher Full Text](#)
46. Cahoon EK, Inskip PD, Gridley G, *et al.*: Immune-related conditions and subsequent risk of brain cancer in a cohort of 4.5 million male US veterans. *Br J Cancer.* 2014; 110(7): 1825–33. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
47. **F** Wulaningsih W, Holmberg L, Garmö H, *et al.*: Investigating the association between allergen-specific immunoglobulin E, cancer risk and survival. *Oncoimmunology.* 2016; 5(6): e1154250. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
48. **F** Fazekas-Singer J, Singer J, Ilieva KM, *et al.*: AllergoOncology: Generating a canine anticancer IgE against the epidermal growth factor receptor. *J Allergy Clin Immunol.* 2018; 142(3): 973–976.e11. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
49. Jensen-Jarolim E, Achatz G, Turner MC, *et al.*: AllergoOncology: the role of IgE-mediated allergy in cancer. *Allergy.* 2008; 63(10): 1255–66. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. **F** Jensen-Jarolim E, Bax HJ, Bianchini R, *et al.*: AllergoOncology - the impact of allergy in oncology: EAACI position paper. *Allergy.* 2017; 72(6): 866–87. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
51. Fu SL, Pierre J, Smith-Norowitz TA, *et al.*: Immunoglobulin E antibodies from pancreatic cancer patients mediate antibody-dependent cell-mediated cytotoxicity against pancreatic cancer cells. *Clin Exp Immunol.* 2008; 153(3): 401–9. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
52. Karagiannis SN, Bracher MG, Hunt J, *et al.*: IgE-antibody-dependent immunotherapy of solid tumors: cytotoxic and phagocytic mechanisms of eradication of ovarian cancer cells. *J Immunol.* 2007; 179(5): 2832–43. [PubMed Abstract](#) | [Publisher Full Text](#)
53. **F** Liao HC, Wu SY, Ou CY, *et al.*: Allergy symptoms, serum total immunoglobulin E, and risk of head and neck cancer. *Cancer Causes Control.* 2016; 27(9): 1105–15. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
54. **F** Nigro EA, Brini AT, Yenagi VA, *et al.*: Cutting Edge: IgE Plays an Active Role in Tumor Immunosurveillance in Mice. *J Immunol.* 2016; 197(7): 2583–8. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
55. Savina A, Amigorena S: Phagocytosis and antigen presentation in dendritic cells. *Immunol Rev.* 2007; 219(1): 143–56. [PubMed Abstract](#) | [Publisher Full Text](#)
56. Platzter B, Elpek KG, Cremasco V, *et al.*: IgE/FcεRI-Mediated Antigen Cross-Presentation by Dendritic Cells Enhances Anti-Tumor Immune Responses. *Cell Rep.* 2015; 10(9): 1487–95. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
57. **F** Crawford G, Hayes MD, Seoane RC, *et al.*: Epithelial damage and tissue γδ T cells promote a unique tumor-protective IgE response. *Nat Immunol.* 2018; 19(8): 859–70. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
58. Karagiannis SN, Wang Q, East N, *et al.*: Activity of human monocytes in IgE antibody-dependent surveillance and killing of ovarian tumor cells. *Eur J Immunol.* 2003; 33(4): 1030–40. [PubMed Abstract](#) | [Publisher Full Text](#)
59. Spicer J: Phase I Study of MOV18 IgE, a First in Class Chimeric IgE Antibody in Patients With Advanced Solid Tumours (NCT02546921). 2015. [Reference Source](#)
60. **F** Crescioli S, Chiaruttini G, Mele S, *et al.*: Engineering and stable production of recombinant IgE for cancer immunotherapy and AllergoOncology. *J Allergy Clin Immunol.* 2018; 141(4): 1519–1523.e9. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Edda Fiebiger

Department of Medicine, Division of Gastroenterology, Hepatology and Nutrition, Boston Children's Hospital, Boston,, Massachusetts, USA

Competing Interests: No competing interests were disclosed.

2. Yui-Hsi Wang

Type 2 Inflammation and Fibrosis Cluster Immunology & Inflammation Research Therapeutic Area, Sanofi, Cambridge, MA, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research