Check for updates

RESEARCH ARTICLE

Screening tuberculosis patients for diabetes mellitus in a routine program setting in Kampala, Uganda: a cross-sectional study [version 1; peer review: 2 approved with reservations]

Joseph Nsonga¹, John Paul Dongo¹, Frank Mugabe², Gerald Mutungi³, Richard Walyomo⁴, Christopher Oundo⁴, Sarah Zalwango⁴, Daniel Okello⁴, Simon Muchuro^{2,5}, Riitta A Dlodlo⁶, Yan Lin ⁶

¹International Union Against Tuberculosis and Lung Disease, Plot 2, Lourdel Road, Nakasero Hill, Kampala, Uganda

²The National Tuberculosis and Leprosy Program, Ministry of Health, Plot 6, Lourdel Road, Nakasero, Kampala, Uganda

³The Non-Communicable Diseases Program, Ministry of Health, Plot 6, Lourdel Road, Nakasero, Kampala, Uganda

⁴Kampala Capital City Authority, City Hall, Plot 1-3, Apollo Kaggwa Road, Kampala, Uganda

⁵University Research Company on USAID Defeat Project, Plot 40 Ntinda II Road, Naguru, Kampala, Uganda

⁶International Union Against Tuberculosis and Lung Disease, 68 Boulevard Saint-Michel, Paris, 75006, France

V1 First published: 17 Jun 2019, 8:872
<https://doi.org/10.12688/f1000research.19279.1>
 Latest published: 10 Oct 2019, 8:872
<https://doi.org/10.12688/f1000research.19279.2>

Abstract

Background: Uganda is located in East Africa and is among the countries with the lowest income globally. The ten health centres in this project serve populations in the under-privileged communities of Kampala. The objective of the study was to implement diabetes mellitus (DM) screening among tuberculosis (TB) patients in a routine program setting with limited resources and high human immunodeficiency virus (HIV) prevalence.

Methods: A descriptive cross-sectional observational study was conducted in ten health centres in Kampala, Uganda. As part of a project to implement DM screening in a routine setting, TB patients were screened for DM by trained health workers. A fasting blood glucose (FBG) value ≥ 7.0 mmol/l was considered to indicate DM. For this study, aggregate data was collected and analysed using SPSS for Windows, version 13.0.

Results: Among 4,590 TB patients registered, 4,016 (88.0%) were screened with random blood glucose (RBG). Of those with RBG ≥ 6.1 mmol/l, 1,093 (83.3%) were screened with FBG. In total, 92 (2.3%) patients were diagnosed with DM and 66 (71.8%) of them were newly diagnosed. The proportion of TB patients screened with FBG in the health centres varied from 58.2% to 100%. The proportion of patients screened with FBG and the prevalence of DM were significantly higher in private health centres compared with public health centres. The

Open Peer Review

Approval Status

	1	2
version 2
(revision)	view	view
10 Oct 2019
version 1
17 Jun 2019	view	view

1. **Razia Kaniz Fatima** , National TB Control Program, Islamabad, Pakistan
2. **Daniel C. Oshi**, The University of West Indies, Mona, Kingston, Jamaica

Any reports and responses or comments on the article can be found at the end of the article.

health centres in peri-urban areas screened more patients with RBG than those in urban areas. Health centres without DM services screened a larger number of patients with RBG and FBG than those with DM services.

Conclusions: It appears feasible to implement screening TB patients for DM in routine program settings with limited resources and high HIV prevalence. Its introduction requires close collaboration between TB and DM services. The challenges identified need government attention and certain institutional and service-related factors need to be better managed at times.

Keywords

Tuberculosis, Diabetes Mellitus, Screening, Uganda

Corresponding author: Yan Lin (ylin@theunion.org)

Author roles: **Nsonga J:** Data Curation, Project Administration, Supervision; **Dongo JP:** Funding Acquisition, Supervision, Validation; **Mugabe F:** Supervision, Validation; **Mutungi G:** Methodology, Resources; **Walyomo R:** Investigation, Validation; **Oundo C:** Investigation, Validation; **Zalwango S:** Project Administration, Resources; **Okello D:** Funding Acquisition, Supervision; **Muchuro S:** Investigation, Methodology; **Dlodlo RA:** Conceptualization, Writing – Review & Editing; **Lin Y:** Conceptualization, Formal Analysis, Methodology, Writing – Original Draft Preparation

Competing interests: No competing interests were disclosed.

Grant information: This project was funded by the World Diabetes Foundation [WDF15-1211].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 Nsonga J *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Nsonga J, Dongo JP, Mugabe F *et al.* **Screening tuberculosis patients for diabetes mellitus in a routine program setting in Kampala, Uganda: a cross-sectional study [version 1; peer review: 2 approved with reservations]** F1000Research 2019, 8:872 <https://doi.org/10.12688/f1000research.19279.1>

First published: 17 Jun 2019, 8:872 <https://doi.org/10.12688/f1000research.19279.1>

Introduction

Although significant progress has been achieved in tuberculosis (TB) care and prevention during the past decades, TB remains a major public health problem and is responsible for more deaths than any other single infectious disease worldwide¹. In 2017 globally, 10.0 million people developed TB and 1.6 million died from it¹. Uganda is among the 30 high TB and human immunodeficiency virus (HIV) burden countries, with an estimated TB prevalence of 253 per 100,000 population (95% CI: 196-317) and 40% HIV co-infection rate^{2,3}.

Along with socio-economic development, urbanization, dietary and lifestyle changes, the prevalence of diabetes mellitus (DM) is escalating in most low- and middle-income countries. Available data suggests that an estimated 425 million people worldwide live with DM and by 2045 this number will grow to 629 million⁴. Although estimates for sub-Saharan Africa are based on limited data with substantial uncertainty, DM appears to be increasing rapidly in many urban centres in the region⁵. In Uganda, some risk factors for DM, such as obesity, were greater among persons of a high socio-economic status in rural areas, though the prevalence of DM in the adult population of the entire country was relatively low (1.4% in a nationwide cross-sectional survey in 2014)^{6,7}.

DM is a well-known risk factor for TB and increases its risk 2-3-fold^{8,9}. Systematic reviews and meta-analyses suggest that DM patients with TB experience worse treatment outcomes compared with patients without DM, with delayed sputum smear conversion, increased risk of drug resistance, treatment failure, death and relapse after successful completion of anti-TB treatment⁸⁻¹². On the other hand, TB patients with previously known DM are at an increased risk of hyperglycemia or worse glycaemic control^{13,14}. Therefore, early detection of DM among TB patients and proper management of both conditions are well known measures to improve treatment outcomes in these patients.

To address this problem in low- and middle-income countries, the World Health Organization (WHO) and the International Union Against Tuberculosis and Lung Disease (The Union) launched a 'Collaborative Framework for the Care and Control of Diabetes and Tuberculosis' in 2011, and recommended bi-directional screening for TB and DM within routine health services¹⁵.

The Union, together with national authorities, implemented the first bi-directional screening of DM and TB within routine health services in China and India, where it was found to be feasible in these settings¹⁶⁻¹⁸. In Uganda, there is no policy on routine screening for DM among TB patients. Whether a similar approach can be implemented in a resource limited setting with a high HIV prevalence remained unknown. To further understand this, a project was implemented from 2016 to 2017 to screen TB patients for DM, supported by the World Diabetes Foundation (WDF). Objectives of this study were to: a) assess whether DM screening can be implemented among TB patients in a routine programmatic health setting in Uganda; b) identify

challenges for DM screening among TB patients and solutions to overcome them; and c) identify institutional and health service related factors that may have an impact on DM screening among TB patients.

Methods

DM screening project

Setting and sites. Kampala is the capital city of Uganda and had an estimated population of 1,607,182 in 2017. In the same year, the Kampala Capital City Authority (KCCA) notified a total of 7,260 TB cases, resulting in a case notification rate of 452/100,000 population³. Treatment success was 84.6%. HIV prevalence was 6.9% nationwide (Uganda Population-based HIV Impact Assessment UPHIA 2016–2017)³.

Health services in the city are provided by the KCCA. Persons with presumed TB are investigated in health centres and diagnosed TB patients receive care in TB clinics. DM services are available mainly in specialised DM clinics at tertiary level health facilities.

Ten health centres, both public and private, in Kampala were selected based on their location in the five administrative divisions of KCCA and representative of the health delivery system in Kampala. These were selected from different communities based on sufficient numbers of registered TB patients, the availability of TB and DM diagnostic and treatment facilities within the same catchment area and willingness of the staff to participate in the project.

Project training, screening algorithm and monitoring. The National TB and Non-Communicable Disease (NCD) Programs' staff, Kampala Capital City Authority (KCCA) and The Union Uganda Office staff attended an inception meeting and training of trainers in February-March 2016. Participants reviewed the association between DM and TB, the need for screening and agreed to use the screening algorithm shown in [Figure 1](#). Cascade training sessions for 40 staff from the selected health centres were then conducted. The Union and KCCA officials performed quarterly monitoring, compiled periodic reports and provided overall supportive supervision for the activities. At the same time, they recorded any challenge encountered during the implementation.

Patients. Patients aged 15 years and over who were seeking health services in the health centres and were newly registered with any form of TB between 1st April 2016 to 31st December 2017 were included in the screening project.

Blood glucose measurement in TB patients. A blood sample was collected for random blood glucose (RBG) using capillary blood immediately after the TB registration in the health centres. A OneTouch glucometer (model SN SAFTM7K8, Lifescan, Johnson & Johnson, New Brunswick, USA) was used with a detection value ranging from 1.7 to 19.1mmol/l. Patients with an RBG value ≥ 6.1 mmol/l were invited to return on the following day, after an overnight fast of at least 10 hours, for measurement of fasting blood glucose (FBG). The FBG cut-off thresholds were

Figure 1. Algorithm for diabetes mellitus (DM) screening among tuberculosis patients in Kampala, Uganda.

based on the World Health Organization (WHO) recommendations as follows: i) FBG ≥ 7.0 mmol/l (≥ 126 mg/dl) indicates presence of DM; ii) FBG of 6.1 – 6.9 mmol/l (from 110 mg/dl to less than 126 mg/dl) indicates impaired glucose tolerance; and iii) FBG < 6.1 mmol/l (< 110 mg/dl) indicates normal blood glucose levels. Health workers provided information on the importance of fasting before returning for a confirmatory test (FBG) to the patients through health education. Reminder phone calls were also provided to all patients who screened positive for RBG to return for confirmatory test.

Study design

This was a descriptive cross-sectional observational study using routine project data from ten health centres in Kampala, Uganda.

Data collection and analysis

District TB and Leprosy Supervisors (DTLs) collected routine programmatic data every six months from the unit TB and DM registers in the 10 implementing health facilities. Final data collection was finished in December 2017. The Union staff reviewed the data and worked together with the health centre staff to trace the missing data (mainly the number of patients with known DM). The DTLs collected aggregated data, which was verified by The Union staff, and double-entered the data into a spreadsheet. Comparisons of characteristics between patients attending different health centres were carried out using the chi square test with odds ratios (OR) and their 95% confidence intervals. Levels of significance were set at 5%.

Statistical analysis was carried out using SPSS for Windows version 13.0 (SPSS Inc., Chicago, IL, USA).

Ethical statement

This project and the research proposal were approved by the National Tuberculosis and Leprosy Program, the Non-Communicable Diseases Program, Ministry of Health of the Government of Uganda, and Kampala Capital City Authority. As the information collected formed part of routine programmatic health service delivery and no individual patient data was collected, review by an ethics committee and patient consent were not considered necessary.

Results

The characteristics of the selected health centres are described in Table 1. Of the 4,590 TB patients notified, 4,016 (88.0%) were screened with RBG²⁵. Of those with RBG ≥ 6.1 mmol/l, 1,093 (83.3%) were screened with FBG. Of the 92 TB patients diagnosed with DM, 66 (71.7%) were newly diagnosed with DM and all, except four patients who died shortly after the TB registration, were referred to DM services. The prevalence of DM in the screened TB patients was 2.3% (Table 2).

The number of registered TB patients in reporting periods one, two, three and four were 372, 1,576, 1,364 and 1,278, respectively. The proportion of patients screened with RBG (number of screened with RBG/number of registered TB patients) were 83.1%, 83.8%, 91.6% and 89.0%, respectively, over the four reporting periods, but the absolute number of TB patients

Table 1. Characteristics of the 10 study health centres, Kampala, Uganda, from 2016–2017.

Characteristics	Kisenyi	Kawaala	Komamboga	Kisugu	Naguru	Kiswa	Murchison Bay	Kitebi	Mengo	Nsambya
Location*	Urban	Urban	Rural	Urban	Urban	Urban	Peri-urban	Peri-urban	Urban	Urban
Type	Public	Public	Public	Public	Public	Public	Public	Public	Private	Private
No. of health workers/volunteers at TB clinic	2/5	2/3	2/2	2/2	2/2	2/2	2/2	2/2	2/2	2/2
No. of health centre beds	32	16	16	16	250	16	100	16	250	250
Average no. of daily patient visits at TB clinic	150	40	20	30	50	40	20	20	40	30
DM services before the study	None	None	None	None	Yes	None	None	None	Yes	Yes

*Urban = central business area; Peri-urban = slightly away from central business area; Rural = furthest away from central business area. TB, tuberculosis; DM, diabetes mellitus

Table 2. Diabetes mellitus screening among TB patients in Kampala, Uganda, from 2016–2017 (10 centres combined).

Registered TB patients screened for and diagnosed with DM	Number
Number of registered TB patients	4,590
Number with known DM	26
Number to be screened with RBG	4,564
Number (%) screened with RBG	4,016 (88.0%)
Number with RBG ≥ 6.1 mmol/l and to be screened with FBG	1,312
Number (%) screened with FBG	1,093 (83.3%)
Number with FBG ≥ 7.0 mmol/l	66
Number (%) of newly diagnosed with DM*	66 (71.7%)
Number (%) with known and newly diagnosed DM	92 (2.3%)

* Percentage of newly diagnosed DM among total number of DM

TB, tuberculosis; DM, diabetes mellitus; RBG, random blood glucose; FBG, fasting blood glucose

screened with RBG in the first reporting period was 7.7%. Of the number of TB patients with RBG ≥ 6.1 mmol/l, the proportion of screened with FBG (number of screened with FBG/number of RBG ≥ 6.1 mmol/l) were 81.4%, 89.8%, 95.7% and 63.9% over the four reporting periods.

The proportion of patients screened with RBG and FBG varied from 78.9% to 94.5%, and from 58.2% to 100.0%, respectively, among the health centres (Table 3). The proportion of patients screened with RBG was significantly higher in the health centres located in peri-urban areas compared with the centres in urban areas (OR 1.50, 95% CI 1.12-1.99, $P=0.006$); and in those without DM services compared with those with DM services (OR 1.35, 95% CI 1.11-1.63, $P=0.002$) (Table 4). Of the patients with RBG ≥ 6.1 mmol/l, the proportion screened with FBG

was significantly higher in private health centres (OR 2.16, 95% CI 1.26-3.69, $P=0.005$) and those without DM services (OR 1.96, 95% CI 1.44-2.67, $P<0.001$) than that in public health centres and in those with DM services.

A total of 92 TB patients were diagnosed with DM, accounting for a prevalence of 2.3%. The prevalence of DM in TB patients was higher in the private health centres (OR 2.26, 95% CI 1.40-3.62, $P=0.001$) and in those with DM services (OR 0.60, 95% CI 0.39-0.93, $P=0.022$).

Discussion

This is the first study to report on screening TB patients for DM in a primary health care setting in Uganda. We included both public and private facilities in urban, peri-urban and rural parts

Table 3. Screening TB patients for DM at the 10 health centres: data combined for the four reporting periods.

	Kisenyi	Kawaala	Komamboga	Kisugu	Naguru	Kiswa	Murchison Bay	Kitebi	Mengo	Nsambya
No. of TB patients registered over the four reporting periods	1,183	593	329	339	555	299	319	324	313	336
No. with known diagnosis of DM	9	3	1	2	3	0	1	0	3	4
No. needing to be screened with RBG	1,174	590	328	337	552	299	318	324	310	332
No. (%) screened with RBG	1,107 (94.3)	466 (79.0)	280 (85.4)	298 (88.4)	468 (84.8)	257 (86.0)	293 (92.1)	292 (90.1)	293 (94.5)	262 (78.9)
No. with RBG ≥ 6.1 mmol/l	346	211	99	94	153	72	76	87	77	98
No. (%) screened with FBG	281 (81.2)	170 (80.6)	91 (91.9)	94 (100.0)	89 (58.2)	70 (97.2)	66 (86.8)	73 (83.9)	72 (93.5)	87 (88.8)
No. with FBG ≥ 7.0 mmol/l referred to DM services	19	5	9	1	6	3	3	3	6	11
No. newly diagnosed with DM*	19	5	9	1	6	3	3	3	6	11
No. (%) with known or newly diagnosed DM	28 (2.5)	8 (1.7)	10 (3.6)	3 (1.0)	9 (1.9)	3 (1.2)	4 (1.4)	3 (1.0)	9 (3.1)	15 (5.7)

* Percentage of newly diagnosed DM among total number of DM

TB = tuberculosis; DM = diabetes mellitus; RBG = random blood glucose; FBG = fasting blood glucose

Table 4. Screening TB patients for DM at the 10 health centres: data combined for the four reporting periods stratified by type, location and available of DM service.

Characteristics of the health centres	No. of TB patients registered	No. (%) screened with RBG	No. with RBG $\geq$ 6.1 mmol/l	No. (%) screened with FBG	No. (%) with known or newly diagnosed DM	% of newly diagnosed DM*
Type						
Public (Reference)	3,941	3,641 (87.8)	1,137	934 (82.2)	68 (2.0)	72.1
Private	649	555 (85.5)	175	159 (90.9)	24 (4.3)	70.8
OR (95% CI)		0.82 (0.65-1.04)		2.16 (1.26-3.69)	2.26 (1.40-3.62)	0.94 (0.34-2.63)
P-value		0.101		0.005	0.001	0.909
Location						
Urban (Reference)	3,618	3,151 (87.1)	1,050	863 (82.2)	75 (2.4)	68.0
Peri-urban	643	585 (91.0)	163	139 (85.5)	7 (1.2)	85.7
OR (95% CI)		1.50 (1.12-1.99)		1.26 (0.79-1.99)	0.50 (0.23-1.08)	2.82 (0.32-24.77)
P-value		0.006		0.334	0.079	0.349
Rural	329	280 (85.4)	99	91 (91.9)	10 (3.6)	90.0
OR (95% CI)		0.85 (0.62-1.17)		2.47 (1.18-5.17)	1.52 (0.78-2.97)	4.24 (0.51-35.36)
P-value		0.307		0.017	0.222	0.182
Available DM service						
Yes (Reference)	1,204	1,023 (85.0)	328	248 (75.6)	33 (3.2)	69.7
None	3,386	2,993 (88.4)	984	845 (85.9)	59 (2.0)	72.9
OR (95% CI)		1.35 (1.11-1.63)		1.96 (1.44-2.67)	0.60 (0.39-0.93)	1.17 (0.46-2.99)
P-value		0.002		<0.001	0.022	0.745

* Percentage of newly diagnosed DM among total number of DM

TB = tuberculosis; DM = diabetes mellitus; RBG = random blood glucose; FBG = fasting blood glucose

of Kampala to assess the feasibility of integrating DM screening into routine TB service.

The overall prevalence of DM in the screened TB patients was 2.3% (92/4,016), which was higher than that in general population and our finding mirrored the findings of previous studies^{6,17,18,20}, although it was lower than that reported by a study among hospitalized TB patients at a large referral hospital in an urban area of Uganda²¹. However, using only FBG to diagnose DM can underestimate the prevalence of DM by as much as 50% when compared with the gold standard test, the oral glucose tolerance test, which is more accurate but unavailable at primary health care level²². In addition, some of the facilities raised the FBG screening threshold to 7.0 mmol/l during the first 6–8 months due to misinterpretation of the screening algorithm. We therefore believe that the actual prevalence of DM among TB patients in Uganda might be higher than we have observed in this study. Of the 92 DM patients detected, 66 (71.7%) had not been diagnosed before this study. This strongly

demonstrated the need for integrating DM screening into routine TB service^{5,23}.

In the first six months, a small number of TB patients were screened with RBG, due to the long engagement of stakeholders followed by the cascade training. One health centre (Nsambya) did not send staff to attend the training and did not screen any patients in the first reporting period due to staff turn-over, a situation that should have been identified by training organizers, and this challenge required KCCA attention. The number of screened TB patients increased rapidly over the rest of the reporting periods, suggesting that the procedure was gradually accepted by staff, and we echo a similar finding in China¹⁷. The proportion of patients screened with FBG declined over the last six-month period due to gluco-strips stock-out at the government medical stores. It was reassuring that the health centre staff largely followed the screening algorithm, although there were certain differences. Kisugu Health Centre ensured FBG screening for everyone with RBG $\geq$ 6.1mmol/l due to strong com-

mitment of the staff. On the contrary, Naguru Health Centre managed to screen only 58.2% of the patients who needed an FBG test even though they had the same number of health workers, although they did have a slightly higher number of TB patient visits. It is necessary to understand reasons for missed opportunities in the DM screening cascade.

Private health centres screened higher proportions of patients with FBG and, subsequently, identified more DM patients than public facilities. Possible reasons for this could be that patients with a higher socio-economic status sought care in private health centres, which have well-motivated staff thanks to better remuneration. Similarly, staff in peri-urban and rural facilities screened higher proportions of patients with RBG and FBG than in urban areas. Murchison Bay is a health centre in peri-urban area operated by the Ministry of Internal Affairs with prisoners being the main source of patients. Given that most patients were confined in prison, they were available for RBG and FBG. In urban health centres there is more frequently congestion and longer waiting times than in rural centres. This may have caused some patients to miss their FBG tests.

An intriguing finding that also requires a further study was that health centres with available DM services at time of the study, a feature that tends to reflect better health services, did not screen more patients with RBG and FBG compared with those without DM services. This was unexpected and not in line with the findings of other studies¹⁶. Low screening at Naguru Health Centre could have contributed to this finding.

There were some service-related barriers identified in the project implementation: 1) lack of a sustainable supply of glucose testing strips, which caused insufficient screening for RBG and FBG; 2) lack of community support mechanisms to help trace patients who failed to return for FBG testing; 3) insufficient number of staff in the health centres caused by staff turn-over; 4) routine DM service was not available in some health centres.

This study had several limitations. Diagnosis of DM relied on FBG alone and we were unable to include oral glucose-tolerance tests, a more accurate test, as part of the screening cascade. This may have resulted in an underestimated prevalence of DM in TB patients²¹. Blood samples for RBG and FBG were taken from capillary blood and measured with glucometers without quality assurance, rather than using venous blood specimens and tested with biochemical analyser. This may have caused a testing bias²⁴. RBG or FBG were measured only once immediately after TB registration, without any confirmatory test at the end of anti-TB treatment, and glycosylated haemoglobin (HbA1c) tests were not carried out, which is influenced less by infection²⁵. Therefore, we do not know whether any TB patient may have been diagnosed as having DM due to infection-induced hyperglycaemia. However, a previous study suggested that taking one blood sample for FBG immediately after TB diagnosis would be appropriate for majority of TB patients¹⁴. Finally, we did not collect individual patient data and perform multivariate analysis and were unable to identify and control potential confounding factors.

The strength of this study is that we implemented DM screening among a large number of TB patients in routine health service using mainly existing resources. Only training, meetings, support supervision, mentorship and provision of glucometers were supported. There were no staff or patient incentives. This is a key point for sustainability and scale up of integration of DM and TB services. Some challenges identified in the study need government attention and this indicates the need for collaboration between TB and non-communicable disease programs.

Conclusion

This study revealed that prevalence of DM in TB patients is higher than that in the general population in Uganda. Screening TB patients for DM in a primary health care setting in a resource-limited country with high HIV-associated TB epidemic appeared feasible. This study identified certain service-related factors that may be barriers and need to be addressed by relevant authorities. Inclusion of bi-directional screening in both TB and DM clinics could be considered. Every year, 44,000-47,000 persons with TB are registered in Uganda and if all were screened for DM, at least 1,748 additional persons with DM could be detected and enrolled to DM care. As early diagnosis and management for the two diseases will improve treatment outcomes and well-being of people, this is an important public health intervention.

Data availability

Underlying data

Figshare: Screening tuberculosis patients for diabetes mellitus in a routine program setting in Kampala, Uganda: a cross-sectional study. <https://doi.org/10.6084/m9.figshare.8248625.v1>¹⁹

This project contains the following underlying data:

- TB_DM DATA _Uganda 2016_2017.xlsx (raw data collected from health centres over the four reporting periods)
- Screening tuberculosis patients for diabetes mellitus in a routine program setting in Kampala, Uganda a cross-sectional study.docx (table of data processes)

Data are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

Grant information

This project was funded by the World Diabetes Foundation [WDF15-1211].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We thank all the staff and patients at the 10 health centres for their participation and collaboration. This study was funded by the World Diabetes Foundation as part of a grant (WDF 15-1211). Thanks are also extended to the Ministry of Health, National TB and Leprosy Program, National Non-Communicable Disease Program, Kampala Capital City Authority, Uganda Diabetes Association and Non-Communicable Disease Alliance.

References

1. World Health Organization. WHO Report: **Global Tuberculosis Control 2018**. 2018. [Reference Source](#)
2. Ministry of Health: **The Uganda National Tuberculosis Prevalence Survey, 2014-2015**. Kampala, Uganda. 2017. [Reference Source](#)
3. Ministry of Health: **National Tuberculosis and Leprosy Program Annual Report 2016-2017**. Kampala, Uganda. 2017. [Reference Source](#)
4. International Diabetes Federation: **IDF Diabetes atlas**. 8th Edition. International Diabetes Federation, Brussels. 2017. [Reference Source](#)
5. Atun R, Davies J, Gale EAM, *et al.*: **Diabetes in sub-Saharan Africa: from clinical care to health policy**. *Lancet Diabetes Endocrinol*. 2017; 5(8): 622–67. [PubMed Abstract](#) | [Publisher Full Text](#)
6. Bahendeka S, Wesonga R, Mutungi G, *et al.*: **Prevalence and correlates of diabetes mellitus in Uganda: a population-based national survey**. *Trop Med Int Health*. 2016; 21(3): 405–416. [PubMed Abstract](#) | [Publisher Full Text](#)
7. Murphy GA, Asiki G, Ekoru K, *et al.*: **Sociodemographic distribution of non-communicable disease risk factors in rural Uganda: a cross-sectional study**. *Int J Epidemiol*. 2013; 42(6): 1740–53. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. Stevenson CR, Forouhi NG, Roglic G, *et al.*: **Diabetes and tuberculosis: the impact of the diabetes epidemic on tuberculosis incidence**. *BMC Public Health*. 2007; 7: 234. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. Jeon CY, Murray MB: **Diabetes mellitus increases the risk of active tuberculosis: a systematic review of 13 observational studies**. *PLoS Med*. 2008; 5(7): e152. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Baker MA, Harries AD, Jeon CY, *et al.*: **The impact of diabetes on tuberculosis treatment outcomes: a systematic review**. *BMC Med*. 2011; 9: 81. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Wang Q, Ma A, Han X, *et al.*: **Hyperglycemia is associated with increased risk of patient delay in pulmonary tuberculosis in rural areas**. *J Diabetes*. 2017; 9(7): 648–655. [PubMed Abstract](#) | [Publisher Full Text](#)
12. Liu Q, Li W, Xue M, *et al.*: **Diabetes mellitus and the risk of multidrug resistant tuberculosis: a meta-analysis**. *Sci Rep*. 2017; 7(1): 1090. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Tabarsi P, Baghaei P, Marjani M, *et al.*: **Changes in glycosylated haemoglobin and treatment outcomes in patients with tuberculosis in Iran: a cohort study**. *J Diabetes Metab Disord*. 2014; 13(1): 123. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
14. Lin Y, Yuan Y, Zhao X, *et al.*: **The change in blood glucose levels in tuberculosis patients before and during anti-tuberculosis treatment in China**. *Glob Health Action*. 2017; 10(1): 1289737. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. The International Union Against Tuberculosis and Lung Disease, World Health Organization: **Collaborative Framework for care and control of Tuberculosis and Diabetes**. Geneva. 2011; WHO/HTM/TB/2011.15. [Reference Source](#)
16. Lin Y, Li L, Mi FL, *et al.*: **Screening patients with diabetes mellitus for tuberculosis in China**. *Trop Med Int Health*. 2012; 17(10): 1302–1308. [PubMed Abstract](#) | [Publisher Full Text](#)
17. Li L, Lin Y, Mi F, *et al.*: **Screening of patients with tuberculosis for diabetes mellitus in China**. *Trop Med Int Health*. 2012; 17(10): 1294–1301. [PubMed Abstract](#) | [Publisher Full Text](#)
18. India Tuberculosis-Diabetes Study Group: **Screening of patients with tuberculosis for diabetes mellitus in India**. *Trop Med Int Health*. 2013; 18(5): 636–45. [PubMed Abstract](#) | [Publisher Full Text](#)
19. Nsonga J: **Screening tuberculosis patients for diabetes mellitus in a routine program setting in Kampala, Uganda: a cross-sectional study**. 2019. <http://www.doi.org/10.6084/m9.figshare.8248625.v1>
20. Lawson L, Muc M, Oladimeji O, *et al.*: **Tuberculosis and diabetes in Nigerian patients with and without HIV**. *Int J Infect Dis*. 2017; 61: 121–125. [PubMed Abstract](#) | [Publisher Full Text](#)
21. Davis K, Richard S, Edrisa M, *et al.*: **Overt diabetes mellitus among newly diagnosed Ugandan tuberculosis patients: a cross sectional study**. *BMC Infect Dis*. 2013; 13: 122. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Mugusi F, Swai AB, Alberti KG, *et al.*: **Increased prevalence of diabetes mellitus in patients with pulmonary tuberculosis in Tanzania**. *Tubercle*. 1990; 71(4): 271–276. [PubMed Abstract](#) | [Publisher Full Text](#)
23. Lonnoth K, Roglic G, Harries AD: **Improving tuberculosis prevention and care through addressing the global diabetes epidemic: from evidence to policy and practice**. *Lancet Diabetes Endocrinol*. 2014; 2(9): 730–39. [PubMed Abstract](#) | [Publisher Full Text](#)
24. World Health Organization, International Diabetes Federation: **Definition and Diagnosis of Diabetes Mellitus and Intermediate Hyperglycaemia**. Geneva, Switzerland: WHO, 2006. [Reference Source](#)
25. World Health Organization: **Use of glycated haemoglobin (HbA1c) in the diagnosis of diabetes mellitus: abbreviated report of a WHO consultation**. Geneva, Switzerland: WHO; 2011; Accessed September 14 2011. [Reference Source](#)

Open Peer Review

Current Peer Review Status: ? ?

Version 1

Reviewer Report 17 September 2019

<https://doi.org/10.5256/f1000research.21133.r53770>

© 2019 Oshi D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Daniel C. Oshi

Department of Community Health and Psychiatry, The University of West Indies, Mona, Kingston, Jamaica

General Comment:

The manuscript is on a very important public health issue, which is at the intersection between communicable and non-communicable diseases control. It is especially relevant because of the contextual issues of high TB and HIV burden in Uganda, the increasing burden of DM as a result of changing lifestyles and nutritional patterns, and limited resources in the country.

Abstract: The abstract is well written and captures the essence of the paper.

Introduction: The Introduction section is well written, supported with key epidemiological facts and figures on TB and DM, globally, in the Sub-Saharan African region and in Uganda. The authors concisely made a strong case for the rationale of the study, from the epidemiological and clinical perspectives and aptly noted the absence of policy on routine screening for TB and DM in health care settings as well as the paucity of evidence to guide the formulation of such a policy. Having created gaps in knowledge and policy that their paper will contribute toward addressing, the authors stated the objectives of their study.

Methods: This section is, on the whole, well written, although it would have been useful to see a justification for the choice of only Kampala Capital City, especially because the research aimed to produce evidence for policy formulation for programmatic screening for TB and DM in the whole country, not just in Kampala Capital City.

Results: This section is also well written. However, there are two issues of concern here. First, since this study was not just an academic exercise but a public health project, the authors should state measures they took, if any, to ensure that patients newly diagnosed with DM and referred to DM clinics actually reached and accessed care at DM clinics. Initial default is a major challenge in such referrals because some of the referred patients fail to go for the needed care. Second, it would have been useful to show the socio-demographic and clinical characteristics of all the patients versus those diagnosed with DM. This would have made this paper (the analysis, results and

discussions) much stronger.

Discussion: This is well written and acknowledged the limitation pointed out above about the non-collection and analysis of patient-level data.

Conclusion: One key conclusion of the study is "Screening TB patients for DM in a primary health care setting in a resource-limited country with high HIV-associated TB epidemic appeared feasible." This conclusion is apparently not supported by the geographical coverage of the study. The study was conducted only in Kampala City Capital. The inclusion of just one rural health centre, which is still part of the KCC is insufficient to be used to draw conclusions regarding rural setting. The geographic spread and the socio-economic characteristics of the locations of almost all the health centres in the study therefore suggest that the conclusion should be drawn to Kampala City Council rather than to the entire country.

Finally, the authors are commended for the effort in implementing this important research project.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Chronic communicable diseases control, social determinants of disease, substance abuse, lifestyles and behavioural change.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 29 Sep 2019

Yan Lin, International Union Against Tuberculosis and Lung Disease, 68 Boulevard Saint-Michel, Paris, France

Reviewer 2:

General Comment:

The manuscript is on a very important public health issue, which is at the intersection between communicable and non-communicable diseases control. It is especially relevant because of the contextual issues of high TB and HIV burden in Uganda, the increasing burden of DM as a result of changing lifestyles and nutritional patterns, and limited resources in the country.

Response: Thank you.

Abstract: The abstract is well written and captures the essence of the paper.

Response: Thank you.

Introduction: The Introduction section is well written, supported with key epidemiological facts and figures on TB and DM, globally, in the Sub-Saharan African region and in Uganda. The authors concisely made a strong case for the rationale of the study, from the epidemiological and clinical perspectives and aptly noted the absence of policy on routine screening for TB and DM in health care settings as well as the paucity of evidence to guide the formulation of such a policy. Having created gaps in knowledge and policy that their paper will contribute toward addressing, the authors stated the objectives of their study.

Response: Many thanks.

Methods: This section is, on the whole, well written, although it would have been useful to see a justification for the choice of only Kampala Capital City, especially because the research aimed to produce evidence for policy formulation for programmatic screening for TB and DM in the whole country, not just in Kampala Capital City.

Response: Point well taken! We have restructured some relevant parts in the method as highlighted in yellow colour to give justification on selection of the health centers. In addition, we have added a sentence in the limitation as "Finally, the selected HCs were only located in Kampala due to financial limitations and transportation barriers. This might have limited the representative value."

Results: This section is also well written. However, there are two issues of concern here. First, since this study was not just an academic exercise but a public health project, the authors should state measures they took, if any, to ensure that patients newly diagnosed with DM and referred to DM clinics actually reached and accessed care at DM clinics. Initial default is a major challenge in such referrals because some of the referred patients fail to go for the needed care.

Response: This is excellent advice, thank you. We have rephrased the last sentences in the first paragraph of the Results as "Ninety-two TB patients were diagnosed with DM, and of them, 66 (71.7%) were newly diagnosed DM and were referred to DM services. All DM patients arrived at the DM clinics and received DM care except four patients who died shortly after TB registration."

Second, it would have been useful to show the socio-demographic and clinical characteristics of all the patients versus those diagnosed with DM. This would have made this paper (the analysis, results and discussions) much stronger.

Response: Thank you for the valuable advice. We agree. To well accept this point, we have rephrased the sentence in the limitation as “We did not collect individual patient data, including socio-demographic data, and perform multivariate analysis and were unable to identify and control potential confounding factors.”

Discussion: This is well written and acknowledged the limitation pointed out above about the non-collection and analysis of patient-level data.

Response: Thanks.

Conclusion: One key conclusion of the study is “Screening TB patients for DM in a primary health care setting in a resource-limited country with high HIV-associated TB epidemic appeared feasible.” This conclusion is apparently not supported by the geographical coverage of the study. The study was conducted only in Kampala City Capital. The inclusion of just one rural health centre, which is still part of the KCC is insufficient to be used to drawn conclusions regarding rural setting. The geographic spread and the socio-economic characteristics of the locations of almost all the health centres in the study therefore suggest that the conclusion should be drawn to Kampala City Council rather than to the entire country.

Finally, the authors are commended for the effort in implementing this important research project.

Response: Thank you for pointing this out. We have revised the sentence in the conclusion as “Screening TB patients for DM in a primary health care setting in Kampala with high HIV-associated TB epidemic appeared feasible.”

Competing Interests: No

Reviewer Report 19 June 2019

<https://doi.org/10.5256/f1000research.21133.r49993>

© 2019 Fatima R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Razia Kaniz Fatima

National TB Control Program, Islamabad, Pakistan

The article is well written and addresses the Global issue of DM screening among TB Patients which is one the common co morbidity recognized with TB that may affect the TB outcome. The points which need to be explained are:

1. The selection criteria of the centres and how public vs private centres were selected moreover was the urban rural representation was also considered or it was random? Moreover it was stated in abstract that health centres without DM screened larger number of patients while in methods it was stated the DM services were mainly in tertiary care hospitals which might create assumptions of large screening in tertiary care hospitals contrary to what was stated in abstract. Table 1 also shows the DM services to be already present in 3 out of 7 centres so this need to be better explained. The average number of daily patients screened was also less in these 3 centres so it might mean that tertiary care where already DM services are present were not part of study?
2. In discussion it was mentioned using only FBG to diagnose DM can underestimate the prevalence of DM by as much as 50% when compared with the gold standard test, the oral glucose tolerance test, which is more accurate but unavailable at primary health care level, so how this should spell out in the policy implication? When the program should plan to do this DM screening of TB patients, more focus should be on the availability of gold standard for FBG using Glucometer, this should also be explained.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Tuberculosis Research & Epidemiology.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 29 Sep 2019

Yan Lin, International Union Against Tuberculosis and Lung Disease, 68 Boulevard Saint-

Michel, Paris, France

Reviewer 1

The article is well written and addresses the Global issue of DM screening among TB Patients which is one the common co morbidity recognized with TB that may affect the TB outcome.

The points which need to be explained are:

- The selection criteria of the centres and how public vs private centres were selected moreover was the urban rural representation was also considered or it was random?

Response: Thank you for this valuable advice. As described in the site and setting, selection of the health centers was based on their location in the five administrative divisions of KCCA and representative of the health delivery system in Kampala as well as willingness of the staff to participate this study. It was however not random selected. We have rephrased the sentence as "These were high volume health facilities selected from different communities based on their ability to manage sufficient numbers of TB patients registered in the previous year and a daily TB patient visits ≥ 20 /day, availability of diagnostic and treatment facilities and willingness of the staff to participate in this study."

- Moreover it was stated in abstract that health centres without DM screened larger number of patients while in methods it was stated the DM services were mainly in tertiary care hospitals which might create assumptions of large screening in tertiary care hospitals contrary to what was stated in abstract. Table 1 also shows the DM services to be already present in 3 out of 7 centres so this need to be better explained. The average number of daily patients screened was also less in these 3 centres so it might mean that tertiary care where already DM services are present were not part of study?

Response: Thank you for the question. The ten health centers recruited in this study are primary health care institutions. There was no any tertiary hospital involved in this study. Of the ten health centers, the health center without DM service screened large number of patients (both RBG and FBG) than those with DM service.

- In discussion it was mentioned using only FBG to diagnose DM can underestimate the prevalence of DM by as much as 50% when compared with the gold standard test, the oral glucose tolerance test, which is more accurate but unavailable at primary health care level, so how this should spell out in the policy implication? When the program should plan to do this DM screening of TB patients, more focus should be on the availability of gold standard for FBG using Glucometer, this should also be explained.

Response: Excellent advice! Thank you. We have added a sentences in the Conclusions as "RBG and FBG are practical tools to be used for screening in routine program settings even though they may underestimate the actual situation".

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research