RESEARCH ARTICLE

Variations in sexual network connectivity may explain dramatic variations in sexually transmitted infection prevalence between populations and over time: a four-country analysis [version 1; peer review: 2 approved with reservations]

Chris Kenyon ^{1,2}

¹HIV/STI Unit, Institute of Tropical Medicine, Antwerp, Antwerp, 2000, Belgium

²Division of Infectious Diseases and HIV Medicine, University of Cape Town, Cape Town, 7700, South Africa

V1 First published: 20 Aug 2020, 9:1009
<https://doi.org/10.12688/f1000research.24968.1>
 Latest published: 30 Aug 2022, 9:1009
<https://doi.org/10.12688/f1000research.24968.2>

Abstract

Background: The incidence of sexually transmitted infections (STIs) has been noted to vary dramatically between population groups and over time. Here, the hypothesis that changes in network connectivity underpin these changes is explored.

Methods: The incidence/prevalence estimates of HIV, herpes simplex virus-2, syphilis, chlamydia, and gonorrhoea, as well as two markers of sexual network connectivity (partner concurrency and multiple partnering) by ethnic group and sexual orientation in Kenya, South Africa, the United Kingdom (UK) and the United States (USA) were extracted from published studies. Pearson's correlation was used to test the association between the markers of network connectivity and the incidence/prevalence of these five STIs. A literature review was performed to evaluate the possible causes of the increases and decreases in syphilis incidence over the past 60 years.

Results: In each country, the five STIs were found to cluster in particular ethnic groups and sexual orientations and to be positively associated with the two markers of network connectivity. Syphilis incidence in the UK and USA was found to increase dramatically in the 1960s/1970s, decline in the 1980s and again increase in the late 1990s. These changes took place predominantly in men who have sex with men, and were preceded by corresponding changes in network connectivity. The large decline in antenatal syphilis prevalence in Kenya and South Africa in the 1990s were likewise preceded by declines in network connectivity.

Conclusions: Although other explanatory variables are not controlled for, the present analysis is compatible with the hypothesis that

Open Peer Review

Approval Status

	1	2
version 2		
(revision)	view	view
30 Aug 2022		
version 1		
20 Aug 2020	view	view

1. **Christine Khosropour** , University of Washington, Seattle, USA

2. **Itzhak Levy**, Tel-Aviv University, Tel-Aviv, Israel
 Sheba Medical Centre, Tel-Aviv, Israel

Any reports and responses or comments on the article can be found at the end of the article.

differential network connectivity is a parsimonious explanation for variations in STI incidence over time and between populations.

Keywords

network connectivity, STI, syphilis, gonorrhoea, herpes simplex virus-2, HIV, chlamydia, epidemiology

This article is included in the **Sociology of Health** gateway.

Corresponding author: Chris Kenyon (chriskenyon0@gmail.com)

Author roles: Kenyon C: Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Kenyon C. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Kenyon C. **Variations in sexual network connectivity may explain dramatic variations in sexually transmitted infection prevalence between populations and over time: a four-country analysis [version 1; peer review: 2 approved with reservations]** F1000Research 2020, 9:1009 <https://doi.org/10.12688/f1000research.24968.1>

First published: 20 Aug 2020, 9:1009 <https://doi.org/10.12688/f1000research.24968.1>

Introduction

The reasons underpinning the dramatic variations in sexually transmitted infection (STI) incidence between different populations are incompletely understood^{1,2}. Not infrequently the incidence of specific STIs varies by an order of magnitude or more between ethnic groups and between groups defined according to sexual orientation^{3–6}. Furthermore, over the past century, there have also been numerous examples of dramatic changes in the incidence of specific STIs. The incidence of syphilis in men who have sex with men (MSM) in the United States (USA), for example, was high in the 1970s, but declined dramatically in the early 1980s only to return to 1970s levels more recently (Figure 1)⁷. In this paper we test the hypothesis that differences in the connectivity of local sexual networks play an important role explaining these cross sectional and longitudinal variations in STI incidence^{8,9}. Although the connectivity of local sexual networks is determined by a number of parameters, we focus on two of the most important - the rate of partner change and the proportion of partnerships that run concurrently^{8,9}. Previous analyses have typically found that one, or a combination of markers of network connectivity, are raised in populations with a higher prevalence of a particular STI^{3,4,9,10}. These studies have however typically been limited to single countries and frequently only investigated variations in a single STI and either considered variations between ethnic groups or sexual orientations but not both^{11–20}. Global country-level analyses have produced a mix bag of results with a small number of studies^{21,22} finding positive associations between markers of network connectivity and STI prevalence, and a larger number finding no association^{23–26}. A possible explanation for this discrepancy is a type of misclassification bias that could be termed a ‘fruit salad bias’^{22,27}. It is increasingly appreciated that each country is made up of a number of sexual networks (defined by sexual orientation, ethnicity, class, religion etc.) that interconnect to

varying degrees^{2,4,9,28}. If the connectivity of these sexual networks is a major determinant of STI prevalence, then it would be most appropriate to compare the association between network connectivity and STI prevalence between sub-populations of sexual networks within a country rather than combining them all into one national population (fruit salad) as a number of studies have done^{10,24,25}.

These insights provided the motivation for the current paper where we aimed to conduct a more systematic analysis of the association between network connectivity and STI prevalence in national subpopulations. We expanded the previously conducted analyses of this association in four ways. Firstly, we assessed this association in four countries: two high income countries (the United Kingdom [UK] and [USA]) where STIs are predominantly concentrated in MSM and certain ethnic groups and two low/middle income countries that have been affected by generalized HIV epidemics (Kenya and South Africa) but where certain ethnic groups have largely escaped this epidemic. Secondly, in each country we assess this association at the levels of both ethnic groups and MSM versus heterosexuals. Thirdly, we assess this association in five STIs (chlamydia, gonorrhoea, HIV, herpes simplex virus-2 [HSV-2] and syphilis). Fourthly, we conduct longitudinal analyses over the past 60 years to assess if increases and decreases in syphilis incidence and prevalence are preceded by corresponding changes in network connectivity.

Methods

Data sources

A search was conducted on PubMed and Google Scholar on 15 February 2020 using the following terms: ‘chlamydia’, ‘gonorrhoea’, ‘HIV’, ‘HSV-2’, ‘syphilis’, ‘concurrency’, ‘multiple partnering’, ‘Kenya’, ‘South Africa’, ‘UK’, ‘United Kingdom’, ‘USA’, ‘United States’, ‘men who have sex with men’, ‘MSM’.

Figure 1. Incidence of primary/secondary syphilis in the United States (MSM and heterosexual men; cases per 100 000/population) between 1963–2013 and cases of primary, secondary and early latent syphilis diagnosed in men and women in England and Wales between 1950 and 2015. Beginning of ‘AIDS’ epidemic and introduction of ‘ART’ – antiretroviral therapy – indicated with arrows.

Cross sectional STI data. The incidence and prevalence estimates of chlamydia, gonorrhoea, HIV, HSV-2, syphilis, concurrency and multiple partnering in Kenya, South Africa, the UK, and USA were extracted from published studies, as detailed in Table 1 and Table 2. When two or more studies were found, we favoured data from studies based on nationally representative samples. In addition, for the cross-sectional analyses, we limited

studies to those based on data from after the year 1990 and favoured the most recent data.

Symptomatic STI: In the case of Kenya and South Africa, we could not find high quality data on STI prevalence, or behaviour reported by sexual orientation. In addition, we could not find reasonable quality data reporting the incidence/prevalence of

Table 1. Description of sources of data for comparison between ethnic groups: prevalence of STIs, multiple partners, concurrency.

Country	Year data collected, reference	Study type, selection procedure, sample size
Kenya		
HIV, HSV-2 & syphilis	2007 ²⁹	The 2007 Kenya AIDS Indicator Survey used a stratified two stage sampling strategy to test a nationally and provincially representative sample of 15 853 15 to 64 year olds.
Male urethral discharge & circumcision	2008 ³⁰	The 2008 Kenya Demographic and Health Survey used a household-based, two-stage stratified sampling approach to recruit 12 677 participants.
Concurrency & multiple partners	2011 ^{31,32}	The Population Services International (PSI) Survey /Kenya 6th HIV Survey conducted in 2011 used a two stage cluster sampling to obtain a provincially representative sample of households from seven of Kenya's eight provinces (the North East was excluded). A total of 3 051 men and women 15-49 year old were included.
South Africa		
HIV	2005 ³³	A two-stage, nationally representative sample of 23 275 persons 2 years old or older. We limited our analysis to the 13 884 individuals aged 15 to 49 years old.
HSV-2	2012 ³⁴	The prevalence of HSV-2 was assessed in the 2012 national antenatal HIV and HSV-2 survey. The prevalence of HSV-2 was estimated for 18 732 women in four provinces - Gauteng, KwaZulu-Natal, Northern Cape and Western Cape. The Focus HerpesSelect 2 @ ELISA IgG Diagnostic kit was used as the diagnostic test.
Syphilis	1991 ³⁵	A sample of 17 318, 15-49 year old women attending antenatal clinics for their first visit were tested for syphilis via the RPR or VDRL test.
Male urethral discharge	1998 ³⁶	The 1998 Demographic and Health Survey employed a 2-stage sampling strategy in South Africa's nine provinces and stratified results into urban and non-urban groups. It was designed to be representative for all provinces and the four major ethnic/racial groups. 6578 men were asked if they had experienced symptoms of a urethral discharge in the last 3 months. Men were not asked questions about their sexual behaviour in this survey.
Concurrency, Multiple partners & circumcision	2003 ^{4,37}	The 2003 Demographic Health Survey (DHS) used a similar study design to the 1998 DHS. The survey sampled 7966 women and 3930 men. All were 15-49 year old.
United Kingdom		
HIV & Male urethral discharge	2010-2012 ¹⁹	National Surveys of Sexual Attitudes and Lifestyles 3 recruited a probability sample of 15 162 women and men aged 16-74 years in Britain. Participants were interviewed with computer-assisted face-to-face and self-completion questionnaires. Urine from a sample of participants aged 16-44 years who reported at least one sexual partner over the lifetime was tested for HIV antibodies.
Chlamydia, gonorrhoea & syphilis	2018 ³⁸	The incidence figures for these three STIs by ethnic group were extracted from the Public Health England report for 2018: Sexually transmitted infections and screening for chlamydia in England. The figures were reported separately for men and women but not combined. We report the results for men. The rank order of each STI incidence by ethnic group did not differ by sex ³⁸ . The results are reported as infections per 100 000 population per year.

Country	Year data collected, reference	Study type, selection procedure, sample size
Concurrency, multiple partners, STI symptoms	2000 ^{39,40}	The second British National Survey of Sexual Attitudes and Lifestyles (NATSAL 2) was a nationally representative sample of 11161 men and women aged 16-44 years ³⁹ . We extracted the relevant variables from a study which broke down the various sexual behaviour variables by ethnic group and sex ³⁹ . Men were asked if they had ever been diagnosed with an STI.
USA		
HIV	2006 ⁴¹	HIV prevalence at the end of 2006 was estimated using information from the national HIV/AIDS Reporting System.
HSV-2	2005-2008 ⁴²	HSV-2 seroprevalence was assessed in a nationally representative sample of 7 293 persons 14-49 years in the NHANES 2005-2008 sample.
Syphilis	2001-2004 ⁴³	Sera from 5767, 18- to 49-year-old participants in the NHANES 2001–2004 were tested for syphilis IgG antibody using an enzyme immunoassay (EIA). Specimens with positive or indeterminate EIAs underwent rapid plasma reagin (RPR) testing.
Gonorrhoea & Chlamydia & Syphilis	2017 ⁴⁴	The incidence estimates for gonorrhoea, chlamydia and syphilis are based on the CDCs 2017 STD Surveillance Report. They are reported as infections per 100 000 population per year.
Concurrency & Multiple partners	1992 ²⁸	The prevalence of concurrency and multiple partners were taken from the 1992 National Health and Social Life Survey (NHSLs). This was a cross-sectional study that used a nationally representative stratified random sample of 3 432 women and men between the ages of 18 and 59.

Abbreviations: STI – sexually transmitted disease, DHS - Demographic and Health Survey, NATSAL - National Survey of Sexual Attitudes and Lifestyles, NHANES - National Health and Nutrition Examination Surveys.

chlamydia or gonorrhoea by ethnic group in these two countries. As a result, we use the proportion of men reporting urethral discharge as a composite proxy for combined chlamydia plus gonorrhoea incidence per ethnic group^{30,36}.

Longitudinal syphilis data. We chose syphilis for the evaluation of longitudinal changes in STI prevalence for a number of reasons, including the fact that syphilis surveillance programmes from the four countries have provided reasonable quality estimates of syphilis incidence for the last 60 years and longer – a far longer period than that for which reasonable incidence estimates are available for other STIs⁴⁵. To a large extent, this is related to the fact that relatively accurate serological tests have been available for the diagnosis of syphilis for longer than other STIs and these have been extensively used to diagnose antenatal syphilis^{45–47}. Antenatal syphilis surveys have been shown to produce reasonable estimates of the prevalence of syphilis in the general population^{45–47}. For Kenya and South Africa, we used large antenatal syphilis prevalence surveys - national in the case of South Africa and limited to the capital in the case of Kenya (Table 1). For the UK and USA, we used incidence estimates based on national case reporting data (Table 1). In the UK, the sexual orientation of individuals with syphilis was not available for much of this period and thus we follow the example of others who compare the ratio of the incidence in men to that in women as a proxy for the incidence in MSM^{1,48,49}.

Multiple partners and concurrency. Studies use a wide variety of definitions of multiple partnering and concurrency.

Concurrency: In all the comparisons of ethnic groups, excluding the UK, the prevalence of concurrency was defined as the percentage of men (15–49 years old) who reported having two or more sex partners at the time of the survey. In the UK, concurrency referred to the percentage of men who reported any overlapping sexual relationships in the past year³⁹. In the comparisons of sexual orientations, for the USA, concurrency was defined as the respondent reporting two or more sexual partnerships with overlapping dates in the prior year⁵⁰. As far as the UK was concerned, this referred to the proportion reporting overlapping sexual partnerships at any point in the past 5 years⁵¹.

Multiple partners: For the ethnic group comparisons, the prevalence of multiple partners was defined as the percentage of men (15–49 years old) who reported having two or more sex partners in the last year. In the UK, this variable was not reported by ethnic group. As a result, multiple partnering here refers to the percentage of men who reported one or more new heterosexual relationship in the prior year. For the MSM vs. heterosexual comparisons, this variable referred to the mean lifetime number of sexual partners reported by respondents in the UK (aged 16–74) and the USA (aged 35–39).

These variations in definitions between studies mean it would be appropriate to compare these variables between studies. Since all the studies we used were limited to specific countries, we only compare subpopulations within each study/country.

Regions vs. ethnic group in Kenya. Many of the national STI surveillance surveys in Kenya do not collect data on

Table 2. Description of sources of data for comparison between men who have sex with men and men who have sex with women: prevalence of STIs, multiple partners and concurrency.

Country	Year data collected, reference	Data source. Study type, selection procedure, sample size
USA		
HIV	2009-12 ⁵²	HIV prevalence estimates for MSM and hetero sexual men were taken from an analysis of the National Health and Nutrition Examination Survey (NHANES) 2009–2012 survey. Men who reported ever having had sex with another man were classified as men who have sex with men (MSM). Men who reported no sex with other men were classified as heterosexual men.
Chlamydia, gonorrhoea and syphilis	2013 ^{7,44}	CDC STI surveillance reports note that with the exception of reported syphilis cases, nationally notifiable STI surveillance data do not routinely include information on sexual partners/orientation. As a result this data is missing for the majority of gonorrhoea and chlamydia cases reported to CDC ⁴⁴ . No estimates are therefore provided for chlamydia or gonorrhoea incidence. The primary and secondary syphilis incidence estimates for 2013 (reported as cases per 100 000 population per year) are taken from a paper that estimated incidence in MSM and heterosexual men based on cases of primary and secondary syphilis reported to the CDC between 1963 and 2013 ⁷ .
Concurrency, multiple partners	1996–2006 ⁵⁰	Sexual behaviour data was taken from a paper that compared sexual behaviour patterns between MSM and male and female heterosexuals aged 18–39 in the USA. Their data was obtained from 4 population-based random digit dialling surveys. A 1996–1998 survey in 4 U.S. cities and 2 Seattle surveys (2003, 2006) provided estimates for MSM; a 2003–2004 Seattle survey provided data about heterosexual men and women. The multiple partner variable refers to the mean lifetime number of partners reported by respondents (aged 35-39 years). Concurrency was defined as the respondent reporting two or more sexual partnerships with overlapping dates in the prior year ⁵⁰ .
United Kingdom		
HIV	2016 ⁵³	HIV prevalence estimates were taken from the Public Health England's report 'HIV in the UK, 2016' ⁵³ . These figures are based on a Bayesian multi-parameter evidence synthesis model, which combines data from multiple sources to provide prevalence estimates for HIV for different sub populations ⁵³ .
Chlamydia, gonorrhoea and syphilis	2018 ³⁸	The incidence figures for these three STIs were taken from Public Health England's report 'Sexually transmitted infections and screening for chlamydia in England, 2018' ³⁸ . This report provides incidence estimates for chlamydia, gonorrhoea and syphilis for heterosexual men, heterosexual women, HIV-infected MSM and 'HIV-uninfected or HIV-undiagnosed MSM' in 2018. We report the results for the first and last two of these categories. There was little difference in the incidence estimates for heterosexual men and women ³⁸ .
Concurrency, multiple partners	2012 ⁵¹	The third British National Survey of Sexual Attitudes and Lifestyles (NATSAL 3) was a nationally representative sample of 15,162 men and women aged 16-74 years conducted in 2012 ⁵¹ . We extracted the relevant variables from a study report which provided prevalence estimates of sexual behaviour for MSM and heterosexual men. The multiple partnering variable referred to the mean lifetime number of sexual partners reported by respondents. The proportion reporting overlapping sexual partnerships at any point in the past 5 years was used as the measure of concurrency.

ethnic group²⁹. There is however a high correlation between region of residence and ethnic group, with large differences in STI incidence and sexual behaviour between regions^{5,31}. We therefore follow the approach used in previous studies and use region of residence as a proxy for ethnic group^{31,54}.

Statistical analysis

We report the incidence/prevalence of the five STIs, multiple partnering and concurrency by ethnic group (all four countries) and MSM vs. heterosexuals in each country with available data (UK and USA). Where this data is provided, we report 95% confidence intervals. We used the Chi-squared test to assess if there was a difference in incidence/prevalence

between the subgroups with the highest and lowest incidence/prevalence values in each country. For continuous variables (lifetime number of partners) the Man-Whitney U test was used. The correlation between the prevalence of the different STIs per subgroup in each country, and between the prevalence of each STI and the two behavioural variables was calculated using Pearson's correlation. Because of the small sample sizes (between 2 and 8), tests of statistical significance were not performed for these calculations. In the case of UK and USA, the STI data is presented as an estimated incidence per 100 000 per year without reporting the number of cases and the total size of the study populations. As a result, we did not test if the association between the prevalence of each STI and

each of the markers of network connectivity is statistically significant. The key differences were however large, which allows substantive inferences to be drawn. Where data was only provided in graphical form, we digitized the data using GetData Graph Digitizer version 2.26. STATA v15.2 was used for all analyses (StataCorp LLC, College Station, TX, USA).

Longitudinal association between network connectivity and syphilis prevalence/incidence

A literature review was performed to evaluate the possible causes of increases and decreases in syphilis incidence in each subpopulation. We used a narrative approach here to evaluate if there was evidence from the literature that increases and decreases in syphilis prevalence were preceded by corresponding changes in markers of network connectivity. The approach adopted was not to perform a systematic review or a quantitative analysis, but rather to assess if there is evidence in the literature that corroborates or refutes the network connectivity hypothesis.

Results

Cross sectional analyses

Kenya

Ethnic group HIV prevalence varied 15-fold between 0.9% (95% confidence interval [CI] 0.3-4%) in the North East and 13.9% (95% CI 11.0-17.9%) in Nyanza (Figure 2; Table 1, Table 4 & Table 7). HSV-2 prevalence varied 5-fold between

15.6% and 76.2% in these same two regions. The prevalence of syphilis varied 5-fold between 0.5% (95% CI 0.2-1.4%) and 2.5% (95% CI 1.8-3.4%) in these same two regions. Likewise, the percentage of men reporting urethral discharge in the prior year varied between 0.0% (95% CI 0.0-0.5%) and 3.5% (95% CI 2.0-5.8%; all $P < 0.001$) in the same two regions. The prevalence/incidence of male urethral discharge, HIV, HSV-2 and syphilis by region were positively correlated with each other ($r = 0.35$ to $r = 0.77$; Figure 2, Table 6).

The region with the lowest HIV prevalence (the North East) was not included in the behavioural survey that provided our data. Other data sources have however found the rate of multiple partnering in this region to be very low⁵. In our data, the region with the second lowest HIV prevalence (Eastern) had the lowest reported prevalence of multiple partnering (12.2%; 95% CI 7.4-19.5%) and concurrency (6.4%; 95% CI 4.2-9.2%; all $P < 0.001$). Nyanza reported the highest prevalence of multiple partnering (30.6%; 95% CI 23.8-38.3%) and the second highest prevalence of concurrency (18.2%; 95% CI 14.9-21.9%).

With the exception of syphilis, which was negatively associated with concurrency prevalence ($r = -0.34$), the prevalence of each STI was positively correlated with both the multiple partnering ($r = 0.38$ to $r = 0.84$) and concurrency ($r = 0.42$ to $r = 0.72$) variables.

Figure 2. Incidence/prevalence of HIV, HSV-2, chlamydia, gonorrhoea, syphilis, male urethral discharge (MUD), partner concurrency and multiple partnering (MP) by ethnic group in Kenya, South Africa, the United Kingdom and the United States of America. For all countries, HIV, HSV-2, male urethral discharge, multiple partners and concurrency prevalence are reported as percentages. The same is true for syphilis in Kenya and South Africa. In the USA and the UK, chlamydia, gonorrhoea and syphilis are reported as cases per 100 000 per year (Point estimates with 95% Confidence Intervals; see Table 1 for sources of data).

South Africa

Ethnic group HIV prevalence varied 40-fold between 0.5% (95% CI 0.3-1.1%) in the White group and 19.9% (95% CI 18.3-21.7%) in the Black group (Figure 1; Table 4). HSV-2 prevalence varied 5-fold from 6.3% to 31.3% between the same groups. The prevalence of syphilis varied 21-fold between 0.4% (95% CI 0.1-0.9%) and 8.3% (95% CI 7.8-8.8%) in these same two groups. Likewise, the percentage of men reporting urethral discharge in the prior year varied 7-fold between 2.0% (95% CI 0.7-3.2%) and 13.2% (95% CI 12.2-14.2%) in the same groups (all $P < 0.001$). The prevalence/incidence of male urethral discharge, HIV, HSV-2 and syphilis by ethnic group were positively correlated with each other ($r = 0.78$ to $r = 0.98$; Figure 2, Table 6).

The Black group reported a 5-fold higher prevalence of concurrency and a 3-fold higher prevalence of multiple partners than the White group (9.2%, 95% CI 7.1-12.0% vs. 2.0%, 95% CI 0.3-12.5% and 16.0%, 95% CI 13.9-18.3% vs. 4.9%, 95% CI 2.1-11.2%, respectively; all $P < 0.001$). The prevalence of each STI was positively correlated with both the multiple partnering ($r = 0.87$ to $r = 1.00$) and concurrency ($r = 0.85$ to $r = 0.99$) variables (Table 6).

USA

Ethnic group HIV prevalence varied 8-fold between 0.2% (95% CI 0.2-0.2%) in the non-Hispanic White group and 1.7%

(95% CI 1.6-1.8%) in the non-Hispanic Black group (Figure 2, Table 4). The incidence of syphilis varied 4-fold between 5 and 24 per 100 000 per year in these same two groups. Likewise, the incidence of chlamydia varied 5-fold and that of gonorrhoea 8-fold between the same groups (all $P < 0.001$). The prevalence/incidence of chlamydia, gonorrhoea, HIV, HSV-2 and syphilis by ethnic group were strongly positively correlated with each other ($r = 0.97$ to $r = 1.00$; Table 6).

The non-Hispanic Black group reported 4-fold higher prevalence of concurrency and a 2-fold higher prevalence of multiple partners than the non-Hispanic Whites group (11.3%, 95% CI 7.5-16.7% vs. 3.1%, 95% CI 2.2-4.3% and 27.1%, 95% CI 23.8-31.4% vs. 15.0%, 95% CI 14.0-16.5%, respectively; all $P < 0.001$). The prevalence of each STI was positively correlated with both the multiple partnering ($r = 0.94$ to $r = 0.98$) and concurrency ($r = 0.79$ to $r = 0.91$) variables.

MSM vs heterosexuals The prevalence of HIV was 36-fold higher (7.2% vs. 0.2%) and the incidence of syphilis 33-fold higher (233 vs. 7/100 000 per year) in MSM compared to heterosexuals (Table 2 & Table 5, Figure 3). The percent reporting concurrency was 3 times higher in MSM than heterosexuals (31.3% vs. 9.7%). Likewise, the mean number of lifetime partners was 7-times higher (67 vs. 10) in MSM.

Figure 3. Incidence/prevalence of HIV, chlamydia, gonorrhoea, syphilis, partner concurrency and multiple partnering by men who have sex with men (MSM) versus heterosexual men in the United Kingdom and the United States of America. HIV and partner concurrency are reported as percentages, multiple partners as the number of lifetime partners and chlamydia gonorrhoea and syphilis as cases per 100 000 per year (Point estimates with 95% Confidence Intervals; see Table 1 for sources of data).

United Kingdom

Ethnic group HIV prevalence varied between 2.8% in the Black Caribbean and Black African groups and 0% in the White and Asian groups (Table 4, Figure 2). The incidence of chlamydia varied 11-fold between the Black Caribbean (1540/100 000/year) and Asian (136/100 000/year) groups. Gonorrhoea incidence varied 8-fold between these same groups (840 and 109/100 000/year, respectively). Likewise, the incidence of syphilis varied 4-fold between 71 and 18 per 100 000 per year in these same two groups. The prevalence/incidence of chlamydia, gonorrhoea, HIV and syphilis by ethnic group were positively correlated with each other ($r=0.71$ to $r=0.98$; Table 6).

The prevalence of concurrency and multiple partnering were approximately 1.5 to 2.5-fold higher in the Black African and Black Caribbean groups than the Asian and White groups. The prevalence of each STI was positively correlated with both the multiple partnering ($r=0.63$ to $r=0.96$) and concurrency ($r=0.37$ to $r=0.91$) variables.

MSM vs heterosexuals The prevalence of HIV was 39-fold higher in MSM outside of London (7.2%) than heterosexual men (0.16%; Table 5, Figure 3). Likewise, the incidence of chlamydia was 12-fold higher (24 vs. 2/100 000/year), gonorrhoea was 68-fold higher (41 vs. 0.6/100 000/year) and syphilis 185-fold higher (74 vs. 0.04/100 000/year) in MSM outside of London than MSW. The incidence of each of these STIs was higher in MSM residing in London. The percentage reporting concurrency was 14 times higher in MSM than heterosexuals (52% vs. 3.8%). Likewise, the mean number of lifetime partners was 8-times higher (111.1 vs. 14.3) in MSM.

Longitudinal analyses

UK and USA. In both the UK and the USA, the incidence of syphilis increased in the 1960s and 1970s but this increase was largely limited to MSM (Figure 1)^{1,7,49}. The incidence remained fairly stable in heterosexuals. Although quantitative data is very limited from this period, a number of analyses attributed this

increased incidence in MSM to behavioural factors, such as increases in rates of partner change^{7,49,55,56}.

In the early 1980s, incidence plummeted in MSM in both countries^{1,7}. The AIDS epidemic played a large role in this decline. Individuals most centrally placed in sexual networks were more likely to die from AIDs, which resulted in a fragmentation of sexual networks^{7,57}. So too reductions in multiple partnering and increases in condom use both served to reduce effective network connectivity^{49,58}. From around 2000, the incidence of syphilis has been increasing in both countries and this increase has been largely limited to MSM^{1,7}. A very similar trend has been evident for MSM in the UK for chlamydia and gonorrhoea where reasonable surveillance data is available and is likely the case in the USA where surveillance data is less detailed as regards sexual orientation^{1,44}. In both countries, numerous lines of evidence suggest that the introduction of effective antiretroviral therapy from 1996 onwards together with other factors resulted in increases in sexual network connectivity, which was in turn responsible for these increases in STI incidence^{1,48}. Reduced deaths from AIDS, increases in rates of partner change and reductions in condom usage all played a role in this reconstitution of network connectivity^{7,48,58,59–61}.

Kenya and South Africa. The prevalence of antenatal syphilis in both Kenya and South Africa was between 5 and 12% in the pre-HIV period (Figure 4; Table 3 and Table 4). A number of analyses attributed this high prevalence of syphilis to a number of factors that included behaviours that translate into dense sexual networks^{45,62,63}. Syphilis prevalence declined rapidly following the AIDS epidemic to below 1% in both countries⁶². Similar, contemporaneous declines in diagnoses of cases of syphilis took place in these countries^{45,64}. Using a range of data sources and study methodologies, a number of analyses have concluded that a combination of connectivity-reducing factors was responsible: reductions in multiple partnering (number of partners and concurrency), delayed sexual debut, increased condom usage and the effect of AIDS mortality removing more centrally placed transmission nodes^{45,62,65–67}.

Figure 4. Prevalence of antenatal syphilis (red circles) and HIV (black squares) in South Africa and Kenya the United States between 1990 and 2012.

Table 3. Description of sources of data for longitudinal changes in syphilis incidence and prevalence.

	Year data collected, reference	Data source. Study type, selection procedure
Kenya	1992-1997 ⁶⁸	81 311 pregnant women from 10 antenatal sentinel sites in Nairobi were screened for syphilis on their first antenatal visit using the RPR only. Results were used to generate annual syphilis prevalence estimates for each year 1992 to 1997. ⁶⁸
South Africa	1938 to 2011 ⁴⁵	Longitudinal syphilis prevalence estimates were taken from a global epidemiology of syphilis over the past century paper. Seventeen studies, all performed in antenatal populations, evaluated the prevalence of syphilis in pregnant women between 1938 and 2011 ⁴⁵
UK	1960 to 2015 ¹	The longitudinal syphilis incidence estimates are for the annual number of new cases of primary, secondary and early latent syphilis diagnosed in men and women in England and Wales between 1960-2015. The data is also based on case reporting surveillance data and is from the Lancet Infectious Diseases Commission: Sexually transmitted infections: challenges ahead ¹ .
USA	1963 to 2013 ⁷	The primary and secondary syphilis incidence estimates for 1963 to 2013 (reported as cases per 100 000 population) are taken from a paper that estimated incidence in MSM and heterosexual men based on cases of primary and secondary syphilis reported to the CDC in this time period ⁷ .

Discussion

Our analysis confirmed previous analyses of dramatic variations in STI prevalence between populations³⁻⁶. A strong tendency towards clustering of STIs within subpopulations was evident. Furthermore, the cross-sectional variations in STI prevalence were almost all positively associated with two markers of network connectivity. In a similar vein, dramatic increases and declines in syphilis incidence could also be explained by corresponding changes in network connectivity. Taken together these findings suggest that network connectivity could represent a parsimonious explanation for both variations in STI incidence between populations and over time.

There are however a number of caveats to bear in mind. Our study evaluated only 4 out of over 200 countries, and the data it used was limited in a number of ways. Not all the data was from nationally representative samples. The data also came from different and limited time periods. The way the behavioural variables were defined varied between studies and in the case of Kenya, region was used as a proxy for ethnic group. For the sake of simplicity, we focused on male sexual behaviours. On the one hand, our analysis is purely ecological and therefore susceptible to the ecological inference fallacy. On the other hand, the prevalence of STIs is to a large extent a function of network level properties. Networks are inherently ecological level entities and thus ecological analyses are appropriate and necessary forms of analysis. We assume that there is a degree of segregation of sexual network by ethnic group or sexual orientation. A high degree of sexual partner homophily by ethnic group has indeed been shown for Kenya, South Africa and the USA^{4,5,28,69,70}. We could find no analyses investigating this type of homophily in the UK. If there was little or none of this type of homophily, this would however result in a homogenization of STI incidence between ethnic groups and would therefore be expected to reduce the strength of the association between STIs and behaviours by ethnic group.

A further crucial weakness is that we did not control for potential confounders. The prevalence of circumcision and condom

use are two examples of factors that could confound our analyses. Other papers have, however, evaluated this question and concluded that these are likely to be only one of many determinants of differential STI prevalence^{3,4,31,65}. For example, in South Africa, the highest STI prevalence ethnic group has been noted to have considerably higher circumcision rates and use of condoms than the two lower STI prevalence populations^{4,65}. Likewise, rates of condom usage are typically higher in MSM than heterosexuals⁵⁰ and as far as we can ascertain, the prevalence of circumcision does not differ significantly between these two populations. A further limitation is that we have not controlled for other differences between MSM and heterosexual sexual networks such as the impact of role-versatility in MSM⁷¹.

These limitations mean that we cannot conclude how generalizable our findings are or even if the associations we found are causal. What we can conclude is that our results are compatible with the network connectivity explanation. We cannot exclude the possibility that other unmeasured variables are the predominant cause.

Our findings are however logical if we recall that the number of sexual contacts per unit time is a key determinant of the rate of spread of an STI². So too increases in concurrency have been shown to lead to non-linear increases in sexual network connectivity^{9,72}. A number of mathematical modelling studies from the USA and South Africa have established that the actual self-reported differences in number of partners and concurrency between ethnic groups in the USA and South Africa were able to explain a large part of the differential spread of HIV in these populations^{3,65}.

A strength of our analysis is its finding that the association between network connectivity and STI prevalence held in four different countries for both comparisons by ethnic group and MSM versus heterosexuals. In the case of Kenya, all the ethnic groups/regions, including those with very low STI rates, were 'black' Africans. The association was found for five different

Table 4. Prevalence/incidence of HIV, HSV-2, chlamydia, gonorrhoea, syphilis, concurrency and multiple partnering by ethnic group in Kenya, South Africa, the United Kingdom and the United States.

Country	Ethnic group	HIV (%)	HSV-2 (%)	Chlamydia [§]	Gonorrhoea [§]	Syphilis ⁺	STI symptoms [#]	Concurrency (%) [†]	Multiple partners (%) [‡]
SA	White	0.5 (0.3-1.1)	6.3	NA	NA	0.4 (0.1-0.9)	2.0 (0.7-3.2)	2.0 (0.3-12.5)	4.9 (2.1-11.2)
	Coloured	3.2 (2.3-4.6)	16.7	NA	NA	6.3 (5.4-7.3)	5.9 (4.2-7.5)	6.5 (2.7-14.8)	8.2 (4.9-13.6)
	Black	19.9 (18.3-21.7)	31.3	NA	NA	8.3 (7.8-8.8)	13.2 (12.2-14.2)	9.2 (7.1-12.0)	16 (13.9-18.3)
Kenya	Nairobi	7.2 (4.2-12.2)	34.8	NA	NA	1.4 (0.8-2.1)	0.91 (0.23-3.5)	16.0 (11.7-20.8)	25.8 (19.8-32.9)
	Central	4.6 (3.2-6.8)	43.8	NA	NA	1.2 (0.7-1.8)	0.88 (0.22-3.3)	14.5 (11.1-18.4)	24.4 (17.3-33.3)
	Coast	4.2 (2.6-7.0)	61.8	NA	NA	1.8 (1.1-2.7)	0.18 (0.04-0.77)	11.7 (8.5-15.6)	19.9 (14.2-27.3)
	Eastern	3.5 (2.1-5.5)	39.9	NA	NA	2.2 (1.5-3.1)	0.24 (0.05-1.0)	6.4 (4.2-9.2)	12.2 (7.4-19.5)
	Nyanza	13.9 (11.0-17.9)	76.2	NA	NA	2.5 (1.8-3.4)	3.45 (2.0-5.8)	18.2 (14.9-21.9)	30.6 (23.8-38.3)
UK	Rift Valley	4.7 (3.1-7.3)	52.6	NA	NA	1.8 (1.2-2.6)	0.54 (0.2-1.6)	10.3 (7.6-13.5)	14.4 (10.5-19.4)
	Western	6.6 (4.9-9.2)	59.9	NA	NA	1.0 (0.5-1.7)	1.52 (0.7-3.2)	18.8 (14.0-24.3)	20.9 (18.4-23.7)
	North Eastern	0.9 (0.3-4.0)	15.6	NA	NA	0.5 (0.2-1.4)	0 (0-0)	NA	NA
	White	0	NA	227	121	22	NA	13.9 (12.7-15.3)	29.6 (28.1-31.1)
	Black Caribbean	2.8	NA	1540	840	71	NA	25.4 (17.6-35.1)	41.6 (31.3-52.7)
USA	Black African	2.8	NA	1011	410	29	NA	34.5 (22.8-48.3)	43.8 (34.1-54.1)
	Asian	0	NA	136	109	18	NA	16.1 (8.0-29.8)	23.2 (16.5-31.5)
	Non-Hispanic white	0.22 (0.21-0.24)	17.6	211	66	5.4	NA	3.1 (2.2-4.3)	15.0 (14.0-16.5)
	Hispanic	0.59 (0.53-0.64)	22.3	404	114	11.8	NA	8.9 (3.6-19.9)	19.9 (15.7-24.7)
	Non-Hispanic black	1.72 (1.61-1.82)	45.9	1176	548	24.2	NA	11.3 (7.5-16.7)	27.1 (23.8-31.4)

NA – Not Available

[§] Incidence in cases per 100 000 population per year⁺ Incidence for Kenya is percent positive of 15–49 year old population, for South Africa is percent antenatal population positive and for UK and USA is cases per 100 000 population per year[#] The definitions used for STI, symptoms, concurrency and multiple partners varied between studies – see [Table 1](#) for details

Table 5. Prevalence/incidence of HIV, HSV-2, chlamydia, gonorrhoea, syphilis, concurrency and multiple partnering by men who have sex with men (MSM) versus heterosexual men the United Kingdom and the United States.

Country	Group	HIV %	Chlamydia [§]	Gonorrhoea [§]	Syphilis [§]	Concurrency % (95% CI)	Multiple partners	Lifetime partners ⁺
USA	MSM	7.2	NA	NA	233	31.3	86%	67
	Heterosexual men	0.2	NA	NA	7	9.7	56%	10
UK	MSM	6.3 (13.4) [#]	24 (80) [#]	41 (108) [#]	7.4 (42.4) [#]	52.4 (43.5-61.1)	24	111.1
	Heterosexual men	0.16	2	0.6	0.04	15.6 (14.5-16.7)	3.8	14.3

NA- Not Available

[§] Incidence in cases per 100 000 population per year.[#] The incidence estimates are provided for MSM living outside of London and in parentheses the incidence figures for MSM living in London.⁺ The mean number of lifetime partners reported by 35–39 years olds in the USA and aged 16–74 in the UK.**Table 6. Associations between incidence/prevalence of HIV, HSV-2, syphilis, gonorrhoea, chlamydia, male urethral discharge, concurrency and multiple partnering by ethnic group in (A) Kenya and South Africa; and (B) the United Kingdom and United States (Pearson's Correlations).**

Section A							
	HIV	HSV-2	Syphilis	Urethral Discharge	Concurrency	Multiple Partners	
Kenya							
HIV	1.00						
HSV-2	0.65	1.00					
Syphilis	0.63	0.39	1.00				
Urethral Discharge	0.77	0.68	0.34	1.00			
Concurrency	0.46	0.43	-0.34	0.72	1.00		
Multiple Partners	0.66	0.38	0.00	0.76	0.84	1.00	
South Africa							
HIV	1.00						
HSV-2	0.96	1.00					
Syphilis	0.78	0.85	1.00				
Urethral Discharge	0.98	0.99	0.90	1.00			
Concurrency	0.86	0.92	0.99	0.95	1.00		
Multiple Partners	0.99	0.99	0.87	0.99	0.93	1.00	
Section B							
	HIV	HSV-2	Syphilis	Chlamydia	Gonorrhoea	Concurrency	Multiple Partners
UK							
HIV	1.00						
HSV-2	NA	NA					
Syphilis	0.71	NA	1.00				
Chlamydia	0.94	NA	0.90	1.00			
Gonorrhoea	0.86	NA	0.97	0.98	1.00		
Concurrency	0.91	NA	0.37	0.73	0.58	1.00	
Multiple Partners	0.96	NA	0.63	0.89	0.78	0.89	1.00
USA							
HIV	1.00						
HSV-2	0.99	1.00					
Syphilis	0.99	0.92	1.00				
Chlamydia	0.99	0.97	0.99	1.00			
Gonorrhoea	0.99	0.99	0.97	0.99	1.00		
Concurrency	0.89	0.68	0.91	0.84	0.79	1.00	
Multiple Partners	0.98	0.89	0.99	0.98	0.95	0.94	1.00

Table 7. Prevalence/incidence of HIV, syphilis, HSV-2, chlamydia, gonorrhoea, urethral discharge, genital ulcers, concurrent partners and lifetime number of partners by ethnic group in four countries and by sexual orientation in two countries. (The precise definition of each variable is provided in the methods section,.)

Country	Ethnic group/ Sexual orientation	Race	HIV Prevalence (%)	Syphilis Prevalence (%)	HSV-2 Prevalence (%)	Concurrency Prevalence (%)	Multiple Partners (%)	Urethral Discharge (%)	Genital Ulcer (%)	Chlamydia Incidence	Gonorrhoea Incidence	Syphilis Incidence/Prevalence	Sex Partners (N)
USA	Ethnic grp	Non-Hispanic white	0	1	12	3	15	6	NA	211	66	5	NA
USA	Ethnic grp	Hispanic	1		10	9	20	9	NA	404	114	12	NA
USA	Ethnic grp	Non-Hispanic black	2	3	39	11	27	32	NA	1176	548	24	NA
SA	Ethnic grp	White	1	0	20	2	5	2	NA	NA	NA	0	NA
SA	Ethnic grp	Coloured	3	6	30	7	8	6	NA	NA	NA	6	NA
SA	Ethnic grp	Black	20	8	61	9	16	13	NA	NA	NA	8	NA
Kenya	Ethnic grp	North East	1	1	16	NA	NA	0	NA	NA	NA	1	NA
Kenya	Ethnic grp	Central	4	1	44	15	24	1	2	NA	NA	1	NA
Kenya	Ethnic grp	Eastern	5	2	40	6	12	0	2	NA	NA	2	NA
Kenya	Ethnic grp	Western	5	1	60	19	21	2	6	NA	NA	1	NA
Kenya	Ethnic grp	Rift Valley	6	2	53	10	14	1	2	NA	NA	2	NA
Kenya	Ethnic grp	Coast	8	2	62	12	20	0	2	NA	NA	2	NA
Kenya	Ethnic grp	Nairobi	9	1	35	16	26	1	1	NA	NA	1	NA
Kenya	Ethnic grp	Nyanza	15	3	76	18	31	3	2	NA	NA	3	NA
UK	Ethnic grp	White	0	2	NA	14	30	11	NA	227	121	22	NA
UK	Ethnic grp	Asian	0	3	NA	16	23	3	NA	136	109	18	NA
UK	Ethnic grp	Black African	3	4	NA	35	44	16	NA	1011	410	29	NA
UK	Ethnic grp	Black Caribbean	3	19	NA	25	42	20	NA	1540	840	71	NA
USA	Sexual orientation	Heterosexual men	0	NA	NA	10	10	NA	NA	NA	NA	7	10
USA	Sexual orientation	MSM	7	NA	NA	31	67	NA	NA	NA	NA	233	67
UK	Sexual orientation	Heterosexual men	0	NA	NA	16	4	NA	NA	2	1	0	14
	Sexual orientation	MSM	6.3	NA	NA	52	24	NA	NA	24	41	7.4	111.1

STIs and in the case of syphilis, was able to explain longitudinal changes in incidence. These findings do not fit with racial theories that biological differences between races explain differential STI spread⁷³. They fit better with sociological explanations that differences in sexual behaviours result in differences in network connectivity and consequent STI incidence^{9,74}. Furthermore, they suggest that behaviour change is possible and can result in rapid reductions in STI incidence. In fact, if network connectivity is a key determinant of STI incidence then it follows that reducing network connectivity would be an optimal way to effect radical STI prevention. Whilst high STI prevalence populations would first need to decide if they wanted to change behaviours that enhance connectivity, there are a number of positive precedents for this. Uganda's 'Zero grazing' campaign was credited with large scaled reductions in multiple partnering and a subsequent decline in HIV prevalence⁷⁵. Similar campaigns amongst MSM in the 1980s had a similar effect^{7,57}.

These insights are relevant in the current setting of increasing incidence of numerous STIs in high income countries¹. A number of analyses have found that a large proportion of these increases are occurring in MSM with high rates of partner change

and particularly in contemporary HIV pre exposure prophylaxis (PrEP) recipients^{1,38,76}. It is not unusual for PrEP recipients to report a median of 12 partners (IQR 6–25) per 3 months⁷⁷. Unsurprisingly these translate into very high STI incidence - the combined incidence of gonorrhoea, chlamydia and syphilis can be up to 169 per 100 person years⁷⁸. Tackling this high incidence via intensive screening and treating without addressing the underlying cause may result in antimicrobial resistance⁷⁹.

Further studies are required to further elucidate the links between norms, structural factors, behaviour, sexual networks and STI incidence. In the interim, the type of evidence reviewed here suggests that reducing STI incidence/prevalence in high prevalence populations to those of low prevalence population may be difficult without reducing network connectivity.

Data availability

Underlying data

All data underlying the results are available as part of the article and no additional source data are required (see [Table 1–Table 3](#) and [Table 7](#)).

References

- Unemo M, Bradshaw CS, Hocking JS, *et al.*: **Sexually transmitted infections: challenges ahead.** *Lancet Infect Dis.* 2017; 17(8): e235–e279.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Aral SO, Leichter JS, Blanchard JF: **Overview: the role of emergent properties of complex systems in the epidemiology and prevention of sexually transmitted infections including HIV infection.** *Sex Transm Infect.* 2010; 86 Suppl 3: iii1–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Morris M, Kurth AE, Hamilton DT, *et al.*: **Concurrent partnerships and HIV prevalence disparities by race: linking science and public health practice.** *Am J Public Health.* 2009; 99(6): 1023–1031.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kenyon C, Buyze J, Colebunders R: **HIV prevalence by race co-varies closely with concurrency and number of sex partners in South Africa.** *PLoS One.* 2013; 8(5): e64080.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kenyon C, Menten J, Vu L, *et al.*: **Male circumcision and sexual risk behaviors may contribute to considerable ethnic disparities in HIV prevalence in Kenya: an ecological analysis.** *PLoS One.* 2014; 9(8): e106230.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Beyrer C, Baral SD, van Griensven F, *et al.*: **Global epidemiology of HIV infection in men who have sex with men.** *Lancet.* 2012; 380(9839): 367–377.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Peterman TA, Su J, Bernstein KT, *et al.*: **Syphilis in the United States: on the rise?** *Expert Rev Anti Infect Ther.* 2015; 13(2): 161–168.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kenyon CR, Delva W: **It's the network, stupid: a population's sexual network connectivity determines its STI prevalence [version 2; peer review: 2 approved].** *F1000Res.* 2018; 7: 1880.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Morris M, Goodreau S, Moody J: **Sexual networks, concurrency and STD/HIV.** In: *Sex Transm Dis.* 4th edn. Edited by Holmes KK. New York: McGraw-Hill Medical. 2008; xxv: 2166.
[Reference Source](#)
- Morris M, Epstein H, Wawer M: **Timing is everything: international variations in historical sexual partnership concurrency and HIV prevalence.** *PLoS One.* 2010; 5(11): e14092.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Koumans EH, Sternberg M, Bruce C, *et al.*: **The prevalence of bacterial vaginosis in the United States, 2001–2004; associations with symptoms, sexual behaviors, and reproductive health.** *Sex Transm Dis.* 2007; 34(11): 864–869.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Garnett GP, Mertz KJ, Finelli L, *et al.*: **The transmission dynamics of gonorrhoea: modelling the reported behaviour of infected patients from Newark, New Jersey.** *Philos Trans R Soc Lond B Biol Sci.* 1999; 354(1384): 787–797.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ghani AC, Swinton J, Garnett GP: **The role of sexual partnership networks in the epidemiology of gonorrhoea.** *Sex Transm Dis.* 1997; 24(1): 45–56.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Helleringer S, Kohler HP, Kalilani-Phiri L: **The association of HIV serodiscordance and partnership concurrency in Likoma Island (Malawi).** *AIDS.* 2009; 23(10): 1285–1287.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kenyon CR, Tsoumanis A, Schwartz IS, *et al.*: **Partner concurrency and HIV infection risk in South Africa.** *Int J Infect Dis.* 2016; 45: 81–87.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Adimora AA, Schoenbach VJ, Bonas DM, *et al.*: **Concurrent sexual partnerships among women in the United States.** *Epidemiology.* 2002; 13(3): 320–327.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kenyon C, Colebunders R, Buve A, *et al.*: **Partner-concurrency associated with herpes simplex virus 2 infection in young South Africans.** *Int J STD AIDS.* 2013; 24(10): 804–812.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Shi R, Devarakonda S, Liu L, *et al.*: **Factors associated with genital human papillomavirus infection among adult females in the United States, NHANES 2007–2010.** *BMC Res Notes.* 2014; 7: 544.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sonnenberg P, Clifton S, Beddows S, *et al.*: **Prevalence, risk factors, and uptake of interventions for sexually transmitted infections in Britain: findings from the National Surveys of Sexual Attitudes and Lifestyles (Natsal).** *Lancet.* 2013; 382(9907): 1795–1806.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Koumans EH, Farley TA, Gibson JJ, *et al.*: **Characteristics of persons with syphilis in areas of persisting syphilis in the United States: sustained transmission associated with concurrent partnerships.** *Sex Transm Dis.* 2001; 28(9): 497–503.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kenyon C, Colebunders R: **Strong association between point-concurrency and national peak HIV prevalence.** *Int J Infect Dis.* 2012; 16(11): e826–827.
[PubMed Abstract](#) | [Publisher Full Text](#)

22. Kenyon C: **Strong associations between national prevalence of various STIs suggests sexual network connectivity is a common underpinning risk factor.** *BMC Infect Dis.* 2017; **17**(1): 682.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Wellings K, Collumbien M, Slaymaker E, *et al.*: **Sexual behaviour in context: a global perspective.** *Lancet.* 2006; **368**(9548): 1706–1728.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Drain PK, Halperin DT, Hughes JP, *et al.*: **Male circumcision, religion, and infectious diseases: an ecologic analysis of 118 developing countries.** *BMC Infect Dis.* 2006; **6**: 172.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Drain PK, Smith JS, Hughes JP, *et al.*: **Correlates of national HIV seroprevalence: an ecologic analysis of 122 developing countries.** *J Acquir Immune Defic Syndr.* 2004; **35**(4): 407–420.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Sawers L, Stillwaggon E: **Concurrent sexual partnerships do not explain the HIV epidemics in Africa: a systematic review of the evidence.** *J Int AIDS Soc.* 2010; **13**: 34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Kenyon C, Buyze J, Schwartz IS: **Strong association between higher-risk sex and HIV prevalence at the regional level: an ecological study of 27 sub-Saharan African countries [version 1; peer review: 2 approved].** *F1000Res.* 2018; **7**: 1879.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Laumann EO: **The social organization of sexuality: Sexual practices in the United States.** University of Chicago Press; 1994.
[Reference Source](#)
29. Department of Health: **Kenya AIDS indicator survey 2007.** In. Nairobi; 2009.
[Reference Source](#)
30. Kenya National Bureau of Statistics (KNBS) and ICF Macro: **Kenya Demographic and Health Survey 2008–09.** In. Calverton: Maryland: KNBS and ICF Macro. 2010.
[Reference Source](#)
31. Kenyon C, Osbak K, Buyze J: **The prevalence of HIV by ethnic group is correlated with HSV-2 and syphilis prevalence in Kenya, South Africa, the United Kingdom and the United States.** *Interdiscip Perspect Infect Dis.* 2014; **2014**: 284317.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Population Services International: **Population Services International (PSI) Survey /Kenya 6th HIV Survey, 2011.** In. Nairobi: PSI; 2012.
33. Shisana O: **South African national HIV prevalence, HIV incidence, behaviour and communication survey, 2005.** Cape Town: HSRC Press; 2005.
[Reference Source](#)
34. South African Department of Health: **The 2012 National Antenatal Sentinel HIV & Herpes Simplex Type-2 Prevalence Survey.** In.: National Department of Health Pretoria; 2013.
[Reference Source](#)
35. Department of Health: **Syphilis screening in women attending antenatal clinics.** *Epidemiological Comments.* 1992; **19**(10): 166–171.
36. Department of Health: **South African Demographic and Health Survey 1998: Report.** In. Pretoria: Department of Health; 1999.
[Reference Source](#)
37. Department of Health, Medical Research Council, OrMacro: **South African Demographic and Health Survey 2003.** In. Pretoria: Department of Health; 2007.
[Reference Source](#)
38. Public Health England: **Sexually transmitted infections and screening for chlamydia in England, 2018.** *Health Protection Report.* 2019; **13**(19).
[Reference Source](#)
39. Fenton KA, Mercer CH, McManus S, *et al.*: **Ethnic variations in sexual behaviour in Great Britain and risk of sexually transmitted infections: a probability survey.** *Lancet.* 2005; **365**(9466): 1246–1255.
[PubMed Abstract](#) | [Publisher Full Text](#)
40. Dave SS, Fenton KA, Mercer CH, *et al.*: **Male circumcision in Britain: findings from a national probability sample survey.** *Sex Transm Infect.* 2003; **79**(6): 499–500.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Centers for Disease Control and Prevention (CDC): **HIV prevalence estimates—United States, 2006.** *MMWR Morb Mortal Wkly Rep.* 2008; **57**(39): 1073–6.
[PubMed Abstract](#)
42. Centers for Disease Control and Prevention (CDC): **Seroprevalence of herpes simplex virus type 2 among persons aged 14–49 years—United States, 2005–2008.** *MMWR Morb Mortal Wkly Rep.* 2010; **59**(15): 456–9.
[PubMed Abstract](#)
43. Gottlieb SL, Pope V, Sternberg MR, *et al.*: **Prevalence of syphilis seroreactivity in the United States: data from the National Health and Nutrition Examination Surveys (NHANES) 2001–2004.** *Sex Transm Dis.* 2008; **35**(5): 507–511.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Centers for Disease Control and Prevention: **Sexually Transmitted Disease Surveillance 2017.** Atlanta: US Department of Health and Human Services. 2018.
[Publisher Full Text](#)
45. Kenyon CR, Osbak K, Tsoumanis A: **The Global Epidemiology of Syphilis in the Past Century - A Systematic Review Based on Antenatal Syphilis Prevalence.** *PLoS Negl Trop Dis.* 2016; **10**(5): e0004711.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. World Health Organization: **Prevalence and incidence of selected sexually transmitted infections, *Chlamydia trachomatis*, *Neisseria gonorrhoeae*, syphilis, and *Trichomonas vaginalis*: methods and results used by the WHO to generate 2005 estimates.** Geneva World Health Organization; 2011.
[Reference Source](#)
47. Guthe T: **The treponematoses as a world problem.** *Br J Vener Dis.* 1960; **36**(2): 67–77.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. Simms I, Fenton KA, Ashton M, *et al.*: **The re-emergence of syphilis in the United Kingdom: the new epidemic phases.** *Sex Transm Dis.* 2005; **32**(4): 220–226.
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Stenger MR, Baral S, Stahlman S, *et al.*: **As through a glass, darkly: the future of sexually transmissible infections among gay, bisexual and other men who have sex with men.** *Sex Health.* 2017; **14**(1): 18–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Glick SN, Morris M, Foxman B, *et al.*: **A Comparison of Sexual Behavior Patterns Among Men Who Have Sex With Men and Heterosexual Men and Women.** *J Acquir Immune Defic Syndr.* 2012; **60**(1): 83–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
51. Mercer CH, Prah P, Field N, *et al.*: **The health and well-being of men who have sex with men (MSM) in Britain: Evidence from the third National Survey of Sexual Attitudes and Lifestyles (Natsal-3).** *BMC Public Health.* 2016; **16**: 525.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
52. Kenyon CR, Tsoumanis A, Schwartz IS: **A population's higher-risk sexual behaviour is associated with its average sexual behaviour—An ecological analysis of subpopulations in Ethiopia, Kenya, South Africa, Uganda and the United States.** *Epidemics.* 2016; **15**: 56–65.
[PubMed Abstract](#) | [Publisher Full Text](#)
53. Kirwan PD CC, Brown AE, Gill ON, *et al.*: **HIV in the UK - 2016 report.** Public Health England, London. 2016.
[Reference Source](#)
54. Johnson K, Way A: **Risk factors for HIV infection in a national adult population: evidence from the 2003 Kenya Demographic and Health Survey.** *J Acquir Immune Defic Syndr.* 2006; **42**(5): 627–636.
[PubMed Abstract](#) | [Publisher Full Text](#)
55. Aral SO, Fenton KA, Holmes KK: **Sexually transmitted diseases in the USA: temporal trends.** *Sex Transm Infect.* 2007; **83**(4): 257–266.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. Sandholzer TA: **Factors Affecting the Incidence and Management of Sexually Transmitted Diseases in Homosexual Men.** In: *Sexually Transmitted Diseases in Homosexual Men.* edn. Springer; 1983; 3–12.
[Publisher Full Text](#)
57. Chesson HW, Dee TS, Aral SO: **AIDS mortality may have contributed to the decline in syphilis rates in the United States in the 1990s.** *Sex Transm Dis.* 2003; **30**(5): 419–424.
[PubMed Abstract](#) | [Publisher Full Text](#)
58. Peterman TA, Heffelfinger JD, Swint EB, *et al.*: **The changing epidemiology of syphilis.** *Sex Transm Dis.* 2005; **32**(10 Suppl): S4–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
59. Crepaz N, Hart TA, Marks G: **Highly active antiretroviral therapy and sexual risk behavior: a meta-analytic review.** *JAMA.* 2004; **292**(2): 224–236.
[PubMed Abstract](#) | [Publisher Full Text](#)
60. Centers for Disease Control and Prevention (CDC): **Resurgent bacterial sexually transmitted disease among men who have sex with men—King County, Washington, 1997–1999.** *MMWR Morb Mortal Wkly Rep.* 1999; **48**(35): 773–7.
[PubMed Abstract](#)
61. Golden MR, Marra CM, Holmes KK: **Update on syphilis: resurgence of an old problem.** *JAMA.* 2003; **290**(11): 1510–1514.
[PubMed Abstract](#) | [Publisher Full Text](#)
62. Kenyon CR, Osbak K, Chico RM: **What underpins the decline in syphilis in Southern and Eastern Africa? An exploratory ecological analysis.** *Int J Infect Dis.* 2014; **29**: 54–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
63. Kenyon CR, Osbak K: **The Prevalence of Syphilis Is Associated with the Prevalence of Male Point-Concurrence: An Ecological Analysis.** *World J AIDS.* 2015; **5**(2): 131–139.
[Publisher Full Text](#)
64. Johnson LF, Coetzee DJ, Dorrington RE: **Sentinel surveillance of sexually transmitted infections in South Africa: a review.** *Sex Transm Infect.* 2005; **81**(4): 287–293.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
65. Johnson LF, Dorrington RE, Bradshaw D, *et al.*: **Sexual behaviour patterns in South Africa and their association with the spread of HIV: insights from a mathematical model.** *Demogr Res.* 2009; **21**: 289–340.
[Publisher Full Text](#)
66. Kenyon CR, Osbak K, Buyze J, *et al.*: **The changing relationship between bacterial STIs and HIV prevalence in South Africa - an ecological study.** *Int J STD AIDS.* 2015; **26**(8): 556–564.
[PubMed Abstract](#) | [Publisher Full Text](#)
67. Kenyon CR, Schwartz I, Buyze J: **Did AIDS mortality decrease the number of lifetime sexual partners in Kenya: an ecological analysis?** *Epidemiol Infect.* 2016; **144**(3): 556–559.
[PubMed Abstract](#) | [Publisher Full Text](#)

68. Temmerman M, Fonck K, Bashir F, *et al.*: **Declining syphilis prevalence in pregnant women in Nairobi since 1995: another success story in the STD field?** *Int J STD AIDS*. 1999; **10**(6): 405–408.
[PubMed Abstract](#) | [Publisher Full Text](#)
69. Laumann EO, Ellingson S, Mahay J, *et al.*: **The sexual organization of the city**. In. Chicago: University of Chicago; 2005.
[Publisher Full Text](#)
70. Laumann EO, Youm Y: **Racial/ethnic group differences in the prevalence of sexually transmitted diseases in the United States: a network explanation**. *Sex Transm Dis*. 1999; **26**(5): 250–261.
[PubMed Abstract](#) | [Publisher Full Text](#)
71. Goodreau SM, Golden MR: **Biological and demographic causes of high HIV and sexually transmitted disease prevalence in men who have sex with men**. *Sex Transm Infect*. 2007; **83**(6): 458–462.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
72. Kretzschmar M, Morris M: **Measures of concurrency in networks and the spread of infectious disease**. *Math Biosci*. 1996; **133**(2): 165–195.
[PubMed Abstract](#) | [Publisher Full Text](#)
73. Kaul R, Cohen CR, Chege D, *et al.*: **Biological factors that may contribute to regional and racial disparities in HIV prevalence**. *Am J Reprod Immunol*. 2011; **65**(3): 317–324.
[PubMed Abstract](#) | [Publisher Full Text](#)
74. Kenyon C, Zondo S: **Why do some South African ethnic groups have very high HIV rates and others not?** *Afr J AIDS Res*. 2011; **10**(1): 51–62.
[PubMed Abstract](#) | [Publisher Full Text](#)
75. Green EC, Halperin DT, Nantulya V, *et al.*: **Uganda's HIV prevention success: the role of sexual behavior change and the national response**. *AIDS Behav*. 2006; **10**(4): 335–346; discussion 347–350.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
76. Ramchandani MS, Golden MR: **Confronting Rising STIs in the Era of PrEP and Treatment as Prevention**. *Curr HIV/AIDS Rep*. 2019; **16**(3): 244–256.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
77. Hoornenborg E, Coyer L, van Laarhoven A, *et al.*: **Change in sexual risk behaviour after 6 months of pre-exposure prophylaxis, use: results from the Amsterdam pre-exposure prophylaxis demonstration project**. *AIDS*. 2018; **32**(11): 1527–1532.
[PubMed Abstract](#) | [Publisher Full Text](#)
78. Beymer MR, DeVost MA, Weiss RE, *et al.*: **Does HIV pre-exposure prophylaxis use lead to a higher incidence of sexually transmitted infections? A case-crossover study of men who have sex with men in Los Angeles, California**. *Sex Transm Infect*. 2018; **94**(6): 457–462.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
79. Kenyon C, Schwartz IS: **Effects of Sexual Network Connectivity and Antimicrobial Drug Use on Antimicrobial Resistance in *Neisseria gonorrhoeae***. *Emerg Infect Dis*. 2018; **24**(7): 1195–1203.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status: ? ?

Version 1

Reviewer Report 18 July 2022

<https://doi.org/10.5256/f1000research.27545.r142231>

© 2022 Levy I. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Itzchak Levy

¹ Tel-Aviv University, Tel-Aviv, Israel

² Sheba Medical Centre, Tel-Aviv, Israel

The study done by Chris Kenyon is very interesting and important. It tries to show the association between the connectivity within a group (which in this work is expressed by the number of partners in a certain time and the concurrency of partners) and the incidence/prevalence of certain sexually transmitted infections (STIs). This is important because understanding sexual networks and how they work and what influences them is very important in the prevention of STIs. It adds to the knowledge we get from the study of social and sexual networks as well to phylogenetic studies of the spread of STIs.

However, since this paper is constructed from many different works that studied different questions (incidence of some STIs, prevalence of STIs, sexual behavior data, etc.) as a clinician I don't understand how the conclusions of the writer can be made and I feel that the input of a statistician would be much more helpful.

It sounds more reasonable to take one or two STIs which are compatible with a dynamic study (e.g. gonorrhea and chlamydia) and to concentrate on a more defined population. However, this study still gives the "wide picture".

HIV in my opinion is not suitable for this study because its prevalence is influenced not only by the incidence but also by some other factors such as the introduction of anti-retroviral therapy (ART) in 1996 and the introduction of better treatments like integrase inhibitors in the 2000s. Incidence may be also influenced by introduction of earlier treatment (treatment as prevention) and introduction of pre exposure prophylaxis (PREP) so it is also less suitable for this study.

The methods section should be somewhat explained better. It is not understandable if the work is a systematic review of the literature, then it would be helpful if you can explain the method by which the specific studies were chosen and why others were not (a flow chart may be helpful).

There is a mixture of studies, some show incidence and some show prevalence – this mixture is confusing.

With such a big population I don't quite understand what is the meaning of the correlation.

There are a lot of prevalence studies which utilize serology like studies 42 and 43 (HSV2 and syphilis). There is no way to know the date of infection in those studies so it is a very big limitation to use them for the purpose of this study which tries to show dynamics.

In Kenya (page 6) ethnicity was a region of residence served as a proxy for ethnicity. I am not sure that this approximation can be done safely for the purpose of this study because it is well established that MSM tend to reside in big cities without regard to their ethnicity so I would leave it as "regional".

In Figure 2 (page 7) The rate of MSM in each ethnic group is not given and this may influence the data and the conclusions.

In Figure 4 legend (page 9) the word the united states should be omitted.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

No

If applicable, is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Infectious Diseases

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 24 Aug 2022

Chris Kenyon, Institute of Tropical Medicine, Antwerp, Antwerp, Belgium

Reply to reviewer 2:

The study done by Chris Kenyon is very interesting and important. It tries to show the association between the connectivity within a group (which in this work is expressed by the number of partners in a certain time and the concurrency of partners) and the incidence/prevalence of certain sexually transmitted infections (STIs). This is important because understanding sexual networks and how they work and what influences them is very important in the prevention of STIs. It adds to the knowledge we get from the study of social and sexual networks as well to phylogenetic studies of the spread of STIs.

However, since this paper is constructed from many different works that studied different questions (incidence of some STIs, prevalence of STIs, sexual behavior data, etc.) as a clinician I don't understand how the conclusions of the writer can be made and I feel that the input of a statistician would be much more helpful.

It sounds more reasonable to take one or two STIs which are compatible with a dynamic study (e.g. gonorrhea and chlamydia) and to concentrate on a more defined population. However, this study still gives the "wide picture".

HIV in my opinion is not suitable for this study because its prevalence is influenced not only by the incidence but also by some other factors such as the introduction of anti-retroviral therapy (ART) in 1996 and the introduction of better treatments like integrase inhibitors in the 2000s. Incidence may be also influenced by introduction of earlier treatment (treatment as prevention) and introduction of pre exposure prophylaxis (PREP) so it is also less suitable for this study.

Reply:

Thank you for these useful comments. The limitations section has been expanded to include the concerns raised by the reviewer (P 23, second paragraph):

Differential uptake of antiretroviral therapy is an important potential determinant of differential HIV prevalence between sub-populations. Once again, we did not control for this.

It should be noted that in the case of South Africa the differential HIV prevalence between ethnic groups was of a similar magnitude in surveys done before ART became available to those following the widespread use of ART.

The methods section should be somewhat explained better. It is not understandable if the work is a systematic review of the literature, then it would be helpful if you can explain the method by which the specific studies were chosen and why others were not (a flow chart may be helpful).

Reply:

As noted above the countries were not selected via a systematic review methodology. The rationale for selecting the studies has been added to the introduction. This limitation has been added to the relevant section of the limitations in the discussion (P 23, 3rd Paragraph).

In addition, the following sentence has been added to the introduction to make it clear how the four countries were chosen (P3, third last sentence).

These countries were chosen due to the familiarity of the author with these countries and the availability of nationally representative datasets with both sexual behaviour and STI prevalence data.

Finally, we have changed the title of the paper from "a four-country analysis" to "a selected four-country analysis."

There is a mixture of studies, some show incidence and some show prevalence – this mixture is confusing.

With such a big population I don't quite understand what is the meaning of the correlation. There are a lot of prevalence studies which utilize serology like studies 42 and 43 (HSV2 and syphilis). There is no way to know the date of infection in those studies so it is a very big limitation to use them for the purpose of this study which tries to show dynamics.

Reply:

It is true that one cannot know the date of infection in studies that define STI incidence/prevalence with HIV, HSV-2 serology or a treponemal test only in the case of syphilis. These variables are then best considered as a measure of cumulative incidence. A number of studies have found that these variables offer a valuable marker of how connected sexual networks are and what their risk is for other STIs (1, 2). As an example, the prevalence of HSV-2 in 20-24 year olds can predict subsequent HIV prevalence (2). The same might be true for emergent STIs such as monkeypox. For most of the studies of syphilis incidence/prevalence used in this paper, syphilis was defined by both a treponemal and non-treponemal test.

In Kenya (page 6) ethnicity was a region of residence served as a proxy for ethnicity. I am not sure that this approximation can be done safely for the purpose of this study because it is well established that MSM tend to reside in big cities without regard to their ethnicity so I would leave it as "regional".

In Figure 2 (page 7) The rate of MSM in each ethnic group is not given and this may influence the data and the conclusions.

Reply:

We provide a justification for the use of regions as a proxy for ethnicity in Kenya in the Methods section (P 11, third paragraph):

In addition, we have added this as a limitation in the discussion section (P 23, second paragraph):

The way the behavioural variables were defined varied between studies and in the case of Kenya, region was used as a proxy for ethnic group

Regions vs. ethnic group in Kenya. *Many of the national STI surveillance surveys in Kenya do not collect data on ethnic group²⁹. There is however a high correlation between region of residence and ethnic group, with large differences in STI incidence and sexual behaviour between regions^{5, 31}. We therefore follow the approach used in previous studies and use region of residence as a proxy for ethnic group^{31, 54}.*

In Figure 4 legend (page 9) the word the united states should be omitted.

Reply:

Thanks for pointing out this error, which has been corrected.

References

1. Kenyon C. Strong associations between national prevalence of various STIs suggests sexual network connectivity is a common underpinning risk factor. BMC Infect Dis. 2017;17(1):682.
2. Kenyon C, Colebunders R, Hens N. Determinants of generalized herpes simplex virus-2 epidemics: the role of sexual partner concurrency. Int J STD AIDS. 2013;24(5):375-82.

Competing Interests: No competing interests

Reviewer Report 08 September 2021

<https://doi.org/10.5256/f1000research.27545.r92342>

© 2021 Khosropour C. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Christine Khosropour

Department of Epidemiology, University of Washington, Seattle, WA, USA

In this review, Dr. Kenyon reviews data on STI/HIV incidence/prevalence and partner concurrency from (mostly) population-based studies in four counties. The results show that STIs cluster within certain groups defined by race/ethnicity and gender of sex partners, and that STIs were correlated with number of sex partners and concurrency. It is helpful to have these data summarized for these countries. The author appropriately addresses the limitations. My main comment has to do with the transparency in how these papers were selected. I think adding in the full search criteria, the full exclusion/inclusion criteria, the number of studies identified and eliminated, etc would be appropriate here. I also had trouble identifying which data supported which conclusions. These and other comments are detailed below.

Minor comments:

1. Could the author specify upfront their use of sex/gender terms throughout the paper? Even just to say that the terms they use reflect those from published papers, where it may have been unclear whether the studies referred to sex at birth or gender identity.
2. The author presents data stratified by "ethnic groups". Could the author provide some description in the methods about how stratification of these groups was operationalized, by country? There is some of that for Kenya already.
3. Page 5 of the PDF, second to last paragraph: The author states: "these variations in definitions between studies mean it would be appropriate to compare these variables across studies". Did the author mean to say "it would be *inappropriate*"?
4. Page 8 of the PDF, first paragraph: I believe the reference should be to Figure 2, not Figure 1.

5. The author uses the terms “heterosexual men” and “men who have sex with women” (Table 2). I think it would be helpful to choose one throughout. There are also places where the author states “heterosexuals” which I assume to be men only but it is a little unclear
6. There appears to be a typo in the Figure 4 title (the phrase “the United States” should not be there)

Major comments:

1. The paper is not described as a systematic review, but was it intended to be? Given that there are summary quantitative estimates provided, it feels like a systematic review. If so, I think it would have been helpful in the results to see how many papers were identified using the search criteria and how many were eliminated, and for what reasons. When I ran this search (using “or” between the terms but also making sure the countries were included) I get >225,000 results. I assume the author used a different order of terms. Could the full order of search terms be included in the paper?
2. Somewhat related to the above, I am having a bit of trouble figuring out how studies were selected. For example, it appears that there were only 3 studies from Kenya published since 1990 that make comparisons on the prevalence of STIs/HIV, multiple partners, or concurrency. Is that the case? Or were these the only population-based studies? I think adding in inclusion/exclusion criteria would be helpful.
3. To calculate chi-square values for incidence/prevalence, did the author pool raw numbers from different studies or use some other method? Would there be potential for violation of an assumption of independence of observations?
4. Figure 3: I do not understand the primary Y axis here (labeled “HIV, syphilis, concurrency, lifetime partners”). Is this the percentage of individuals who reported living with HIV, being diagnosed with syphilis, reporting concurrency, or reporting over a certain number of partners?
5. The discussion focuses on the relationship between STIs and concurrency. Unless I misunderstand the data, it seems that these data are mostly from Table 6, which shows the correlation of STIs and concurrency and multiple partners. To me, it does not look like the associations are particularly strong in Kenya, and it does not really hold for syphilis in the UK or gonorrhea in the UK. Are there other data from which the author is drawing to make these conclusions?
6. Page 14 of the PDF, first paragraph: I would change the phrase: “These findings do not fit with racial theories that biological differences...” to “These findings do not fit with *racist* theories that biological differences...”
7. Page 14 of the PDF, first paragraph: The author suggests reducing network connectivity as a way to decrease STI spread. I do not understand how that would be operationalized. Is the author referring to reducing concurrency? I am not familiar with the Zero grazing campaign; perhaps that could be explained more to help the reader understand the concept of “breaking up” network connectivity

8. In the conclusion, I think it may be worth adding a comment about investment in STI vaccines being a plausible solution to curbing STI rates. Probably a more viable solution than behavior change.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

No

If applicable, is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: STI/HIV epidemiology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 24 Aug 2022

Chris Kenyon, Institute of Tropical Medicine, Antwerp, Antwerp, Belgium

Reply to reviewer 1:

1 In this review, Dr. Kenyon reviews data on STI/HIV incidence/prevalence and partner concurrency from (mostly) population-based studies in four countries. The results show that STIs cluster within certain groups defined by race/ethnicity and gender of sex partners, and that STIs were correlated with number of sex partners and concurrency. It is helpful to have these data summarized for these countries. The author appropriately addresses the limitations. My main comment has to do with the transparency in how these papers were selected. I think adding in the full search criteria, the full exclusion/inclusion criteria, the number of studies identified and eliminated, etc would be appropriate here. I also had trouble identifying which data supported which conclusions. These and other comments are detailed below.

Reply:

The following sentence has been added to the introduction to make it clear how the four countries were chosen (P3, third last sentence). These countries were chosen due to the familiarity of the author with these countries and the availability of nationally representative datasets with both sexual behaviour and STI prevalence data.

Minor comments:

- Could the author specify upfront their use of sex/gender terms throughout the paper? Even just to say that the terms they use reflect those from published papers, where it may have been unclear whether the studies referred to sex at birth or gender identity.

Reply: A new section “Terminology” has been added to the Methods section to address this point:

Sex, gender, ethnicity and race: Where this was specified, each data source used a slightly different definition of sex, gender, ethnicity and race. Wherever possible, we used the same terms used in these original sources.

- The author presents data stratified by “ethnic groups”. Could the author provide some description in the methods about how stratification of these groups was operationalized, by country? There is some of that for Kenya already.

Reply: A new section to the methods section has been added to address this question (P11, penultimate paragraph).

- Page 5 of the PDF, second to last paragraph: The author states: “these variations in definitions between studies mean it would be appropriate to compare these variables across studies”. Did the author mean to say “it would be *inappropriate*”

Reply: Thanks for pointing this typo out. It has been corrected.

- Page 8 of the PDF, first paragraph: I believe the reference should be to Figure 2, not Figure 1.

Reply: Thanks for pointing this error out. It has been corrected.

- The author uses the terms “heterosexual men” and “men who have sex with women” (Table 2). I think it would be helpful to choose one throughout. There are also places where the author states “heterosexuals” which I assume to be men only but it is a little unclear

Reply: The phrase “men who have sex with women” has been deleted everywhere and replaced with “heterosexual men.”

- There appears to be a typo in the Figure 4 title (the phrase “the United States” should not be there)

Reply: Thanks for pointing this error out. It has been corrected.

Major comments:

- The paper is not described as a systematic review, but was it intended to be? Given that there are summary quantitative estimates provided, it feels like a systematic review. If so, I think it would have been helpful in the results to see how many papers were identified using the search criteria and how many were eliminated, and for what reasons. When I ran this search (using “or” between the terms but also making sure the countries were included) I get >225,000 results. I assume the author used a different order of terms. Could the full order of search terms be included in the paper?

Reply: As noted above the countries were not selected via a systematic review methodology. The rationale for selecting the studies has been added to the introduction. This limitation has been added to the relevant section of the discussion (P 23, 3rd Paragraph).

- Somewhat related to the above, I am having a bit of trouble figuring out how studies were selected. For example, it appears that there were only 3 studies from Kenya published since 1990 that make comparisons on the prevalence of STIs/HIV, multiple partners, or concurrency. Is that the case? Or were these the only population-based studies? I think adding in inclusion/exclusion criteria would be helpful.

Reply:

Where possible surveys using nationally representative samples were used. Where this data was not available other sources were used such as detailed in Table 2. This information has been added to the methods section.

- To calculate chi-square values for incidence/prevalence, did the author pool raw numbers from different studies or use some other method? Would there be potential for violation of an assumption of independence of observations?

Reply: Chi-squared tests were performed in the raw numbers per group. It is acknowledged that this approach is of limited utility and if the reviewer prefers, these statistical tests could be removed.

- Figure 3: I do not understand the primary Y axis here (labeled “HIV, syphilis, concurrency, lifetime partners). Is this the percentage of individuals who reported living with HIV, being diagnosed with syphilis, reporting concurrency, or reporting over a certain number of partners?

Reply: These are defined in the Figure 3 legend as follows: HIV and partner concurrency prevalence are reported as percentages, multiple partners as the number of lifetime partners and chlamydia, gonorrhoea and syphilis incidence as cases per 100 000 per year.

- The discussion focuses on the relationship between STIs and concurrency. Unless I misunderstand the data, it seems that these data are mostly from Table 6, which shows the correlation of STIs and concurrency and multiple partners. To me, it does not look like the associations are particularly strong in Kenya, and it does not really

hold for syphilis in the UK or gonorrhea in the UK. Are there other data from which the author is drawing to make these conclusions?

Reply: The data referred to is that from Table 6. I agree with the reviewer's comments and have added this concern to the list of limitations (P23, third paragraph): Not all the associations between multiple partnering/ concurrency and STI incidence/prevalence were strongly positive. In fact, one association (between syphilis and concurrency by ethnic group in Kenya) was weakly negative (Table 6).

- Page 14 of the PDF, first paragraph: I would change the phrase: "These findings do not fit with racial theories that biological differences..." to "These findings do not fit with *racist* theories that biological differences..."

Reply: This change has been made.

- Page 14 of the PDF, first paragraph: The author suggests reducing network connectivity as a way to decrease STI spread. I do not understand how that would be operationalized. Is the author referring to reducing concurrency? I am not familiar with the Zero grazing campaign; perhaps that could be explained more to help the reader understand the concept of "breaking up" network connectivity

Reply: This section has been expanded as suggested.

- In the conclusion, I think it may be worth adding a comment about investment in STI vaccines being a plausible solution to curbing STI rates. Probably a more viable solution than behavior change.

Reply: Thank you for this point, which has been added to the last sentence of the paper.

Competing Interests: No competing interests

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research