REVIEW

Chromatin remodelling comes into focus [version 1; peer review: 4 approved]

Ramasubramian Sundaramoorthy, Tom Owen-Hughes

Centre for Gene Regulation and Expression, School of Life Sciences, University of Dundee, Dundee, Dundee, DD1 5EH, UK

V1 First published: 20 Aug 2020, 9(Faculty Rev):1011
<https://doi.org/10.12688/f1000research.21933.1>

Latest published: 20 Aug 2020, 9(Faculty Rev):1011
<https://doi.org/10.12688/f1000research.21933.1>

Abstract

ATP-dependent chromatin remodelling enzymes are molecular machines that act to reconfigure the structure of nucleosomes. Until recently, little was known about the structure of these enzymes. Recent progress has revealed that their interaction with chromatin is dominated by ATPase domains that contact DNA at favoured locations on the nucleosome surface. Contacts with histones are limited but play important roles in modulating activity. The ATPase domains do not act in isolation but are flanked by diverse accessory domains and subunits. New structures indicate how these subunits are arranged in multi-subunit complexes providing a framework from which to understand how a common motor is applied to distinct functions.

Keywords

chromatin remodelling, SWI/SNF complex, nucleosome structure, INO80, SWR1, CHD1, SMARCA, BAF.

Open Peer Review

Approval Status

	1	2	3	4
version 1 20 Aug 2020				

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Gregory Bowman**, Johns Hopkins University, Baltimore, USA
2. **Craig L. Peterson**, University of Massachusetts Medical School, Worcester, USA
3. **Blaine Bartholomew**, The University of Texas MD Anderson Cancer Center, Science Park, Smithville, USA
4. **Alexander Brehm**, University of Marburg, Marburg, Germany

Any comments on the article can be found at the end of the article.

Corresponding author: Tom Owen-Hughes (t.a.owenhughes@dundee.ac.uk)

Author roles: Sundaramoorthy R: Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; Owen-Hughes T: Conceptualization, Funding Acquisition, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Medical Research Council research grant MR/S021647/1.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2020 Sundaramoorthy R and Owen-Hughes T. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Sundaramoorthy R and Owen-Hughes T. **Chromatin remodelling comes into focus [version 1; peer review: 4 approved]** F1000Research 2020, 9(Faculty Rev):1011 <https://doi.org/10.12688/f1000research.21933.1>

First published: 20 Aug 2020, 9(Faculty Rev):1011 <https://doi.org/10.12688/f1000research.21933.1>

Introduction

In addition to packaging DNA within nuclei, chromatin provides a means of segmenting the genome into distinct chromatin states that ensure that transcription is regulated correctly in time and space¹. Conversion between chromatin states involves changes to chromatin at different levels, including post-translational modifications to histones and DNA, recruitment of chromatin-binding factors and direct changes to the structure of nucleosomes as a result of the action of chromatin remodelling ATPases.

ATP-dependent chromatin remodelling enzymes are molecular machines that act to reconfigure nucleosomes so as to enable gene regulation in response to developmental and environmental signals. Interest in this fundamental process has heightened with the finding that many subunits of these complexes are mutated at high frequencies in cancers and neurological disorders.

The involvement of chromatin remodelling ATPases in this process was first evident from the finding that mutations in the genes that encode components of these complexes affect processes such as mating type switching (SWI) and sucrose fermentation (SNF) in budding yeast². Many of the genes identified in these screens were found to associate as multi-subunit SWI/SNF complexes that had the ability to reconfigure chromatin³. Subsequently, budding yeast were found to encode a related complex, remodels the structure of chromatin (RSC), in which many (but not all) subunits are paralogs of those found in SWI/SNF⁴. Multicellular organisms also encode related complexes. In *Drosophila*, the Brahma complex falls within the trithorax group of developmental regulators^{5–7}. In mammalian cells, three forms of SWI/SNF-related complexes have been identified^{6–8} (Table 1). Homozygous loss of genes encoding most subunits results in early developmental defects; in humans, components of the complexes are frequently mutated in cancers⁹. More recently, it has emerged that mutations to complex

components are also detected in neurological disorders¹⁰. Intriguingly, different subunits are found to be mutated both in cancers of different tissues and in neurological disorders.

Biochemical characterisation of the complexes indicated that they can disrupt nucleosome structure^{3,11}. Consistent with this, the complexes are linked to the maintenance of accessible chromatin structure at promoters in yeast and at enhancers in mammalian cells^{12–18}. The complexes contain a catalytic subunit that is related to ancient ATP-dependent DNA translocases and that acts to drive DNA across the surface of nucleosomes. These specialised ATPase domains are found in an extended family of some 20 yeast and 40 human proteins that regulate DNA–protein interactions¹⁹. In the context of SWI/SNF complexes, ATPase subunits act in the repositioning, destabilisation and dissociation of histones from DNA²⁰. Until recently, a structural framework on which the mechanism of action can be built has been lacking.

Here, we summarise recent insights into the structure of ATP-dependent remodelling enzymes. These show that remodelling enzymes share an ATPase module that acts on DNA within nucleosomes and that this motor domain is tuned to different purposes within distinct multi-subunit complexes.

Snf2 ATPases: a motor domain for nucleosome reorganisation

The ATPase domains found in the yeast Snf2 protein are also present in an extended family of chromatin remodelling enzymes. Crystal structures of Rad54 and Chd1 proteins illustrated that each domain is made up of folds related to those found in bacterial RecA^{21–23}. More recently, cryogenic electron microscopy (cryo-EM) has been used to obtain structures of yeast Snf2, human (SNF2H) and yeast ISWI (imitation SWI) proteins and yeast Chd1 proteins in complex with nucleosomes^{24–29} (Figure 1). These studies show that each enzyme is capable of binding to

Table 1.

Yeast SWISNF	Yeast RSC	Human BAF	Human PBAF	
Snf2/Swi2	Sth1	SMARCA2/SAMRCA4	SMARCA4	SMARCA4
Swi1/Adr6	Rsc9	ARID1A /ARID1B	ARID2	GLTSCR1 / BICRAL1
Swi3	Rsc8	SMARCC1/SMARCC2	SMARCC1/SMARCC2	SMARCC1
Snf12/Swp73	Rsc6	SMARCD1/D2/D3	SMARCD1/D2/D3	SMARCD1
Snf5	Sfh1	SMARCB1	SMARCB1	
	Rsc2/4/58		PBRM /Brd7	BRD9
Swp82	Rsc7	DPF1/2/3	PHF10	DPF3
Arp7/9	Arp7/9	ACTL6A/β-Actin	ACTL6A/β-Actin	ACTL6A/β-Actin
	HTL1			
	Rsc3/30			
		SMARCE1	SMARCE1

Figure 1. Structure of Snf2-related enzymes SNF2H, Snf2 and Chd1 bound to nucleosome. (A) Structural model of *Saccharomyces cerevisiae* Snf2 ATPase domain fragment bound to a nucleosome at super helical location 2 (SHL $\pm$ 2) (PDB ID 5XOY)²⁵. The Snf2-related ATPase lobe1 and 2 are coloured in marine and blue spheres. The nucleosomal DNA is shown in surface representation, the backbone phosphate atom is highlighted in red sphere, and the dyad (SHL 0) of the nucleosome is marked. The histones are shown in black surface representation. The histone H4 tail that interacts with the lobe2 of ATPase domain is coloured green. (B) Structural model of human ISWI remodeller SNF2H bound to nucleosome (PDB ID 6NE3)²⁹. The nucleosome and the ATPase lobes are presented in the same colours as in frame A. (C) Structural model of Chd1 bound to nucleosomes. The structure of Chd1–nucleosome complex resolved at 4.5-Å resolution using cryogenic electron microscopy (PDB ID 6FTX)²⁸. The Chd1 ATPase shown in spheres is characterised by N-terminal tandem chromodomains and a C-terminal SANT-SLIDE-containing DNA-binding domain (DNABD). The Chd1 DNA-binding domain coloured in slate was found to be located at the edge of the nucleosome in the boundary between nucleosomal and linker DNA. The Chd1 ATPase lobes drawn in marine and blue are bound at the SHL 2 location distal to the linker DNA, and the chromodomains drawn in yellow at the SHL 1 location. Similar to the Snf2 and Snf2h, the histone H4 tail interacts with the ATPase lobe2. The dyad of the nucleosome is marked. Two turns of nucleosomal DNA prised from the surface of the histone octamer upon Chd1 binding.

nucleosomes two helical DNA turns away from their centres. Each of these enzymes engages with nucleosomes predominantly through contacts with DNA, and there are relatively few contacts with the histone components of nucleosomes. One exception observed in each structure is that the N-terminal region of histone H4 contacts the second ATPase domain. This region on the H4 tail is required for full activity of the SNF2H, Chd1 and Snf2 proteins and may act to ensure that full activity is reached only when the ATPase domains are correctly docked on nucleosomes. Higher-resolution studies provide enough detail to detect changes in the positioning of individual DNA bases in different nucleotide-bound states. Remarkably, binding in the presence of ADP results in the propagation of a distortion to DNA, predominantly on one strand, across some 55 base pairs of the octamer surface²⁴. As distinct distortions to DNA are

observed in different nucleotide-bound states, it is possible to envision how co-ordinated small movements could drive DNA across the nucleosome surface^{12,13}.

ATPase subunits are found within larger proteins. In the case of Chd1, chromodomains and a DNA-binding domain adjacent to the ATPase domain contribute to the nucleosome-bound state (Figure 1C) and are sufficient to generate an enzyme active in chromatin remodelling. This does raise the question of why many remodelling ATPases are found as components of much larger multi-subunit complexes.

The INO80 and SWR1 enzymes

A first close-up view of multi-subunit remodelling ATPases came from the structural characterisation of the INO80 and SWR1

complexes^{31–33} (Figure 2). Surprisingly, the ATPase domains of these complexes engage with nucleosomes at different

locations. The Ino80 ATPase domains interact with DNA at the edge of the nucleosomes causing unwrapping of outer turns of

Figure 2. Structure of INO80 and SWR1 complexes. (A) Schematic showing the signature ATPase subunit of the Ino80 and Swr1 remodelers. The ATPase is characterised by a split ATPase domain with a large insert region that associates with RuvB proteins and an N-terminal HSA domain that associates with ARP proteins. (B) Structural model of INO80–nucleosome complex resolved at 4.3-Å resolution using cryogenic electron microscopy (cryo-EM) (PDB ID 6FML)³². The subunits that form the INO80–nucleosome complex are labelled. The RuvB hexamer is shown in grey, and the Ino80 insert region that threads through RuvB hexamer in yellow. The nucleosome is shown in surface representation, and the DNA backbone phosphate atoms are highlighted as red spheres. The dyad of the nucleosome SHL 0 is labelled. The Ino80 HSA-Arp8-Actin-Arp4 module (PDB ID 5NBN)³⁵ is docked into EM density observed in this region³³ (coloured light blue). From this location, the Arp8-Actin-Arp4 module potentially interacts with nucleosomal linker DNA and these subunits do indeed cross-link with DNA in this region³⁹. Arp5 is shown in orange, Ies6 in pink and Ies6 purple. Ino80 ATPase lobe1 and 2 are coloured in marine and dark blue respectively. (C) Structural model of SWR1–nucleosome complex resolved at 3.6-Å resolution using cryo-EM (PDB ID 6FML)³¹. The RuvB hexamer is shown in light grey, and the Swr1 insert region that threads through RuvB hexamer in yellow sphere. This extended helical region protrudes through the RuvB hexamer and makes contact with the Arp6 subunit (orange). The Swc6 and Swc2 subunits are coloured pink and purple. The nucleosome and ATPase domains are represented as in frame B.

DNA, whereas in Swr1 they interact at the off-centre site also observed for Snf2 and Chd1. Despite these differences, the Swr1 and Ino80 proteins share distinctive large insertion regions separating the two ATPase domains. In both cases, the insert regions are intertwined with a hetero-hexamer of RuvB proteins, a second type of conserved ATPase belonging to the AAA+ group that are found in both INO80 and SWR1 complexes. The Ino80 and Swr1 insert domains appear to represent a specific adaption of the Snf2-related ATPase domains to function in concert with hetero-hexameric RuvB-related proteins. Mysteriously, ATP hydrolysis by the RuvB proteins is not required for Swr1 histone exchange activity; as a result, they have been proposed to function as a scaffold³¹. The INO80 and SWR1 complexes also contain hetero-dimers of proteins that contain actin folds. These include the Arp5/Ies6 hetero-dimer in the INO80 complex and the Arp6/Swc6 hetero-dimer in the SWR1 complex. These dimers interact on the side of the nucleosome opposite to that occupied by the ATPase domains. In the case of the SWR1 complex, the Arp6/Swc6 interacts with outer turns of nucleosomal DNA at super-helical location (SHL) 6 and histone H2A/H2B dimer surface and generates an unwrapping of one turn of nucleosomal DNA^{14,15}. In the case of Arp5/Ies6, the interaction is internal to the nucleosome. INO80 and SWR1 also contain a second grouping of actin fold proteins: the Arp8/Arp4/actin module. This module has been studied in isolation and is known to interact with a helicase-SANT-associated (HSA) domain present N-terminal to the ATPase domains of many remodelling ATPases where it acts as a linker DNA-binding module that regulates activity¹⁶⁻¹⁹. Density for the Arp8/Arp4/actin module is not observed in one structure³², but weak density is observed in the other³³ (Figure 2B). From this location, the module is well placed to interact with extranucleosomal linker DNA. Consistent with this, this region cross-links to linker DNA and regulates the coupling between ATP-hydrolysis and nucleosome repositioning³⁹.

In summary, while the INO80 and SWR1 complexes share insert regions that mediate interactions with RuvB hetero-hexamers, they engage with nucleosomes in different orientations. The two complexes also have distinct biochemical activities. INO80 is capable of repositioning nucleosomes^{20,21}, whereas SWR1 directs replacement of histone H2A/H2B dimers with the variant histone H2A.Z/H2B dimers⁴². The molecular mechanisms underlying these distinct outcomes remain to be determined.

SWI/SNF complexes: multi-tools for chromatin remodelling

As with the INO80 and SWR1 complexes, cryo-EM has been applied to determine the structures of the yeast SWI/SNF and RSC complexes. Initially, lower-resolution structures indicated that these complexes are globular with a C-shaped central cavity of appropriate dimensions to accommodate a nucleosome²²⁻²⁵. Very recent higher-resolution structures of budding yeast RSC and SWI/SNF and human BAF complexes reveal a distinct organisation in which the Rsc8, Swi3 or human SMARCC proteins form a dimeric hub at the core of the complex⁴⁶⁻⁵⁰ (Figure 3A-D). Yeast Rsc8, Swi3 and the human SMARCC1 and SMARCC2 proteins all contain N-terminal SWIRM domains, Zinc

finger-binding modules, a SANT domain and C-terminal dimerisation domains. Within the complexes, the long dimerisation helices interact in a parallel orientation reminiscent of the Fos/Jun dimerisation module⁵¹ (Figure 3D). Furthermore, the N-terminal SWIRM domains are not arranged symmetrically but adopt distinct conformations (Figure 3D). This asymmetry is likely imposed by the interaction of the two SWIRM domains with tandem repeats within the Sfh1/Snf5/SMARCB1 protein (Figure 3D). The Rsc8/Swi3/SMARCC dimerisation interface is also contacted and likely stabilised by Rsc6/Snf12/SMARCD1 and the N-terminal region of the ATPase subunit Sth1/Snf2/SMARCA4. The core formed by the Rsc8/Swi3/SMARCC dimers and Rsc6/Snf12/SMARCD1 is conserved from yeast to humans and these subunits are present within the three major forms of SWI/SNF-related complex present in humans (Table 1). This core likely serves as a platform from which accessory modules adapt the complex for distinct functions. Within the RSC, SWI/SNF and BAF complexes, five distinct modules are appended to the core (Figure 3):

- i) Adjacent to the Rsc8 dimerisation helices, the armadillo repeats of the Rsc9/Swi1/ARID1A protein are visible. The ARID1A and ARID2 proteins are distinguishing features of the BAF and PBAF forms of mammalian complex positioned some distance from the ATPase domains and likely provide as-yet-uncharacterised complex-specific functionality.
- ii) The tandem bromodomain-containing Rsc2 and Rsc4 proteins are located close to the C-terminus of Rsc8. In mammalian PBAF complex, the polybromo protein PBRM1 is likely to interact at a similar location.
- iii) In between these regions, the HSA domain of Sth1/Snf2/SMARCA4 protrudes *en route* to the motor domains which engage with the nucleosome in an orientation similar to that observed with the isolated Snf2 protein. The HSA domain interacts with the actin fold subunits Arp7 and Arp9 and, together with Rtt102, has been crystallised independently³⁶. This HSA-ARP module links the ATPase domains to the central core based around the Rsc8 dimer. This module is shared in SWI/SNF, RSC, INO80 and SWR1 complexes¹⁶⁻¹⁹.
- iv) The nucleosomal linker DNA extends back towards the Rsc8 core and may make contact with a DNA-binding domain which, at present, is poorly resolved but likely to be composed of the Rsc3 and Rsc30 subunits.
- v) The Sfh1/Snf5/SMARCB1 protein C-terminus extends back forming a distinct nucleosome-binding module located such that it is placed to interact with the acidic patch region on the lateral surface of the nucleosome. This contact is shared in SWI/SNF and RSC complexes that also have the ability to both reposition and evict nucleosomes but is not present in SNF2H and Chd1 complexes which only reposition nucleosomes. Furthermore, mutation of the C-terminal region of Sfh1 that interacts with the acidic patch specifically decreases the ability of RSC complexes to evict nucleosomes

Figure 3. Structure of the RSC and human SWI/SNF (BAF) complex bound to a nucleosome. (A) The structure of the RSC–nucleosome complex (PDB ID 6K15, 6KW3, 6KW4)⁴⁶. The structure can be considered a core or hub module made up principally of the Rsc8 and Rsc6 subunits which are coloured teal. From this, additional modules with distinct functionality extend in different directions: the armadillo repeat-containing Rsc9 is shown in orange; Sth1 which contacts the nucleosome lateral surface, the pin module, is shown in marine; the two tandem bromodomain-containing subunits Rsc2 and Rsc4, together with Rsc58, are coloured pink; the main ATPase Sth1 is coloured red; and Arp7–Rtt102–Arp9 are coloured yellow. The bound nucleosome is shown in similar representation as in Figure 1 and Figure 2. (B) The structure of the SWI/SNF chromatin remodelling complex PDB 6UXW⁴⁸. Homologous subunits (Table 1) are coloured similarly to those in RSC. The Swi3, Snf12 and Snf6 constituents of the core module are coloured in teal, the armadillo repeat-containing Swi1 protein in orange, the Arp module in yellow, the Snf5 pin module in marine, the Snf2 ATPase in red and the nucleosome as in frame A. (C) The structure of Arid1A-containing human BAF complex (PDB 6LTH, 6LTJ)⁴⁷. The complex has a geometry similar to that of the yeast RSC and SWI/SNF complexes. The core of the BAF complex is composed of the SMARCC2 dimer, the HMG domain-containing SMARCE1 protein and the SWIB domain-containing SMARCD1 protein and is shown in teal. The SWIRM domain of the SMARCC2 dimer mediates interaction with the tandem RPT domain of SMARCB1 protein shown in marine. The nucleosome acidic patch is contacted by the SMACB1 C-terminal region, and the pin module is similar to that observed in the RSC and SWI/SNF–nucleosome complexes. SMARCA4 ATPase is shown in red bound to the nucleosome at SHL 2 location and the N-terminal region of ATPase and the pre-HSA domain anchored onto the core of the complex. The Arid1A armadillo repeat domain drawn in orange occupies the central cavity formed by the L-shaped SMARCC2 dimer. The ARP module drawn in yellow surface constitutes the Act16A/β-actin dimer associated with the ATPase HSA domain and forms a bridge between the SMARCA4 ATPase and the core module. (D) Expanded view of SMARCC dimer coloured cyan and dull blue. SMARCC2 has N-terminal SWIRM and SANT domains. SMARCB1 tandem repeat domains (RPT1 and RPT2), shown in blue and marine, interact with each of the SWIRM domains from the dimer of SMARCC2. Similarly, the SANT domain clamps the N-terminal region of ATPase SMARCA4 shown in cartoon representation and coloured in red.

with little effect on ATPase activity⁴⁶. This contact is conserved in human forms of SWI/SNF complex⁴⁷ and mutations in this region observed in patients with

Coffin–Siris syndrome also affect the ability of complexes to reconfigure nucleosomes⁵². It is possible that contact with this region confers specificity for histone variants⁵³.

- vi) The structures of nucleosome-bound complexes are in all cases relatively low and combined with higher-resolution structures of nucleosome-free complexes. This means that it is not possible to determine whether detail observed in the higher-resolution structures of the Snf2 ATPase fragment bound to a nucleosome, including the location of the histone H4 tail, hold true in the context of nucleosome-bound complexes.

Overall, the structure can be considered a central hub from which modules representing sites of association for ARID-containing proteins, multi-bromodomain-containing subunits, the ATPase domains, a linker DNA-binding domain and a module that interacts with the lateral surface of the nucleosome are projected in different directions (Figure 3A–C). The need to accommodate these modules, each likely to be associated with distinct functionalities, goes some way to explaining the larger size of SWI/SNF complexes.

Mutations to SWI/SNF components in human disease

The major features of the RSC nucleosome structure are conserved in the yeast SWI/SNF complex and many of the subunits have equivalents in human forms of SWI/SNF complex (Table 1). A major difference in humans is that there are two genes most similar to Rsc8 in humans, *SMARCC1* and *SMARCC2*. These can be present as either hetero- or homo-dimers within SWI/SNF complexes, but the published structure contains only *SMARCC2*.

Many features of the modular organisation of the complex are consistent with previous observations. For example, SWI/SNF-related complexes remain substantially intact following loss of most subunits^{7,26–28}. This is consistent with the idea that individual modules are largely independent. The exception is the core of the complex which, through acting as a scaffold

for association of many components, plays a more significant role in complex integrity. Consistent with this, perturbing *SMARCC1* and *SMARCC2* levels results in substantial degradation of SWI/SNF complexes^{7,29,30}. Similarly, loss of all *SMARCD* isoforms (equivalent to Rsc6) results in disruption of core complexes and recovery of a residual core module⁷. Subunits that are not assembled correctly are subject to ubiquitinylation and proteasome-mediated degradation^{29,30}. Differences in the potency of this surveillance system may explain some differences in the effects of deleting subunits in specific cell types. For example, the peripheral association of the Sfh1/Snf5/SMARCB1 subunits would suggest that these are not required for complex integrity. Consistent with this, SWI/SNF complexes remain largely intact following deletion of the Snf5 subunit with only Swp82 and Taf14 dissociating^{26,27}. Deletion of the human homolog *SMARCB1* does not severely compromise complex integrity in HEK293T cells⁷ but does in rhabdoid cell lines¹⁷. Thus, subtle effects on subunit associations may have different effects on complex integrity in different tissue types. The partial or complete dissociation of complexes following loss of subunits is, of course, relevant to diseases in which these subunits are lost. However, it does not necessarily inform on the pathway by which complexes are assembled which may be distinct. To characterise how complexes assemble requires study of the order in which nascently translated peptides associate, as observed for assembly of SAGA complexes⁵⁹.

It is estimated that 20% of all human cancers contain a mutation to at least one subunit of one form of SWI/SNF complex⁹. However, the ARID1A, PBRM1 and SMARCA4 subunits are mutated at highest frequencies and these are mutated at much higher rates in tumours of some tissues in comparison with others (Table 2). It is notable that the *SMARCC1* and *SMARCC2* subunits that form the core of the complex and are essential for its integrity⁵⁸ are mutated at substantially lower rates (Table 2 and Figure 4A). It is possible that mutations that severely compromise

Table 2.

Gene ID	Module ¹	Missense-Mutation	Truncating-mutations ¹	Mutations/bp ²	Predicted oncogenic ³
Arid1A	ARID	1110	2231	0.487381473	231
ARID2	ARID	878	648	0.277202543	74
Arid1B		960	334	0.192903995	24
SAMRAC4	ATPASE	1306	335	0.332119004	32
SMARCA2	ATPASE	489	74	0.11802935	1
SMARCC1	CORE	257	61	0.095927602	NA
SMARCC2	CORE	337	93	0.118066996	NA
SMARCD1	CORE	160	42	0.130744337	NA
SMARCD2	CORE	119	30	0.093534212	NA
SMARCD3	CORE	103	16	0.082125604	NA
SMARCE1	CORE	95	24	0.096512571	NA

Gene ID	Module ¹	Missense-Mutation	Truncating-mutations ¹	Mutations/bp ²	Predicted oncogenic ³
SMARCB1	PIN	233	97	0.285714286	15
ACTL6A	ARP	105	31	0.105672106	NA
ACTB	ARP	275	23	0.264888889	NA
PBRM1	BROMO	661	721	0.272745214	35
DPF1		99	31	0.114035088	NA
DPF2		136	35	0.145780051	NA
DPF3		139	31	0.149911817	NA
PHF10		100	54	0.103078983	NA
BRD7		147	91	0.121863799	NA

¹This indicates the module the protein product of gene is assigned to in [Figure 4](#)

²This is the sum of missense and truncating mutations identified from cBioPortal non-redundant studies divided by the coding sequence length in base pairs.

³This indicates number of mutations associated with annotation indicating a potential oncogenic function.

A Distribution of mutations from tumor samples across modules of human BAF complex

B Distribution of mis sense mutations in the SMARCA4 protein

Figure 4. Depiction of different modules of BAF/PBAF complex and mutation rates. (A) A schematic representation of the major modules of human BAF and PBAF remodelling enzymes. Placement of modules is based on structures shown in [Figure 3](#); the subunits included in each module are listed in [Table 2](#). Note that the bromo module is not present in BAF forms of complex. Subunits are coloured by mutation frequency which is also represented as a graph. Mutation frequency is the sum of truncating and missense mutations in the coding regions of genes encoding each subunit and recovered from cBioportal⁶¹ divided by the coding length in base pairs. Truncating mutations and mutations annotated as likely to be oncogenic are more highly enriched in peripheral modules and depleted from the core region ([Table 2](#)). **(B)** Sites of missense mutations within SMARCA4 (red) are shown in black. Mutations are greatly enriched in the ATPase domains in comparison with the HSA domain that interacts with ARP proteins or the N-terminal region which is folded into the core region. Mutated sites obtained from non-redundant studies cohort at cBioportal⁶¹.

complex integrity are disfavoured. Instead, mutations to a subset of subunits that affect one aspect of SWI/SNF functionality are those mutated at high rates (Table 2 and Figure 4A). The ARID1A component which shares armadillo repeats with Rsc9 and Swi1 represents a discrete region adjacent to the core of the complex which remains intact following loss of ARID1A^{28,31}. Similarly, the PBRM1 component contains six bromodomains and is likely related to the Rsc2, Rsc4 and Rsc58 subunits which occupy a location at one extremity of the core region. Deletion of PBRM1 does not affect association of other subunits^{7,28}. Mutations to SMARCB1 which is related to the yeast *SFH1* and *SNF5* genes are relatively rare but debilitating as they drive malignant rhabdoid tumours⁶⁰. The relatively low frequency of SMARCB1 mutations may be related to the fact it has no obvious partially redundant paralogs and is present in two forms of human SWI/SNF-related complex. SMARCA4 is mutated at relatively high frequency, but unlike other subunits, these mutations are predominantly missense mutations clustering to the ATPase domains and appear to act in a dominant fashion¹⁶. In contrast, the high frequency of mutations in ATPase domains, the HSA and N-terminal region of SMARCA4 is mutated at relatively low frequency (Figure 4B). As a result, the distribution of mutation within the SMARCA4 module re-enforces the notion that loss of specific aspects of the function of the complexes—in

this case, ATPase activity—drives cancer, rather than mutations that affect the core region and destabilise the entire complex.

This first wave of structures provides a first insight into the layout of different classes of chromatin remodelling enzyme. Though spectacular, substantial proportions of many subunits are not defined, meaning that it is not possible to assign functions to the bromodomain and armadillo repeat-containing subunits that define different forms of complex. In addition, the current structures represent snapshots of moving machines. To build a complete picture of their function will require views of different stages of the reactions they drive. It will also be critical to determine where and how their activity is regulated by contacts with interaction partners. Nonetheless, insight into the complexes is proceeding at a dramatic pace and provides a structural framework encompassing many of the subunits and domains present.

Abbreviations

HSA, helicase-SANT-associated; Ies, inositol eighty subunit; INO80, INOitol requiring; RSC, remodels the structure of chromatin; SMARCC, SWI/SNF-related matrix-associated actin-dependent regulator of chromatin; SWI/SNF, mating type SWItching/sucrose non-fermenting; SWR1, SWI-related 1; cryo-EM, cryogenic electron microscopy.

References

- Filion GJ, van Bommel JG, Braunschweig U, et al.: **Systematic protein location mapping reveals five principal chromatin types in *Drosophila* cells.** *Cell*. 2010; **143**(2): 212–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Winston F, Carlson M: **Yeast SNF/SWI transcriptional activators and the SPT/SIN chromatin connection.** *Trends Genet*. 1992; **8**(11): 387–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Côté J, Quinn J, Workman JL, et al.: **Stimulation of GAL4 derivative binding to nucleosomal DNA by the yeast SWI/SNF complex.** *Science*. 1994; **265**(5168): 53–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cairns BR, Henry NL, Kornberg RD: **TFG/TAF30/ANC1, a component of the yeast SWI/SNF complex that is similar to the leukemogenic proteins ENL and AF-9.** *Mol Cell Biol*. 1996; **16**(7): 3308–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Tamkun JW, Deuring R, Scott MP, et al.: **brahma: A regulator of *Drosophila* homeotic genes structurally related to the yeast transcriptional activator SNF2SWI2.** *Cell*. 1992; **68**(3): 561–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wang W, Xue Y, Zhou S, et al.: **Diversity and specialization of mammalian SWI/SNF complexes.** *Genes Dev*. 1996; **10**(17): 2117–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mashtalir N, D'Avino AR, Michel BC, et al.: **Modular Organization and Assembly of SWI/SNF Family Chromatin Remodeling Complexes.** *Cell*. 2018; **175**(5): 1272–1288.e20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Alpsoy A, Dykhuizen EC: **Glioma tumor suppressor candidate region gene 1 (GLTSCR1) and its paralog GLTSCR1-like form SWI/SNF chromatin remodeling subcomplexes.** *J Biol Chem*. 2018; **293**(11): 3892–903.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Kadoch C, Crabtree GR: **Mammalian SWI/SNF chromatin remodeling complexes and cancer: Mechanistic insights gained from human genomics.** *Sci Adv*. 2015; **1**(5): e1500447.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sokpor G, Xie Y, Rosenbusch J, et al.: **Chromatin Remodeling BAF (SWI/SNF) Complexes in Neural Development and Disorders.** *Front Mol Neurosci*. 2017; **10**: 243.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Imbalzano AN, Kwon H, Green MR, et al.: **Facilitated binding of TATA-binding protein to nucleosomal DNA.** *Nature*. 1994; **370**(6489): 481–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ocampo J, Chereji RV, Eriksson PR, et al.: **The ISW1 and CHD1 ATP-dependent chromatin remodelers compete to set nucleosome spacing in vivo.** *Nucleic Acids Res*. 2016; **44**(10): 4625–35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kelso TWR, Porter DK, Amaral ML, et al.: **Chromatin accessibility underlies synthetic lethality of SWI/SNF subunits in ARID1A-mutant cancers.** *eLife*. 2017; **6**: e30506.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Brahma S, Henikoff S: **RSC-Associated Subnucleosomes Define MNase-Sensitive Promoters in Yeast.** *Mol Cell*. 2019; **73**(2): 238–249.e3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Kubik S, O'Duibhir E, de Jonge WJ, et al.: **Sequence-Directed Action of RSC Remodeler and General Regulatory Factors Modulates +1 Nucleosome Position to Facilitate Transcription.** *Mol Cell*. 2018; **71**(1): 89–102.e5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
- Hodges HC, Stanton BZ, Cermakova K, et al.: **Dominant-negative SMARCA4 mutants alter the accessibility landscape of tissue-unrestricted enhancers.** *Nat Struct Mol Biol*. 2018; **25**(1): 61–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Wang X, Lee RS, Alver BH, et al.: **SMARCB1-mediated SWI/SNF complex function is essential for enhancer regulation.** *Nat Genet*. 2017; **49**(2): 289–95.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)

18. Langer LF, Ward JM, Archer TK: **Tumor suppressor SMARCB1 suppresses super-enhancers to govern hESC lineage determination.** *eLife*. 2019; 8: e45672.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
19. Flaus A, Martin DMA, Barton GJ, et al.: **Identification of multiple distinct Snf2 subfamilies with conserved structural motifs.** *Nucleic Acids Res*. 2006; 34(10): 2887–905.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
20. Clapier CR, Iwasa J, Cairns BR, et al.: **Mechanisms of action and regulation of ATP-dependent chromatin-remodelling complexes.** *Nat Rev Mol Cell Biol*. 2017; 18(7): 407–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
21. Hauk G, McKnight JN, Nodelman IM, et al.: **The chromodomains of the Chd1 chromatin remodeler regulate DNA access to the ATPase motor.** *Mol Cell*. 2010; 39(5): 711–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
22. Thomä NH, Czyzewski BK, Alexeev AA, et al.: **Structure of the SWI2/SNF2 chromatin-remodeling domain of eukaryotic Rad54.** *Nat Struct Mol Biol*. 2005; 12(4): 350–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Dürr H, Körner C, Müller M, et al.: **X-ray structures of the *Sulfolobus solfataricus* SWI2/SNF2 ATPase core and its complex with DNA.** *Cell*. 2005; 121(3): 363–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
24. Li M, Xia X, Tian Y, et al.: **Mechanism of DNA translocation underlying chromatin remodelling by Snf2.** *Nature*. 2019; 567(7748): 409–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
25. Liu X, Li M, Xia X, et al.: **Mechanism of chromatin remodelling revealed by the Snf2-nucleosome structure.** *Nature*. 2017; 544(7651): 440–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Farnung L, Vos SM, Wigge C, et al.: **Nucleosome-Chd1 structure and implications for chromatin remodelling.** *Nature*. 2017; 550(7677): 539–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
27. Sundaramoorthy R, Hughes AL, Singh V, et al.: **Structural reorganization of the chromatin remodeling enzyme Chd1 upon engagement with nucleosomes.** *eLife*. 2017; 6: e22510.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Sundaramoorthy R, Hughes AL, El-Mkami H, et al.: **Structure of the chromatin remodelling enzyme Chd1 bound to a ubiquitinated nucleosome.** *eLife*. 2018; 7: e35720.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Armache JP, Gamarra N, Johnson SL, et al.: **Cryo-EM structures of remodeler-nucleosome intermediates suggest allosteric control through the nucleosome.** *eLife*. 2019; 8: e46057.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
30. Bowman GD, Deindl S: **Remodeling the genome with DNA twists.** *Science*. 2019; 366(6461): 35–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
31. Willhoft O, Ghoneim M, Lin CL, et al.: **Structure and dynamics of the yeast SWR1-nucleosome complex.** *Science*. 2018; 362(6411): eaat7716.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
32. Ayala R, Willhoft O, Aramayo RJ, et al.: **Structure and regulation of the human INO80-nucleosome complex.** *Nature*. 2018; 556(7701): 391–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
33. Eustermann S, Schall K, Kostrewa D, et al.: **Structural basis for ATP-dependent chromatin remodelling by the INO80 complex.** *Nature*. 2018; 556(7701): 386–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
34. Zhou CY, Stoddard CI, Johnston JB, et al.: **Regulation of Rvb1/Rvb2 by a domain within the INO80 Chromatin Remodeling Complex Implicates the Yeast Rvbs as Protein Assembly Chaperones.** *Cell Rep*. 2017; 19(10): 2033–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. Knoll KR, Eustermann S, Niebauer V, et al.: **The nuclear actin-containing Arp8 module is a linker DNA sensor driving INO80 chromatin remodeling.** *Nat Struct Mol Biol*. 2018; 25(9): 823–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
36. Szerlong H, Hinata K, Viswanathan R, et al.: **The HSA domain binds nuclear actin-related proteins to regulate chromatin-remodeling ATPases.** *Nat Struct Mol Biol*. 2008; 15(5): 469–76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Schubert HL, Wittmeyer J, Kasten MM, et al.: **Structure of an actin-related subcomplex of the SWI/SNF chromatin remodeler.** *Proc Natl Acad Sci U S A*. 2013; 110(9): 3345–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Cao T, Sun L, Jiang Y, et al.: **Crystal structure of a nuclear actin ternary complex.** *Proc Natl Acad Sci U S A*. 2016; 113(32): 8985–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Brahma S, Ngubo M, Paul S, et al.: **The Arp8 and Arp4 module acts as a DNA sensor controlling INO80 chromatin remodeling.** *Nat Commun*. 2018; 9(1): 3309.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
40. Willhoft O, McCormack EA, Aramayo RJ, et al.: **Crosstalk within a functional INO80 complex dimer regulates nucleosome sliding.** *eLife*. 2017; 6: e25782.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Krientein N, Wal M, Watanabe S, et al.: **Genomic Nucleosome Organization Reconstituted with Pure Proteins.** *Cell*. 2016; 167(3): 709–721.e12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Mizuguchi G, Shen X, Landry J, et al.: **ATP-driven exchange of histone H2AZ variant catalyzed by SWR1 chromatin remodeling complex.** *Science*. 2004; 303(5656): 343–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
43. Smith CL, Horowitz-Scherer R, Flanagan JF, et al.: **Structural analysis of the yeast SWI/SNF chromatin remodeling complex.** *Nat Struct Biol*. 2003; 10(2): 141–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
44. Dechassa ML, Zhang B, Horowitz-Scherer R, et al.: **Architecture of the SWI/SNF-nucleosome complex.** *Mol Cell Biol*. 2008; 28(19): 6010–21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
45. Chaban Y, Ezeokonkwo C, Chung WH, et al.: **Structure of a RSC-nucleosome complex and insights into chromatin remodeling.** *Nat Struct Mol Biol*. 2008; 15(12): 1272–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. Ye Y, Wu H, Chen K, et al.: **Structure of the RSC complex bound to the nucleosome.** *Science*. 2019; 366(6467): 838–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
47. He S, Wu Z, Tian Y, et al.: **Structure of nucleosome-bound human BAF complex.** *Science*. 2020; 367(6480): 875–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
48. Han Y, Reyes AA, Malik S, et al.: **Cryo-EM structure of SWI/SNF complex bound to a nucleosome.** *Nature*. 2020; 579(7799): 452–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
49. Patel AB, Moore CM, Greber BJ, et al.: **Architecture of the chromatin remodeler RSC and insights into its nucleosome engagement.** *eLife*. 2019; 8: e54449.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
50. Wagner FR, Dienemann C, Wang H, et al.: **Structure of SWI/SNF chromatin remodeler RSC bound to a nucleosome.** *Nature*. 2020; 579(7799): 448–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
51. van Dam H, Castellazzi M: **Distinct roles of Jun: Fos and Jun: ATF dimers in oncogenesis.** *Oncogene*. 2001; 20(19): 2453–64.
[PubMed Abstract](#) | [Publisher Full Text](#)
52. Valencia AM, Collings CK, Dao HT, et al.: **Recurrent SMARCB1 Mutations Reveal a Nucleosome Acidic Patch Interaction Site That Potentiates mSWI/SNF Complex Chromatin Remodeling.** *Cell*. 2019; 179(6): 1342–1356.e23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
53. Cakiroglu A, Clapier CR, Ehrensberger AH, et al.: **Genome-wide reconstitution of chromatin transactions reveals that RSC preferentially disrupts H2AZ-containing nucleosomes.** *Genome Res*. 2019; 29(6): 988–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
54. Dutta A, Sardiu M, Gogol M, et al.: **Composition and Function of Mutant Swi/Snf Complexes.** *Cell Rep*. 2017; 18(9): 2124–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
55. Sen P, Luo J, Hada A, et al.: **Loss of Snf5 Induces Formation of an Aberrant SWI/SNF Complex.** *Cell Rep*. 2017; 18(9): 2135–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. Schick S, Rendeiro AF, Runggatscher K, et al.: **Systematic characterization of BAF mutations provides insights into intracomplex synthetic lethality in human cancers.** *Nat Genet*. 2019; 51(9): 1399–410.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
57. Chen J, Archer TK: **Regulating SWI/SNF subunit levels via protein-protein interactions and proteasomal degradation: BAF155 and BAF170 limit expression of BAF57.** *Mol Cell Biol*. 2005; 25(20): 9016–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
58. Narayanan R, Pirouz M, Kerimoglu C, et al.: **Loss of BAF (mSWI/SNF) Complexes Causes Global Transcriptional and Chromatin State Changes in**

Forebrain Development. *Cell Rep.* 2015; **13**(9): 1842–54.
[PubMed Abstract](#) | [Publisher Full Text](#)

59. Kamenova I, Mukherjee P, Conic S, *et al.*: **Co-translational assembly of mammalian nuclear multisubunit complexes.** *Nat Commun.* 2019; **10**(1): 1740.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) |
[Faculty Opinions Recommendation](#)

60. Versteeg I, Sévenet N, Lange J, *et al.*: **Truncating mutations of hSNF5/IN11 in aggressive paediatric cancer.** *Nature.* 1998; **394**(6689): 203–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
61. Cerami E, Gao J, Dogrusoz U, *et al.*: **The cBio cancer genomics portal: An open platform for exploring multidimensional cancer genomics data.** *Cancer Discov.* 2012; **2**(5): 401–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Alexander Brehm**

Institute for Molecular Biology and Tumour Research, University of Marburg, Marburg, Germany

Competing Interests: No competing interests were disclosed.

2. **Blaine Bartholomew**

Department of Epigenetics and Molecular Carcinogenesis, The University of Texas MD Anderson Cancer Center, Science Park, Smithville, TX, USA

Competing Interests: No competing interests were disclosed.

3. **Craig L. Peterson**

Program in Molecular Medicine, University of Massachusetts Medical School, Worcester, MA, USA

Competing Interests: No competing interests were disclosed.

4. **Gregory Bowman**

Department of Biophysics, Johns Hopkins University, Baltimore, MD, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research