STUDY PROTOCOL

Protocol for a controlled, randomized, blind, clinical trial to assess the effects of transcranial direct current stimulation associated with balance training using games in the postural balance of elderly people [version 1; peer review: 2 approved]

Andre Issao Kunitake , João Carlos Ferrari Corrêa , Klaine Silva Nascimento, Bianca Barioni Cardoso de Oliveira, Natalia Maciel Muniz, Soraia Micaela Silva, Fernanda Ishida Corrêa

Universidade Nove de Julho, Nove de Julho University, São Paulo, São Paulo, 01504-001, Brazil

V1 First published: 21 Aug 2020, 9:1018
<https://doi.org/10.12688/f1000research.25164.1>
 Latest published: 23 Feb 2021, 9:1018
<https://doi.org/10.12688/f1000research.25164.2>

Abstract

Aims: This study aims to evaluate the additional effect of transcranial direct current stimulation (tDCS) on training postural balance with the use of video games in aged.

Methods: This is a blinded, randomized, controlled clinical trial protocol, with elderly people of both genders. Participants will be randomized into three training groups: Group 1 (videogame balance training), group 2 (videogame balance training associated with active tDCS), group 3 (videogame balance training associated with sham tDCS). The training will be carried out twice a week for four weeks, totaling eight sessions, and will be performed with the Nintendo Wii videogame console, using games that stimulate the postural balance associated with tDCS, with anode applied over the left dorsolateral prefrontal cortex and cathode on the contralateral supraorbital region at 2 mA for 20 minutes. The postural balance will be assessed using the Mini Test of the Balance Assessment System and posturography. Evaluations will be carried out before and after eight training sessions and 30 days after the end of treatment.

Discussion: Some studies show favorable results from the use of video games in improving postural balance in the elderly; however, their effect does not remain long-term. TDCS associated with other therapies can potentiate and prolong the effects of these therapies owing to its ability to stimulate neurotrophins important for neurogenesis, facilitating tasks that require attention, and helping to consolidate learning and memory. The effect of the two associated techniques on balance have not yet been tested in this population.

Registration: Brazilian Registry of Clinical Trials ID [U1111-1213-4266](#); registered on 15 October, 2018.

Open Peer Review

Approval Status

	1	2
version 2 (revision) 23 Feb 2021		 view
version 1 21 Aug 2020	 view	 view

1. **Fernando Zanela da Silva Arêas** , Federal University of Espírito Santo (UFES), Vitória, Brazil

2. **Rodrigo Vitória** , Oregon Health and Science University, Portland, USA

Any reports and responses or comments on the article can be found at the end of the article.

Keywords

Elderly, postural balance, transcranial direct current stimulation, video game

Corresponding author: Andre Issao Kunitake (issaokun@gmail.com)

Author roles: **Kunitake AI:** Conceptualization, Methodology, Project Administration, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Corrêa JCF:** Funding Acquisition, Resources, Software; **Nascimento KS:** Investigation, Project Administration, Supervision; **Oliveira BBCd:** Investigation, Project Administration, Supervision; **Muniz NM:** Investigation, Project Administration, Supervision; **Silva SM:** Data Curation, Formal Analysis, Validation; **Corrêa FI:** Conceptualization, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This study is funded and supported by the Higher Education Personnel Improvement Coordination in conjunction with Nove de Julho University (Opinion number 2.962.837), Brazil, São Paulo.

Copyright: © 2020 Kunitake AI *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Kunitake AI, Corrêa JCF, Nascimento KS *et al.* **Protocol for a controlled, randomized, blind, clinical trial to assess the effects of transcranial direct current stimulation associated with balance training using games in the postural balance of elderly people [version 1; peer review: 2 approved]** F1000Research 2020, 9:1018 <https://doi.org/10.12688/f1000research.25164.1>

First published: 21 Aug 2020, 9:1018 <https://doi.org/10.12688/f1000research.25164.1>

Introduction

Aging is a physiological occurrence that has several consequences, including an increased risk of falls^{1,2}. The incidence of falls increases proportionally to age, from 28% to 35% in the elderly over 65 years and from 32% to 42% in those over 75 years. Falls can lead to musculoskeletal injuries, and in more severe cases, they can lead to death^{1,3,4}.

Preventive therapies for the risk of falls can be performed with the use of technological resources, such as videogames. Studies using videogames have shown positive effects in improving the postural control, balance, and functional capacity of the elderly when practiced regularly⁵. They have been widely used by therapists with high acceptance in clinical practice because they are motivating, challenging, and recreational, showing good evidence to improve postural control in the elderly⁶⁻⁹.

Another technique that has aroused interest in clinical practice is transcranial direct current stimulation (tDCS), which consists of low-intensity current, and can be applied to the scalp by means of rubber electrodes, thus promoting changes in the potential of resting membrane, such as depolarization by the anode electrode or hyperpolarization by the cathode; modulating cortical excitability^{10,11}. Anodic tDCS, when applied over

the long term, can stimulate the brain-derived neurotrophic factor (BDNF), which is an important protein for stimulating neuroplasticity, improving attention, consolidating learning, and memory. Thus, its use has aroused interest for the elderly, in order to reduce cognitive and motor declines¹²⁻¹⁵.

However, the protocols for the use of tDCS associated with tasks that involve balance, in order to improve balance are not yet defined, and are being applied in different dosages and locations¹⁶⁻¹⁸, therefore, further studies are necessary to verify the benefit of tDCS in association with other therapies. For that reason, the objective of this study is to assess whether tDCS can enhance and prolong the effects of videogame training on improving postural balance in the elderly. This study hypothesizes that the association of tDCS with virtual reality therapy is more effective in improving balance in the elderly than therapy with videogames alone, and that the addition of tDCS will prolong the effects of balance training.

Methods

Study design

This is a protocol for a randomized, controlled blind, longitudinal clinical trial, as shown in [Figure 1](#). This project will be carried out at University Nove de Julho in São Paulo, Brazil.

TIMEPOINT	STUDY PERIOD										
	Enrolment	Allocation	Post-allocation								Close-out
	$-t_1$	0	1	2	3	4	5	6	7	8	t_9
ENROLMENT:	X										
Eligibility screen	X										
Informed consent	X										
Ethics committee approval and Trial Registration	X										
Allocation		X									
OUTCOMES											
Postural Balance		X									X
Static Postural Balance		X									X
Quality of Life		X									X
Depressive symptoms		X									X
Side effects questionnaire (tDCS)											
Blinding and Satisfaction											
INTERVENTIONS:											
Video Game Training (VGT)			←								→
VGT associated with active tDCS			←								→
VGT associated with sham tDCS			←								→

Figure 1. Study enrollment schedule, outcomes, and interventions. tDCS: transcranial direct current stimulation.

Eligibility criteria

The inclusion criteria are people between 60 and 80 years old, of both genders, able to stand or walk without the aid of auxiliary devices, with reduced balance (score ≤ 25 points) evaluated using the Mini-BESTest¹⁹. Exclusion criteria are cognitive impairment (scores ≤ 14 points) corrected for education, measured by the Mini-Mental State Examination (MMSE)²⁰, presence of lower limb fractures or amputations, neurological diseases, cardiovascular diseases that limit the performance of exercises, untreated acute and chronic respiratory diseases and the presence of pain that limits movement; the presence of contraindications to the use of tDCS.

Consent to participate

The informed consent form will be explained to all participants. The volunteers who participate in this study, must sign the form (supplementary file 1), which guarantees the secrecy and confidentiality of data, free access to the final data, explanations of any kind related to the study and possible compensation for those that suffer from participation in the study.

Ethics approval

The protocol was approved in May 2018 by Human Research Ethics Committee of the University Nove de Julho, São Paulo, Brazil (Opinion number 2.962.837), and Brazilian Clinical Trials Registry (ReBec), number: U1111-1213-4266, in accordance with Resolution 466/12 of the National Health Council of Brazil.

Recruitment and allocation

Participants in this study will be community elders elected according to the criteria established for allocation. They will be randomized into 3 groups:

Group 1: Control Group (Balance training with video game);

Group 2: Active group (balance training with video games associated with active tDCS);

Group 3: Sham Group (Balance training with video games associated with sham tDCS).

A researcher not involved in evaluations or training will be responsible for allocating participants via www.randomization.com.

Sample size

The sample size was obtained using G*Power 3.0.10 software, based on the outcomes from a pilot study. The calculation was carried out by the two-way repeated measures analysis of variance (ANOVA), considering the mean value (before and after training, and 30-day follow-up) for the control groups (25.46) active (27.46) and sham (24.66) and the pooled standard deviation (SD_{pooled}) (2.89), with $\alpha = 0.05$, $\beta = 0.2$ (80% of power) and the effect size of 0.40. A total of 15 individuals were determined to be required for each group (total sample: 45 individuals). Considering possible dropouts and to ensure a sample size that will demonstrate the effect of the intervention,

the sample will be expanded by 25%, resulting in 19 individuals in each group, thus totalling 54 participants.

Outcome assessments

All evaluations will be carried out on three occasions: pre-intervention, after eight treatment sessions and 30 days after the end of the training (follow-up). The training will be held twice a week, for 4 weeks, totalling 8 sessions.

Postural balance. The assessment of postural balance will be performed by the Mini BESTest Scale, which consists of 14 functional tasks, such as sitting and getting up from a chair, standing up, balancing on tiptoes, and on one foot, overcoming walking obstacles, and double activities task²¹.

Performance can range from 0 to 28 points. Each test can be performed for up to three attempts and the best result will be obtained. If adaptations are necessary to accomplish this, a point will be deducted from the maximum score obtained. This is a very reliable instrument, with an intraclass correlation (ICC) of 0.84²².

Static postural balance. Posturography data will be collected using the Wii Balance Board (Nintendo, Kyoto, Japan), which is a validated instrument for posturography evaluation²³. The software for postural assessment is available at <http://www.rehabtools.org/sway.html>.

The evaluation protocol will be performed in two conditions, standing with eyes open and then with eyes closed. These two postures are reliable for measuring body sway, with eyes open with (ICC: 0.77) and with eyes closed (ICC: 0.89)²³.

To standardize the collections, the initial position of the feet on the evaluation platform will be marked and the same measurement repeated in all evaluations. The tests will last 1 minute, the initial 30 seconds will be to establish the patient's suitability and the final 30 seconds will be to collect the posturography data.

Potential confusion factors

Symptoms of depression. The symptoms of depression will be evaluated and classified according to severity, using the Beck Depression Inventory (BDI), a questionnaire composed of 21 questions, which can be self-applied. Scores from 0 to 10 indicate an absence of depression; 11 to 18 indicate mild depression; 19 to 29 indicate moderate depression and 30 to 63 is considered severe depression²⁴. In studies with the elderly, the BDI score shows a degree of reliability $r = 0.78$ ²⁵. Subsequently, it will be assessed whether the emotional state interfered with the results.

Quality of life (QOL). QOL will be measured by the World Health Organization's Quality of Life Instrument, Bref version²⁶, which is composed of 26 questions, scored from 1 to 5, with the worst and the best scores being 26 and 130, respectively. In studies with the elderly, it showed a high degree of reliability

to mediate quality of life in the domain of physical capacity (0.89), psychological well-being (0.95), social relationships (0.81), and the environment (0.93).

The use of medication

The information related to continuous-use medication will be monitored through the application of a questionnaire prepared by the author. Such information will be used to characterize the sample.

Intervention procedures

Balance training with video games. The balance training sessions will be performed using only the video games in group 1 (control), group 2 will be the balance training with the video games associated with the active tDCS and group 3 will be the balance training with the video games associated with the tDCS sham.

Video game training will be carried out with the Nintendo Wii and Wii Fit Plus, placed in a slide projector to enlarge the image. The sequence of games will always be the same, and the participant should play an average of 5 to 7 minutes each game, passing the stage according to their learning.

The sessions will be held twice a week, for four weeks, totalling eight sessions. Each session will last 20 minutes. The selected games are table tilt, penguin slide and ski slalom, which will be performed standing on the Wii Board Balance and require movements that stimulate balance, such as the anterior, posterior and lateral tilt of the body, without moving (Figure 1).

tDCS

tDCS will be performed with a tDCS device, the NeuroConn DC_STIMULATOR PLUS, from Germany, by means of two non-metallic surface electrodes, cathode 35 cm² (5 × 7 cm) and anode 25 cm² (5 × 5 cm), both wrapped in sponges moistened in a saline solution.

The intensity will be 2mA with a 20-second ramp up and down for a 20-minute period of stimulation. The montage will be anodal, unbalanced bilateral bipolar, anode positioned over the dorsolateral prefrontal cortex of the dominant hemisphere (F3 or F4) and the cathode electrode on the supraorbital region contralateral to the anode (Figure 2), according to the criteria of the 10–20 electroencephalogram system²⁷.

Assessment of the potential adverse effects

At the end of each session, a questionnaire on the adverse effects of tDCS will be applied to the participant²⁶ (see Extended data, Assessment of ETCC Adverse Events form).

Blinding

The NeuroConn DC-STIMULATOR PLUS device has settings that allow the selection of the active stimulation or sham mode, by inserting codes. A researcher not involved in the procedures will program the equipment with the code to which the individual will be allocated. The stimulus mode will not be

Figure 2. Positioning the participant on the board for the execution of the therapy with Nintendo Wii associated with active stimulation or sham. Source: author's own, consent received for publication.

perceived by the external (supplementary functioning of the device, therefore, neither the researcher who will apply the intervention nor the individual will know what treatment will be applied (double-blind).

Participants in the group that will only perform training with videogames will not be blind to treatment; however, a researcher will be responsible for the exclusive training of this group and will not be aware of the procedures with the tDCS of the other groups. All assessments will be made by researchers who have not participated in the training of the elderly, and will, therefore, be blind to the intervention. A blinding questionnaire will be applied to guarantee its reliability and satisfaction with the treatment received (supplementary files 3 and 4), without the researcher knowing the training carried out by the participant.

Statistical analysis

All statistical tests will be performed using SPSS (V22, IBM Corporation, New York, USA). Initially, we will perform the test of normality of the sample using the Shapiro-Wilk

test, considering the significance level defined as a value of $\alpha < 0.05$. The parametric data will be expressed as a mean $\pm$ SD (standard deviation), and the nonparametric data as a median (IQR); the categorical data will be described as absolute values and as a percentage of the total sample.

The variables of the data of the Mini-BESTest, Posturography, obtained pre-intervention, post-intervention and 30-day follow-up for the three groups (control, active and sham) will be calculated by the two-way repeated measures analysis of variance (ANOVA) for the parametric data and the Friedman's test for the nonparametric data.

Finally, to verify whether there is a correlation between BESTest values and the symptoms of depression and quality of life, Pearson's R correlation (parametric) or Spearman correlation (nonparametric) for the three groups in three evaluated times will be performed, verifying the degree of correlation (r) and its significance (p).

Trial status

The protocol was approved in May 2018 by the Brazilian Clinical Trials Registry (ReBec) U1111-1213-4266. Submission of the manuscript occurred after this period, with the collection taking place and the completion is expected to happen in December 2020. The outcomes of this study will be subsequently published in a journal of interest.

Discussion

The incidence of falls has become an aggravating problem with the increase in the number of elderly people and life expectancy. The costs of falling are also high; prevention is an effective and inexpensive alternative when compared to any other procedure^{28,29}.

Preventive therapies can be performed with the use of video games, since they can be used to improve postural control and balance in the elderly, helping to prevent falls. Besides, they are well accepted for being motivating and recreational and can be easily performed at home⁷⁻¹⁰.

However, there are several types of training protocols, which can vary from 4 to 20 weeks²². For some elderly people, this long process of therapy can be tiring; however, for the consolidation of learning and memory to occur, repetitive training is necessary. According to²⁹, the learning and memory process is directly related to the number of repetitions performed by the elderly, with a better effect in the long run. In this sense, the simultaneous application of tDCS can be a resource that can assist in this process of consolidating learning, improving attention for its ability to stimulate neuroplasticity. For the elderly, it can be an important tool in the rehabilitation process, as it is known that, with advancing age, they present cognitive and motor decline¹¹⁻¹⁴.

However, one of the limitations of using tDCS for the elderly is that few studies have investigated its effects on this population; some showing positive effects^{16,18,30,31}, while others do not^{17,32,33}. Protocols combining the two resources have already been carried out in children with cerebral palsy³⁴, Parkinson's disease³⁵, spinal cord injury³⁶ and stroke victims^{37,38}. However, studies that associated tDCS and video games in the elderly was not found.

Conclusion

This article presents a randomized, controlled and blind protocol developed to show the effect of the combination of transcranial direct current stimulation (tDCS) associated with training using video games in the elderly. The outcomes obtained will be published and their evidences may contribute for new training alternatives in the elderly.

Data availability

Underlying data

No underlying data are associated with this article.

Extended data

Harvard Dataverse: A PROTOCOL OF A CONTROLLED, RANDOMIZED, BLIND, CLINICAL TRIAL, TO CHECK THE EFFECTS OF TRANSCRANIAL DIRECT CURRENT STIMULATION (tDCS) ASSOCIATED WITH BALANCE TRAINING USING GAMES IN THE POSTURAL BALANCE OF ELDERLY PEOPLE. <https://doi.org/10.7910/DVN/OJBWHJ>

File 'Supplementary File.docx' contains the following extended data:

- Model informed consent form.
- Assessment of ETCC Adverse Events form.
- Blinding questionnaire.
- Satisfaction questionnaire.

Reporting guidelines

Harvard Dataverse: SPIRIT checklist for 'Protocol for a controlled, randomized, blind, clinical trial to assess the effects of transcranial direct current stimulation associated with balance training using games in the postural balance of elderly people'. <https://doi.org/10.7910/DVN/OJBWHJ>

Data are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

Acknowledgments

The authors of this study thank the University Nove de Julho and its collaborators for supporting the present study.

References

1. da Saúde M, Motta LB da: **Envelhecimento e Saúde da Pessoa idosa**. 1st ed. Brasil: Ministerio da Saúde; BRASIL. 2006.
[Reference Source](#)
2. da Saúde M: **Atenção à Saúde da Pessoa Idosa e Envelhecimento**. Brasil. 2010.
3. Horak FB, Henry SM, Shumway-Cook A: **Postural perturbations: new insights for treatment of balance disorders**. *Phys Ther*. 1997; **77**(5): 517–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Ferreira OGL, Maciel SC, Costa SMG, et al.: **Envelhecimento Ativo e Sua Relação Com a Independência Funcional**. *Texto e Context Enferm*. 2012; **21**: 513–8.
[Publisher Full Text](#)
5. Donath L, Rössler R, Faude O: **Effects of Virtual Reality Training (Exergaming) Compared to Alternative Exercise Training and Passive Control on Standing Balance and Functional Mobility in Healthy Community-Dwelling Seniors: A Meta-Analytical Review**. *Sport Med*. 2016; **46**(9): 1293–309.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Bisson E, Contant B, Sveistrup H, et al.: **Functional Balance and Dual-Task Reaction Times in Older Adults Are Improved by Virtual Reality and Biofeedback Training**. *Cyberpsychol Behav*. 2007; **10**(1): 16–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Cho GH, Hwangbo G, Shin HS: **The Effects of Virtual Reality-based Balance Training on Balance of the Elderly**. *J Phys Ther Sci*. 2014; **26**(4): 615–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. Robinson J, Dixon J, Macsween A, et al.: **The effects of exergaming on balance, gait, technology acceptance and flow experience in people with multiple sclerosis: a randomized controlled trial**. *BMC Sports Sci Med Rehabil*. 2015; **7**: 8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. Tsang WWN, Fu ASN: **Virtual reality exercise to improve balance control in older adults at risk of falling**. *Hong Kong Med J*. 2016; **22** Suppl 2: 39–42.
[PubMed Abstract](#)
10. Lefaucheur JP, Antal A, Ayache SS, et al.: **Evidence-based guidelines on the therapeutic use of transcranial direct current stimulation (tDCS)**. *Clin Neurophysiol*. 2017; **128**(1): 56–92.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Nitsche MA, Paulus W: **Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation**. *J Physiol*. 2009; **527** Pt 3(Pt 3): 633–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Cespón J, Rodella C, Rossini PM, et al.: **Anodal transcranial direct current stimulation promotes frontal compensatory mechanisms in healthy elderly subjects**. *Front Aging Neurosci*. 2017; **9**: 420.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Fritsch B, Reis J, Martinowich K, et al.: **Direct current stimulation promotes BDNF-dependent synaptic plasticity: potential implications for motor learning**. *Neuron*. 2010; **66**(2): 198–204.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
14. Nitsche MA, Cohen LG, Wassermann EM, et al.: **Transcranial direct current stimulation: State of the art 2008**. *Brain Stimul*. 2008; **1**(3): 206–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Coffman BA, Trumbo MC, Clark VP: **Enhancement of object detection with transcranial direct current stimulation is associated with increased attention**. *BMC Neurosci*. 2012; **13**: 108.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Manor B, Zhou J, Harrison R, et al.: **Transcranial Direct Current Stimulation May Improve Cognitive-Motor Function in Functionally Limited Older Adults**. *Neurorehabil Neural Repair*. 2018; **32**(9): 788–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Kaminski E, Hoff M, Rjosk V, et al.: **Anodal Transcranial Direct Current Stimulation Does Not Facilitate Dynamic Balance Task Learning in Healthy Old Adults**. *Front Hum Neurosci*. 2017; **11**: 16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Yosephi MH, Ehsani F, Zoghi M, et al.: **Multi-session anodal tDCS enhances the effects of postural training on balance and postural stability in older adults with high fall risk: Primary motor cortex versus cerebellar stimulation**. *Brain Stimul*. 2018; **11**(6): 1239–1250.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Magnani PE, Genovese MB, Porto JM, et al.: **Use of the BESTest and the Mini-BESTest for Fall Risk Prediction in Community-Dwelling Older Adults Between 60 and 102 Years of Age**. *J Geriatr Phys Ther*. 2019; **1**.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. Bertolucci PHF, Brucki SMD, Campacci SR, et al.: **O Mini-Exame do Estado Mental em uma população geral. Impacto da escolaridade**. *Arg Neuropsiquiatr*. 1994; **52**: 1–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Potter K, Brandfass K: **The Mini-Balance Evaluation Systems Test (Mini-BESTest)**. *J Physiother*. 2015; **61**(4): 225.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Anson E, Thompson E, Ma L, et al.: **Reliability and Fall Risk Detection for the BESTest and Mini-BESTest in Older Adults**. *J Geriatr Phys Ther*. 2019; **42**(2): 81–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Clark RA, Bryant AL, Pua Y, et al.: **Validity and reliability of the Nintendo Wii Balance Board for assessment of standing balance**. *Gait Posture*. 2010; **31**(3): 307–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Beck AT, Ward CH, Mendelson M, et al.: **An inventory for measuring depression**. *Arch Gen Psychiatry*. 1961; **4**: 561–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Snyder AG, Stanley MA, Novy DM, et al.: **Measures of depression in older adults with generalized anxiety disorder: a psychometric evaluation**. *Depress Anxiety*. 2000; **11**(3): 114–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Bikson M, Grossman P, Thomas C, et al.: **Safety of transcranial Direct Current Stimulation: Evidence Based Update 2016**. *Brain Stimul*. 2016; **9**(5): 641–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Manor B, Zhou J, Jor'dan A, et al.: **Reduction of Dual-task Costs by Noninvasive Modulation of Prefrontal Activity in Healthy Elders**. *J Cogn Neurosci*. 2016; **28**(2): 275–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Schoene D, Lord SR, Delbaere K, et al.: **A Randomized Controlled Pilot Study of Home-Based Step Training in Older People Using Videogame Technology**. *PLoS One*. 2013; **8**(3): e57734.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Van Ooteghem K, Frank JS, Horak FB: **Practice-related improvements in posture control differ between young and older adults exposed to continuous, variable amplitude oscillations of the support surface**. *Exp Brain Res*. 2009; **199**(2): 185–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Zhou J, Lo OY, Lipsitz LA, et al.: **Transcranial direct current stimulation enhances foot sole somatosensation when standing in older adults**. *Exp Brain Res*. 2018; **236**(3): 795–802.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Ehsani F, Samaei A, Zoghi M, et al.: **The effects of cerebellar transcranial direct current stimulation on static and dynamic postural stability in older individuals: a randomized double-blind sham-controlled study**. *Eur J Neurosci*. 2017; **46**(12): 2875–2884.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Doeltgen SH, Young J, Bradnam LV: **Anodal Direct Current Stimulation of the Cerebellum Reduces Cerebellar Brain Inhibition but Does Not Influence Afferent Input from the Hand or Face in Healthy Adults**. *Cerebellum*. 2016; **15**(4): 466–74.
[PubMed Abstract](#) | [Publisher Full Text](#)
33. Nilsson J, Lebedev AV, Lövdén M: **No significant effect of prefrontal tDCS on working memory performance in older adults**. *Front Aging Neurosci*. 2015; **7**: 230.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Grecco LAC, Duarte NAC, Zanon N, et al.: **Effect of a single session of transcranial direct-current stimulation on balance and spatiotemporal gait variables in children with cerebral palsy: A randomized sham-controlled study**. *Brazilian J Phys Ther*. 2014; **18**(5): 419–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. Schabrun SM, Lamont RM, Brauer SG: **Transcranial direct current stimulation to enhance dual-task gait training in Parkinson's disease: A pilot RCT**. *PLoS One*. 2016; **11**(6): e0158497.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Soler MD, Kumru H, Pelayo R, et al.: **Effectiveness of transcranial direct current stimulation and visual illusion on neuropathic pain in spinal cord injury**. *Brain*. 2010; **133**(9): 2565–77.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Viana RT, Laurentino GEC, Souza RJP, et al.: **Effects of the addition of transcranial direct current stimulation to virtual reality therapy after stroke: a pilot randomized controlled trial**. *NeuroRehabilitation*. 2014; **34**(3): 437–46.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Kim YJ, Ku J, Cho S, et al.: **Facilitation of corticospinal excitability by virtual reality exercise following anodal transcranial direct current stimulation in healthy volunteers and subacute stroke subjects**. *J Neuroeng Rehabil*. 2014; **11**: 124.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 01 December 2020

<https://doi.org/10.5256/f1000research.27770.r74514>

© 2020 **Vitório R.** This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Rodrigo Vitório

Department of Neurology, Oregon Health and Science University, Portland, OR, USA

This interesting ongoing randomized controlled trial examines whether anodal tDCS applied over the dorsolateral prefrontal cortex can enhance and/or prolong the effects of videogame training on balance in older adults. The authors report existing approval by the local ethics committee and ongoing training/intervention and data collection. The study is registered at ensaioclinicos.gov.br. Methods are sound. I have only a few suggestions that I believe will make the manuscript clearer.

- Please mention stimulation type (i.e., "anodal" or "anodic") and targeted area (i.e., dorsolateral prefrontal cortex) in the title and aims (abstract and main text).
- Please clarify why the study includes a "control group" (no electrodes positioning) and a "sham group" (placebo tDCS).
- Please provide further details about the sham tDCS (e.g., will any current intensity be applied? For how long? Etc.)
- Please avoid the use of "elderly". Consider using "older adults/people" throughout the entire manuscript.
- The primary/secondary outcome measures should be explicitly identified.

Is the rationale for, and objectives of, the study clearly described?

Yes

Is the study design appropriate for the research question?

Yes

Are sufficient details of the methods provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Not applicable

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Movement Science and Neuroscience, with experience in conducting clinical trials and reporting their findings.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 21 Dec 2020

Andre Issao Kunitake, University Nove de Julho, São Paulo, Brazil

Dear reviewer. We appreciate your time reading our study and making your considerations. I adapted the article taking into account everything that was requested, to enrich and make this article more quality.

Best regards

Competing Interests: The authors of this study declare that they have no conflicts of interest.

Reviewer Report 02 November 2020

<https://doi.org/10.5256/f1000research.27770.r71407>

© 2020 da Silva Arêas F. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Fernando Zanela da Silva Arêas

Center of Health Sciences, Discipline of Physical Therapy, Federal University of Espírito Santo (UFES), Vitória, Brazil

The paper is well delineated, and well written - it is clear, objective and relevant. The protocols follow all the ethics criteria for reproduction. Also, the literature does not have works that investigate the approach with this methodologic quality.

Does the project have some finance?

This is a good project. I recommend indexing.

Is the rationale for, and objectives of, the study clearly described?

Yes

Is the study design appropriate for the research question?

Yes

Are sufficient details of the methods provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Neurorehabilitation, neuromodulation, traumatic brain injury

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research