RESEARCH ARTICLE

Assessment of the phytochemical, antioxidant and antibacterial activities of *Heteromorpha arborescens* (Spreng.) Cham & Schltdl. leaf extracts [version 1; peer review: 2 approved]

Taiwo Oluwafunmilola Abifarin , Gloria Aderonke Otunola ,
Anthony Jide Afolayan

Medicinal Plants and Economic Development (MPED) Research Centre, Department of Botany, University of Fort Hare, Alice 5700, Eastern Cape, South Africa

V1 First published: 01 Sep 2020, 9:1079
<https://doi.org/10.12688/f1000research.25197.1>
 Latest published: 01 Sep 2020, 9:1079
<https://doi.org/10.12688/f1000research.25197.1>

Abstract

Background: *Heteromorpha arborescens* (Spreng.) Cham. and Schltdl (Apiaceae) is widely used traditionally for the treatment of a wide range of diseases in Southern and Eastern Africa. Although previous studies have reported the biological activities of hexane, ethyl acetate and methanol extracts of *H. arborescens* leaves, there is no scientific information on the phytochemical contents, antioxidant and antibacterial activities of acetone, ethanol, aqueous and blanched extracts. This study is therefore aimed to investigate and compare the phytochemical contents, antioxidant and antibacterial activities of acetone, ethanol, aqueous and blanched extracts of *H. arborescens* leaves.

Methods: Phytochemical analysis for the total phenolic, flavonoid, proanthocyanidin, alkaloid and saponin contents of all the fractions were determined by spectroscopic methods, while the free radical scavenging potential of the extracts were evaluated using DPPH, ABTS radical scavenging and total antioxidant capacity assays. Micro dilution method was used to determine the Minimum Inhibitory Concentrations (MIC) of *H. arborescens* leaf extracts against *Bacillus pumilus*, *Staphylococcus epidermidis*, *Staphylococcus aureus*, *Escherichia coli* and *Klebsiella pneumoniae*.

Results: Total phenol content of the extracts ranged between 15.10 mg GAE/g- 42.50 mg GAE/g, proanthocyanidin was 459-8402.1 mg QE/g, and flavonoid content of 109.24-235.79 mg QE/g. In addition, alkaloids (5.59%) and saponins (23.33%) were present in significant amounts. Based on the IC₅₀ values, the ethanol extract exhibited the highest total antioxidant activity (0.013 mg/mL) with highest inhibition against DPPH and ABTS radicals (0.06 and 0.049 mg/mL respectively).

Open Peer Review

Approval Status

	1	2
version 1		
01 Sep 2020	view	view
1. Abimbola Oloye , Federal University Of Agriculture Abeokuta, Ogun State, Nigeria		
2. Balasubramani Ravindran , Kyonggi University, Suwon, South Korea		
Any reports and responses or comments on the article can be found at the end of the article.		

Considerable antibacterial activities were observed in the acetone, ethanol and blanched extracts with MIC values ranging from 1.563-12.5 mg/mL; however, the aqueous extract was inactive against all the bacteria strains.

Conclusion: The study suggests that *H. arborescens* leaves could be a valuable source of bioactive compounds. Although the blanching process significantly decreased polyphenolic contents and antioxidant activities of the extracts, it increased the antibacterial compounds.

Keywords

Phytochemicals, *Heteromorpha arborescens*, antioxidant, antibacterial, bioactives, blanching, traditional, medicines.

Corresponding author: Gloria Aderonke Otunola (gotunola@ufh.ac.za)

Author roles: Abifarin TO: Data Curation, Formal Analysis, Methodology, Project Administration, Resources, Writing – Original Draft Preparation; Otunola GA: Funding Acquisition, Investigation, Project Administration, Writing – Review & Editing; Afolayan AJ: Funding Acquisition, Supervision, Visualization, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This research was financially supported by Govan Mbeki Research and Development Center, University of Fort Hare (C127).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2020 Abifarin TO *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Abifarin TO, Otunola GA and Afolayan AJ. **Assessment of the phytochemical, antioxidant and antibacterial activities of *Heteromorpha arborescens* (Spreng.) Cham & Schltdl. leaf extracts [version 1; peer review: 2 approved]** F1000Research 2020, 9:1079 <https://doi.org/10.12688/f1000research.25197.1>

First published: 01 Sep 2020, 9:1079 <https://doi.org/10.12688/f1000research.25197.1>

Introduction

Plants from the Apiaceae family are generally known to be rich sources of phytochemicals and antioxidants and are commonly used as food, flavoring agents and medicine¹. These bioactive constituents may contribute significantly to the protection of humans from a wide range of diseases²⁻⁴. *Heteromorpha arborescens* (Spreng.) Cham. and Schltld otherwise referred to as parsley tree in English, “wildepitersielie” in Afrikaans and “umbangandlala” in Xhosa is a large shrub or small tree that belongs to the Apiaceae family³.

The leaves vary from simple to compound and the flowers are small, greenish to yellowish in colour, occurring in compound umbels, with waxy bark which is smooth or glossy in texture⁵. The species is generally considered as a medicinally important plant throughout Africa since almost all its parts are traditionally used for the treatment of different ailments⁶. In South Africa, the leaves and roots are used for blood purification, diabetes and shortness of breath⁷⁻⁹. The bark, leaves and roots are also used for respiratory problems in Kenya, Lesotho and Tanzania⁹. The leaves are consumed as vegetables in Kenya¹⁰, while the roots are given to malnourished children in Botswana and Swaziland¹¹. The plant is also used to treat abdominal pains, dysmenorrhea, nervous and mental disorders, as well as a vermicide in children¹².

The healing ability of the plant is attributed to the presence of bioactive compounds, such as tannins, phenols, alkaloids, saponin, flavonoids, and proanthocyanidin. Although very scanty, previous scientific reports exist on the phenolic contents, antioxidant^{13,14} and antibacterial activities³ of hexane, ethyl acetate and methanol extracts of *H. arborescens* leaves. To the best of our knowledge no study exists on comparison of polyphenolic contents, antioxidant and antibacterial activities of blanched, aqueous, acetone and ethanol extracts of *H. arborescens* leaves. The present study was therefore conducted to determine the phytochemical contents, antioxidant and antibacterial activities of blanched, aqueous, acetone and ethanol extracts of *H. arborescens* leaves.

Methods

Ethical approval

Ethical approval was granted by the University of Fort Hare Animal and Plant Use Research Ethics Committee, South Africa with protocol number OTA011SABI01/19/E.

Chemical and reagents

All reagents and chemicals including gallic acid, rutin, quercetin, aluminum chloride (AlCl_3), ferric chloride ($\text{FeCl}_3 \cdot 7\text{H}_2\text{O}$), Folin-Ciocalteu, oxalic acid, trichloroacetic acid (TCA), sulphuric acid (H_2SO_4), hydrochloric acid (HCl), Na_2CO_3 , ammonium molybdate, potassium ferricyanide ($\text{K}_3\text{Fe}(\text{CN})_6$), catechin, 2,2-diphenyl-1-picrylhydrazyl (DPPH) and 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid diammonium salt) (ABTS) are products of Sigma-Aldrich, South Africa. Other chemicals such as anhydrous sodium carbonate, sodium nitrite (NaNO_2), ethanol, methanol, n-butanol, sodium acetate, butylated hydroxytoluene (BHT), diethyl ether, glacial acetic acid, sulfanilic acid, potassium persulphate, sodium nitroprusside were also

purchased from the same company. All the reagents used were of analytical grade. The bacterial strains including; *Klebsiella pneumoniae* ATCC 13883, *Staphylococcus aureus*, ATCC 29213, *Pseudomonas aeruginosa* ATCC 19582, *Escherichia coli* 25922, *Bacillus pumilus* ATCC 14884, *Staphylococcus epidermis* ATCC 12228 were obtained from the Department of Biochemistry and Microbiology, University of Fort Hare, South Africa.

Plant material

Fresh leaves of *H. arborescens* were obtained in the month of June, 2019 from an area located on latitude $32^\circ 47' 50.4''$ S, $26^\circ 52' 41.8''$ E along Hogsback road, Alice Town, Eastern Cape, South Africa. The plant was authenticated by a taxonomist at the University of Fort Hare and a voucher specimen was preserved in the Giffen Herbarium and assigned number Abif2019/03.

Preparation of plant extracts

Fresh *H. arborescens* leaves were oven dried at 40°C and crushed to coarse powder. 100 g of the powdered plant material was extracted separately in 500 mL acetone, ethanol and water for 24 hours on an orbital shaker. An equal weight of the fresh leaves was immersed into 500 mL of hot water (80°C) for 5 min to simulate home cooking and ground with a blender. The extracts were filtered under pressure using a Buchner funnel and Whatman filter paper (150 mm); the acetone and ethanol extracts were concentrated to dryness using a rotary evaporator while the aqueous extract and blanched sample were subjected to freeze drying. The extracts were thereafter stored at 4°C until further analysis.

Phytochemical screening

Determination of total phenolic content. The total phenolic content was determined by spectrometry using Folin-Ciocalteu reagent. Briefly, 0.5 mL of extract (1mg/mL) and gallic acid (0.02 to 0.1mg/mL) were put separately in test tubes. Thereafter, 2.5mL of 10% Folin-Ciocalteu reagent was added into the test tubes, after which 2.5 mL of 7.5% sodium carbonate solution was added to the mixture, stirred and incubated at 40°C for 30 min. The absorbance was then measured at 760 nm using a Hewlett Packard VR-2000 spectrophotometer. All samples were analysed in triplicate. Total phenolic content was expressed as milligrams of gallic acid equivalent per gram (mg GAE/g) with the standard curve: $y = 10.875x + 0.1025$, $R^2 = 0.996$.

Where R is the determined coefficient, x is the concentration, and y is the absorbance.

Determination of flavonoids content. The amount of flavonoids present in the extract was determined using the aluminium chloride (AlCl_3) colorimetric assay, as described by Majouli *et al.*¹⁵. A mixture of 0.5 mL of extract was added to an equal volume of 2% AlCl_3 solution. The mixture was incubated for 10 min and vigorously shaken, after which the absorbance was measured at 420 nm. Varying concentrations of the standard (quercetin) were also prepared with the same method. All samples were analysed in triplicates. Flavonoid content was

expressed as milligrams of quercetin equivalent per gram (mg QE/g) with the standard curve: $y = 2.9422x - 0.4438$, $R^2 = 0.913$

Where R is the determined coefficient, x is the concentration, and y is the absorbance.

Determination of proanthocyanidin (condensed tannins).

Proanthocyanidin content was determined as described by Sagbo *et al.*¹⁶. Briefly, a mixture of 3mL of 4% vanillin-methanol solution and 1.5mL of HCl was added to 0.5 mL of each extract. The solution was stirred and incubated at 27°C for 15 min, after which absorbance was measured at 500 nm. All samples were analysed in triplicate and the amount of proanthocyanidin was expressed as mg/g dry weight of quercetin equivalent (mg QE/g) of the extract with the standard curve: $y = 0.0252x + 0.0482$, $R^2 = 0.9005$.

Where R is the determined coefficient, x is the concentration, and y is the absorbance.

Determination of alkaloid content.

Alkaloid content of *H. arborescens* leaves was determined as previously described by Abifarín *et al.*¹⁷. 0.5 g of the pulverized leaves was mixed with 200mL of 10% acetic acid in ethanol. The mixture was covered, incubated at room temperature for 4 h, filtered and concentrated to about a quarter of its original volume in a water bath. To the extract, concentrated ammonium hydroxide was added in drops till precipitation was complete. After the solution was allowed to settle, precipitates obtained were washed with dilute ammonium hydroxide and then filtered. The residue was dried in an oven (40°C), weighed and the alkaloid content was determined using the following formula:

$$\% \text{ Alkaloid} = \text{weight of precipitate} / \text{initial weight of sample} \times 100.$$

Determination of saponin content.

Saponin content of *H. arborescens* leaves was determined as previously described by Unuofin *et al.*¹⁸. Briefly, 0.5 g of pulverized *H. arborescens* leaves was measured into 50 mL of 20% ethanol prepared in distilled water. The mixture was heated in a hot water bath for 4 h at 55°C. The mixture was filtered and the residue extracted again with another 50 mL of 20% ethanol. The two filtrates were combined and reduced to 20 mL over a hot water bath (90°C). The concentrated solution obtained was poured into a 250 mL separating funnel containing 20 mL of diethyl ether. The aqueous layer was collected while the ether layer was discarded. 20 mL of n-butanol was added to the filtrate and then washed thrice with 10 mL of 5% sodium chloride. The mixture was heated in an oven (40°C) to constant weight. The percentage saponin content of the sample was calculated using the following formula:

$$\% \text{ Saponins} = \text{weight of final filtrate} / \text{weight of sample} \times 100$$

Antioxidant activity

DPPH radical scavenging activity. DPPH radical scavenging activity for each plant extract was determined as previously described by Ohikhen *et al.*¹⁹. Briefly, a reaction mixture

containing 2.5 mL of DPPH solution (0.13 mM) and 2.5 mL of each plant extract or standard (rutin and BHT) dissolved in methanol at varying concentrations (0.005, 0.01, 0.02, 0.04, 0.08 mg/mL), stirred and kept in the dark for 30 min. The absorbance was measured at 517 nm and DPPH radical scavenging activity was calculated as:

$$\% \text{ DPPH scavenging activity} = [(Abs \text{ DPPH} - Abs \text{ Sample}) / Abs \text{ DPPH}] \times 100$$

Where Abs control is the absorbance of DPPH radical + methanol; Abs sample is the absorbance of DPPH radical + sample extract/standard.

ABTS radical scavenging assay.

ABTS scavenging activity of the different plant extracts was determined as described by Unuofin *et al.*²⁰. A mixture was prepared by reacting 7 mM ABTS solution and 2.45 mM $K_2S_2O_8$ (1:1), which was kept in the dark for 12 hours to produce a bluish green coloration. The solution was adjusted with methanol until an absorbance of 0.700 ± 0.01 at 734 nm was obtained. 1mL of plant extract was allowed to react with 1 mL of the working solution and the absorbance was measured at 734 nm after 7 min. The ABTS scavenging capacity of the extracts were compared with that of BHT and rutin and percentage inhibition was calculated as:

$$ABTS \text{ radical scavenging activity } (\%) = [(Abs \text{ control} - Abs \text{ sample}) / (Abs \text{ control})] \times 100.$$

Where Abs control is the absorbance of ABTS radical + methanol; Abs sample is the absorbance of ABTS radical + sample extract/standard.

Total antioxidant capacity (phosphomolybdenum) assay.

The total antioxidant capacity was determined by the method described by Abifarín *et al.*¹⁷. Briefly, 0.3 mL of the extracts and standard (0.025-0.4 mg/mL) were measured into separate test tubes and each was dissolved in 3 mL of reagent solution (0.6 M sulfuric acid, 4 mM ammonium molybdate, and 28 mM sodium phosphate). The test tubes were incubated at 90°C in a water bath for 90 min, allowed to cool to room temperature and the absorbance was measured at 695 nm. Rutin and BHT were used as standards. The percentage inhibition (%TAC) was calculated as:

$$\%TAC = [(Absorbance \text{ of sample} - Absorbance \text{ of control}) / (absorbance \text{ of sample})] \times 100$$

Determination of antibacterial activity

Microorganisms and media. The bacteria used in this study were chosen primarily on the basis of their importance as opportunistic microorganisms for humans with diabetes mellitus. The bacterial strains include: *Klebsiella pneumoniae* ATCC 13883, *Staphylococcus aureus*, ATCC 29213, *Pseudomonas aeruginosa* ATCC 19582, *Escherichia coli* 25922, *Bacillus pumilus* ATCC 14884, *Staphylococcus epidermidis* ATCC 12228.

Minimum Inhibitory Concentration (MIC). The MIC of the extracts were evaluated against selected Gram positive (*Bacillus*

pumilus, *Staphylococcus aureus* and *Staphylococcus epidermidis*) and Gram negative bacteria (*Klebsiella pneumoniae*, *Pseudomonas aeruginosa* and *Escherichia coli*) by microdilution method, as described by Mohsenipour *et al.*²¹. The bacterial concentration in the inoculum was standardized at 0.5 McFarland turbidity scale, (1×10^8 CFU/ml).

Firstly, sterile round bottom 96-well plates were filled with 100 μ l of distilled water. 100 μ l stock fractions of 50 mg/mL of plant extract and standard (Erythromycin) were then added into the first row. The fractions were serially diluted to make varying concentrations of the plant extracts (0.78125–12.5 mg/mL) and standard (0.0625–1 μ g/mL) and then 50 μ l of inoculums was added into the wells. Thereafter, 100 μ l sterile nutrient broth culture medium plus 50 μ l of the culture of each organism was added into each well and the inoculated micro plates were incubated at 37°C for 24 h. 1% DMSO was used as the negative control while 100 μ l of inoculums only was used as the growth hormone. The plates were incubated for 24 h at 37°C, and subsequently 40 μ l of 0.4 mg/mL INT dye was added to each well. The plates were gently agitated and incubated for another 30 min. The MICs were then determined as the lowest concentrations at which there was no indication of colour change to pink (no bacterial growth).

Statistical analyses

All data were expressed as mean $\pm$ standard deviation (SD) of triplicates and subjected to one-way analysis of variance (ANOVA). Where the data showed significant difference ($p < 0.05$) among the extracts, a mean separation was done using Fischer's Least Significant Difference. MINITAB 17 statistical package was used for analysis.

Results

Phytochemical contents

The total phenol, proanthocyanidin and flavonoid contents of *H. arborescens* leaf extracts are presented in Table 1. The highest phenol content was found in the ethanol extract (42.50 mg GAE/g), followed by the aqueous extract (34.24 mg GAE/g), blanch extract (21.42 mg GAE/g) and acetone extract (15.10 mg GAE/g). Ethanol extract (8402.1 mg QE/g) showed the highest proanthocyanidin content followed by acetone extract (4923 mg QE/g DW), blanch extract

(928.6 mg QE/g) and aqueous extract (459 mg QE/g). The concentration of flavonoid was highest in the ethanol extract (235.79 mg QE/g), followed by acetone extract (230.52 mg QE/g), aqueous extract (154.95 mg QE/g) and blanch extract (109.24 mg QE/g). Quantitative determination of the alkaloid and saponin contents of *H. arborescens* leaves also revealed low alkaloid content (5.59%) with considerable amount of saponin (23.33%).

Antioxidant activities

ABTS radical scavenging activity. The ABTS radical scavenging activities of the extracts and standards (BHT and rutin) are presented in Figure 1. The samples exhibited significant ABTS radical scavenging activities, which increased in a concentration dependent manner. Although the standards showed higher ABTS inhibitory potentials than the extracts, the highest inhibitory capacity of the extracts was observed in the ethanol and aqueous extracts. Based on the IC_{50} (Table 2), ABTS scavenging activity of the samples was in the order: BHT > rutin > ethanol & aqueous > acetone > blanch.

Total antioxidant capacity. The total antioxidant capacity of the extracts and standards (BHT and rutin) are shown in Figure 2. There was a dose dependent increase in total antioxidant capacity of the samples, with the ethanol extract showing the highest antioxidant capacity when compared with other extracts. Based on the IC_{50} (Table 2), the values of the total antioxidant capacity of the standards and extracts were in the order: BHT > ethanol > rutin > aqueous > acetone > blanch.

DPPH scavenging activity. There was a dose-dependent increase in percentage inhibitory activity of the extracts and standards against DPPH radical (Figure 3). Ethanol extracts showed the highest DPPH radical inhibitory activity; even though the standards inhibited best when compared with the extracts. With respect to the IC_{50} (Table 2) the inhibitory capacity of the standards and extracts were in the order: rutin > BHT > ethanol > aqueous > blanch > acetone.

Minimum inhibitory concentration. Antibacterial activity of erythromycin (standard) and extracts of *H. arborescens* leaves are presented in Table 3. While all Gram-positive and Gram-negative bacteria strains tested were resistant to the

Table 1. Total phenol, proanthocyanidin, flavonoid, alkaloids and saponin contents of *H. arborescens* leaves.

Extracts	Phenol (mg GAE/g)	Proanthocyanidins (mg QE/g)	Flavonoids (mg QE/g)	Alkaloids (%)	Saponins (%)
Aqueous	34.24 $\pm$ 0.46 ^a	459 $\pm$ 49.9 ^a	154.95 $\pm$ 9.99 ^a		
Acetone	15.10 $\pm$ 1.38 ^b	4923 $\pm$ 579 ^b	230.52 $\pm$ 0.59 ^b		
Blanch	21.42 $\pm$ 0.28 ^c	928.6 $\pm$ 90.9 ^c	109.24 $\pm$ 0.17 ^c		
Ethanol	42.50 $\pm$ 0.69 ^d	8402.1 $\pm$ 89.5 ^d	235.79 $\pm$ 4.52 ^d		
Pulverized	*	*	*	5.59 $\pm$ 0.45	23.33 $\pm$ 2.3

The results are expressed as mean $\pm$ standard deviation (n=3). Values with different superscripts are significantly different ($P < 0.05$) across the different extracts. * - not determined.

Figure 1. ABTS radical scavenging activity of extracts of *H. arborescens* leaf extracts and standards (rutin and BHT). Results are expressed as mean $\pm$ standard deviation (n=3). Columns with different letters are significantly different ($P < 0.05$) across the different samples.

Table 2. IC_{50} values of various solvent extracts of *H. arborescens* leaf extracts and standards.

Extracts/standard	DPPH (mg/mL)	ABTS (mg/mL)	TAC (mg/mL)
Aqueous	0.1	0.049	0.024
Blanched	0.12	0.109	0.152
Acetone	0.21	0.081	0.046
Ethanol	0.06	0.049	0.013
Rutin	0.026	0.024	0.017
BHT	0.031	0.012	0.012

aqueous extract at all the concentrations tested, the blanched, acetone, and ethanol extracts showed considerable antibacterial activities with MICs ranging from 1.563-12.5 mg/mL. Ethanol extract showed the best antibacterial activity against *B. pumilus*, *S. epidermidis*, *S. aureus* and *P. aeruginosa*, exhibiting a low MIC of (1.563 and 3.125 mg/mL). Generally speaking, the Gram-negative bacteria were observed to be more resistant against the extracts tested.

Discussion

Plant extracts have been reported to have numerous protective roles due to their phytochemical contents, which contributes to a large extent towards their antioxidant and antimicrobial

activities²². Oxidative stress, which leads to the production of free radicals in the body, plays a major role in the development of chronic and degenerative ailments such as cancer, arthritis, aging, autoimmune disorders, cardiovascular and neurodegenerative diseases²³. Antioxidants play a vital role in deactivating and scavenging of the free radicals in order to reduce the risk of these chronic diseases²⁴.

In the present study, the phytochemical, antioxidant and antibacterial activities of the acetone, ethanol, aqueous and blanched extracts of *H. arborescens* leaves were assessed. Phytochemical constituents varied significantly in all the extracts, which revealed the presence of phenols, flavonoids

Figure 2. Total antioxidant capacity of *H. arborescens* leaf extracts and standards (Rutin and BHT). Results are expressed as mean $\pm$ standard deviation (n=3). Columns with different letters are significantly different ($P < 0.05$) across the different samples.

Figure 3. DPPH scavenging activity of *H. arborescens* leaf extracts and standards (Rutin and BHT). Results are expressed as mean $\pm$ standard deviation (n=3). Columns with different letters are significantly different ($P < 0.05$) across the different samples.

and proanthocyanidin in considerable amounts. The variation in the phytochemical constituents could be attributed to differences in the extracting capabilities of different solvents. Phenols, a major phytochemical found in plants have been widely studied because of their ability to reduce oxidative stress-related degenerative diseases, including cancer. Flavonoids are a group of hydroxylated phenolics, and their beneficial effects are linked to their antioxidant, antibacterial, anticancer,

anti-inflammatory^{25,26} anti-hypertensive, and cardio protective activities^{27,28}. Proanthocyanidins are a class of compounds belonging to the flavonoid family, with protective effects against tissue damage and cancer, and they also improve blood circulation by strengthening the capillaries, arteries and veins²⁹.

Ethanol has been reported to be suitable for extraction of compounds with a wide range of polarity while water is suitable

Table 3. Minimum Inhibitory Concentration (MIC) values of *H. arborescens* leaf extracts against selected bacterial strains.

Samples	Bacterial strains (MIC) mg/mL					
	<i>B. pumilus</i> ^a	<i>S. epidermidis</i> ^a	<i>S. aureus</i> ^a	<i>E.coli</i> ^b	<i>K. pneumoniae</i> ^b	<i>P. aeruginosa</i> ^b
Aqueous	Na	Na	Na	Na	Na	Na
Blanched	12.5	6.25	12.5	12.5	12.5	12.5
Acetone	6.25	1.563	3.125	12.5	6.25	6.25
Etdanol	3.125	3.125	1.563	6.25	6.25	12.5
Erythromycin (µg/mL)	0.125	0.0625	0.0625	0.0625	0.125	0.125

Bacillus pumilus, *Staphylococcus epidermidis*, *Staphylococcus aureus*, *Escherichia coli*, *Klebselia pneumoniae*.

Bacteria ^a- Gram positive. Bacteria ^b- Gram negative. Na- not active at tde highest concentration tested.

for very polar compounds³⁰. Consequently, a higher quantity of phenolic compounds, proanthocyanidin and flavonoids were observed in the ethanol samples when compared to the acetone, blanched and aqueous extracts; hence, better activity. This is in agreement with previous reports that ethanol is more suitable for the extraction of phenolic compounds in plants^{30–32}. Furthermore, high phenolic compounds were indicated in ethyl acetate and methanol extracts, suggesting the reason for its high antioxidant activity¹³. Flavonoid content of acetone extract of *H. arborescens* was comparable to results obtained by Elisha *et al.*¹³; however, total phenolic content was much higher as opposed to the current study.

Our results indicate that blanching resulted in a significant decrease in total phenols and flavonoids with significant increase in proanthocyanidin content. This correlates with similar observations by Korus *et al.*³³, Jaiswal *et al.*³⁴ and Ironi *et al.*³⁵. Losses of polyphenolic compounds upon blanching was also observed in some cruciferous vegetables, such as spinach³⁶, kale, broccoli³⁷ and cauliflower^{38,39}. The decrease in polyphenolic contents could be attributed to the loss of water soluble vitamins and nitrogen compounds during the blanching process³⁴. Contrary to the findings of this study, some researchers have claimed that the heating process may alter the cell membrane, causing a release of some membrane-bound phytochemicals which may increase bioavailability as observed by Jimenez-Monreal *et al.*⁴⁰. Similarly, Ma *et al.*⁴¹ observed an increase in polyphenolic contents of *Daucus carota* L. juice after blanching (at 95°C for 3 min).

Saponins possess medicinal properties such as anti-inflammatory, antibacterial, anticancer and cytotoxic activities⁴². Alkaloids have also been shown to exhibit antioxidant properties by reducing oxidative damage induced by hydrogen peroxide⁴³. The amount of alkaloids and saponin exhibited by *H. arborescens* leaves is promising in playing the aforementioned biological roles.

Further assessment of the antioxidant activities of the extracts revealed the free radical scavenging potential of the extracts,

which varied among the methods used. This variation was due to the fact that antioxidants are able to neutralize free radicals by different modes of action such as transition metal chelation, singlet oxygen quenchers, and donation of hydrogen⁴⁴.

The total antioxidant capacity and scavenging activity of the extracts against DPPH+ and ABTS+ were compared to the standards (BHT and rutin) and the extracts showed considerable antioxidant activity. Since positive correlation between polyphenolic compounds and antioxidant activities have been reported^{43,45,46}, the highest antioxidant activity exhibited by the ethanol extract could be attributed to its high polyphenolic compounds, as shown in this study. In terms of the relatively higher DPPH and ABTS inhibitory potential of the ethanol extract, these results are in agreement with antioxidant activity of other Apiaceae species such as: *Coriandrum sativum*⁴⁷, *Ammi majus* L.⁴⁸, *Daucus carota* L.⁴⁹, *Ferula gummosa*⁵⁰, *Seseli libanotis* (L.) Koch⁵¹, *Foeniculum vulgare* and *Anethum graveolens*¹. This strengthens the suggestion that the most important bioactive potentials are observed in the plant extracts with high amounts of phenolics and flavonoids. This finding therefore provides some scientific verification for the biological activities of *H. arborescens* leaves.

In recent years, more attention has been given to the development of antibacterial drugs from medicinal plants instead of synthetic ones. The antibacterial activities of plant extracts have been confirmed against an array of both Gram-positive and Gram-negative bacteria⁵². Previous studies have established that the crude, hexane, ethyl acetate, chloroform and butanol extracts reveal antibacterial activities against *E. coli*, *S. aureus* and *P. aeruginosa*. However, *P. aeruginosa* was more susceptible to three out of the extracts with the butanol fraction showing the highest antibacterial activity⁵³. The resistance of all the bacterial strains to the aqueous extract at all concentrations tested and the significant antibacterial activity of the blanched extract obtained in the present study, could be attributed to poor solubility of the antibacterial compounds in cold water and high extractive power of boiled water.

In addition, the higher antibacterial activity observed in the acetone and ethanol extracts could be due to relatively higher flavonoid and phenolic contents present⁵⁴.

Our findings are in agreement with McGaw *et al.*⁵⁵ who reported that in the evaluation of antibacterial activities of hexane, ethanol and aqueous leaf extracts of *H. arborescens* against *Bacillus subtilis*, *S. aureus*, *E. coli*, and *K. pneumoniae*, only ethanol extract exhibited antibacterial activities. These findings correlate with observation made by Wigmore *et al.*⁵⁶ that aqueous extracts of plants showed less antibacterial activity when compared with other solvents. Furthermore, in agreement with our findings, the aqueous extract was inactive against all Gram-positive bacteria used. However, contrarily, the aqueous extract was active against *S. aureus* and *S. epidermidis*³.

Conclusion

Our results clearly indicated that the extraction of phenolic compounds and their antioxidant capacity is highly dependent on the solvent of extraction. The study also revealed that *H. arborescens* aqueous, blanched and ethanol leaf extracts possess various levels and concentrations of phytochemical constituents, which may be essential for human health. A positive correlation between polyphenolic contents, antioxidant

and antibacterial activities in extracts of *H. arborescens* leaves was established, which indicates that certain phenolic compounds may be responsible for high antioxidant activity. It was also observed that the blanching process (at 80° C) significantly decreased polyphenolic content and antioxidant activities but increased antibacterial activity of *H. arborescens* leaves. This study therefore agrees with its reported use in traditional medicine for the treatment of some bacterial infections and diseases.

Data availability

Underlying data

Figshare: Absorbance values for antioxidant activity.xlsx, <https://doi.org/10.6084/m9.figshare.12654479.v1>⁵⁷.

This project contains the following underlying data:

- Replicate absorbance values for DPPH, TAC and ABTS
- Replicate phytochemical analysis results
- MIC values of antibacterial activity of the extracts

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC0 1.0 Public domain dedication).

References

1. Bagdassarian VLC, Bagdassarian KS, Atanassova MS: **Phenolic profile, antioxidant and antimicrobial activities from the Apiaceae family (dry seeds).** *Mint J Pharm Medi Scie.* 2013; 2 (4): 26–31.
[Reference Source](#)
2. Fotina AA, Fisnin VI, Surai, PF: **Recent developments in usage of natural antioxidants to improve chicken meat production and quality.** *Bulg J Agric Sci.* 2013; 19(5): 889–896.
[Reference Source](#)
3. Nkomo M, Kambizi L: **Antimicrobial activity of Gunnera perpensa and Heteromorpha arborescens var. abyssinica.** *J Med Pl Res.* 2009; 3(12): 1051–1055.
[Reference Source](#)
4. Abifarini TO, Ogunola GA, Afolayan AJ: **Cytotoxicity evaluation and anti-inflammatory potentials of Cucumis africanus L.f. leaves.** *Med Plants-Inter J Phytomed Relat Indus.* 2020; 12(1): 48–52.
[Publisher Full Text](#)
5. Palgrave MC: **Keith Coates Palgrave Trees of Southern Africa.** 3rd ed. Cape Town. Struik Publishers (Pty) Ltd. 2002.
[Reference Source](#)
6. Maroyi A: **Heteromorpha arborescens: a review of its botany, medicinal uses, and pharmacological properties.** *Asian Journal of Pharmaceutical and Clinical Research.* 2018; 11(11): 75–82.
[Publisher Full Text](#)
7. Mushi MJ, Mbembo ZH: **Experience of Tanzanian traditional healers in the management of non-insulin dependent diabetes mellitus.** *Pharm Biol.* 2002; 40(7): 552–560.
[Publisher Full Text](#)
8. Schmidt E, Lotter M, McClelland W: **Trees and Shrubs of Mpumalanga and Kruger National Park.** Johannesburg, Jacana; 2002.
[Reference Source](#)
9. Van BE Wyk, Van Oudtshoorn N: **Medicinal Plants of South Africa.** Pretoria: Briza Publications. 2013.
10. Bussmann RW: **Ethnobotany of the Samburu of Mt. Nyiru, South Turkana, Kenya.** *J Ethnobiol Ethnomed.* 2006; 2(1): 35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Setshogo MP, Mbereki CM: **Floristic diversity and uses of medicinal plants sold by street vendors in Gaborone, Botswana.** *African Journal of Plant Science and Biotechnology.* 2011; 5(1): 69–74.
[Reference Source](#)
12. Nkomo M, Nkeh-Chungag BN, Kambizi L, *et al.*: **Investigation of the antinociceptive and anti-inflammatory properties of Heteromorpha arborescens (Apiaceae).** *Afr J Tradit Complement Altern Med.* 2011; 8(4): 412–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Sunil C, Kumar V, Van Staden J: **In vitro alpha-glucosidase inhibitory, total phenolic composition, antiradical and antioxidant potential of Heteromorpha arborescens (Spreng.) Cham. & Schldl. leaf and bark extracts.** *South African Journal of Botany.* 2019; 1(124): 380–386.
[Publisher Full Text](#)
14. Elisha IL, Dzoyem JP, McGaw LJ, *et al.*: **The anti-arthritis, anti-inflammatory, antioxidant activity and relationships with total phenolics and total flavonoids of nine South African plants used traditionally to treat arthritis.** *BMC Complement Altern Med.* 2016; 16(1): 307.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Majouli K, Hamdi A, Hilla MB: **Phytochemical analysis and biological activities of Hertia cheirifolia L. roots extracts.** *Asian Pac J Trop Med.* 2017; 10(12): 1134–1139.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Sagbo IJ, Afolayan AJ, Bradley G: **Antioxidant, antibacterial and phytochemical properties of two medicinal plants against the wound infecting bacteria.** *Asian Pacific Journal of Tropical Medicine.* 2017; 7(9): 817–825.
[Publisher Full Text](#)
17. Abifarini TO, Afolayan AJ, Ogunola GA: **Phytochemical and antioxidant activities of Cucumis africanus L.f. A wild vegetable of South Africa.** *J Evid Based Integr Med.* 2019; 24: 1–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Unuofin JO, Ogunola GA, Afolayan AJ: **Polyphenolic content, antioxidant and antimicrobial activities of Vernonia mespilifolia Less. used in folk medicine in the Eastern Cape Province, South Africa.** *J Evid Based Integr Med.* 2018; 23: 1–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Ohikhen FU, Wintola OA, Afolayan AJ: **Quantitative phytochemical constituents and antioxidant activities of the mistletoe, Phragmanthera capitata (Sprengel) Balle extracted with different solvents.** *Pharmacognosy Res.* 2018; 10(1): 16–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Unuofin JO, Ogunola GA, Afolayan AJ: **Phytochemical screening and in vitro evaluation of antioxidant and antimicrobial activities of Kedrostis africana (L.) Cogn.** *Asian Pacific Journal of Tropical Biomedicine.* 2017; 7(10): 901–908.
[Publisher Full Text](#)
21. Mohsenipour Z, Hassanshahian M: **The effects of Allium sativum extracts on**

- biofilm formation and activities of six pathogenic bacteria. *Jundishapur J Microbiol.* 2015; 8(8): e18971.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Kuspradini H, Wulandari I, Putri AS, *et al.*: Phytochemical, antioxidant and antimicrobial properties of *Litsea angulata* extracts [version 2; peer review: 3 approved]. *F1000 Res.* 2018; 7: 1839.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 23. Shinde A, Ganu J, Naik P: Effect of free radicals & antioxidants on oxidative stress: a review. *Journal of Dental and Allied Sciences.* 2012; 1(2): 63–66.
[Publisher Full Text](#)
 24. Guleria S, Tikku AK, Singh G, *et al.*: In vitro antioxidant activity and phenolic contents in methanol extracts from medicinal plants. *J Plant Biochem Biotechnol.* 2013; 22(1): 9–15.
[Publisher Full Text](#)
 25. Olajuyigbe OO, Afolayan AJ: Phenolic content and antioxidant property of the bark extracts of *Ziziphus mucronata* Willd. subsp. *mucronata* Willd. *BMC Complement Altern Med.* 2011; 11(1): 130.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 26. Khanam UKS, Oba S, Yanase E, *et al.*: Phenolic acids, flavonoids and total antioxidant capacity of selected leafy vegetables. *J Funct Foods.* 2012; 4(4): 979–987.
[Publisher Full Text](#)
 27. Abushouk AI, Salem AMA, Abdel-Daim MM: *Berberis vulgaris* for cardiovascular disorders: a scoping literature review. *Iran J Basic Med Sci.* 2017; 20(5): 503–510.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 28. Uddin MS, Hossain MS, Al Mamun A, *et al.*: Phytochemical analysis and antioxidant profile of methanolic extract of seed, pulp and peel of *Baccaurea ramiflora* Lour. *Asian Pac J Trop Med.* 2018; 11(7): 443–450.
[Reference Source](#)
 29. Di Majo D, La Guardia M, Giammanco S, *et al.*: The antioxidant capacity of red wine in relationship with its polyphenolic constituents. *Food Chem.* 2008; 111(1): 45–49.
[Publisher Full Text](#)
 30. Sun C, Wu ZS, Wang Z, *et al.*: Effect of ethanol/water solvents on phenolic profiles and antioxidant properties of Beijing Propolis extracts. *Evid Based Complement Alternat Med.* 2015; 2015: 595393.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 31. Wen TN, Prasad KZ, Yang B, *et al.*: Bioactive substance contents and antioxidant capacity of raw and blanched vegetables. *Innovative Food Science & Emerging Technologies.* 2010; 11(3): 464–469.
[Publisher Full Text](#)
 32. Jimoh MO, Afolayan AJ, Lewu FB: Antioxidant and phytochemical activities of *Amaranthus caudatus* L. harvested from different soils at various growth stages. *Sci Rep.* 2019; 9(1): 12965.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 33. Korus A, Lisiewska Z: Effect of preliminary processing and method of preservation on the content of selected antioxidative compounds in kale (*Brassica oleracea* L. var. *acephala*) leaves. *Food Chem.* 2011; 129(1): 149–154.
[Publisher Full Text](#)
 34. Jaiswal AK, Gupta S, Abu-Ghannam N: Kinetic evaluation of colour, texture, polyphenols and antioxidant capacity of Irish York cabbage after blanching treatment. *Food Chem.* 2012; 131(1): 63–72.
[Publisher Full Text](#)
 35. Ironi EA, Akintunde JK, Agboola SO, *et al.*: Blanching influences the phenolics composition, antioxidant activity, and inhibitory effect of *Adansonia digitata* leaves extract on α -amylase, α -glucosidase, and aldose reductase. *Food Science & Nutrition.* 2012; 5(2): 233–242.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 36. Ismail A, Lee WY: Influence of cooking practice on antioxidant properties and phenolic content of selected vegetables. *Asia Pac J Clin Nutr.* 2004; 13: S162–S162.
[Reference Source](#)
 37. Gliszczynska-Swiglo A, Ciska E, Pawlak-Lemańska K, *et al.*: Changes in the content of health-promoting compounds and antioxidant activity of broccoli after domestic processing. *Food Addit Contam.* 2006; 23(11): 1088–1098.
[PubMed Abstract](#) | [Publisher Full Text](#)
 38. Zhang D, Hamauzu Y: Phenolics, ascorbic acid, carotenoids and antioxidant activity of broccoli and their changes during conventional and microwave cooking. *Food Chem.* 2004; 88(4): 503–509.
[Publisher Full Text](#)
 39. Sikora E, Cieślak E, Leszczynska T, *et al.*: The antioxidant activity of selected cruciferous vegetables subjected to aquathermal processing. *Food Chem.* 2008; 107(1): 55–59.
[Publisher Full Text](#)
 40. Jiménez-Monreal AM, García-Diz L, Martínez-Tomé M, *et al.*: Influence of cooking methods on antioxidant activity of vegetables. *J Food Sci.* 2009; 74(3): H97–H103.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. Ma T, Tian C, Luo J, *et al.*: Influence of technical processing units on polyphenols and antioxidant capacity of carrot (*Daucus carot* L.) juice. *Food Chem.* 2013; 141(3): 1637–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
 42. Ashour AS, El Aziz MMA, Melad ASG, *et al.*: A review on saponins from medicinal plants: chemistry, isolation, and determination. *J Nanomed Res.* 2019; 8(1): 6–12.
[Publisher Full Text](#)
 43. Kibiti CM, Afolayan AJ: Preliminary phytochemical screening and biological activities of *Bulbine abyssinica* used in the folk medicine in the Eastern Cape Province, South Africa. *Evid Based Complement Alternat Med.* 2015; 2015: 617607.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 44. Cioffi G, D'Auria M, Braca A, *et al.*: Antioxidant and free-radical scavenging activity of constituents of the leaves of *Tachigalla paniculata*. *J Nat Prod.* 2002; 65(11): 1526–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 45. Gomes da Cruz Silva ME, Guimarães AL, Oliveira AP, *et al.*: HPLC-DAD analysis and antioxidant activity of *Hymenaea martiana* Hayne (Fabaceae). *J Chem Pharm Res.* 2012; 4(2): 1160–1166.
[Reference Source](#)
 46. Adefuye AO, Ndip RN: Phytochemical analysis and antibacterial evaluation of the ethyl acetate extract of the stem bark of *Bridelia micrantha*. *Pharmacogn Mag.* 2013; 9(33): 45–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 47. Matejić JS, Džamić A, Mihajilov-Krstev T, *et al.*: Total phenolic content, flavonoid concentration, antioxidant and antimicrobial activity of methanol extracts from three *Seseli* L. taxa. *Central European Journal of Biology.* 2012; 7(6): 1116–1122.
[Publisher Full Text](#)
 48. Al-Hadhrami RMS, Hossain MA: Evaluation of antioxidant, antimicrobial and cytotoxic activities of seed crude extracts of *Ammi majus* grown in Oman. *Egyptian Journal of Basic Applied Sciences.* 2016; 3(4): 329–334.
[Publisher Full Text](#)
 49. Chatatikun M, Chiabchalard A: Phytochemical screening and free radical scavenging activities of orange baby carrot and carrot (*Daucus carota* Linn.) root crude extracts. *Journal of Chemical and Pharmaceutical Research.* 2013; 5(4): 97–102.
[Reference Source](#)
 50. Ebrahimzadeh MA, Nabavi SM, Nabavi SF, *et al.*: Antioxidant activity of hydroalcoholic extract of *Ferula gummosa* Boiss roots. *Eur Rev Med Pharmacol Sci.* 2011; 15(6): 658–64.
[PubMed Abstract](#)
 51. Shah MR, Satardeka KV, Barve SS: Screening of phenolic content, antioxidant and in-vitro eye irritation activities from Apiaceae family (dry seeds) for potential cosmetic applications. *International Journal of Pharmaceutical Science Research.* 2014; 5(10): 4366–4374.
[Reference Source](#)
 52. Bazargani MM, Rohloff J: Antibiofilm activity of essential oils and plant extracts against *Staphylococcus aureus* and *Escherichia coli* biofilms. *Food control.* 2016; 61: 156–64.
[Publisher Full Text](#)
 53. Adamu M, Bagla VP, Eloff JN: Fractionation of *Heteromorpha arborescens* var. *abyssinica* (Apiaceae) leaf extracts based on polarity leads to a marked change in cytotoxicity that may yield a commercially useful product. *South African Journal of Botany.* 2016; 103: 36–40.
[Publisher Full Text](#)
 54. Rathee P, Chaudhary H, Rathee S, *et al.*: Mechanism of action of flavonoids as anti-inflammatory agents: A review. *Inflamm Allergy Drug Targets.* 2009; 8(3): 229–235.
[PubMed Abstract](#) | [Publisher Full Text](#)
 55. McGaw LJ, Jäger AK, Van Staden J, *et al.*: Antibacterial, anthelmintic and anti-moebic activity in South African medicinal plants. *J Ethnopharmacol.* 2000; 72(1–2): 247–263.
[PubMed Abstract](#) | [Publisher Full Text](#)
 56. Wigmore SM, Naiker M, Bean DC: Antimicrobial activity of extracts from native plants of temperate Australia. *Pharmacogn Commn.* 2016; 6(2): 80–84.
[Reference Source](#)
 57. Abifarint T, Otunola GA, Afolayan AJ: Absorbance values for antioxidant activity. *xlsx. figshare.* Dataset. 2020.
<http://www.doi.org/10.6084/m9.figshare.12654479.v1>

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 02 October 2020

<https://doi.org/10.5256/f1000research.27806.r71376>

© 2020 Ravindran B. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Balasubramani Ravindran

Department of Environmental Energy and Engineering, Kyonggi University, Suwon, South Korea

- **Title:** The title reflects sufficiently, the work done and reported by the authors.
- **Abstract:** Adequate
- **Introduction:** This is adequate with current and relevant citations.
- **Methods:** Methods used were sufficient and reproducible by others.
- **Results** were well presented with Tables and Figures where appropriate. These were also properly captioned and can stand alone.
- **Discussion** projected the results and made appropriate reference to similar works by other researchers.
Replace part of lines 4 and 5 "Oxidative stress which leads to the production of free radicals in the body," with: Oxidative stress which occurs as a result of imbalance between production and accumulation of free radicals especially reactive oxygen species (ROS),
- **Conclusion** is adequate and reflected the findings of the study.
- I approve that the article be indexed.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Phytochemistry and microbiology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 17 September 2020

<https://doi.org/10.5256/f1000research.27806.r70661>

© 2020 Oloye A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abimbola Oloye

Department of Veterinary Surgery and Theriogenology, College of Veterinary Medicine, Federal University Of Agriculture Abeokuta, Ogun State, Nigeria

The work prescribed an ethno-medical alternative to treating oxidative stress and perhaps bacterial infections which are largely considered safer, easy to access, and cheaper when compared to most orthodox prescriptions.

The wideness of the spread of *H. arborescens* especially in southern and eastern Africa makes it fit for consideration as an alternative to some chemically compounded drugs.

The conclusions drawn about the leaf extracts of this plant by the authors, especially in establishing a relationship between the phenolic content of the extracts and their potencies as antioxidant and antibacterial, were appropriate. However, I wished the authors could give consideration to seasonal variation in concentration statuses of polyphenolics and other bioactive compounds of the plant's leaves. The month of June during which the study was carried out in the Eastern Cape, South Africa is the winter period, it would be revealing repeating the study in summer (between November and April) and drawing comparisons.

Studying the effect of blanching was a brilliant one. Indigenes mostly blanch their veggies in the process of making them into edibles or in preparing them as oral medicines for the locals. Blanching, as against chemical extraction, deactivates bioactive compounds hence reduces the

therapeutic effects associated with the plant. This has been implied in the study. Interestingly, however, the antibacterial effects of the plant seemed enhanced with blanching. The authors attributed this to enhanced extractive proficiency of water at boiling water temperature. In addition to this, however, I also think the heat-stable bioactive antibacterial factor should be identified and pinned down.

The authors should reconcile the justification for the study enumerated in the abstract with the one stated in the introductory section of the manuscript. The assertion that no literature enumerating the phytochemical contents, antioxidative and antibacterial properties of acetone, ethanol and aqueous exist may not be totally correct. There are few studies on these. (See Citations).^{1,2} However, they may be correct about the blanched extract. The justification in the introductory section of the main text appears more appropriate.

The concentrations of saponin and alkaloids determined in the study were said to be enough to elicit the bioactivities associated with the two compounds. Is there any empirical reference that corroborates this assertion especially considering the fact that the authors considered the alkaloid concentration as low?

A minor correction: The MIC of Ethanol extract against *P. aeruginosa* was 12.5 and this did not fall within the range of 1.563- 3.125 stated in the result. This, therefore, revealed an ethanol extract that had a weak antibacterial effect on *P. aeruginosa*. The authors should take a look at this.

References

1. McGaw L, Jäger A, van Staden J: Antibacterial, anthelmintic and anti-amoebic activity in South African medicinal plants. *Journal of Ethnopharmacology*. 2000; **72** (1-2): 247-263 [Publisher Full Text](#)
2. Maroyi A: HETEROMORPHA ARBORESCENS: A REVIEW OF ITS BOTANY, MEDICINAL USES, AND PHARMACOLOGICAL PROPERTIES. *Asian Journal of Pharmaceutical and Clinical Research*. 2018; **11** (11). [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Reproductive toxicology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research