REVIEW

Obesity and Bone [version 1; peer review: 2 approved]

Joseph Proietto

Department of Medicine, Austin Health, University of Melbourne, Repatriation Hospital, Heidelberg Heights 3081, Victoria, Australia

V1 First published: 09 Sep 2020, 9(Faculty Rev):1111
<https://doi.org/10.12688/f1000research.20875.1>
 Latest published: 09 Sep 2020, 9(Faculty Rev):1111
<https://doi.org/10.12688/f1000research.20875.1>

Abstract

Obesity and osteoporosis are both common conditions with high rates of morbidity and mortality. There is a relationship between obesity and bone. There are multiple factors that influence the risk of fracture, including the quality of bone, the risk of falls, and the padding around the bone. These multiple factors partly explain the finding that obesity protects against fractures in some sites while increasing the risk in other parts of the body. While it is well known that increased weight builds bone, there are several mechanisms related to the obese state that make the bone more fragile. These include the increased production of bone marrow fat cells at the expense of bone-forming osteoblasts, an increase in inflammatory cytokines leading to the activation of bone-resorbing osteoclasts, mutations in the *FTO* gene, and obesity-induced increased osteoblast senescence. Surprisingly, the relationship between bone and obesity is not unidirectional; there is now evidence that osteocytes are able to regulate body weight by acting as weighing machines.

Keywords

Osteoporosis, Body weight regulation, Inflammation, *FTO* gene, Bone marrow, Osteoblast, osteoclast, osteocyte

Open Peer Review

Approval Status

	1	2
version 1		
09 Sep 2020		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Maria Felicia Faienza**, University "A. Moro" of Bari, Bari, Italy
2. **Jasminka Z. Ilich**, Florida State University, Tallahassee, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Joseph Proietto (j.proietto@unimelb.edu.au)

Author roles: Proietto J: Data Curation

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Proietto J. This is an open access article distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Proietto J. **Obesity and Bone [version 1; peer review: 2 approved]** F1000Research 2020, 9(Faculty Rev):1111
<https://doi.org/10.12688/f1000research.20875.1>

First published: 09 Sep 2020, 9(Faculty Rev):1111 <https://doi.org/10.12688/f1000research.20875.1>

Introduction

Both obesity and osteoporosis are common conditions leading to increased morbidity and mortality. Cardiovascular disease, cancer, type 2 diabetes, obstructive sleep apnoea, osteoarthritis, psychosocial problems, hypertension, fatty liver, and infertility are also all caused by or aggravated by obesity. Of these, it has been shown that cardiovascular disease and cancer account for the highest mortality¹. Osteoporosis increases the risk of fractures and results in increased mortality² and morbidity, including increased pain and reduced mobility, giving a reduction in quality of life³. The very first link between obesity and bone was the common belief that an individual with obesity was heavier because he was “big boned”⁴. Although this is a common myth, science has revealed a surprising bidirectional relationship between excess fat and bone. Furthermore, Ilich and colleagues have proposed the existence of osteosarcopenic obesity linking obesity and osteoporosis to muscle weakness⁵. In their formulation, they discuss possible mechanisms, such as mesenchymal cell development, linking these three conditions. Osteosarcopenic obesity occurs in old age, but it has been shown that even preschool obesity is associated with an increase in fracture risk in teenage children⁶. This review will briefly discuss how excess fat damages bone and how bone contributes to the regulation of body weight.

How does obesity influence bone?

Weight-bearing exercise increases bone density⁷, so pressure from excessive weight may strengthen bone⁸. Consistent with this, significant weight loss achieved with bariatric surgery results in bone loss⁹, which can be mitigated by exercise¹⁰. However, despite more bone mass, obesity confers an increased risk of osteoporosis¹¹ and bone fractures¹², at least in some sites. Compston and colleagues¹³ followed 52,939 post-menopausal women for 3 years. During that time, 3,628 reported a fracture. BMI was protective for hip with a hazard ratio (HR) of 0.8 (95% confidence interval [CI] 0.71–0.90), spine with a HR of 0.83 (95% CI 0.76–0.92), and for wrist with a HR of 0.88 (95% CI 0.83–0.94) (HR with 95% CI for each increase in BMI of 5 kg/m²). However, excess weight increased the risk of ankle fractures (HR 1.05 [1.02–1.07]) ($P < 0.01$). For pelvic and rib fractures, they found a non-linear relationship with inverse associations at low BMI/body weight ($P = 0.05$) and positive associations at high values ($P = 0.03$). Interestingly, Liu and colleagues in a cross-sectional study involving 471 women divided into normal, overweight, and obese groups found that there is a threshold effect of fat on bone density. They concluded that while body fat below 33% has a positive relationship with bone density, for body fat above 33% the relationship between fat mass and bone is negative for most skeletal sites¹⁴. So extra weight builds bone and extra fat padding protects the bone during a fall. So why is there an increased rate of fracture in some sites? It appears that obesity causes bone to be of poor quality.

How can obesity damage bone?

Several mechanisms have been proposed for the deleterious effects of obesity on bone:

1. Replacement of osteoblasts by fat cells in bone marrow
2. Increased inflammation present in obesity
3. Mutations in the fat mass and obesity-associated (*FTO*) gene leading to weight gain and bone fragility
4. Increased metabolism and accelerated senescence in stromal stem cells

Replacement of osteoblasts by fat cells in bone marrow

Clinically, it has been found that situations leading to bone loss such as old age, thiazolidinedione treatment, and prolonged corticosterone treatment are associated with increased fat in bone marrow¹⁵. Adipocytes and osteoblasts are both derived from the same bone marrow-derived mesenchymal cell precursor (Figure 1). Increasing adipocyte differentiation decreases osteoblast differentiation and vice versa^{16,17}. The signals that cause the stem cells to differentiate into one or the other include the transcriptional activator PPAR γ , which can move the differentiation towards the adipocyte lineage¹⁸. Other triggers include leptin¹⁹ and the Wntless-related Integration site (Wnt) signalling glycoproteins. Wnt signalling is an ancient controller of cell fate determination, cell migration, and more during embryogenesis²⁰. It has been found that Wnt activation stimulates osteoblast formation and inhibits adipocyte differentiation. Secreted frizzled-related protein 1 (sFRP-1), an inhibitor of Wnt signalling, is increased in adipocyte stem cells²¹. sFRP-1 has been shown to be increased in mild obesity, leading to increased adipocyte formation, but interestingly is reduced in severe obesity, possibly to limit further adipocyte formation²².

Increased inflammation present in obesity

Obesity is now known to be a state of low-grade inflammation associated with elevated levels of cytokines^{23,24}. Cytokines have been shown to increase bone resorption. They do this primarily by enhancing osteoclast activity. Monocyte chemoattractant protein-1 (MCP-1), tumour necrosis factor (TNF)-related apoptosis-inducing ligand (TRAIL), receptor activator on nuclear factor κ B ligand (RANKL), interleukin 6 (IL-6), TNF α , and TNF super family 14 (TNFSF14)/LIGHT activate genes involved in the formation of bone-resorbing osteoclasts, as reviewed by Faienza and colleagues²⁵.

Mutations in the *FTO* gene leading to weight gain and bone fragility

The *FTO* gene has been linked to overweight and obesity²⁶. *FTO* is an RNA demethylase and has also been linked to increased risk of osteoporosis²⁷. Deletion of *FTO* in mice using the CRE/LOX system in osteoblasts leads to increased death of osteoblasts and bone loss. It has been shown that *FTO* functions to increase the stability of mRNA-encoding proteins that protect osteoblasts from genotoxic damage²⁸. Thus, those individuals whose obesity is caused by a mutation in the *FTO* gene are at increased risk of osteoporosis because of increased susceptibility of osteoblasts to cell death.

Increased metabolism and accelerated senescence in stromal stem cells

The next possible mechanism for obesity-related bone fragility is proposed obesity-induced hypermetabolism leading

Figure 1. Ontogeny of fat, osteoblast, and osteoclast cells in bone marrow.

to senescence in osteoblasts. Tencerova and colleagues²⁹ studied 54 healthy men whom they divided into lean (BMI $22.9 \pm 0.3 \text{ kg/m}^2$), overweight (BMI $28.0 \pm 0.4 \text{ kg/m}^2$), and obese (BMI $36.1 \pm 0.8 \text{ kg/m}^2$). Using DXA scanning, they showed that the men with obesity had an increase in total hip bone mineral density but no difference in lumbar spine compared with lean men. Subjects with obesity also had a reduction in bone turnover. Because the presence of inflammation leads to increased bone turnover, the authors proposed that this means that the bone marrow is protected from the generalised increase in inflammation seen in obesity. They obtained bone marrow-derived mesenchymal stem cells (BM-MSCs) from these men and performed RNA sequencing. They found that 129 genes were upregulated and 168 genes were downregulated in men with obesity compared to lean men. Gene ontology analysis showed that the genes that were upregulated in obesity are genes involved in cell differentiation while those that were downregulated were genes for glucose metabolism and response to hypoxia. Further analysis of glucose metabolism genes showed an increase in oxidative phosphorylation while genes for anaerobic glycolysis were downregulated. Interestingly, they also found upregulation of PPAR γ and other markers of differentiation into fat cells in obese compared with lean BM-MSC cultured cells. Obesity is known to be associated with insulin resistance in peripheral fat cells, muscle, and liver. In contrast, insulin resistance was not found in BM-MSCs from subjects with obesity. The BM-MSCs were then divided into those expressing the insulin and leptin receptors and those not expressing them. They showed that those expressing insulin and leptin receptors were more likely to differentiate into adipocytes while those that were negative for these receptors were more likely to differentiate into osteoblasts. It had previously been shown that osteoblast-like cells prefer glycolysis for ATP production while adipocyte-like cells prefer mitochondrial oxidative phosphorylation³⁰. High levels of oxidative phosphorylation

can increase the production of toxic reactive oxygen species (ROS), which may drive senescence³¹. When they compared insulin receptor- and leptin receptor-positive BM-MSCs from their patients, they found that these cells had an increase in the expression of senescence-associated genes when compared to cells that were negative for these receptors. Cultures of BM-MSCs from men with obesity contained a higher number of senescent cells and increased ROS production. It was concluded that enhanced insulin signalling and enrichment of adipocyte progenitor cells in the marrow of individuals with obesity lead to increased oxidative phosphorylation and increase ROS production, leading to a senescent phenotype, which results in poor bone formation.

How does bone influence body weight?

It appears that bone can assess body weight. Jansson and colleagues³² placed fluid-filled capsules into the abdomens of Sprague-Dawley rats and C57BL6 mice with diet-induced obesity to equal approximately 15% of their body weight. Control rodents had empty capsules inserted into their abdominal cavities (3% of their body weight). Increased loading suppressed body weight in both rats and mice with the difference in weight compared to control rodents already present from day 2 post-surgery. After 14 days of loading, 80% of the increased load was counteracted by reduced body weight. At the end of the experiment, body weight plus capsule weight was similar in the load and control rodents.

MRI imaging on day 14 showed that increased loading reduced the amount of white adipose tissue with no difference in brown adipose tissue. Not surprisingly, leptin levels were lower, consistent with the reduced adipose tissue mass. Weight loss was due to reduced food intake in both rats and mice. There were no changes in VO_2 max, respiratory quotient, or muscle activity, suggesting that all of the effects of capsule loading

were due to a reduction in energy intake. This was further confirmed by the fact that pair fed rodents decreased their body weight to the same extent as load rodents.

The effects of loading to suppress weight were shown to be independent of leptin when they showed that leptin-deficient ob/ob mice respond similarly to wild-type mice to capsule load. In any case, the reduction of leptin should have led to an increase in energy intake, not in the observed reduction in intake, so what is the signal?

It was known that osteocytes can sense dynamic short-term high-impact bone loading, so the authors hypothesised that osteocytes may also sense chronic static bone loading due to increased body weight. To test this hypothesis, mice deficient in osteocytes were loaded with the capsules. It was found that the control mice lost weight as shown before but the equally loaded osteocyte-deficient mice did not. So, load sensing is mitigated by the osteocyte.

The following question then arose: how does the osteocyte send the signal of increased load to the hypothalamus? In previous

studies, dynamic short-term high-impact loading was shown to reduce the expression of *Sost*, the gene coding for sclerostin. However, there was no difference in sclerostin levels. There was also no change in carboxylated and undercarboxylated forms of osteocalcin, fibroblast growth factor 23, and lipocalin 2. In addition, deficiency of ghrelin, glucagon-like peptide 1, the melanocortin 4 receptor, or the oestrogen receptor did not remove the effect. This leaves the possibility of a neuronal signal from osteocytes and the hypothalamus. To begin to investigate this, the authors measured urinary noradrenaline/creatinine ratio and serum choline level but found no differences. It was concluded that the osteocyte can sense chronic weight load and can send an unknown signal that can fairly accurately correct body weight.

Conclusion

There is a bidirectional complex relationship between obesity and bone. The advances in understanding how obesity leads to increased fragility in bone may lead to new treatments to improve bone strength. Finally, finding the signal that osteocytes send to the hypothalamus may lead to new approaches in the treatment of obesity.

References

1. Abdelaal M, Le Roux CW, Docherty NG: **Morbidity and mortality associated with obesity.** *Ann Transl Med.* 2017; **5**(7): 161.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
2. Bolland MJ, Grey AB, Gamble GD, et al.: **Effect of osteoporosis treatment on mortality: A meta-analysis.** *J Clin Endocrinol Metab.* 2010; **95**(3): 1174–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Lips P, van Schoor NM: **Quality of life in patients with osteoporosis.** *Osteoporos Int.* 2005; **16**(5): 447–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Stuart-Jones V: **Is 'obese' the new 'big-boned'?** *British Journal of Cardiac Nursing.* 2010; **5**: 361.
[Publisher Full Text](#)
5. Ilich JZ, Kelly OJ, Inglis JE, et al.: **Interrelationship among muscle, fat, and bone: Connecting the dots on cellular, hormonal, and whole body levels.** *Ageing Res Rev.* 2014; **15**: 51–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Lane JC, Butler KL, Poveda-Marina JL, et al.: **Preschool Obesity Is Associated With an Increased Risk of Childhood Fracture: A Longitudinal Cohort Study of 466,997 Children and Up to 11 Years of Follow-up in Catalonia, Spain.** *J Bone Miner Res.* 2020; **35**(6): 1022–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
7. Etherington J, Harris PA, Nandra D, et al.: **The effect of weight-bearing exercise on bone mineral density: A study of female ex-elite athletes and the general population.** *J Bone Miner Res.* 1996; **11**(9): 1333–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Felson DT, Zhang Y, Hannan MT, et al.: **Effects of weight and body mass index on bone mineral density in men and women: The Framingham study.** *J Bone Miner Res.* 1993; **8**(5): 567–73.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Ablett AD, Boyle BR, Avenell A: **Fractures in Adults After Weight Loss from Bariatric Surgery and Weight Management Programs for Obesity: Systematic Review and Meta-analysis.** *Obes Surg.* 2019; **29**(4): 1327–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
10. Murai IH, Roschel H, Dantas WS, et al.: **Exercise Mitigates Bone Loss in Women With Severe Obesity After Roux-en-Y Gastric Bypass: A Randomized Controlled Trial.** *J Clin Endocrinol Metab.* 2019; **104**(10): 4639–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
11. Greco EA, Fornari R, Rossi F, et al.: **Is obesity protective for osteoporosis? Evaluation of bone mineral density in individuals with high body mass index.** *Int J Clin Pract.* 2010; **64**(6): 817–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Kim KC, Shin DH, Lee SY, et al.: **Relation between obesity and bone mineral density and vertebral fractures in Korean postmenopausal women.** *Yonsei Med J.* 2010; **51**(6): 857–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Compston JE, Flahive J, Hosmer DW, et al.: **Relationship of Weight, Height, and Body Mass Index With Fracture Risk at Different Sites in Postmenopausal Women: The Global Longitudinal Study of Osteoporosis in Women (GLOW).** *J Bone Miner Res.* 2014; **29**(2): 487–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
14. Liu PY, Ilich JZ, Brummel-Smith K, et al.: **New insight into fat, muscle and bone relationship in women: Determining the threshold at which body fat assumes negative relationship with bone mineral density.** *Int J Prev Med.* 2014; **5**(11): 1452–63.
[PubMed Abstract](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
15. Abdallah BM, Kassem M: **New factors controlling the balance between osteoblastogenesis and adipogenesis.** *Bone.* 2012; **50**(2): 540–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Beresford JN, Bennett JH, Devlin C, et al.: **Evidence for an inverse relationship between the differentiation of adipocytic and osteogenic cells in rat marrow stromal cell cultures.** *J Cell Sci.* 1992; **102**(Pt 2): 341–51.
[PubMed Abstract](#)
17. Nuttall ME, Gimble JM: **Controlling the balance between osteoblastogenesis and adipogenesis and the consequent therapeutic implications.** *Curr Opin Pharmacol.* 2004; **4**(3): 290–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Siersbaek R, Nielsen R, Mandrup S: **PPARGgamma in adipocyte differentiation and metabolism--novel insights from genome-wide studies.** *FEBS Lett.* 2010; **584**(15): 3242–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Cirmanová V, Bayer M, Stárka L, et al.: **The effect of leptin on bone: An evolving concept of action.** *Physiol Res.* 2008; **57** Suppl 1: S143–51.
[PubMed Abstract](#)

20. Komiya Y, Habas R: **Wnt signal transduction pathways.** *Organogenesis*. 2014; **4**(2): 68–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Taipaleenmäki H, Abdallah BM, AlDahmash A, *et al.*: **Wnt signalling mediates the cross-talk between bone marrow derived pre-adipocytic and pre-osteoblastic cell populations.** *Exp Cell Res*. 2011; **317**(6): 745–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
22. Lagathu C, Christodoulides C, Tan CY, *et al.*: **Secreted frizzled-related protein 1 regulates adipose tissue expansion and is dysregulated in severe obesity.** *Int J Obes (Lond)*. 2010; **34**(12): 1695–705.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Hotamisligil GS, Arner P, Caro JF, *et al.*: **Increased adipose tissue expression of tumor necrosis factor-alpha in human obesity and insulin resistance.** *J Clin Invest*. 1995; **95**(5): 2409–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Ferrante AW: **Obesity-induced inflammation: A metabolic dialogue in the language of inflammation.** *J Intern Med*. 2007; **262**(4): 408–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Faienza MF, D'Amato G, Chiarito M, *et al.*: **Mechanisms Involved in Childhood Obesity-Related Bone Fragility.** *Front Endocrinol (Lausanne)*. 2019; **10**: 269.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
26. Frayling TM, Timpson NJ, Weedon MN, *et al.*: **A Common Variant in the FTO Gene Is Associated with Body Mass Index and Predisposes to Childhood and Adult Obesity.** *Science*. 2007; **316**(5826): 889–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
27. Guo Y, Liu H, Yang TL, *et al.*: **The fat mass and obesity associated gene, FTO, is also associated with osteoporosis phenotypes.** *PLoS One*. 2011; **6**(11): e27312.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Zhang Q, Riddle RC, Yang Q, *et al.*: **The RNA demethylase FTO is required for maintenance of bone mass and functions to protect osteoblasts from genotoxic damage.** *Proc Natl Acad Sci U S A*. 2019; **116**(36): 17980–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
29. Tencerova M, Frost M, Figeac F, *et al.*: **Obesity-Associated Hypermetabolism and Accelerated Senescence of Bone Marrow Stromal Stem Cells Suggest a Potential Mechanism for Bone Fragility.** *Cell Rep*. 2019; **27**(7): 2050–2062.e6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
30. Guntur AR, Gerencser AA, Le PT, *et al.*: **Osteoblast-like MC3T3-E1 Cells Prefer Glycolysis for ATP Production but Adipocyte-like 3T3-L1 Cells Prefer Oxidative Phosphorylation.** *J Bone Miner Res*. 2018; **33**(6): 1052–65.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
31. Manolagas SC: **From estrogen-centric to aging and oxidative stress: A revised perspective of the pathogenesis of osteoporosis.** *Endocr Rev*. 2010; **31**(3): 266–300.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Jansson JO, Palsdottir V, Hägg DA, *et al.*: **Body weight homeostat that regulates fat mass independently of leptin in rats and mice.** *Proc Natl Acad Sci U S A*. 2018; **115**(2): 427–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Jasminka Z. Ilich**

Institute for Successful Longevity, Florida State University, Tallahassee, FL, USA

Competing Interests: No competing interests were disclosed.

2. **Maria Felicia Faienza**

Pediatric Section, Department of Biomedical Sciences and Human Oncology, University "A. Moro" of Bari, Bari, Italy

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research