Check for updates

REVIEW

Novel therapeutic choices in immune aplastic anemia [version 1; peer review: 2 approved]

Phillip Scheinberg

Division of Hematology, Hospital A Beneficência Portuguesa, São Paulo, Brazil

V1 First published: 10 Sep 2020, 9(Faculty Rev):1118
<https://doi.org/10.12688/f1000research.22214.1>

Latest published: 10 Sep 2020, 9(Faculty Rev):1118
<https://doi.org/10.12688/f1000research.22214.1>

Abstract

Aplastic anemia (AA) in its severe form has historically been associated with high mortality. With limited supportive care and no effective strategy to reverse marrow failure, most patients diagnosed with severe AA (SAA) died of pancytopenia complications. Since the 1970s, hematopoietic stem cell transplantation (HSCT) and immunosuppressive therapy (IST) have changed SAA's natural history by improving marrow function and pancytopenia. Standard IST with horse anti-thymocyte globulin plus cyclosporine produces a hematologic response rate of 60 to 70%. In the long term, about one-third of patients relapse, and 10 to 15% can develop cytogenetic abnormalities. Outcomes with either HSCT or IST are similar, and choosing between these modalities relies on age, availability of a histocompatible donor, comorbidities, and patient preference. The introduction of eltrombopag, a thrombopoietin receptor agonist, improved SAA outcomes as both salvage (second-line) and upfront therapy combined with IST. As a single agent, eltrombopag in doses up to 150 mg daily improved cytopenias in 40 to 50% in those who failed initial IST, which associated with higher marrow cellularity, suggesting a pan-stimulatory marrow effect. When eltrombopag was combined with IST as upfront therapy, overall (about 90%) and complete responses (about 50%) were higher than observed extensively with IST alone of 65% and 10%, respectively. Not surprisingly, given the strong correlation between hematologic response rates and survival in SAA, most (>90%) were alive after a median follow-up of 18 months. Longer follow-up and real-world data continue to confirm the activity of this agent in AA. The use of eltrombopag in different combinations and doses are currently being explored. The activity of another thrombopoietin receptor agonist in AA, romiplostim, suggests a class effect. In the coming years, the mechanisms of their activity and the most optimal regimen are likely to be elucidated.

Keywords

aplastic anemia, bone marrow failure, pancytopenia, antithymocyte globulin

Open Peer Review

Approval Status

	1	2
version 1		
10 Sep 2020		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Carlo Dufour**, G. Gaslini Children's Research Hospital, Genova, Italy
2. **Maria A. Pereda**, UH Rainbow Babies & Children's Hospital, Cleveland, USA
Jignesh Dalal, UH Rainbow Babies & Children's Hospital, Cleveland, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Phillip Scheinberg (scheinbp@bp.org.br)

Author roles: Scheinberg P: Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: The author is a speaker and advisory at Novartis.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Scheinberg P. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Scheinberg P. **Novel therapeutic choices in immune aplastic anemia [version 1; peer review: 2 approved]** F1000Research 2020, 9(Faculty Rev):1118 <https://doi.org/10.12688/f1000research.22214.1>

First published: 10 Sep 2020, 9(Faculty Rev):1118 <https://doi.org/10.12688/f1000research.22214.1>

The initial treatment choice is an important decision point in the management of aplastic anemia (AA), especially in its severe form. The progression of pancytopenia can be gradual but often presents acutely in a patient who weeks earlier was otherwise doing well. In these cases, the clinical presentation is associated with complications of low blood counts, namely infections (from neutropenia), weakness (from anemia), or bleeding (from thrombocytopenia) or a combination of these. These patients often are admitted for transfusion support and evaluation. A hypocellular bone marrow points to marrow failure, and if no other causes are identified in the setting of severe pancytopenia, severe AA (SAA) is diagnosed. Exposures to drugs, chemicals, and toxins often are interrogated but rarely are the direct cause of marrow failure. A history of autoimmune disease or infections (viral) may suggest an association. But the most important differential diagnosis is with myelodysplastic syndrome, which can mimic AA^{1,2}.

Once AA is diagnosed, it is important to differentiate between congenital and immunologic forms of the disease. Younger age of onset (childhood), family history, and physical stigmata of distinct forms of hereditary AA (dyskeratosis congenita, Fanconi anemia, and Shwachman–Diamond syndrome) can be indicative of a hereditary problem³. Genetics, telomere length, and clastogenic testing often are applied in this evaluation. However, older age of onset, absence of family history, and physical stigmata do not rule out alterations in the telomere length or in the genetics, which are associated with hereditary syndromes^{4,5}. Nonetheless, most cases of AA, especially when severe, are immune-mediated and treated with either hematopoietic stem cell transplantation (HSCT) or immunosuppressive therapy (IST)⁶. The degree of pancytopenia defines the severity of AA⁷.

For younger (<40 years) patients with an HLA-matched sibling, allogeneic HSCT is preferred^{5,6}. HLA disparities and the fact that risks with HSCT increase with age are reasons why IST is preferred for patients who are older than 40 or who lack a histocompatible sibling (or both)⁸. In some settings where a matched unrelated donor (MUD) can be readily available (<6 weeks), a MUD HSCT can be considered as the first therapy in children⁹. However, the uncertainty of finding a histocompatible donor in registries (especially in ethnic minorities), the time for identifying such a donor (in many places this can be several months), and applying supportive care only for a patient with severe pancytopenia limit the generalizability of this approach. A haploidentical platform with post-transplant cyclophosphamide has shown encouraging results in younger patients, but experience so far has relied primarily on smaller retrospective studies with limited follow-up¹⁰. The optimal conditioning has not yet been defined, and to date, this approach is considered experimental and should be applied in a clinical protocol. In the aggregate, most cases of SAA worldwide are treated with non-transplant options because of the lack of a suitable HLA-matched donor, older age, comorbidities, patient choice, or lack of access to transplant. The standard IST consists of horse anti-thymocyte globulin (h-ATG) plus cyclosporine (CsA), which is associated with hematologic recovery and transfusion independence in 60 to 70% of cases¹¹.

Rabbit anti-thymocyte globulin (r-ATG) is more widely available, especially outside the US, but produces inferior outcomes; hematologic recoveries occur in about 30 to 40% of cases^{11,12}. Several attempts have been made to improve outcomes beyond h-ATG/CsA but were unsuccessful. The addition of a third drug to h-ATG/CsA, such as granulocyte colony-stimulating factor (G-CSF), mycophenolate mofetil, or sirolimus, did not improve outcomes, and substitution of h-ATG by r-ATG, cyclophosphamide, or alemtuzumab was equally ineffective because of inferior response rates or excessive toxicities or both¹. The strategy that has been more fruitful was not by intensifying immunosuppression but by adding a pan-stimulatory marrow agent of the class of the thrombopoietin receptor agonists (Tpo-RAs)¹³. This class is represented by eltrombopag, romiplostim, and avatrombopag, which were developed for immune thrombocytopenia. Most of the experience with Tpo-RA in marrow failure has been with eltrombopag, which also has approval in SAA.

About 10 years ago, the first trial of eltrombopag in SAA was initiated in patients who had failed first-line therapy with IST. In a 43-patient study, hematologic response was observed in about 40 to 50% of patients with bi- and tri-lineage count recoveries observed¹⁴. Eltrombopag, in these earlier studies, was given stepwise with the initial 50 mg dose that was escalated every 2 weeks up to 150 mg for a total of 16 weeks. Response usually was noted at the higher dose range. This initial cohort was expanded to 83 patients, and the latter 40 patients received eltrombopag at a starting dose of 150 mg from day 1 up to 6 months. In this 40-patient cohort, the response rate remained at about 50% but more multilineage (robust) recoveries were observed¹⁵. The possibility of discontinuation of eltrombopag was noted in the earlier trials, and the robustness of blood count recovery appeared to be associated with successful discontinuation without the need for further therapy. Relapses, when they occurred, were responsive to the resumption of eltrombopag. The mechanism by which eltrombopag was active in SAA was hypothesized to be via c-mpl (the Tpo-RA receptor) signaling and stimulation of early progenitor cells which expressed the receptor. *In vitro* and *in vivo* data supported this possibility. Of concern, however, has been the possible stimulation of abnormal clones in the marrow given the temporal association between the initiation of eltrombopag and the appearance of cytogenetic abnormalities (usually in the first 6 months)¹⁵. The cumulative incidence to date has not shown an increased rate of higher-risk clonal evolution, but longer follow-up is necessary to better define this risk. More recent data support that off-target effects might positively contribute to its immunomodulatory and marrow stimulatory effects¹⁶. Likely, the overall net effect of target and off-target mechanisms contributes to its effectiveness.

With the single-agent activity observed in second line, eltrombopag was added to standard h-ATG/CsA in front line in subsequent studies. In a three-cohort 92-patient study, eltrombopag was added to standard h-ATG/CsA in slightly different regimens to decrease toxicity and maximize effectiveness¹⁷. The combination was well tolerated and the better overall results were observed in the third cohort which had eltrombopag initiated concomitantly

with h-ATG/CsA on day 1 and continued for 6 months. In this group, the overall response rate was 94% and the complete response rate was 58%, which differ from the vast historical experience of h-ATG/CsA alone of an overall response rate of 60 to 70% and a complete response rate of about 10%. Overall survival after a median follow-up of 18 months was higher than 90 to 95% given the very intimate association between hematologic response and survival after IST in SAA⁷. Relapse in the upfront trial occurred at an unexpectedly higher rate (close to 50%), which nearly halfway into the study led to an amendment that allowed for continued long-term use of CsA beyond 6 months (instead of its discontinuation at 6 months). This led to a significant reduction of relapses to about 10 to 15%. The risk for clonal cytogenetic evolution was an important consideration in the design of the trial, prompting periodic bone marrow evaluations. Follow-up to fully determine this risk has been relatively short, although with a median follow-up of over 2 years to date, this risk has been in accordance with what has been observed historically with non-eltrombopag-containing

regimens in SAA⁵. These data led to the approval of eltrombopag in front line with IST with benefit primarily in the adult population⁵.

The strategy of stem cell stimulation with eltrombopag in SAA had more success than other approaches of intensifying immunosuppression or stimulation of more committed marrow progenitor cells (G-CSF, erythropoietin, and so on). Alternative mechanisms of action for eltrombopag—which includes immunomodulatory effects, evasion of inhibitory signals, and iron chelation which could be beneficial in SAA—are emerging^{16,18,19}. Studies outside the US have confirmed the activity of eltrombopag in SAA^{20–24}. More recently, alternative Tpo-RAs such as romiplostim have been investigated in AA as second-line therapy also showing activity^{25,26}. It is anticipated that romiplostim will be further developed in SAA. Markers of sustained response to eltrombopag could be useful in stratifying patients who are more likely to benefit from it^{27,28}. Longer-term follow-up and different combinations and doses are needed to better define the role and optimal delivery of the Tpo-RAs in AA in the future.

References

- Scheinberg P, Young NS: **How I treat acquired aplastic anemia.** *Blood.* 2012; **120**(6): 1185–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Durrani J, Maciejewski JP: **Idiopathic aplastic anemia vs hypocellular myelodysplastic syndrome.** *Hematology Am Soc Hematol Educ Program.* 2019; **2019**(1): 97–104.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Scheinberg P: **Recent Advances and Long-Term Results of Medical Treatment of Acquired Aplastic Anemia: Are Patients Cured?** *Hematol Oncol Clin North Am.* 2018; **32**(4): 609–18.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Calado RT, Young NS: **Telomere diseases.** *N Engl J Med.* 2009; **361**(24): 2353–65.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Young NS: **Aplastic Anemia.** *N Engl J Med.* 2018; **379**(17): 1643–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Bacigalupo A: **How I treat acquired aplastic anemia.** *Blood.* 2017; **129**(11): 1428–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
- Rosenfeld S, Follmann D, Nunez O, et al.: **Antithymocyte globulin and cyclosporine for severe aplastic anemia: Association between hematologic response and long-term outcome.** *JAMA.* 2003; **289**(9): 1130–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Giammarco S, de Latour RP, Sica S, et al.: **Transplant outcome for patients with acquired aplastic anemia over the age of 40: Has the outcome improved?** *Blood.* 2018; **131**(17): 1989–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
- Dufour C, Veys P, Carraro E, et al.: **Similar outcome of upfront-unrelated and matched sibling stem cell transplantation in idiopathic paediatric aplastic anaemia. A study on behalf of the UK Paediatric BMT Working Party, Paediatric Diseases Working Party and Severe Aplastic Anaemia Working Party of EBMT.** *Br J Haematol.* 2015; **171**(4): 585–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
- ElGohary G, El Fakih R, de Latour R, et al.: **Haploidentical hematopoietic stem cell transplantation in aplastic anemia: A systematic review and meta-analysis of clinical outcome on behalf of the severe aplastic anemia working party of the European group for blood and marrow transplantation (SAAWP of EBMT).** *Bone Marrow Transplant.* 2020.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
- Scheinberg P, Nunez O, Weinstein B, et al.: **Horse versus rabbit antithymocyte globulin in acquired aplastic anemia.** *N Engl J Med.* 2011; **365**(5): 430–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Marsh JC, Bacigalupo A, Schrezenmeier H, et al.: **Prospective study of rabbit antithymocyte globulin and cyclosporine for aplastic anemia from the EBMT Severe Aplastic Anaemia Working Party.** *Blood.* 2012; **119**(23): 5391–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Olmes MJ, Scheinberg P, Calvo KR, et al.: **Eltrombopag and improved hematopoiesis in refractory aplastic anemia.** *N Engl J Med.* 2012; **367**(1): 11–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Desmond R, Townsley DM, Dumitriu B, et al.: **Eltrombopag restores trilineage hematopoiesis in refractory severe aplastic anemia that can be sustained on discontinuation of drug.** *Blood.* 2014; **123**(12): 1818–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Winkler T, Fan X, Cooper J, et al.: **Treatment optimization and genomic outcomes in refractory severe aplastic anemia treated with eltrombopag.** *Blood.* 2019; **133**(24): 2575–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Scheinberg P: **Activity of eltrombopag in severe aplastic anemia.** *Blood Adv.* 2018; **2**(21): 3054–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Townsley DM, Scheinberg P, Winkler T, et al.: **Eltrombopag Added to Standard Immunosuppression for Aplastic Anemia.** *N Engl J Med.* 2017; **376**(16): 1540–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Zhao Z, Sun Q, Sokoll LJ, et al.: **Eltrombopag mobilizes iron in patients with aplastic anemia.** *Blood.* 2018; **131**(21): 2399–402.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Fattizzo B, Cavallaro F, Milesi G, et al.: **Iron mobilization in a real life cohort of aplastic anemia patients treated with eltrombopag.** *Am J Hematol.* 2019; **94**(9): E237–E239.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
- Ecsedi M, Lengline É, Knol-Bout C, et al.: **Use of eltrombopag in aplastic anemia in Europe.** *Ann Hematol.* 2019; **98**(6): 1341–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
- Lengline E, Drenou B, Peterlin P, et al.: **Nationwide survey on the use of**

- eltrombopag in patients with severe aplastic anemia: A report on behalf of the French Reference Center for Aplastic Anemia.** *Haematologica*. 2018; **103**(2): 212–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
22. Yamazaki H, Ohta K, Iida H, *et al.*: **Hematologic recovery induced by eltrombopag in Japanese patients with aplastic anemia refractory or intolerant to immunosuppressive therapy.** *Int J Hematol*. 2019; **110**(2): 187–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
  23. Hwang YY, Gill H, Chan TSY, *et al.*: **Eltrombopag in the management of aplastic anaemia: Real-world experience in a non-trial setting.** *Hematology*. 2018; **23**(7): 399–404.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
  24. Konishi A, Nakaya A, Fujita S, *et al.*: **Evaluation of eltrombopag in patients with aplastic anemia in real-world experience.** *Leuk Res Rep*. 2019; **11**: 11–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
  25. Scheinberg P: **Stem cell stimulation continues to pay off in aplastic anaemia.** *Lancet Haematol*. 2019; **6**(11): e543–e544.
[PubMed Abstract](#) | [Publisher Full Text](#)
  26. Lee JW, Lee SE, Jung CW, *et al.*: **Romiplostim in patients with refractory aplastic anaemia previously treated with immunosuppressive therapy: A dose-finding and long-term treatment phase 2 trial.** *Lancet Haematol*. 2019; **6**(11): e562–e572.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
  27. Zhao X, Feng X, Wu Z, *et al.*: **Persistent elevation of plasma thrombopoietin levels after treatment in severe aplastic anemia.** *Exp Hematol*. 2018; **58**: 39–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
  28. Fattizzo B, Kulasekararaj AG, Hill A, *et al.*: **Clinical and morphological predictors of outcome in older aplastic anemia patients treated with eltrombopag.** *Haematologica*. 2019; **104**(11): e494–e496.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Maria A. Pereda**

Pediatric Hematology and Oncology, UH Rainbow Babies & Children's Hospital, Cleveland, OH, USA

Jignesh Dalal

Pediatric Hematology and Oncology, UH Rainbow Babies & Children's Hospital, Cleveland, OH, USA

Competing Interests: No competing interests were disclosed.

2. **Carlo Dufour**

Hematology Unit, G. Gaslini Children's Research Hospital, Genova, Italy

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research