Check for updates

RESEARCH ARTICLE

Green tea influence on iron overload in thalassemia intermedia patients: a randomized controlled trial [version 1; peer review: 2 approved]

Hayder Al-Momen ¹, Hussein Khudhair Hussein¹, Zaid Al-Attar²,
Mohammed Jalal Hussein¹

¹Department of Pediatrics, Al-Kindy College of Medicine, University of Baghdad, Baghdad, 00964, Iraq

²Department of Pharmacology, Al-Kindy College of Medicine, University of Baghdad, Baghdad, 00964, Iraq

V1 First published: 16 Sep 2020, 9:1136
<https://doi.org/10.12688/f1000research.25553.1>

Latest published: 16 Sep 2020, 9:1136
<https://doi.org/10.12688/f1000research.25553.1>

Abstract

Background: Although iron chelation therapies have been available for many years for thalassemia intermedia patients, iron accumulation remains the major cause of death. Therefore, the need for additional chelation options is in demand. This randomized controlled study aimed to understand the effects of green tea on iron balance in thalassemia intermedia patients.

Methods: Using a random selection method, 141 thalassemia intermedia patients were initially screened for inclusion in this trial; only 68 patients included after applying exclusion criteria. Two equal groups were generated (n=34/group): green tea (three cups/day after meals) + usual treatment (deferasirox iron chelator and on demand blood transfusion); and control (only usual treatment). The study lasted for a period of 12 months. Patients failing to comply to the trial methodology were excluded, leaving a final total of 29 patients in the green tea group and 28 patients in the control group. Liver iron concentration, and serum ferritin were assessed at baseline and 12 months, while hemoglobin levels were assessed monthly.

Results: At baseline, both groups were matched regarding general demographics. At 12 months, the net drop of liver iron concentration in the green tea group (7.3 mg Fe/g dry weight) was significantly higher than the control group (4.6 mg Fe/g dry weight) (p<0.05). This was also seen with serum ferritin; net reduction in green tea and control groups were 1289 ng/ml and 871 ng/ml, respectively (p<0.05). Hemoglobin levels were slightly higher in the green tea group compared with the control group, but this was not significant.

Conclusions: Regular green tea consumption had a significant capability to improve iron deposition in thalassemia intermedia patients who already undergo deferasirox iron chelation therapy.

Trial registration: UMIN-CTR Clinical Trials Registry, [UMIN000040841](https://clinicaltrials.gov/ct2/show/study?term=UMIN000040841) (retrospectively registered June 21, 2020).

Open Peer Review

Approval Status

	1	2
version 1
16 Sep 2020	view	view
1. Alaa Alwan , Mustansiriyah University, Baghdad, Iraq		
2. Hassan Abolghasemi , Baqiyatallah University of Medical Sciences, Tehran, Iran		
Any reports and responses or comments on the article can be found at the end of the article.		

Keywords

Deferasirox, Liver iron concentration, Serum ferritin, Thalassemia.

This article is included in the **All trials matter** collection.

Corresponding author: Hayder Al-Momen (hayder77almusawi@gmail.com)

Author roles: **Al-Momen H:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Hussein HK:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Al-Attar Z:** Conceptualization, Formal Analysis, Funding Acquisition, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Hussein MJ:** Conceptualization, Formal Analysis, Investigation, Methodology, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Al-Momen H *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Al-Momen H, Hussein HK, Al-Attar Z and Hussein MJ. **Green tea influence on iron overload in thalassemia intermedia patients: a randomized controlled trial [version 1; peer review: 2 approved]** F1000Research 2020, 9:1136 <https://doi.org/10.12688/f1000research.25553.1>

First published: 16 Sep 2020, 9:1136 <https://doi.org/10.12688/f1000research.25553.1>

Introduction

Thalassemia intermedia (TI) is part of the thalassemia syndromes, which are the commonest genetic disorders worldwide, and is a member of the non-transfusion-dependent thalassemia (NTDT) group. By definition, β -thalassemia intermedia is a hemoglobin disorder characterized by a reduction in the expression of the β -globin gene^{1,2}. Iron overload in TI has more than one cause, the most important is increased gut absorption. In addition, ineffective erythropoiesis and hemolysis may increase the need for blood transfusions with the eventual result of iron overload^{3,4}. High morbidity rates usually accompany iron accumulation in NTDT and TI and has a significant impact on free radicals and oxidative stress tissue damage. Iron chelators are the best solution to alleviate these pathologies^{5,6}.

Deferasirox is the main oral iron chelation therapy according to international guidelines, but it is not free from side effects, leading to issues surrounding compliance, resulting in excessive body iron. Accordingly, recent research aims to assist, improve, or even substitute the use of iron chelating drugs using certain plants^{7,8}. The common tea plant *Camellia sinensis* has been utilized globally as a tea drink in various subtypes based on the manufacturing process: black (fermented), green (non-fermented), and oolong (semi-fermented). Tea belongs to the group of common beverages worldwide and it is considered as the number one beverage (after water) in Iraq and some other parts of the world^{9,10}.

Regarding green tea (GT), polyphenols are the main ingredient, representing 24–36% of dry weight. Catechins are the mainstay of these polyphenol compounds, which have inhibitory effects on iron (heme and non-heme) absorption across gut cells. This may lead to extraordinary negative iron balance in humans who consume GT as part of their daily diet^{11,12}.

Consequently, this randomized controlled trial aimed to assess the iron accumulation levels in TI patients treated with deferasirox with additional GT intake using serum ferritin (SF) and liver iron concentration (LIC). Also, hemoglobin (Hb) levels were assessed regularly.

Methods

Trial design

This is a randomized controlled (parallel-group; 1:1) study performed prospectively at the Baghdad Hereditary Anemia Center at Ibn Al-Baladi Hospital in Baghdad, Iraq, which is the largest thalassemia center in the country. The study was conducted over a period from 1st March, 2019 until 29th February, 2020.

Patients

All β -thalassemia intermedia patients aged ≥ 10 years old who were regularly managed by the Baghdad Hereditary Anemia Center were entered in a baseline screen. No sample size was calculated as all patients were considered.

Patients with chronic illnesses such as diabetes mellitus, heart failure, asthma, renal failure, liver disease, and any

hematological problem other than thalassemia, were excluded from the study. Patients that failed to have full compliance with blood transfusion and iron chelation therapy according to physicians' instructions in the last three months before baseline recruitment were also excluded.

At the initial screening, 141 patients were eligible. In total, 73 patients were excluded as they did not meet the inclusion criteria, leaving 68 patients who were randomly divided into two groups with matched age, sex, and BMI. Patients were asked to take part in the study during routine check-up appointments.

Outcomes

Iron overload status through measurements of SF and LIC at baseline and end of the study period, in addition to regular monthly assessment of Hb levels.

Interventions

The patients were randomly divided into two groups (n=34/group): GT group and control group.

GT groups: Patients in the GT group were instructed to drink a cup of GT after meals (within an hour) three times daily. Patients were given GT tea bags from Lipton™ containing 2 grams of GT (*Camellia sinensis*), which was collected from Kenya and Indonesia and made in January 2019, in Jebel Ali, United Arab Emirates by Unilever Gulf FZE (P.O. Box 17055, Dubai). Patients were instructed to infuse the tea bag in a hot cup of water (150°C) for 2–3 minutes, according to the manufacturer's instructions. Patients were undergoing deferasirox treatment and blood transfusion (when needed as decided by the attending physician) as their usual treatment.

Control group: Patients in the control group were instructed to drink one cup or more of either warm or cold drinking water after each meal, and to avoid any beverage other than water for three hours after each meal daily. Patients were undergoing deferasirox treatment and blood transfusion (when needed as decided by the attending physician) as their usual treatment.

Patients of both groups were asked to report their daily meals through a phone call (twice a week) to ensure low-iron food consumption, which might be considered as a confounding factor. Follow-up visits were performed at least every month with the patients attending physician.

During the study interval (12 months), participants failing to follow these instructions for three days a month (a total of nine drinking events during 30 days, which represents 10%) were excluded from analysis.

Randomization

Simple randomization was performed to allocate participants into GT and control groups in an equal pattern with matched age, sex, and BMI. Researchers have given members of each group a unique sequential number which was applied to all laboratory blood samples and MRI results. The researchers have enrolled and assigned participants to GT and control

groups. The attending physicians and staff of the laboratory and MRI units were blinded to the study groups.

Data collection

Full medical history and examination with the decision for blood transfusion requirement and iron chelation therapy (deferasirox 20–40 mg/kg/day) were made by the attending physician during each follow-up visit (at least once a month) during the study period. Body mass index (BMI (kg/m²)) was calculated at the start of the study for all patients. Also, packed red blood cells (RBC) transfusion volume during the previous year, calculated just before the start of study period, was estimated for all participants.

The following measurements were taken at baseline and 12 months for all patients:

- Serum ferritin (SF) readings (ng/ml), determined using a MINI VIDAS analyzer (bioMérieux S.A., Lyon, France) through an electrofluorescence assay method;
- Liver iron concentration (LIC) values (mg Fe/dry gram weight), determined using R2 magnetic resonance imaging (MRI; Siemens MRI machine with 3 Tesla and Medis suite software);
- Hemoglobin levels (g/dl) (documented monthly), determined using microprocessor hemoglobin meter (model number LE182, and manufactured by GPC Medical Ltd.)

Statistical analysis

Data are expressed as mean±standard deviation (SD) unless otherwise indicated. Shapiro-Wilk test was used for normal distribution. Pearson's chi-square test (for discrete or nominal variables) or Student t-tests (for continuous variables) were used to compare the characteristics between GT and control

groups. When $p < 0.05$, results were assumed to be significant. SPSS 22.0 for Windows (IBM Corp., NY, USA) was utilized in the analysis.

Ethical considerations

Ethical approval was obtained from Ethical and Scientific Committee of Al-Kindy College of Medicine at University of Baghdad (No.: 367). Baghdad Hereditary Anemia Center is affiliated with Al-Kindy College of Medicine at University of Baghdad. Written informed consent for participation was taken from all participants or their caregivers/legal guardians for those aged <18 years. All work was done according to the most recent version of the Declaration of Helsinki. The trial was registered retrospectively on UMIN-CTR Clinical Trials Registry [UMIN000040841](#) (June 21, 2020) according to the University of Baghdad recommendations.

Results

Due to non-compliance, the GT group had 29 patients (five patients excluded) and the control group had 28 patients (six patients excluded) at the end of the study (Figure 1).

The two groups had comparable (non-significant) general demographic data at baseline (Table 1). Iron overload (measured by SF or LIC) and volume of packed red blood cells (RBC) transfusion during the previous year calculated just before the start of study period showed no significant differences between groups. Deferasirox mean doses for the GT (36.89 ± 7.24 mg/kg/day) and control (37.21 ± 1.13 mg/kg/day) groups were high and near to the maximum approved doses, but were not significantly different between the groups.

LIC was significantly reduced ($P < 0.05$) after 12 months in both the GT group (baseline, 19.4 ± 2.3 Fe/gram/dry weight;

Figure 1. Flowchart of participants.

Table 1. Participant characteristics at baseline.

Variable	Green tea group (n=29)	Control group (n=28)	P value
Age, years (mean±SD)	13.62±4.27	13.47±2.58	0.81 ^a
Male:female	15:14 (51.72%:48.28%)	14:14 (50%)	
Body mass index, kg/m ² (mean±SD)	18.45±4.91	19.11±3.40	0.63 ^a
Deferasirox dose, mg/kg/day (mean±SD)	36.89±7.24	37.21±1.13	0.75 ^a
Pre-transfusion, hemoglobin g/dl (mean±SD)	8.21±3.36	8.16±4.15	0.86 ^a
Packed red blood cell transfusion volume during the year of pre-study period, ml/kg (mean±SD)	71.84±9.97	73.42±9.53	0.41 ^a
Splenectomized, n (%)	7 (24.14)	7 (25.07)	0.90 ^b
Serum ferritin, ng/mL (mean±SD)	2617±549	2684±164	0.57 ^a
Liver iron concentration, mg Fe/g dry weight (mean±SD)	19.4±2.3	19.8±1.4	0.84 ^a

a: Student t-tests, b: Pearson's chi-square test.

12 months, 12.1±1.3 Fe/gram/dry weight), and control groups (baseline, 19.8±1.7 Fe/gram/dry weight; 12 months, 15.2±1.6 Fe/gram/dry weight) (Figure 2). Similarly, SF levels were significantly decreased at the end of study period ($P<0.05$) in the GT (baseline, 2617±549 ng/ml; 12 months, 1328±127 ng/ml) and control (baseline, 2684±164 ng/ml; 12 months, 1813±235 ng/ml) groups.

Deferasirox use was comparable at the start and end of the study period for both the GT (baseline, 36.89±7.24 mg/kg/day; 12 months, 36.24±6.83 mg/kg/day) and control (baseline, 37.21±1.13 mg/kg/day; 12 months, 37.15±3.78 mg/kg/day) groups.

The mean volume of transfused packed RBC during the study period was estimated after 12 months from the baseline and revealed no statistical significance between GT (70.64±7.84 ml/kg) and control (72.14±8.35 ml/kg) patients. The net reduction of LIC (Fe/gram/dry weight) in GT patients was 7.3 (baseline, 19.4; 12 months, 12.1), while for control patients the net result was 4.6 (baseline, 19.8; 12 months, 15.2). A significant reduction ($P<0.05$) of LIC was found in the GT group in comparison with the control cases. The drop of serum ferritin in GT patients after 12 months [1289 ng/ml (2617-1328)] was significantly ($P<0.05$) higher than that of the control group [871 ng/ml (2684-1813)] (Table 2).

Hemoglobin readings gradually increased in both the GT and control groups over the study period. GT patients had higher Hb levels after 12 months than the control group (8.4 versus 8.2), but this was not statistically significant ($P=0.37$) (Figure 3).

Discussion

Iron overload is a major consequence of TI. Physicians have several challenges when using medications that eliminate

excessive iron, including type, dose, expected adverse effects, and the best time to start an iron chelator. Therefore, the need for a naturally mediated iron chelator based on plants is increasing along with the increase of thalassemia patients around the world¹³⁻¹⁵. Green tea (GT) seems to be one of the best available options due to its natural properties to counteract iron absorption in addition to other promising advantages^{16,17}.

There are several ways to estimate the overload of iron in thalassemia patients, such as the widely available test of SF, and the most accurate non-invasive recent method of LIC using the R2 MRI technique^{18,19}.

Patients with TI usually have high iron content loading in their tissues as evidenced by other studies from different areas of the world^{13,20,21}. Our sample of patient had even higher iron content, which is in line with other reports from Middle East^{3,14}. This might be due to compliance deterioration, background cultural beliefs, possible side effects of iron chelating drugs, and the frequent use of mimic medications in this part of the world²².

GT consumption in our sample was associated with decreased ferritin levels mirroring better iron chelation. This was also noticed by other works done on both humans and animals with transfusion dependent thalassemia²³⁻²⁵. SF could be affected by the antioxidant properties of GT^{26,27}. However, a recent study that added curcumin to GT for thalassemia patients discovered a slight increase of SF after two months, with the reasons for this unknown to the authors²⁸.

A more sophisticated method of iron accumulation assessment in TI is LIC, and in this study, we found a significant drop in the GT group compared to the control group. This has

Figure 2. Liver iron concentration (LIC) at the start and end of study period in the green tea (GT) and control groups.

Table 2. Relationship between specific events within the study groups.

Event	Green tea group (n=29)	Control group (n=28)	P value
Packed red blood cell transfusion volume (ml/kg) over 12 months (mean±SD)	70.64±7.84	72.14±8.35	0.37 ^a
Net drop of mean liver iron concentration (mg Fe/g dry weight) at the end of the study	7.3	4.6	< 0.05 ^a
Net drop of mean serum ferritin (ng/ml) at the end of the study	1289	871	< 0.05 ^a

^a: Student t-tests.

only been previously reported in animal studies, and an online search conducted by the present authors failed to reveal human trials¹⁵. It should be noted that iron deficiency has been documented in healthy individuals consuming large amounts of GT²⁹.

Surprisingly, in the GT group, TI patients developed slightly higher hemoglobin readings than controls. Again, no previously published data was found to verify this, but it might be related to better iron removal and effects of antioxidant characteristics

of GT, which could be associated with improved hematopoiesis and less red blood cells hemolysis. However, GT is historically associated with anemia in a healthy population, especially in high doses³⁰.

Although this trial was conducted in the largest thalassemia center within Iraq country borders, its small sample size could be considered as a limitation. We recommend other interested scientists throughout the world to involve more patients with more variables and risk factors to deeply investigate the effects

Figure 3. Hemoglobin (Hb) levels in the green tea (GT) and control groups throughout the study (measured monthly).

of GT in TI and discover more plant-based iron chelation therapies with better effects.

Conclusion

Thalassemia intermedia patients having usual medical care, including blood transfusion and deferasirox oral iron chelation therapy, have significantly superior levels of iron chelation and slightly better hemoglobin levels when consuming GT regularly compared with a control group.

Data availability

Underlying data

Mendeley Data: Green tea influence on iron overload in thalassemia intermedia, <http://dx.doi.org/10.17632/p7558gcrmw.2>³¹.

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

Reporting guidelines

Mendeley Data: CONSORT checklist for 'Green tea influence on iron overload in thalassemia intermedia patients: a randomized controlled trial', <http://dx.doi.org/10.17632/9c5bx4p65c.1>³².

Acknowledgements

Many thanks for the enormous encouragements, and great deal of support offered by the medical and management staff of Baghdad Hereditary Anemia Center at Ibn Al-Baladi Hospital.

References

- Ben Salah N, Bou-Fakhredin R, Mellouli F, *et al.*: **Revisiting beta thalassemia intermedia: past, present, and future prospects.** *Hematology*. 2017; **22**(10): 607–16.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Asadov C, Alimirzoeva Z, Mammadova T, *et al.*: **β -Thalassemia intermedia: a comprehensive overview and novel approaches.** *Int J Hematol*. 2018; **108**(1): 5–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Yassin MA, Soliman AT, De Sanctis V, *et al.*: **Final height and endocrine complications in patients with β -thalassemia intermedia: our experience in non-transfused versus infrequently transfused patients and correlations with liver iron content.** *Mediterr J Hematol Infect Dis*. 2019; **11**(1): e2019026.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Vichinsky E: **Non-transfusion-dependent thalassemia and thalassemia intermedia: epidemiology, complications, and management.** *Curr Med Res Opin*. 2016; **32**(1): 191–204.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Al-Momen H: **Iron chelation therapy in sickle cell/beta thalassemia syndrome, a 2 years' extension study.** *Al-Kindy College Medical Journal*. 2017; **13**(1): 76–81.
[Reference Source](#)
- Kontoghiorghe CN, Kontoghiorghe GJ: **Efficacy and safety of iron-chelation therapy with deferoxamine, deferiprone, and deferasirox for the treatment of iron-loaded patients with non-transfusion-dependent thalassemia syndromes.** *Drug Des Devel Ther*. 2016; **10**: 465–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Daher AM, Al-Momen H, Jasim SK: **Deferasirox in thalassemia: a comparative study between an innovator drug and its copy among a sample of Iraqi patients.** *Ther Adv Drug Saf*. 2019; **10**: 2042098619880123.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

8. Moayed B, Gharagozloo M, Esmaeil N, *et al.*: **A randomized double-blind, placebo-controlled study of therapeutic effects of silymarin in β -thalassemia major patients receiving desferrioxamine.** *Eur J Haematol.* 2013; **90**(3): 202–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Jasim SK, Al-Momen H, Al-Asadi F: **Maternal Anemia Prevalence and Subsequent Neonatal Complications in Iraq.** *Open Access Maced J Med Sci.* 2020; **8**(B): 71–5.
[Reference Source](#)
10. Shafik SS, Rejah BK: **Measurement of the uranium concentration in different types of tea used in Iraqi kitchen.** *Iraqi Journal of Science.* 2014; **55**(3A): 1039–43.
[Reference Source](#)
11. Xing L, Zhang H, Qi R, *et al.*: **Recent advances in the understanding of the health benefits and molecular mechanisms associated with green tea polyphenols.** *J Agric Food Chem.* 2019; **67**(4): 1029–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Nawab A, Farooq N: **Review on green tea constituents and its negative effects.** *J Pharm Innov.* 2015; **4**(1): 21–4.
[Reference Source](#)
13. Taher AT, Porter JB, Viprakasit V, *et al.*: **Defining serum ferritin thresholds to predict clinically relevant liver iron concentrations for guiding deferasirox therapy when MRI is unavailable in patients with non-transfusion-dependent thalassaemia.** *Br J Haematol.* 2015; **168**(2): 284–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Al-Momen H, Jasim SK, Hassan QA, *et al.*: **Relationship between liver iron concentration determined by R2-MRI, serum ferritin, and liver enzymes in patients with thalassemia intermedia.** *Blood Res.* 2018; **53**(4): 314–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Saewong T, Ounjaijean S, Munde Y, *et al.*: **Effects of green tea on iron accumulation and oxidative stress in livers of iron-challenged thalassemic mice.** *Med Chem.* 2010; **6**(2): 57–64.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Chacko SM, Thambi PT, Kuttan R, *et al.*: **Beneficial effects of green tea: a literature review.** *Chin Med.* 2010; **5**: 13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Ke X, Qin N, Zhang T, *et al.*: **Highly Augmented Antioxidant and Anticancer Effect of Biocompatible MIL-100 (Fe)@ SiO 2-Immobilized Green Tea Catechin.** *J Inorg Organomet Polym Mater.* 2020; **30**(3): 935–42.
[Publisher Full Text](#)
18. Al-Momen HH, Obed AA, Mahdi ZK, *et al.*: **Isdeferiasirox as Effective as Desferrioxamine in Treatment of Iron Overload in Patients with Thalassemia Major?**
19. Bharadwaj NS, Jain R, Bhatia P, *et al.*: **Assessment of Iron Overload and Role of Genetic Modifiers in Patients with Beta Thalassemia Intermedia.** *Pediatric Hematology Oncology Journal.* 2019; **4**(2): S39.
20. Saliba AN, Musallam KM, Cappellini MD, *et al.*: **Serum ferritin values between 300 and 800 ng/mL in nontransfusion-dependent thalassemia: A probability curve to guide clinical decision making when MRI is unavailable.** *Am J Hematol.* 2017; **92**(3): E35–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Sarhan MM, El Asmy AE, Mohamed GM: **Magnetic Resonance Imaging Versus Serum Ferritin Levels in Detection of Liver and Cardiac Iron Overload in Non-Transfusion Dependent Thalassemia.** *Pediatr Therapeut.* 2017; **7**(03): 10–4.
[Publisher Full Text](#)
22. Mattar M, Alwan AF, Boukhalal H, *et al.*: **On the use of substandard medicines in hematology: An emerging concern in the Middle East and North Africa region.** *Eur J Intern Med.* 2018; **48**: e40–1.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Ounjaijean S, Thephinlap C, Khansuwan U, *et al.*: **Effect of green tea on iron status and oxidative stress in iron-loaded rats.** *Med Chem.* 2008; **4**(4): 365–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Badiee MS, Nili-Ahmadabadi H, Zeinvand-Lorestani H, *et al.*: **Green tea consumption improves the therapeutic efficacy of deferoxamine on iron overload in patients with (Beta)-thalassemia major: a randomized clinical study.** *InBiological Forum.* 2015; **7**(2): 383. [Research Trend](#).
[Reference Source](#)
25. Soeizi E, Rafrat M, Asghari-Jafarabadi M, *et al.*: **Effects of Green Tea on Serum Iron Parameters and Antioxidant Status in Patients with β -Thalassemia Major.** *Pharm Sci.* 2017; **23**(1): 27–36.
[Publisher Full Text](#)
26. Bou-Abdallah F, Paliakkara JJ, Melman G, *et al.*: **Reductive mobilization of iron from intact ferritin: mechanisms and physiological implication.** *Pharmaceuticals (Basel).* 2018; **11**(4): 120.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Peluso I, Serafini M: **Antioxidants from black and green tea From dietary modulation of oxidative stress to pharmacological mechanisms.** *Br J Pharmacol.* 2017; **174**(11): 1195–208.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Koonyosying P, Tantiworawit A, Hantrakool S, *et al.*: **Consumption of a green tea extract-cucurmin drink decreases blood urea nitrogen and redox iron in β -thalassemia patients.** *Food Funct.* 2020; **11**(1): 932–943.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Fan FS: **Iron deficiency anemia due to excessive green tea drinking.** *Clin case Rep.* 2016; **4**(11): 1053–1056.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Rani S, Rasyid R, Desmawati D: **High intake of green tea decreased hemoglobin and hematocrit levels in Rattus novergicus strain wistar albino.** *International Journal of Research in Medical Sciences.* 2018; **6**(11): 3688.
[Publisher Full Text](#)
31. Al-Momen H, Hussein KH, Al-Attar Z, *et al.*: **“Green tea influence on iron overload in thalassemia intermedia ”.** Mendeley Data, v2, 2020.
<http://www.doi.org/10.17632/p7558gcrmw.2>
32. Al-Momen H, Al-Musawi H, Al-Attar Z, *et al.*: **CONSORT Checklist for (Green tea influence on iron overload in thalassemia intermedia patients).** Mendeley Data, V1, 2020.
<http://www.doi.org/10.17632/9c5bx4p65c.1>

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 25 September 2020

<https://doi.org/10.5256/f1000research.28201.r71683>

© 2020 Abolghasemi H. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Hassan Abolghasemi

Faculty of Medicine, Baqiyatallah University of Medical Sciences, Tehran, Iran

In this article the impact of green tea on reduction of liver iron concentration have been studied in a randomized clinical trials by authors. They showed liver iron was decreased in the case groups compared with control group significantly. The level of hemoglobin was not significantly changed at the end of trial.

The trial was done appropriately.

The only problem with the study was the level of ferritin at the start of trial which was higher than usual for thalassemia intermedia and the dose of Deferasirox. It seems that some of the patients were transfusion dependent.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: pediatric hematology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 24 September 2020

<https://doi.org/10.5256/f1000research.28201.r71684>

© 2020 Alwan A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Alaa Alwan

Department of Clinical Hematology, National Center of Hematology, Mustansiriyah University, Baghdad, Iraq

- It is interested subject which has covered important issue in beta thalassemia.
- The introduction is well covered with recent references.
- Materials and method done appropriately especially with use of statistical methods and randomization.
- Regarding the results, they are clear and in my opinion it is obvious.
- The discussion part is well written and it is good to see comparison with other studies.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: benign and malignant hematology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research