CASE REPORT

Case Report: Complete heart block as a manifestation of cardiac metastasis of oral cancer [version 1; peer review: 1 approved with reservations]

Andrianto Andrianto , Eka Prasetya Budi Mulia , Denny Suwanto, Dita Aulia Rachmi, Mohammad Yogiarto

Department of Cardiology and Vascular Medicine, Airlangga University - Dr. Soetomo General Hospital, Surabaya, East Java, 60286, Indonesia

V1 First published: 15 Oct 2020, 9:1243
<https://doi.org/10.12688/f1000research.26438.1>
Latest published: 11 Nov 2020, 9:1243
<https://doi.org/10.12688/f1000research.26438.2>

Abstract

Metastatic tumors of the heart presenting with complete heart block (CHB) is an extremely uncommon case. There are no available guidelines in managing CHB in terminal cancer. Permanent pacemaker implantation in such cases is a challenge in terms of clinical utility and palliative care.

We report a case of a 24-year-old man suffering from tongue cancer presenting with CHB. A intracardiac mass and moderate pericardial effusion were present, presumed as the metastatic tumor of tongue cancer. We implanted a temporary pacemaker for his symptomatic heart block and cardiogenic shock, and pericardiocentesis for his massive pericardial effusion. We decided that a permanent pacemaker would not be implanted based on the low survival rate and significant comorbidities.

Multiple studies report a variable number of cardiac metastasis incidence ranging from 2.3% to 18.3%. It is rare for such malignancies to present with CHB. The decision to implant a permanent pacemaker is highly specific based on the risks and benefits of each patient. It needs to be tailored to the patient's functional status, comorbid diseases, prognosis, and response to conservative management.

Keywords

tongue cancer, cardiac metastasis, complete heart block, case report

Open Peer Review

Approval Status

	1	2
version 2 (revision) 11 Nov 2020	 view	 view
version 1 15 Oct 2020	 view	

1. **Radityo Prakoso** , Universitas Indonesia, Jakarta, Indonesia
2. **Pieter A.F.M. Doevendans** , Universitas Indonesia, Jakarta, Indonesia

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Andrianto Andrianto (andrianto@fk.unair.ac.id)

Author roles: **Andrianto A:** Conceptualization, Funding Acquisition, Writing – Review & Editing; **Mulia EPB:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Suwanto D:** Data Curation, Formal Analysis, Funding Acquisition, Investigation, Resources, Software, Writing – Original Draft Preparation; **Rachmi DA:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Yogiarto M:** Funding Acquisition, Investigation, Resources, Supervision, Validation, Visualization

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Andrianto A *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Andrianto A, Mulia EPB, Suwanto D *et al.* **Case Report: Complete heart block as a manifestation of cardiac metastasis of oral cancer [version 1; peer review: 1 approved with reservations]** F1000Research 2020, 9:1243 <https://doi.org/10.12688/f1000research.26438.1>

First published: 15 Oct 2020, 9:1243 <https://doi.org/10.12688/f1000research.26438.1>

Introduction

Cardiac metastasis is the least common presentation in malignant cancer. Primary cardiac tumors are also rare (on postmortem analysis commonly between 0.01% to 0.1%). However, the frequency of secondary metastatic tumors to the pericardium, myocardium, great vessels, and coronary arteries are between 0.7% to 3.5% in the general population and up to 9.1% in patients with a history of malignancies¹.

Complete heart block (CHB) as the primary clinical presentation of heart metastasis is very unusual². There are currently no guidelines for the management of CHB in terminal stage of cancer.

We report a case of CHB caused by cardiac metastasis and review the literature to further help the management of our patient.

Case presentation

A 24-year old Asian man was admitted the cardiology department with CHB and hypotension. The patient was a chef and

had a history of tongue cancer for six months and had undergone 30 cycles of radiotherapy. A week before, the patient came to the emergency department (ED) because of oral bleeding and general weakness. The patient denied any history of cardiovascular disease.

The patient was presented with chest discomfort and general weakness. He was hypotensive and bradycardic with blood pressure of 80/40 mmHg, regular heart rate of 44 beats per minute, respiratory rate of 18 breaths per minute, and oxygen saturation of 97% on room air. Chest auscultation was clear, and no murmurs heard. Electrocardiogram showed CHB with a junctional escape rhythm at 44 bpm (Figure 1). Echocardiography showed normal left ventricle kinetic, normal left ventricular ejection fraction (62%), and normal right ventricle systolic function. There were moderate pericardial effusion and intracardiac masses (2.1 × 0.9 cm and 1.8 × 0.8 cm) in the right atrial and septal leaflet of tricuspid. Hyperechoic areas in the annulus of tricuspid, lateral wall of right atrium and right ventricle, and interventricular septum were also

Figure 1. Electrocardiography on the first day of consultation showed complete heart block.

found in an echocardiogram (Figure 2 and Figure 3). Laboratory finding revealed anemia (hemoglobin 8.5 g/dL; normal range 13.3-16.6 g/dL), leukocytosis (white blood count 18,470/mL; normal range 3,370-10,000/mL), hypoalbuminemia (albumin 2.6 g/dL; normal range 3.4-5.0 g/dL), hypokalemia (potassium 3.4 mmol/L; normal range 3.5-5.1 mmol/L), and hypercalcemia (calcium 16.2 mg/dL, corrected calcium 16.9 mg/dL; normal range 8.6-10.3 mg/dL).

Figure 2. Echocardiography showed the presence of a mass in the right atrium right atrial and septal leaflet of tricuspid (yellow arrow). Hyperechoic areas were found in the annulus of tricuspid, lateral wall of right atrium and right ventricle, and interventricular septum (red arrows).

Previous magnetic resonance imaging (MRI) of head and neck, six months before this admission, revealed malignant tongue mass (staging AJCC 2010 of lip and oral cavity mass: T4N1Mx) and bilateral nasal cavity thickening. Multi-slice computed tomography (CT) scan of the head revealed an enhancing mass 1×1.3×1.5 cm at the base of the tongue and multiple lymph node enlargements subcentimeter in the upper and lower paratracheal. Histopathology examination of tongue biopsy confirmed poorly differentiated squamous cell carcinoma.

A temporary pacemaker was immediately implanted as the patient showed symptomatic heart block and cardiogenic shock. CHB persisted despite corrected electrolyte imbalance. On the 14th day of admission, the patient developed pleural effusion, and worsening pericardial effusion with echocardiogram showed massive pericardial effusion and sign of tamponade. Further chest X-ray evaluation on the 14th day of treatment (which includes electrolyte imbalance correction, supportive treatment of general weakness condition, anemia, hypoalbuminemia, and infection) showed left parahilar ground glass appearance with suspicions of lung metastasis and pleural effusion (Figure 4). Pericardiocentesis was then performed with pericardial fluid showing hemorrhagic typical for malignant disease.

Because the patient showed general improvement after supportive treatment for seven days and no symptoms of CHB, it was decided that the temporary pacemaker would be extracted. Considering the poor prognosis of this cancer, risk

Figure 3. Pericardial effusion was found in (A) anterior, posterior, (B) inferior, (C) base, and (D) left-lateral of the heart.

Figure 4. Chest X-ray on the 14th day of treatment showed left parahilar ground glass appearance with suspicion of lung metastasis, pleural effusion. Pericardial fluid pigtail was already inserted for drainage of pericardial effusion.

of permanent pacemaker (PPM) implantation, and severe comorbidities, we decided not to implant a PPM after acquiring the patient's and his family's consent. The patient died due to respiratory failure and septic shock 20 days after showing the first symptoms of cardiac metastasis.

Discussion

From the literature, we acquired a total of 14 articles regarding heart block as a manifestation of cardiac metastasis, as summarized in Table 1²⁻¹⁵. Oral cavity, uterus, and thyroid are the most common primary cancers that metastasize to the heart. Squamous cell carcinoma was the primary histologic finding. Heart metastasize may be present with clinically silent symptoms to an alarming presentation of hemoptysis and syncope. Locations of metastasis were mostly in the right ventricle, supporting the hypothesis of the hematologic spread of cancer cells. PPM implantation was performed in 10 cases, yet only one case reported a significant lifespan after PPM implantation.

The prevalence of cardiac metastasis, in general, is arguably low. However, multiple studies reported a variable number of cardiac metastasis incidence ranging from 2.3% to 18.3%. The prevalence of malignancy originating from the oral cavity was 5.3%. The involvement of pericardium made up two-thirds of all cardiac metastasis. Myocardium and endocardium involvement each made up one-third of all cardiac metastasis. Only 5% involved the endocardium. The most common site of metastasis for squamous cell carcinomas is epicardium (41.4%)¹⁶.

Myocardial infiltration by cancer cells may present with arrhythmias, such as atrial flutter or fibrillation, premature beats, or ventricular arrhythmias. Conduction system involvement may induce various degree of atrioventricular blocks¹⁶.

The presence of right atrial mass in our patient supports the possibility of hematologic spreading of metastatic cancer cells into the endocardium. Pericardial effusion may represent metastasize or inflammatory reaction toward the malignancy. The presence of CHB suggests the infiltration of the heart conduction system.

Valves are an uncommon site for metastasis because of the absence of vessels in the physiological valvular stroma and the constant cusp motion. Bussani *et al.* reported, out of over a thousand of postmortem examination, there was only one case of valve involvement¹⁶. The mass in the septal leaflet of tricuspid valves appeared in echocardiography examination in this patient showed valve involvement of cardiac metastasis.

Asymptomatic in the early stages, cardiac metastasis could lead to a wide range of signs and symptoms, such as cardiac failure, conduction disturbances, angina, and pain as it progresses. Disruption of the cardiac conduction system by cardiac metastases can lead to lethal arrhythmias, including atrial fibrillation with a rapid ventricular response, CHB, or ventricular fibrillation¹. From the literature, we acquired 14 cases reporting CHB as a manifestation of cardiac metastasis originating from various malignancies, three of which are from the oral cavity.

Cardiac masses in our case presented with features favoring tumor, such as echo density similar to myocardium, normal wall motion, valvular lesion, no history suggestive of coronary artery disease, and a clinical history of oral cancer as primary site suspected to metastasize to the heart.

Cardiac MRI is the best imaging modality and, along with positron emission tomography scanning, are mostly used in investigating the extent of infiltration by malignant cells^{2,17}. Patients with specific cardiac devices, such as pacemaker and defibrillators, would be disqualified from undergoing MRI, an important consideration given the frequency at which arrhythmia complicates cardiac metastasis¹⁷. The temporary pacemaker implanted during the early stages of CHB made MRI impractical for our patient.

In our patient, CHB was initially thought to be the result of electrolyte imbalance. However, as the electrolyte was restored to its normal level without any improvement of CHB, it suggested that CHB was caused by infiltration of the metastatic cell to the conduction system of the heart. The presence of hemorrhagic pericardial effusion supports the suggestion of pericardial metastasis. Regardless of the lack of histological confirmation, we suggest that this case was cardiac metastasis diagnosed antemortem.

Table 1. Literature review of case reports regarding heart block caused by cardiac metastasis.

Article	Age (years)	Primary site of cancer	Type of cancer cell	Location of cardiac metastasis	Sign / symptoms	ECG findings	Other cardiac manifestation	PPM implantation	Type of PPM	Reversibility of heart block	Treatment of cardiac metastasis	Diagnostic tools to detect cardiac metastasis	Lifespan after cardiac manifestation
Buckberg and Fowler, 1961 ³	42	bronchogenic carcinoma	adeno carcinoma	interventricular septum, anterior wall of left ventricle	shortness of breath, cough with blood	complete heart block	intracardiac mass	no	No	no	none	postmortem autopsy	6 weeks
Clifford et al., 2003 ⁴	64	lymph	follicular small cell cleaved lymphoma	interventricular septum and RV anterior wall	nausea, diaphoresis, and dizziness	complete heart block with a ventricular escape	RV mass	yes	dual chamber	yes	chemotherapy	echocardiogram	2 years, still alive
Mocini et al., 2005 ⁵	44	lung	malignant neoplasm	interventricular septum, left ventricular wall	syncope	complete heart block	mild pericardial effusion	yes	dual chamber	no	none	postmortem autopsy	3 days
Ferraz et al., 2006 ¹⁰	63	uterus	squamous cell carcinoma	interventricular septum and RV	fatigue and dyspnea on mild exertion	complete heart block	pericardial effusion, RV mass	yes	permanent atrioventricular epimyocardial pacemaker	no	surgery for RV mass	echocardiogram, CT	4 months
Ozyuncu et al., 2006 ¹¹	56	right thigh	malignant melanoma	interventricular, interatrial septum	nausea, vomiting, presyncope	complete heart block	mitral regurgitation, pericardial effusion	yes	VDD	no	chemo-immunotherapy	CT thorax, echocardiogram	2 months, still alive
Knowles et al., 2007 ¹²	42	right maxillary sinus	right maxillary sinus	not stated	syncope	complete heart block with a ventricular escape	pericardial effusion	no	no	yes	chemotherapy	echocardiogram, CT	1 year
Rathi et al., 2008 ¹³	67	skin	malignant melanoma	all myocardial walls	asymptomatic	complete heart block, junctional escape	pericardial effusion	no	no	intermittent complete heart block	chemotherapy	echocardiogram, MRI	not stated

Article	Age (years)	Primary site of cancer	Type of cancer cell	Location of cardiac metastasis	Sign / symptoms	ECG findings	Other cardiac manifestation	PPM implantation	Type of PPM	Reversibility of heart block	Treatment of cardiac metastasis	Diagnostic tools to detect cardiac metastasis	Lifespan after cardiac manifestation
Lin <i>et al.</i> , 2015 ¹⁴	74	thyroid	papillary thyroid carcinoma	RVOT	exertional dyspnea and palpitations	first-degree atrioventricular block and subsequently a new intermittent complete atrioventricular block	RVOT mass	yes	not stated	no	none	CTA, C MRI, PET Scan	not stated (not long after discharge)
Yoneda <i>et al.</i> , 2016 ¹⁵	53	gingival	squamous cell carcinoma	atrial septum, left ventricle, AV node	cough, syncope	complete heart block, ventricular fibrillation	none	no	No	no	chemotherapy	postmortem autopsy	4 weeks
Park <i>et al.</i> , 2016 ¹⁶	54	right leg	leiomyosarcoma	interventricular septum	dizziness and dyspnea	complete heart block and idioventricular escaped rhythm of bifascicular block morphology	VT, intraventricular mass	yes	dual chamber	no	palliative chemotherapy	MRI	3 months
Yoshihiro <i>et al.</i> , 2017 ¹⁷	57	thyroid	squamous cell carcinoma	interventricular septum	cough and shortness of breath	complete heart block	RV mass	yes	not stated	no	chemoradiotherapy	FDG-PET	25 days
Karsai <i>et al.</i> , 2007 ¹⁸	56	lung	adenocarcinoma	interventricular septum	dull pain, presyncope	complete heart block	LV mass	yes	not stated	no	none	echocardiogram, CT, postmortem autopsy	19 days
Kumar <i>et al.</i> , 2018 ¹⁹	28	tongue	squamous cell carcinoma	interventricular septum	syncope	complete heart block	none	yes	not stated	no	none	PET Scan	5 days
Cho <i>et al.</i> , 2018 ²⁰	70	oral cavity	squamous cell carcinoma	interventricular septum	dizziness	complete heart block with a ventricular escape	none	yes	DDD	no	palliative chemotherapy	FDG PET	not stated

Predominantly, after a reversible or transient cause of bradycardia is excluded, cardiac pacing indication is decided by bradycardia severity, instead of its etiology. The European Society of Cardiology (ESC) guidelines state that some types of persistent bradycardia require permanent pacing. In acquired AV block, pacing is indicated in patients with second-degree type 2 or third-degree AV block regardless of symptoms (class I)¹⁸. Similar to the ESC guidelines, the American College of Cardiology/American Heart Association/Heart Rhythm Society guidelines also state that patients with transient or reversible causes of AV block should receive medical and supportive treatment if necessary, including temporary transvenous pacing, prior to confirmation of the need for permanent pacing (class of Recommendation/COR I). In addition, for patients with acquired CHB not associated with physiologic or reversible etiology, permanent pacing is recommended regardless of symptoms (COR I)¹⁹.

In our case, we assume that the CHB was persistent after trying to correct possible external causes, such as hypercalcemia and hypokalemia, and after a sufficient waiting period. Moreover, it is still also unknown whether CHB would resolve after cancer treatment^{2,20}. Our literature review showed that there are only two cases of metastatic CHB that are reversible after undergoing chemotherapy for cardiac metastasis^{8,12}, while several other cases showed that CHB is irreversible^{2-7,9-11,13-15}. The short life expectancy in patients with metastatic CHB also makes it difficult to follow up on CHB reversibility. Our literature review showed only one metastatic CHB case reported a significant lifespan more than two years after PPM implantation⁸, while other cases reported a lifespan of no longer than one year^{2-7,9-15}.

Our patient was expected to continue the radiotherapy cycle. Radiotherapy (RT) itself can induce pacemaker malfunction. Software impairments are the primary manifestation of malfunction during RT. This will perhaps lead to pacemaker reset, leaving only device basic function. Radiation dosage

appears to contribute less to inducing pacemaker malfunctions than the beam energy of the RT²¹.

For patients with indications for permanent pacing, but accompanied with significant comorbidities or who are anticipated having a shortened lifespan because of terminal progressive illness, the implantation of a PPM should not be performed if it is unlikely to deliver significant clinical benefits or if it hinders the main therapy for the patient's goal of care. Even though pacemaker implantation risks are rather low, the risk-benefit ratio is not favorable if the possible benefit is also quite low¹⁹. Thus, after discussing with our patient and his family, we decided not to implant a PPM after acquiring the patient's and his family's consent.

Conclusion

CHB in a patient with oral cancer should increase the physician's suspicion of cardiac metastasis. It is rare for such malignancies to present with CHB. The decision to implant a permanent pacemaker is highly specific based on the risks and benefits of each patient. It needs to be tailored to the patient's functional status, comorbid diseases, prognosis, and response to conservative management. Even though the pacemaker implantation risks are rather low, the risk-benefit ratio is not favorable if the possible benefit is also quite low. A permanent pacemaker was not implanted in our patient because of the poor prognosis, severe comorbidities, and low expected lifespan.

Consent

Written informed consent for publication of their clinical details and/or clinical images was obtained from the patient's parent.

Data availability

All data underlying the results are available as part of the article and no additional source data are required.

References

- Goldberg AD, Blankstein R, Padera RF, et al.: **Tumors Metastatic to the Heart.** *Circulation.* 2014; **128**(16): 1790–1794.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kumar D, Mankame P, Sabnis G, et al.: **A case report: metastatic complete heart block.** *Eur Heart J Case Rep.* 2018; **2**(4): yty131.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Buckberg GD, Fowler NO: **Complete Atrioventricular Block due to Cardiac Metastasis of Bronchogenic Carcinoma.** *Circulation.* 1961; **24**: 657–661.
- Park YM, Shin JO, Kim M, et al.: **Cardiac metastasis of leiomyosarcoma complicated with complete atrio-ventricular block and ventricular tachycardia.** *Korean Circ J.* 2016; **46**(2): 260–263.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Yoshihiro T, Tsuchihashi K, Kusaba H, et al.: **Cardiac metastasis of squamous cell carcinoma of the thyroid gland with severe disseminated intravascular coagulation: A case report.** *Mol Clin Oncol.* 2017; **6**(1): 91–95.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kasai T, Kishi K, Kawabata M, et al.: **Cardiac metastasis from lung adenocarcinoma causing atrioventricular block and left ventricular outflow tract obstruction.** *Chest.* 2007; **131**(5): 1569–1572.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cho JY, Kim KH, Park H, et al.: **Complete atrioventricular block as an initial manifestation of recurrent oral cavity cancer: a case report.** *BMC Cardiovasc Disord.* 2018; **18**(1): 142.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Clifford SM, Guerra SM, Mangion JR: **Massive metastatic intracardiac lymphoma presenting with complete heart block with resolution following chemotherapy.** *Echocardiography.* 2003; **20**(2): 201–202.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mocini D, Longo R, Colivicchi F, et al.: **A complete atrioventricular block secondary to myocardial metastases of lung cancer . A case report.** *Ital Hear J.* 2005; **6**(11): 931–932.
[PubMed Abstract](#)
- Ferraz JGG, Martins ALM, De Souza JF, et al.: **Metastatic Tumor of Squamous Cell Carcinoma From Uterine Cervix to Heart: Ante-Mortem Diagnosis.** *Arq Bras Cardiol.* 2006; **87**(4): e104–7.
[PubMed Abstract](#) | [Publisher Full Text](#)

11. Ozyuncu N, Sahin M, Altin T, *et al.*: **Cardiac metastasis of malignant melanoma : a rare cause of complete atrioventricular block.** *Europace.* 2006; **8**(7): 545–548.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Knowles JW, Elliott AB, Brody J: **A case of complete heart block reverting to normal sinus rhythm after treatment for cardiac invasive Burkitt's lymphoma.** *Ann Hematol.* 2007; **86**(9): 687–690.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Rathi VK, Williams RB, Yamrozik J, *et al.*: **Cardiovascular magnetic resonance of the charcoal heart.** *J Cardiovasc Magn Reson.* 2008; **10**(1): 37.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
14. Lin CK, Cochet A, Lewi JE: **Complete Heart Block in a Patient With Metastatic Papillary Thyroid Carcinoma.** *Fed Pract.* 2015; **32**(1): 34–35.
[PubMed Abstract](#) | [Free Full Text](#)
15. Yoneda T, Kase K, Amino Y, *et al.*: **A case of gingival cancer with pulmonary metastases that developed complete atrioventricular block and ventricular fibrillation as a result of myocardial metastases.** *Clin Case Rep.* 2016; **4**(12): 1075–1081.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Bussani R, Abbate A, Silvestri F: **Cardiac metastases.** *J Clin Pathol.* 2007; **60**(1): 27–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Lichtenberger JP, Reynolds DA, Keung J, *et al.*: **Metastasis to the Heart: A Radiologic Approach to Diagnosis With Pathologic Correlation.** *AJR Am J Roentgenol.* 2016; **207**(4): 764–772.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Brignole M, Auricchio A, Baron-Esquivias G, *et al.*: **2013 ESC Guidelines on cardiac pacing and cardiac resynchronization therapy.** *Rev Esp Cardiol (Engl Ed).* 2014; **67**(1): 58.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Kusumoto FM, Schoenfeld MH, Barrett C, *et al.*: **2018 ACC/AHA/HRS Guideline on the Evaluation and Management of Patients With Bradycardia and Cardiac Conduction Delay: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society.** *Circulation.* 2019; **140**(8): e382–e482.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. Fotouhi Ghiam A, Dawson LA, Abuzeid W, *et al.*: **Role of palliative radiotherapy in the management of mural cardiac metastases: who when and how to treat? A case series of 10 patients.** *Cancer Med.* 2016; **5**(6): 989–996.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Zaremba T, Jakobsen AR, Sogaard M, *et al.*: **Radiotherapy in patients with pacemakers and implantable cardioverter defibrillators: A literature review.** *Europace.* 2016; **18**(4): 479–491.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status: ?

Version 1

Reviewer Report 27 October 2020

<https://doi.org/10.5256/f1000research.29188.r73101>

© 2020 Prakoso R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Radityo Prakoso

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia

Some typos and grammars should be evaluated. In the presentation, what does author mean about "improvement"? Why did TPM extracted by improvement but the complete heart block was actually not improving? I mean its 7th days were they sure that the worsening condition not due to the low cardiac output because of too low of heart rate? Maybe the author could add some suggestions of using medication to increase heart rate if there were significant reasons to "must take" the TPM?

Conclusion

So what does the author advise if one day there were similar patients came with complete heart block? Should the author suggest to implant the TPM or maybe using medication to increase the heart rate, such as sympathomimetic amines.

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Yes

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Partly

Is the case presented with sufficient detail to be useful for other practitioners?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Pediatric Cardiology and Congenital Heart Disease

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 31 Oct 2020

Andrianto Andrianto

Some typos and grammars should be evaluated. In the presentation, what does author mean about "improvement"? Why did TPM extracted by improvement but the complete heart block was actually not improving? I mean its 7th days were they sure that the worsening condition not due to the low cardiac output because of too low of heart rate? Maybe the author could add some suggestions of using medication to increase heart rate if there were significant reasons to "must take" the TPM?

Answer: Thank you for your comment. We have revised come typos and grammar. We also added a clearer definition in general condition, supportive treatment including dopamine use, and reason the TPM was extracted in section *case presentation* paragraf 4-6.

Conclusion

So what does the author advise if one day there were similar patients came with complete heart block? Should the author suggest to implant the TPM or maybe using medication to increase the heart rate, such as sympathomimetic amines.

Answer: Thank you for your comment. We have added our suggestion of TPM use in such cases in section *conclusion* paragraf 1.

Competing Interests: None declared

Comments on this article

Version 1

Reader Comment 29 Oct 2020

Mohammad Saifur Rohman, Cardiology and vascular medicine, faculty of medicine, Brawijaya University, Malang, Indonesia

Very interesting and might be considered to be indexed.

This is an extremely rare case of complete heart block which might related or not to tongue cancer metastatic. Since, no histopathology of biopsy available taken from intracardiac mass or

myocardium post mortem. To confirm whether the same result as tongue biopsy showing poorly differentiated squamous cell carcinoma.

Pericardiocentesis showed a hemorrhagic typical for malignant disease, however the authors did not show the cytology analysis results taken from pericardiocentesis procedure. The bloody fluid of pericardial effusion might come from elsewhere metastatic organ other than the cardiac.

Therefore, it is pretty much convincing if the mass biopsy and pericardial fluid cytology are available.

Competing Interests: NO conflict of interest

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research