REVIEW

Recent advances in the treatment of pancreatic cancer

[version 1; peer review: 2 approved]

Marc T Roth , Dana B Cardin, Jordan D Berlin

Department of Hematology and Oncology, Vanderbilt-Ingram Cancer Center, Nashville, TN, USA

V1 First published: 21 Feb 2020, 9(F1000 Faculty Rev):131
<https://doi.org/10.12688/f1000research.21981.1>

Latest published: 21 Feb 2020, 9(F1000 Faculty Rev):131
<https://doi.org/10.12688/f1000research.21981.1>

Abstract

Pancreatic ductal adenocarcinoma is one of the deadliest solid tumor malignancies and is projected to become a leading cause of cancer-related death in coming years. Improving quality of life and survival amongst these patients will require new ideas and novel therapies in a multidisciplinary approach. This review will cover the most recent advances in the comprehensive treatment of pancreatic cancer and place them within a historical context when necessary. Treatment of all disease stages will be discussed, but the focus is on systemic therapy as novel drugs and new treatment combinations enter the clinic. This will include more aggressive chemotherapy in earlier disease stages, approved uses for immunotherapy, and targetable mutations. In addition, negative trials of importance and controversial topics will be noted.

Keywords

pancreatic cancer, review

Open Peer Review**Approval Status**

	1	2
version 1 21 Feb 2020

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Thomas Seufferlein**, Ulm University, Ulm, Germany
2. **Jörg Kleeff**, Martin-Luther-University Halle-Wittenberg, University Medical Center Halle, Halle, Germany

Any comments on the article can be found at the end of the article.

Corresponding author: Marc T Roth (marc.roth@vumc.org)

Author roles: Roth MT: Conceptualization, Writing – Original Draft Preparation; Cardin DB: Conceptualization, Writing – Review & Editing; Berlin JD: Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Roth MT *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Roth MT, Cardin DB and Berlin JD. **Recent advances in the treatment of pancreatic cancer [version 1; peer review: 2 approved]** F1000Research 2020, 9(F1000 Faculty Rev):131 <https://doi.org/10.12688/f1000research.21981.1>

First published: 21 Feb 2020, 9(F1000 Faculty Rev):131 <https://doi.org/10.12688/f1000research.21981.1>

Introduction

Pancreatic cancer has one of the highest mortalities of all malignancies. It is estimated that 56,770 diagnoses were made and 45,750 deaths occurred in 2019 in the US¹. One analysis projects that pancreatic cancer will surpass breast, prostate, and colorectal cancers as the leading cause of cancer-related death in the US by the year 2030². These estimates highlight the urgent need for new and innovative treatment options for patients with this deadly disease. Indeed, numerous efforts are under way to alter the disease course in order to keep pace with the improved outcomes seen in other malignancies. The aims of this review are to highlight these advancements and to revisit the historical basis for current treatment options in pancreatic ductal adenocarcinoma (PDAC).

Chemotherapy

Chemotherapy remains a cornerstone of treatment for PDAC in all stages of disease. Recent data and US Food and Drug Administration (FDA) approvals are divided into three major treatment categories: adjuvant, unresectable/metastatic, and neoadjuvant/induction treatments.

Adjuvant setting

The role of adjuvant chemotherapy in PDAC was established in multiple trials, including ESPAC-1, CONKO-001, and ESPAC-3^{3,4}. ESPAC-4 compared the combination of gemcitabine/capecitabine with gemcitabine monotherapy in the adjuvant setting⁵. This study, published in 2017, found that the rates of median overall survival (mOS) were 28.0 months in the combination arm and 25.5 months in the monotherapy arm (hazard ratio [HR] 0.82, 95% confidence interval [CI] 0.68–0.98; $P = 0.032$). GI PRODIGE 24, published in 2018, compared adjuvant modified FOLFIRINOX—infusion 5-fluorouracil (5-FU), folinic acid, irinotecan, and oxaliplatin—with gemcitabine⁶. The term “modified” refers to the reduction in irinotecan dosing from 180 to 150 mg/m² and the omission of the 5-FU bolus. These changes were made after concerns with the triplet regimen arose during a protocol-specified safety analysis. The mOS rates were an unprecedented 54.4 months for modified FOLFIRINOX and 35.0 months in the gemcitabine arm. It is important to note that those eligible for modified FOLFIRINOX included patients who had fully recovered from surgery and had a performance status of 0 to 1 and adequate hematologic and hepatologic function, making it important to assess the fitness of adjuvant therapy candidates. Gemcitabine/capecitabine and modified FOLFIRINOX are the current standards of care in the adjuvant setting for appropriate patients with resected PDAC.

One notable negative study (APACT, ClinicalTrials.gov Identifier: NCT01964430) compared gemcitabine and nab-paclitaxel (Abraxane®, Celgene Corporation, Summit, NJ, USA) (G/A) with gemcitabine monotherapy. The trial did not meet its primary endpoint of improvement in disease-free survival when compared with gemcitabine alone⁷.

Unresectable/metastatic disease

Many of the cytotoxic regimens discussed previously were first studied in patients with metastatic disease. The ACCORD trial,

published in 2011, compared FOLFIRINOX with gemcitabine in metastatic PDAC (mPDAC), resulting in mOS rates of 11.1 months in the triplet arm and 6.8 months with monotherapy (HR 0.57, 95% CI 0.45–0.73; $P < 0.001$)⁸ and leading to a new standard of care for patients with advanced disease with a good performance status. The MPACT trial, published in 2013, evaluated G/A in mPDAC, finding improved mOS rates of 8.5 months in the combination arm and 6.7 months with gemcitabine alone (HR 0.72, 95% CI 0.617–0.835; $P < 0.001$)⁹, again adding a new regimen to the standard-of-care options for patients with advanced disease. In 2015, the NAPOLI-1 trial compared 5-FU/nanoliposomal irinotecan (nal-IRI) with either drug as monotherapy in metastatic PDAC previously treated with gemcitabine¹⁰. In this phase III randomized controlled trial (RCT), 417 patients were enrolled. The mOS rates were 6.1 months in the combination arm and 4.2 months in the 5-FU arm (HR 0.67, 95% CI 0.49–0.92; $P = 0.012$). Of note, mOS in the nal-IRI arm was 4.9 months. This regimen is currently listed as category 1 in the second-line setting for metastatic disease by the National Comprehensive Cancer Network (NCCN)¹¹ and is important as a rare randomized trial in this setting, especially one demonstrating the potential benefit of salvage chemotherapy in a difficult-to-treat patient population. Most recently, the combination of gemcitabine, nab-paclitaxel, and cisplatin was tested in a phase Ib/II trial in metastatic PDAC with a primary endpoint of complete response rate (CRR) (25% needed for significance)¹². Although this benchmark was not met (CRR 8%), the triplet did yield an overall response rate (ORR) of 71% and mOS of 16.4 months (95% CI 10.2–25.3). The triplet continues to be tested in the neoadjuvant setting and in advanced biliary cancer.

Although cytotoxic chemotherapy is responsible for the largest survival advantages seen in the metastatic setting, novel agents are actively being investigated. Pegvorhialuronidase alfa (PEGPH20), which degrades hyaluronan in the extracellular matrix, was the subject of much interest in recent years until development was ceased in late 2019 by Halozyme Therapeutics. Although phase II data showed an improvement in progression-free survival (PFS) when combined with G/A¹³, the triplet failed to meet its primary endpoint of OS in a phase III placebo-controlled RCT (11.2 versus 11.5 months, HR 1.00; $P = 0.9692$)¹⁴. AVENGER 500 (ClinicalTrials.gov Identifier: NCT03504423) is a phase III RCT investigating modified FOLFIRINOX with or without CPI-613¹⁵, a tricarboxylic acid cycle inhibitor. Accrual should be complete by mid-2020, and results are highly anticipated after showing promise in the phase I setting (ORR 61%, CRR 17%)¹⁶.

Neoadjuvant and induction treatment

Since establishing the efficacy of chemotherapy in the adjuvant setting and advanced disease, its movement into the pre-operative space has been a logical progression. The rationale behind this includes potentially increasing the R0 resection rate, administering more therapy prior to surgery when it is better tolerated, and providing time to evaluate the biology of a patient's individual disease. In addition, as criteria for resection change and procedures become more extensive, conversion or downstaging therapy with chemotherapy in the borderline

resectable pancreatic cancer/locally advanced pancreatic cancer (BRPC/LAPC) space is becoming commonplace. Multiple institutional and small multi-center trials have been conducted to determine the optimal chemotherapy regimen or utility of radiation (or both) in this space, although currently there is no established standard of care in this arena. The spectrum of radiographic staging and sequence of treatment in these trials can make extrapolation into common clinical practice difficult.

Several ongoing trials are investigating pre-operative chemotherapy for upfront resectable PDAC. One such study is the phase II/III Prep-02/JSAP-05 (UMIN000009634), which compared gemcitabine/S1 followed by resection with upfront surgery, and OS was its primary endpoint¹⁷. The mOS rates were 36.7 months in the chemotherapy arm and 26.6 months with upfront surgery alone (HR 0.72, 95% CI 0.55–0.94; $P = 0.015$). Resection rate, R0 rate, and morbidity were similar between the two arms. Further studies should be performed in this space to inform generalization to the broader PDAC population.

Current practice for many institutions in the US for fit patients with borderline resectable PDAC is neoadjuvant modified FOLFIRINOX with consideration of radiation therapy (RT) followed by resection if appropriate. This stems from the phase I single-arm ALLIANCE trial A021101 (published in 2016), in which patients who received the above regimen achieved an mOS of 21.7 months (95% CI 15.7–not reached) and a 68% resection rate (95% CI 49–88%)¹⁸. The confirmatory trial is ongoing (A021501, discussed below). NEOLAP (ClinicalTrials.gov Identifier: NCT02125136) is the first randomized trial comparing FOLFIRINOX and G/A in LAPC. In that study, patients were given two cycles of G/A induction and then randomly assigned to either two additional cycles of G/A or four cycles of FOLFIRINOX. Final results presented at the European Society for Medical Oncology annual congress in 2019 were notable for conversion rates of 30.6% for G/A and 45.0% for FOLFIRINOX, although this difference was not considered statistically significant (odds ratio 0.54, 95% CI 0.26–1.13; $P = 0.135$)¹⁹. The highly anticipated phase II SWOG 1505 (ClinicalTrials.gov Identifier: NCT02562716) trial moves the comparison of G/A with FFX entirely into the upfront resectable setting with a “pick the winner” design using 2-year OS as the primary endpoint²⁰. Other ongoing studies will add to the growing body of evidence in this arena. NEONAX (ClinicalTrials.gov Identifier: NCT02047513), a phase II randomized study, has recently completed accrual. That trial compares two pre-operative and four post-operative cycles of G/A with six cycles of adjuvant G/A.

Radiation

Evidence regarding the role of RT in pancreatic cancer has been conflicting, although its use has been of great interest. Cross-trial comparisons are even more difficult with RT given that the dose and mode of radiation delivery have changed over the years. LAP07 was one of the largest randomized trials to evaluate the role of gemcitabine and erlotinib with or without chemoradiation in LAPC. The trial was stopped early for futility as no significant survival difference was noted

between the chemoradiotherapy- and chemotherapy-alone arms²¹. Few definitive practice-changing trials have been conducted recently, but two studies hoping to clarify the role of RT in this space are ongoing. The ALLIANCE trial A021501 (ClinicalTrials.gov Identifier: NCT02839343) is comparing neoadjuvant FOLFIRINOX followed by stereotactic body RT (SBRT) with FOLFIRINOX alone followed by resection (if possible) in borderline resectable PDAC²². Interestingly, the radiation arm has since closed because of futility. Results of the interim analysis have yet to be reported but will be an intriguing addition to an ongoing discussion. The PREOPANC-1 (NTR3709) trial is a phase III RCT comparing upfront resection to gemcitabine-based chemoradiation followed by resection, and both arms are receiving adjuvant gemcitabine. Though preliminary, data suggest an improved OS (17.1 versus 13.5 months, HR 0.71; $P = 0.047$) and R0 resection rate (65% versus 31%; $P < 0.001$) with pre-operative chemoradiation compared with chemotherapy alone²³.

In 2019, the American Society for Radiation Oncology released new guidelines for the treatment of pancreatic cancer²⁴. These recommended the consideration of conventionally fractionated RT or SBRT in high-risk adjuvant settings, as neoadjuvant downstaging for BRPC/LAPC, and for LAPC as part of definitive treatment, all admittedly with low to moderate quality of evidence. It should be noted that in contrast to other data presented regarding efficacy of chemotherapy in PDAC, the use of radiation is not founded on solid evidence as of yet.

Immunotherapy

PDAC has been largely excluded from the recent success stories in the field of immunotherapy (IO) relative to many other malignancies. Single- and double-agent IO, as well as combinations with cytotoxic chemotherapy and radiation among others, have been studied but have not shown meaningful clinical benefit²⁵. There are multiple hypotheses for the underlying mechanism of resistance to IO in PDAC. These include a lack of effector cells in the tumor microenvironment combined with an immunosuppressive infiltrate, dense stroma impairing migration of effector cells, and immune checkpoint signaling²⁶. Many studies are under way to discover the key to overcoming these barriers. It is important to highlight that IO, though used in many other malignancies, is not approved for the treatment of PDAC outside of clinical trials.

Currently, the only exception to this is in PDAC with microsatellite instability (MSI). In May 2017, the FDA granted its first tissue/site-agnostic approval to pembrolizumab for the treatment of MSI-high or deficient mismatch repair (dMMR) solid tumors on the basis of data from five single-arm trials. One enrolled 86 patients, including eight patients with PDAC²⁷. That study had co-primary endpoints of immune-related PFS and ORR. In the intention-to-treat population, the ORR was 53% (95% CI 42–64%) and PFS rates at 1 and 2 years were 64% and 53%, respectively. Among patients with PDAC, the ORR was 62%, and two patients (25%) achieved a complete response. Updated results from the KEYNOTE-158 study are less promising. Among 22 patients with PDAC, the ORR was 18.2%,

the median PFS (mPFS) was 2.1 months, and the mOS was 4.0 months²⁸.

Targeted therapy

Recent large-scale molecular profiling efforts have attempted to uncover genomic subtype prevalence and corresponding therapeutic relevance in PDAC. One such effort found that 14% of evaluated patients harbored mutations (either germline or somatic) in *BRCA1*, *BRCA2*, and *PALB2*²⁹. Similarly, initial results from the Know Your Tumor initiative showed that 27% of evaluated patients had highly actionable mutations, which when paired with matched therapy resulted in a longer mPFS than that of patients without matched therapy³⁰. A number of other genes are frequently mutated in PDAC (*KRAS*, *TP53*, *CDKN2A*, *SMAD4*, *MLL3*, *TGFBR2*, *ARID1A*, and *SF3B1*) but as of yet do not have therapeutic indications though may provide detail on prognosis³¹. Neurotrophic receptor tyrosine kinase (*NTRK*) gene fusions have been found in a small fraction of PDAC, and some estimates are as high as 6%³². Current options for targeted treatment in PDAC are discussed below.

NTRK fusions

There are two new treatment options for patients found to have fusion proteins in the *NTRK* gene detected by next-generation sequencing or fluorescence *in situ* hybridization. Larotrectinib³³ was granted accelerated approval in November 2018 on a tissue-agnostic basis for unresectable or metastatic solid tumors harboring the gene fusion. This approval was based on data from three single-arm clinical trials: LOXO-TRK-14001, SCOUT, and NAVIGATE³⁴. In total, 55 patients (children and adults) were enrolled between the trials. At the time of assessment, the primary endpoint of ORR was 75% (95% CI 67–90%) and the mPFS had not been reached after a median follow-up duration of 9.9 months. It should be noted that only one patient with pancreatic cancer was included, although that person was among the responders. The drug was generally well tolerated; grade 3 and higher adverse events (AEs) occurred in less than 5% of patients. The most common side effects were aspartate aminotransferase/alanine aminotransferase (AST/ALT) abnormalities, fatigue, and vomiting.

Entrectinib³⁵ was granted accelerated approval for the same indication in August 2019 on the basis of three additional single-arm trials: ALKA-372-001, STARTRK-1, and STARTRK-2³⁶. Between the two phase I studies, 60 patients harboring a gene rearrangement in *NTRK*, *ROS1*, or *ALK* were enrolled. Given their early phase, only preliminary efficacy data are available, but in the three evaluable patients with *NTRK* fusions, the ORR was 100% (95% CI 44–100%), although none had PDAC. The phase II study (STARTRK-2) is under

way. The most common side effects were fatigue, dysgeusia, and paresthesias.

Homologous recombination deficiency

Mutations in homologous recombination deficiency (HRD) genes (including *BRCA1/2*) have been correlated with defective DNA repair²⁹. Given the inherent genomic instability in this setting, various interventions to exploit this and induce apoptosis have been studied in PDAC. Patients with germline mutations in *BRCA1/2* appear to have an increased sensitivity to platinum agents³⁷, and the NCCN recommends treatment of this subpopulation with gemcitabine plus cisplatin in locally advanced and metastatic disease¹¹.

A more recent mechanism to leverage this pathway involves poly(adenosine diphosphate-ribose) polymerase (PARP) inhibition. The POLO trial was a phase III RCT evaluating olaparib (a PARP inhibitor) maintenance versus placebo in patients with metastatic PDAC and germline *BRCA1/2* mutations whose disease did not progress after first-line platinum-based chemotherapy³⁸. The primary endpoint was PFS. mPFS rates were 7.4 months in the treatment arm and 3.8 in the control arm (HR 0.53, 95% CI 0.35–0.82; $P = 0.004$) but this did not equate to an OS advantage during the interim analysis (mOS 18.9 versus 18.1 months, respectively; HR for death, 0.91; 95% CI 0.56–1.46; $P = 0.68$). Rates of grade 3 and higher AEs were 40% in the treatment group and 23% in the placebo group, and the most common AEs were fatigue, nausea, diarrhea, and abdominal pain. In December 2019, the FDA approved olaparib for use in those who had germline *BRCA1/2* mutations, a good performance status, and no disease progression after at least 4 to 6 months of chemotherapy. Further studies are ongoing with novel drug combinations targeting a wider range of DNA damage repair pathways in PDAC.

Summary

Patients with pancreatic cancer today have increased options for treatment relatively speaking, but survival for this disease is dismal and falling behind in a rapidly advancing field. Clinical trial design must evolve to address this need, and trial participation should be considered whenever possible. Established questions such as the role of radiation in pancreatic cancer and the precise populations that will benefit remain unanswered. New issues regarding resistance to IO and the role of targeted therapy have also been presented. Progress is being made, however, from pharmaceutical innovation to an increasing number of studies including the elderly and those with a poor performance status and refractory disease. With such efforts, there is hope that more effective treatment strategies are within reach.

References

1. Siegel RL, Miller KD, Jemal A, et al.: **Cancer statistics, 2019.** *CA Cancer J Clin.* 2019; **69**(1): 7–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
2. Rahib L, Smith BD, Aizenberg R, et al.: **Projecting cancer incidence and deaths to 2030: the unexpected burden of thyroid, liver, and pancreas cancers in the United States.** *Cancer Res.* 2014; **74**(11): 2913–21.
[PubMed Abstract](#) | [Publisher Full Text](#)

3. Neoptolemos JP, Stocken DD, Friess H, *et al.*: **A randomized trial of chemoradiotherapy and chemotherapy after resection of pancreatic cancer.** *N Engl J Med.* 2004; **350**(12): 1200–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Oettle H, Neuhaus P, Hochhaus A, *et al.*: **Adjuvant chemotherapy with gemcitabine and long-term outcomes among patients with resected pancreatic cancer: the CONKO-001 randomized trial.** *JAMA.* 2013; **310**(14): 1473–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. **F** Neoptolemos JP, Palmer DH, Ghaneh P, *et al.*: **Comparison of adjuvant gemcitabine and capecitabine with gemcitabine monotherapy in patients with resected pancreatic cancer (ESPAC-4): a multicentre, open-label, randomised, phase 3 trial.** *Lancet.* 2017; **389**(10073): 1011–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
6. **F** Conroy T, Hammel P, Hebbar M, *et al.*: **FOLFIRINOX or Gemcitabine as Adjuvant Therapy for Pancreatic Cancer.** *N Engl J Med.* 2018; **379**(25): 2395–406.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
7. Tempero MA, Reni M, Riess H, *et al.*: **APACT: phase III, multicenter, international, open-label, randomized trial of adjuvant nab-paclitaxel plus gemcitabine (nab-P/G) vs gemcitabine (G) for surgically resected pancreatic adenocarcinoma.** *JCO.* 2019; **37**: 4000.
[Publisher Full Text](#)
8. **F** Conroy T, Desseigne F, Ychou M, *et al.*: **FOLFIRINOX versus gemcitabine for metastatic pancreatic cancer.** *N Engl J Med.* 2011; **364**(19): 1817–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
9. **F** Von Hoff DD, Ervin T, Arena FP, *et al.*: **Increased survival in pancreatic cancer with nab-paclitaxel plus gemcitabine.** *N Engl J Med.* 2013; **369**(18): 1691–703.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
10. Wang-Gillam A, Li CP, Bodoky G, *et al.*: **Nanoliposomal irinotecan with fluorouracil and folinic acid in metastatic pancreatic cancer after previous gemcitabine-based therapy (NAPOLI-1): A global, randomised, open-label, phase 3 trial.** *Lancet.* 2016; **387**(10018): 545–57.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. **F** Tempero MA: **NCCN Guidelines Updates: Pancreatic Cancer.** *J Natl Compr Canc Netw.* 2019; **17**(5.5): 603–5.
[PubMed Abstract](#) | [F1000 Recommendation](#)
12. **F** Jameson GS, Borazanci E, Babiker HM, *et al.*: **Response Rate Following Albumin-Bound Paclitaxel Plus Gemcitabine Plus Cisplatin Treatment Among Patients With Advanced Pancreatic Cancer: A Phase 1b/2 Pilot Clinical Trial.** *JAMA Oncol.* 2020; **6**: 125.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
13. **F** Hingorani SR, Zheng L, Bullock AJ, *et al.*: **HALO 202: Randomized Phase II Study of PEGPH20 Plus Nab-Paclitaxel/Gemcitabine Versus Nab-Paclitaxel/Gemcitabine in Patients With Untreated, Metastatic Pancreatic Ductal Adenocarcinoma.** *JCO.* 2018; **36**(4): 359–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
14. Halozyme Announces HALO-301 Phase 3 Study Fails To Meet Primary Endpoint. Halozyme Therapeutics. 2019; Accessed January 6, 2020.
[Reference Source](#)
15. CPI-613® (devimistat) [package insert]. Newark, NJ: Rafael Pharmaceuticals, Inc. [Reference Source](#)
16. **F** Alistar A, Morris BB, Desnoyer R, *et al.*: **Safety and tolerability of the first-in-class agent CPI-613 in combination with modified FOLFIRINOX in patients with metastatic pancreatic cancer: a single-centre, open-label, dose-escalation, phase 1 trial.** *Lancet Oncol.* 2017; **18**(6): 770–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
17. Unno M, Motoi F, Matsuyama Y, *et al.*: **Randomized phase I/II trial of neoadjuvant chemotherapy with gemcitabine and S-1 versus upfront surgery for resectable pancreatic cancer (Prep-02/JSAP-05).** *JCO.* 2019; **37**: 189.
[Publisher Full Text](#)
18. Katz MH, Shi Q, Ahmad SA, *et al.*: **Preoperative Modified FOLFIRINOX Treatment Followed by Capecitabine-Based Chemoradiation for Borderline Resectable Pancreatic Cancer: Alliance for Clinical Trials in Oncology Trial A021101.** *JAMA Surg.* 2016; **151**(8): e161137.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Kunzmann V, Algül H, Goekkurt E, *et al.*: **Conversion rate in locally advanced pancreatic cancer (LAPC) after nab-paclitaxel/gemcitabine- or FOLFIRINOX-based induction chemotherapy (NEOLAP): Final results of a multicenter randomised phase II AIO trial.** *Ann Oncol.* 2019; **30**: v253.
[Publisher Full Text](#)
20. Sohal D, McDonough SL, Ahmad SA, *et al.*: **SWOG S1505: A randomized phase II study of perioperative mFOLFIRINOX versus gemcitabine/nab-paclitaxel as therapy for resectable pancreatic adenocarcinoma.** *JCO.* 2018; **36**(4_suppl): TPS547-TPS547.
[Publisher Full Text](#)
21. **F** Hammel P, Huguet F, van Laethem JL, *et al.*: **Effect of Chemoradiotherapy vs Chemotherapy on Survival in Patients With Locally Advanced Pancreatic Cancer Controlled After 4 Months of Gemcitabine With or Without Erlotinib: The LAP07 Randomized Clinical Trial.** *JAMA.* 2016; **315**(17): 1844–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
22. Katz MHG, Ou FS, Herman JM, *et al.*: **Alliance for clinical trials in oncology (ALLIANCE) trial A021501: preoperative extended chemotherapy vs. chemotherapy plus hypofractionated radiation therapy for borderline resectable adenocarcinoma of the head of the pancreas.** *BMC Cancer.* 2017; **17**(1): 505.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. van Tienhoven G, Versteijne E, Suker M, *et al.*: **Preoperative chemoradiotherapy versus immediate surgery for resectable and borderline resectable pancreatic cancer (PREOPANC-1): A randomized, controlled, multicenter phase III trial.** *JCO.* 2018; **36**(18_suppl): LBA4002-LBA4002.
[Publisher Full Text](#)
24. **F** Palta M, Godfrey D, Goodman KA, *et al.*: **Radiation Therapy for Pancreatic Cancer: Executive Summary of an ASTRO Clinical Practice Guideline.** *Pract Radiat Oncol.* 2019; **9**(5): 322–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
25. **F** Henriksen A, Dyhl-Polk A, Chen I, *et al.*: **Checkpoint inhibitors in pancreatic cancer.** *Cancer Treat Rev.* 2019; **78**: 17–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
26. **F** Upadhrasta S, Zheng L: **Strategies in Developing Immunotherapy for Pancreatic Cancer: Recognizing and Correcting Multiple Immune "Defects" in the Tumor Microenvironment.** *J Clin Med.* 2019; **8**(9): pii: E1472.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
27. **F** Le DT, Durham JN, Smith KN, *et al.*: **Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade.** *Science.* 2017; **357**(6349): 409–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
28. **F** Marabelle A, Le DT, Ascierto PA, *et al.*: **Efficacy of Pembrolizumab in Patients With Noncolorectal High Microsatellite Instability/Mismatch Repair-Deficient Cancer: Results From the Phase II KEYNOTE-158 Study.** *J Clin Oncol.* 2020; **38**(1): 1–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
29. Waddell N, Pajic M, Patch AM, *et al.*: **Whole genomes redefine the mutational landscape of pancreatic cancer.** *Nature.* 2015; **518**(7540): 495–501.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. **F** Pishvaian MJ, Bender RJ, Halverson D, *et al.*: **Molecular Profiling of Patients with Pancreatic Cancer: Initial Results from the Know Your Tumor Initiative.** *Clin Cancer Res.* 2018; **24**(20): 5018–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
31. **F** Biankin AV, Waddell N, Kassahn KS, *et al.*: **Pancreatic cancer genomes reveal aberrations in axon guidance pathway genes.** *Nature.* 2012; **491**(7424): 399–405.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
32. **F** Dreyer SB, Chang DK, Bailey P, *et al.*: **Pancreatic Cancer Genomes: Implications for Clinical Management and Therapeutic Development.** *Clin Cancer Res.* 2017; **23**(7): 1638–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
33. Vitkravi® (larotrectinib) [package insert]. Whippany, NJ: Bayer Healthcare Pharmaceuticals, Inc. 2019.
34. **F** Drilon A, Laetsch TW, Kummar S, *et al.*: **Efficacy of Larotrectinib in TRK Fusion-Positive Cancers in Adults and Children.** *N Engl J Med.* 2018; **378**(8): 731–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
35. Rozlytrek™ (entrectinib) [package insert]. South San Francisco, CA: Genentech, Inc. 2019.
36. **F** Drilon A, Siena S, Ou SI, *et al.*: **Safety and Antitumor Activity of the Multitargeted Pan-TRK, ROS1, and ALK Inhibitor Entrectinib: Combined Results from Two Phase I Trials (ALKA-372-001 and STARTRK-1).** *Cancer Discov.* 2017; **7**(4): 400–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
37. Golan T, Kanji ZS, Epelbaum R, *et al.*: **Overall survival and clinical characteristics of pancreatic cancer in BRCA mutation carriers.** *Br J Cancer.* 2014; **111**(6): 1132–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. **F** Golan T, Hammel P, Reni M, *et al.*: **Maintenance Olaparib for Germline BRCA-Mutated Metastatic Pancreatic Cancer.** *N Engl J Med.* 2019; **381**(4): 317–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Jörg Kleeff

Department of Visceral, Vascular and Endocrine Surgery, Martin-Luther-University Halle-Wittenberg, University Medical Center Halle, Halle, Germany

Competing Interests: No competing interests were disclosed.

2. Thomas Seufferlein

Department of Internal Medicine I, Ulm University, Ulm, Germany

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research