SYSTEMATIC REVIEW

REVISED Rapid systematic review of systematic reviews: what befriending, social support and low intensity psychosocial interventions, delivered remotely, may reduce social isolation and loneliness among older adults and how? [version 2; peer review: 2 approved with reservations]

Previously titled: "Rapid systematic review of systematic reviews: what befriending, social support and low intensity psychosocial interventions, delivered remotely, are effective in reducing social isolation and loneliness among older adults? How do they work?"

Elisabeth Boulton ^{1,2*}, Dylan Kneale ^{3*}, Claire Stansfield³, Paul Heron⁴, Katy Sutcliffe³, Brenda Hayanga ³, Alex Hall^{1,2}, Peter Bower ^{1,2}, Dymrna Casey ⁵, Dawn Craig⁶, Simon Gilbody ⁴, Barbara Hanratty⁶, Dean McMillan⁴, James Thomas ³, Chris Todd ^{1,2}

¹National Institute for Health Research (NIHR) Older People and Frailty Policy Research Unit, School of Health Sciences, University of Manchester, Manchester, UK

²Manchester Academic Health Science Centre, Manchester, UK

³EPPI-Centre, UCL Social Research Institute, University College London, London, UK

⁴University of York and Hull York Medical School (HYMS), York, UK

⁵School of Nursing & Midwifery, National University of Ireland, Galway, Galway, Ireland

⁶National Institute for Health Research (NIHR) Older People and Frailty Policy Research Unit, Population Health Sciences Institute, Newcastle University, Newcastle-upon-Tyne, UK

* Equal contributors

V2 First published: 25 Nov 2020, 9:1368
<https://doi.org/10.12688/f1000research.27076.1>
 Latest published: 28 Apr 2021, 9:1368
<https://doi.org/10.12688/f1000research.27076.2>

Abstract

Background: During the COVID-19 pandemic 'social distancing' has highlighted the need to minimise loneliness and isolation among older adults (aged 50+). We wanted to know what remotely delivered befriending, social support and low intensity psychosocial interventions may help to alleviate social isolation and loneliness and how they work.

Methods: We followed a systematic 'review of reviews' approach. Searches of 11 databases from the fields of health, social care, psychology and social science were undertaken during April 2020.

Open Peer Review

Approval Status

	1	2
version 2 (revision) 28 Apr 2021	view

version 1 25 Nov 2020	view	view

Reviews meeting our PICOS criteria were included if they focussed on the evaluation of remote interventions to reduce levels of social isolation or loneliness in adults aged 50+ and were critically appraised using AMSTAR2. Narrative synthesis was used at a review and study level to develop a typology of intervention types and their effectiveness. Intervention Component Analysis (ICA) and Qualitative Comparative Analysis (QCA) were used at a study level to explore the characteristics of successful interventions.

Results: We synthesised evidence from five systematic reviews and 18 primary studies. Remote befriending, social support and low intensity psychosocial interventions took the form of: (i) supported video-communication; (ii) online discussion groups and forums; (iii) telephone befriending; (iv) social networking sites; and (v) multi-tool interventions. The majority of studies utilised the first two approaches, and were generally regarded positively by older adults, although with mixed evidence around effectiveness. Focussing on processes and mechanisms, using ICA and QCA, we found that the interventions that were most successful in improving social support: (i) enabled participants to speak freely and to form close relationships; (ii) ensured participants have shared experiences/characteristics; (iii) included some form of pastoral guidance.

Conclusions: The findings highlight a set of intervention processes that should be incorporated into interventions, although they do not lead us to recommend specific modes of support, due to the heterogeneity of interventions.

Keywords

social isolation, loneliness, remote interventions, review of reviews

This article is included in the [Sociology of Health](#) gateway.

This article is included in the [Coronavirus](#) collection.

1. **Syed Ghulam Sarwar Shah** , Oxford
University Hospitals NHS Foundation Trust,
Oxford, UK
John Radcliffe Hospital, Oxford, UK
2. **Jennifer Anne Chipps**, University of the
Western Cape, Cape Town, South Africa

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding authors: Elisabeth Boulton (elisabeth.boulton@manchester.ac.uk), Dylan Kneale (D.Kneale@ucl.ac.uk)

Author roles: **Boulton E:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Kneale D:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Stansfield C:** Data Curation, Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Heron P:** Data Curation, Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Sutcliffe K:** Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Hayanga B:** Formal Analysis, Investigation, Writing – Original Draft Preparation, Writing – Review & Editing; **Hall A:** Writing – Original Draft Preparation, Writing – Review & Editing; **Bower P:** Funding Acquisition, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Casey D:** Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Craig D:** Funding Acquisition, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Gilbody S:** Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Hanratty B:** Funding Acquisition, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **McMillan D:** Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Thomas J:** Funding Acquisition, Methodology, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Todd C:** Funding Acquisition, Methodology, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the National Institute for Health Research Policy Research Programme. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2021 Boulton E *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Boulton E, Kneale D, Stansfield C *et al.* **Rapid systematic review of systematic reviews: what befriending, social support and low intensity psychosocial interventions, delivered remotely, may reduce social isolation and loneliness among older adults and how? [version 2; peer review: 2 approved with reservations]** F1000Research 2021, 9:1368 <https://doi.org/10.12688/f1000research.27076.2>

First published: 25 Nov 2020, 9:1368 <https://doi.org/10.12688/f1000research.27076.1>

REVISED Amendments from Version 1

We have made the following changes to the current submission, and have worked on enhancing the clarity of the presentation based on the suggestions of the reviewers by:

- updating the reference list and highlighting that relevant systematic reviews have been published in the area after the publication of this review;
- explaining some of the methods more clearly including some of the study selection details and the implications of a rapid design;
- explaining in more detail the findings of the review in relation to the effectiveness of remote interventions;
- providing clearer definitions of social isolation and loneliness;
- updating the terminology used;
- making changes to the presentation of the QCA results to add clarity to the findings.

We thank both reviewers for helping us to enhance the clarity of the manuscript.

Any further responses from the reviewers can be found at the end of the article

Introduction

During the height of the first wave of the COVID-19 pandemic, millions of people aged 70 years and over were advised to avoid social contact with those outside their household¹, with older age identified as a risk factor for poorer COVID-19 prognosis². Older adults are more likely to have long-term illness or disability, to live alone and to be widowed, all of which are risk factors for loneliness³. Social isolation and loneliness adversely affect quality of life, wellbeing and mental health, and are associated with physical ill health and mortality⁴. Social distancing and restrictions on face-to-face contact increase the risk of social isolation and loneliness. The requirement for older adults to restrict their activities during the COVID-19 pandemic puts a spotlight on the need to understand how to minimise the impact of loneliness and isolation using remotely-delivered approaches. Here we use a broad definition of 'older' adult, defined as those aged 50+, which captures those in middle age who may be nearing or experiencing age-related transitions, such as retirement or unpaid caring, or living with age-related long term conditions.

In the voluntary and community sector, many existing services are shifting to providing remote support, often via the telephone. In England, the call during March 2020 for NHS Volunteer Responders included roles to make 'regular phone calls to check on people isolating at home'⁵. Fulfilment of such roles requires that:

- (i) the programmes and interventions staffed by these volunteers are effective and have minimal adverse consequences for older adults; and
- (ii) the volunteers making phone calls and providing other forms of support are adequately trained and supported to fulfil these roles, with training based on evidence of how the

intervention should be delivered and the key processes that generate successful interventions.

This review focusses on interventions that seek to ameliorate loneliness or social isolation, or both. We conceptualise loneliness as an emotional response by individuals when there is a 'deficit between their desired and actual quality and quantity of social engagement and relationships'^{6, p64*}. Social isolation reflects the number of social contacts that people have⁷, and people who are socially isolated tend to have social networks of low density that are not maintained through frequent engagement⁸. Both loneliness and social isolation are conceptually distinct from living alone, the latter having limited utility as a proxy for either social isolation or loneliness⁹. However, we recognise that defining social isolation and loneliness is challenging, particular as researchers have used terms involving social relationships, including social isolation, loosely¹⁰. Furthermore, while we recognise social isolation and loneliness as distinct concepts, here we explore both simultaneously as the COVID-19 pandemic and measures adopted to mitigate its spread have exacerbated both isolation and loneliness.

A number of evidence reviews have highlighted the diverse range of interventions to alleviate loneliness amongst older adults in a variety of settings^{11–14}. In the main, these have been face-to-face interventions, either in groups or between individuals. During the height of the first wave of COVID-19 pandemic, and during subsequent waves, these interventions were of limited utility as lockdown regulations in many countries confined the vast majority of the population to their homes, except for essential outings. In this period all opportunities for face-to-face social contact outside the home were curtailed, and visiting friends and family for social contact prohibited. Even as these regulations were eased social distancing has restricted opportunities for social interaction, by restricting face-to-face connections and physical contact. During this period there has been considerable growth in the use of remote communication tools including telephones, videoconferencing, or other internet 'chat' facilities.

This rapid review examines evidence specifically on whether befriending, social support, and low intensity psychosocial interventions delivered remotely can reduce social isolation or loneliness among older adults. Specifically, the aims are to:

- (i) Identify existing systematic reviews on befriending, social support, and low intensity psychosocial interventions delivered remotely for older adults.
- (ii) Synthesise review-level findings on the nature and effectiveness of these interventions.
- (iii) Generate new understandings on how interventions work and which core components and processes are associated with successful interventions, using the innovative methods of Intervention Component Analysis and Qualitative Comparative Analysis.
- (iv) Map the review-level and study-level evidence to better understand evidence gaps.

This paper is an abridged summary of a full report, available elsewhere, containing further details¹⁵. The rapid review was conducted in a short timescale (four weeks for the main body of work), and adopted a review of reviews approach to meet these timescales.

Methods

We followed a systematic ‘review of reviews’ methodology to synthesise evidence from related (but differing) interventions for social isolation and loneliness, to help inform decisions about different approaches¹⁶. Although broad frameworks for conducting overviews exist¹⁷, specific guidance that can be used in a directive way to carry out overviews is lacking. Nevertheless, we followed elements of practice recommended by Caird and colleagues (2015)¹⁸ in balancing some of the challenges of conducting reviews of reviews with the need to produce policy-relevant evidence at speed in the context of the COVID-19 pandemic, and the present review was developed over a four-week period in April-May 2020. We also examined how existing reviews of reviews in the area, and particularly a review conducted by Chipps and colleagues (2017)¹⁹, navigated the challenge around differences between the scope of a systematic review and the scope of the review of the reviews.

For the purposes of this review, we define ‘older adults’ as those aged 50 years and above.

The review adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) checklist for the reporting of systematic reviews²⁰. A protocol was agreed before data extraction and published on the [EPPI-Centre website](#).

Search strategy

Searches of 11 bibliographic databases and online resources across the fields of health, social care, psychology and social science were carried out on 23rd–24th April 2020. We searched: Applied Social Sciences Index and Abstracts (ASSIA)(Proquest), Emerging Sources Citation Index (Web of Science), Database of Promoting Health Effectiveness Reviews (DoPHER), Epistemonikos, Medline (OVID), NHS Evidence, PsycInfo (OVID), Social Policy and Practice (OVID), Social Sciences Citation Index (Web of Science), Social Systems Evidence and Sociological Abstracts (Proquest).

The search terms reflected four concepts that needed to be present in each of the study citations:

- 1) **Population:** older and middle-aged populations aged 50+ years in the community and in residential settings.
- 2) **Interventions** that enable remote delivery: technology, remote communication, telephone, helplines, self-help, bibliotherapy.
- 3) **Outcomes:** loneliness, social isolation (or measures focussed on a particular aspect e.g. social support and social contact).

- 4) **Study design:** systematic reviews.

An example search history for Medline is presented in the full report¹⁵ and as *Extended data*²¹.

Inclusion and exclusion criteria

Inclusion and exclusion criteria were also based on the Population, Intervention, Comparator, Outcome and Study Design (PICOS) framework:

Population: We included reviews on ‘older’ adults age 50+ (see protocol for further details²²). Participants could be located in a variety of settings in the community or residential care, although reviews of interventions delivered to older adults in hospital settings were excluded. Studies included older adults who were socially isolated, lonely, or who were otherwise at risk of loneliness or isolation.

While we identified all reviews on older adults, we only synthesised evidence from reviews focussed on populations of older adults. We did not synthesise evidence from reviews focussed exclusively on particular groups of older adults, specifically older caregivers (see 15), although interventions including caregivers are well represented in the evidence presented. The decision to synthesise evidence from a subset of reviews was in line with the rapid timescales of the review (see Figure 1).

Intervention: Included reviews examined interventions that sought to reduce levels of social isolation or loneliness, through strengthening individuals’ social contacts and social relationships (e.g. befriending and social support), or through low intensity psychosocial interventions (e.g. internet-delivered CBT - iCBT), using remote methods and technologies. Interventions were delivered on a one-to-one basis (e.g. befriending), or as remote group-based interventions (e.g. discussion groups). We did not include interventions that examined the use of social robots, pets or virtual pets, or reviews solely focused on the use of technology for educational or training purposes.

Comparator/control: We included reviews that included studies with most forms of control group (randomised and non-randomised) and those without a control group (pre-post designs). Reviews on the implementation of interventions, including qualitative evidence syntheses were also included.

Outcomes: Included reviews measured social isolation or loneliness as a primary outcome.

As outlined earlier, we conceptualise loneliness as an emotional response by individuals when there is a ‘deficit between their desired and actual quality and quantity of social engagement and relationships^{6, p64}. Social isolation reflects the number of social contacts that people have⁷, and people who are socially isolated tend to have social networks of low density that are not maintained through frequent engagement⁸. Based on previous reviews ⁸, we expected various measures of loneliness and

Figure 1. Flow of studies through the review.

social isolation to be reported, including measures that ranged in the degree of subjectivity. Despite a number of tools for measuring social isolation being available¹⁰, many of the available comprehensive measures of social isolation in particular are underutilised within many intervention studies (for example 23), and we planned to draw on narrower measures where necessary and appropriate, for example reports of contact with family and friends and levels of social support. This is in accordance with conceptualisations of social isolation elsewhere, for example with social support being viewed as indicative of individuals' everyday social worlds and a key indicator of social isolation in several studies (see, for example 24).

Study design: We defined systematic reviews as those that met at least four of the following criteria²⁵:

1. Were inclusion/exclusion criteria reported?
2. Was the search adequate?
3. Were the included studies synthesised?
4. Was the quality of the included studies assessed?
5. Were sufficient details about the individual included studies presented?

We did not include any other reviews of reviews, but used these to identify additional systematic reviews. We included unpublished manuscripts. We did not place any restrictions on date of publication, although only reviews in English were selected.

As we were expecting some heterogeneity in the question being addressed by reviews, and expected this to be reflected in the design of primary studies included within reviews, we did not specify that source systematic reviews had to be confined to a particular study design. In line with previous reviews in the field, we expected studies measuring quantitative outcomes to be composed of single-group pre-post studies, non-randomised comparison studies, and randomised comparison group studies.

Study selection and data extraction

We exported search records to EPPI-Reviewer web²⁶ and de-duplicated the records. Title and abstract screening was undertaken independently by three reviewers (DK, EB, PH) following joint screening of 204 citations (10%) to ensure consistency (with levels of agreement reached 93%). For records included for full-text screening, each record was examined in duplicate, and reviewers met online to reconcile any differences. Reasons for exclusion are reported in Figure 1.

Systematic reviews in this area often include a mix of eligible and ineligible interventions. In line with previous overviews¹⁹, and in addition to the criteria outlined above, systematic reviews were included if they:

- (i) contained only or a majority of interventions within scope; or
- (ii) contained separate evidence tables, or defined sections of evidence tables, presenting evidence on interventions within scope; or
- (iii) contained separate synthesis sections presenting evidence on interventions within scope.

Interventions in scope were befriending, social support, and low intensity psychosocial interventions, delivered remotely, to reduce social isolation and loneliness among older adults. We did not include reviews where only a single study within the review met our criteria. Individual studies reported within systematic reviews were identified as relevant, using the same inclusion criteria as above (albeit applied at the study, not review level) and after agreement of two reviewers.

Data were extracted by two reviewers and any differences agreed in online reconciliation meetings. We extracted the following data from reviews:

- Lead author and team;
- Year of publication;
- Number of primary studies included in the review;
- Primary study design(s) (e.g., RCT studies, qualitative studies);
- Aims of review and main topic focus; (e.g. if focussed on social isolation/loneliness);
- Target population (e.g., if focussed on particular group e.g. bereaved older adults);
- Participant characteristics (e.g., age, gender);

- Intervention approaches in primary studies (e.g., type of remote intervention);

- Synthesised outcomes/key findings relating to social isolation and/or loneliness; secondary outcomes relating to implementation and adverse effects;

- Quality assessment characteristics and rating.

Once eligible reviews had been identified, primary papers were extracted if they met the criteria in line with the review inclusion and exclusion criteria stated above. That is:

Population: older adults, located in the community or residential care, socially isolated or lonely, or at risk of social isolation or loneliness.

Intervention: interventions that sought to reduce levels of social isolation or loneliness, through strengthening individuals' social contacts and social relationships. Interventions were delivered on a one-to-one basis (e.g. befriending), or as remote group-based interventions (e.g. discussion groups).

Comparator/control: studies with most forms of control group (randomised and non-randomised) and those without a control group (pre-post designs).

Outcomes: measures of social isolation or loneliness, including measures focussed on a particular aspect, such as social support.

Critical appraisal

Included systematic reviews were critically appraised using AMSTAR-2²⁷ by two reviewers (DK/BH and EB/PH). Criteria were summed and categories of quality created based on the AMSTAR-2 assessment (low risk of bias: equivalent to high confidence in AMSTAR-2; unclear: equivalent to moderate confidence; and high risk of bias: equivalent to low or critically low confidence).

The quality of the primary studies was reported where it had been assessed by the review authors. Not all reviews included a quality assessment of their included studies.

Data synthesis

Descriptive analysis of reviews and studies. We produced textual descriptions of the reviews and their findings and presented this in tabular form to develop a preliminary understanding of the evidence. The results also helped to populate an evidence map (see later synthesis).

Narrative synthesis of the evidence. A narrative synthesis was conducted to examine review-level and study-level findings. The narrative synthesis focussed on the outcomes of befriending, social support, and low intensity psychosocial interventions delivered remotely. Building on the descriptive analysis, we followed guidance outlined elsewhere²⁸.

Intervention Component Analysis and Qualitative Comparative Analysis. We drew on two complementary synthesis methods – Intervention Component Analysis (ICA) and Qualitative Comparative Analysis (QCA) – and applied these to primary studies

contained within the reviews that presented quantitative results, to understand how interventions ‘worked’. The first approach, ICA²⁹, is an inductive approach developed in response to the poor reporting of intervention processes. It involves (a) inductively coding the nature of intervention features (i.e. components) and (b) using trialists’ informally-reported experiences of implementing the intervention (i.e. information usually located in introduction and discussion sections of trial reports, which is usually not incorporated into analysis)²⁹. This information is then used in conducting the QCA.

The second approach, QCA, is applied to numeric data and is based on set-theory³⁰. QCA is employed as a solution to the challenge of analysing data containing a small number of studies (known as cases in QCA terminology), each with an extensive array of factors that may trigger a given outcome³¹. This ‘small N-many variables’ challenge is similar to that faced by systematic reviewers, and Thomas and colleagues provide one of the first examples where QCA was utilised within a systematic review to understand configurations of intervention components that were aligned with ‘successful’ interventions³². Studies were eligible for QCA if they reported quantitative findings (see *Results*). We identified studies as belonging to both ‘condition sets’ (i.e. belonging to a distinct set of studies distinguished by the presence or absence of different characteristics or processes) and ‘outcome sets’ (i.e. belonging to a group of studies differentiated by whether they are considered most effective or least effective). Ultimately, we were interested in establishing which condition sets ‘overlapped’ with successful outcome sets. The goal of QCA is to identify the simplest expression of characteristics/processes that lead to effective interventions; to find the simplest expression we drew on Boolean minimisation. We followed standards of good practice that have been laid out elsewhere in conducting the QCA³³. Further explanation of the approach is provided alongside the results.

A fundamental element of QCA is the selection of an appropriate theory to base the analysis on, and to help identify suitable evidence to extract as part of the ICA (described in the methods section). To understand which processes might be important to incorporate in interventions – regardless of specific mode of delivery (i.e. videoconferencing or internet chat group) – we drew on Robert Weiss’s³⁴ ‘Fund of Sociability’ theoryⁱ. The theory is intended to capture assumptions, content, and functions of social ties that can help to support developing social relationships. The theory specifies six characteristics of social interactions and relationships that are necessary for well-being and the avoidance of loneliness³⁴. **Table 1** outlines the six categories, their definitions and how we interpreted them in relation to the interventions in the QCA.

ⁱ This theory also served as the conceptual framework underpinning one of the included studies 49. Weinert C, Cudney S, Hill WG. Rural women, technology, and self-management of chronic illness. *The Canadian journal of nursing research*= *Revue canadienne de recherche en sciences infirmières* 2008;40(3):114.

Results

Review and study characteristics

The literature searches identified 2,715 citations. After duplicates were removed, 2,057 citations were screened at title and abstract level, identifying 75 possible studies for inclusion. Full texts were obtained for all 75 records, with nine potential reviews identified and five included for synthesisⁱⁱ (see **Figure 1**). Not all of the primary studies within these five reviews met our inclusion criteria (see *Methods*) and from the 112 primary studies included across the five reviews, we identified 18 studies as eligible for synthesis.

Review populations. The reviews covered a range of populations, using different definitions and age thresholds for ‘older adults’, with a combined age range of 50–95. The settings were not always clearly stated, but were primarily older adults’ own homes, nursing homes, or supported living facilities, in North America, Europe and Taiwan. Whilst some reviews contained studies focused on the general older adult population, others included studies of people with multiple chronic conditions, specific conditions (such as Alzheimer’s Disease, or breast cancer), or in a particular geographical area.

Review study designs. RCTs, quasi-experimental cohort studies, survey studies, and qualitative (semi-structured interviews and focus groups) were all represented. Three of the five reviews conducted quality appraisals on the included studies^{35–37}, one evaluated only the effectiveness of the technologies within the studies, not the quality of the study itself³⁸, and one did not report any quality appraisal³⁹ⁱⁱⁱ.

The reviews contained studies reporting interventions using various technologies to deliver remote befriending, social support or low intensity psychosocial interventions including those in scope (e.g. video-communication and telephone befriending) and those out of scope (e.g. computer training and internet training). There was a range of different outcome measures within the reviews, although all contained some measure of loneliness or social isolation.

ⁱⁱ Four of the reviews focussed exclusively on interventions to support caregivers: 40. Corry M, Neenan K, Brabyn S, *et al.* Telephone interventions, delivered by healthcare professionals, for providing education and psychosocial support for informal caregivers of adults with diagnosed illnesses. *Cochrane Database of Systematic Reviews* 2019(5); 41. Elvish R, Lever S-J, Johnstone J, *et al.* Psychological interventions for carers of people with dementia: A systematic review of quantitative and qualitative evidence. *Counselling and Psychotherapy Research* 2013;13(2):106–25.; 42. Hopwood J, Walker N, McDonagh L, *et al.* Internet-based interventions aimed at supporting family caregivers of people with dementia: systematic review. *Journal of medical Internet research* 2018;20(6):e216.; and 43. Lins S, Hayder-Beichel D, Ruecker G, *et al.* Efficacy and experiences of telephone counselling for informal carers of people with dementia. *Cochrane database of systematic reviews* 2014(9). In view of the need for rapid evidence synthesis, we excluded these reviews from the synthesis, which only included the reviews focussed on the general older population (this included care givers in some instances).

ⁱⁱⁱ Note that while Beneito-Montagut and colleagues’ study (2018) was a self-defined ‘review of the literature’, it was deemed to sufficiently meet the criteria of a systematic review for this review of reviews.

Table 1. Six categories of the fund of sociability theory.

Category	Definition in Weiss 1969	Application in Qualitative Comparative Analysis
1. Intimacy (which we describe as close relationships in our narrative)	<i>'An effective emotional integration in which individuals can express their feelings freely and without self-consciousness' (p.38)</i>	Intervention supports participants to express feelings freely and without self-consciousness (e.g. opportunities for unstructured discussions with peers)
2. Interaction	<i>'Participants share concerns, either because of similar situations ("we are in the same boat") or because they are striving for similar objectives' (p.39)</i>	Target population has shared experience (e.g. being a carer, stroke survivor etc.) and shared characteristics (e.g. women only, people of similar age / SES etc.)
3. Nurturance	<i>'Opportunity for nurturant behavior ... absence of this function may be signalled by a sense that one's life is unfulfilled, meaningless, and empty of purpose' (p.39)</i>	Intervention values / encourages participant sharing of experiences for others benefit (e.g. group discussions / bulletin boards invite participants to share experiences)
4. Self-worth (control)	<i>'Relationships that attest to an individual's competence in some role' (p.39)</i>	Intervention enhances sense of competence by offering control over design / delivery (e.g. participants determine frequency of discussion groups / identify topics for discussion)
5. Availability	<i>'Assistance that is not limited in time and extent' (p.40)</i>	Intervention is available continuously and provides opportunities for asynchronous and 'real-time' interactions (e.g. website information resources (continuous), discussion board (asynchronous), videoconferencing / 'live-chat' (real-time))
6. Support	<i>'This function might be characterized as guidance, and may be provided by mental-health professionals such as social workers or psychiatrists or by ministers and priests, among others.' (p.40)</i>	Services include some form of pastoral care (e.g. light-touch oversight of a discussion forum by professionals or opportunities for participants to contact professionals for advice)

Risk of bias assessment of included reviews. All of the reviews were deemed to be of low or critically low quality (displayed as having a high risk of bias in Table 2). Although all had reasonably clearly defined PICO components and had conducted reasonably comprehensive search strategies, the majority had failed to prepare a protocol, and many failed to justify the choice of study selection. This latter concern was particularly problematic where authors had included studies of various designs.

Primary study characteristics. Befriending, social support and low intensity psychosocial interventions reported in the 18 primary studies fell into five categories reflecting modes of delivery:

- Supported videoconferencing to alleviate loneliness⁴⁴⁻⁴⁸.
- Telephone befriending to reduce social isolation^{49,50}.
- Online discussion groups/forums to reduce social isolation and/or loneliness, or to improve/maintain social connectedness⁵⁰⁻⁵⁷.
- Supported use of social networking sites for mitigating social isolation and loneliness⁵⁸.
- Multi-tool interventions (PC, training, messaging, chat groups) to reduce loneliness and/or social isolation, or increase social connectedness^{54,57,59,60}.

Further primary study characteristics, including their populations, details of implementation, methodological details including how the outcome was measured, and outcomes as found in Table 1.

Narrative synthesis of findings

Supported videoconferencing to alleviate loneliness. Four reviews included a total of three qualitative studies, three quantitative studies, and one mixed-methods study, on supported video-communication interventions. Five studies involved supporting older adults to communicate with family members^{45-48,61}, with the other two reporting on the videoconferencing element of the ACTION service in Sweden and Norway^{44,56}. The qualitative evidence suggests the interventions were generally regarded positively by older adults, with increased feelings of connection with their family members⁴⁵. The quantitative evidence showed some evidence of decreases in feelings of loneliness and increases in social support scores. Two quantitative studies^{47,48} found reduced feelings of loneliness at one week, three months and 12 months, although this achieved statistical significance in only one study^{iv47}. Torp *et al.*'s⁵⁶ mixed-methods pilot cohort study, employing questionnaires and focus groups,

^{iv} It was also unclear whether the intraclass correlation had been accounted for in calculating measures of effect.

Table 2. Characteristics of included reviews and primary studies.

Included review	Type of review and study numbers	Review focus / aim	AMSTAR2 rating	Primary studies included in our review	Population and setting	Study design and intervention	Measure of loneliness /social isolation	Outcomes	Quality tool used in review (rating)
Beneito-Montagut et al. (2018).	Literature review that included a systematic review of 25 studies	To review previous research that investigates the relationship between internet use for communication and social isolation and loneliness, including its effects on social relationships in later life.	Critically Low	Ballarneye et al. (2010)	Participants: n=60 older adults, age range 69-85. 3 males, 1 female. Living at home, Australia.	Pilot project: to evaluate use of social network site (SNS) in the Age-Active Ageing (A3) One-on-one tutoring. In how to use the social networking site, supported throughout the intervention by project team member.	Qualitative: Connectedness (proxy for isolation)	The utilisation of a SNS has the potential to reduce loneliness and improve mental health (especially in the evening) and connectedness.	None
				Tsai & Tsai (2011)	Participants: n=90 older adults, age over 60 yrs Experimental (n=40; females=22) Control (n=50; females=30). Nursing home residents, Taiwan.	Quasi-experimental: Supported video-conferencing. Experimental group received at least 5 minutes/week for 3 months of video-conference interaction with their family members in addition to usual family visits. Videoconferencing supported by trained research assistant. Comparison group received regular family visits only.	Loneliness (UCLA Loneliness Scale)	Alleviated perceived loneliness at 3, 6, and 12m after the intervention. Experimental group had significantly lower mean loneliness and depressive status scores at 3m (-5.40, P < .001), -2.64, P < .001, respectively), 6m (-6.47, P < .001), -4.33, P < .001, and 12m (-6.27, P = .001, -4.40, P < .001) compared with baseline than those in the comparison group. Experimental group had significantly lower mean scores at 3m (-0.44, P = .03), 6m (-0.44, P = .03), and 12m (-0.44, P = .03) compared with baseline than those in the comparison group. Mean change in emotional social support at 3m (0.74, P < .001) and 12m (0.61, P = .02), and in appraisal support at 3m (0.74, P = .001).	None
								No significant differences between experimental and control group regarding informational social support.	
Included review	Type of review and study numbers	Review focus / aim	AMSTAR2 rating	Primary studies included in our review	Population and setting	Study design, intervention, comparison	Measure of loneliness /social isolation	Outcomes	Quality tool used in review (rating)
Bennett (2015)	Systematic review (part of doctoral dissertation) 10 studies	To answer what impact video-communication has on older adults existing relationships and their social relationships when it is used to communicate with friends and relatives.	Critically Low	Demiris et al. (2008)	Participants: n=4 older adults, age 65+. Nursing home residents, USA.	Qualitative: Supported videophone intervention. Both residents and family members were asked to conduct a videocal at least once/week for four weeks. They discussed and completed a form after each video call to document the quality of the quality. Participants supported by long-term care facility staff to use videophone.	Qualitative: Thematic synthesis	Themes reported included; participants being included into family interactions and feeling part of the family; participants valuing seeing the other person's facial expressions, videoconferencing reduced feelings of isolation, videoconferencing had the potential for emotional conversations and concerns with using the technology.	None
				Schwinderhammer, (2013)	Participants: n=80 older adults, age 65+ Intervention condition: n=40, mean age 85.42 (5.88). Control condition: n=40, mean age 86.82 (5.28). Nursing home residents, USA.	Quasi-experimental within-between repeated measures: Supported video-conferencing intervention. Experimental group received ten Skype sessions over a 14-week period, 10-30 minutes each time. Supported by researcher. Control group had regular care.	UCLA Loneliness Scale	The results indicate the intervention group felt a significantly decreased level of loneliness after videoconferencing with family members than before, while the control group did not significantly change their loneliness level from pre-intervention to post-intervention.	None
				Tsai and Tsai (2010)	Participants: n=34 older adults, age range=60-95. Nursing home residents, Taiwan.	Qualitative: Supported video-conferencing intervention. Videoconferencing programme once/week for three weeks. Contact with spouse, child or grandchild. The programme was either MSN or SKYPE, using a large laptop. Residents assisted to use technology by a trained research assistant, who spent at least five minutes/week with residents at the appointment time.	Qualitative: Thematic synthesis	Themes included: enriched life, second best option for visiting, life adjustments and true picture of family life.	None
				Tsai and Tsai (2011) (Duplicate - see description in Beneito-Montagut (2018))					None
				Tsai et al. (2010)	Participants: n=57 older adults, (experimental group, n= 24) Mean age = 78 and control group, n = 33; Mean age = 74). Nursing home residents, Taiwan.	Quasi-experimental: Supported video-conferencing intervention. Experimental group received at least 5 minutes/week for 3 months of video-conference interaction with their family members in addition to usual family visits. Videoconferencing was supported by a trained research assistant. Comparison group received regular family visits.	Depressive symptoms, loneliness and social support (Social support behavior scale (Hsling, 1999)), via self-report questionnaires, administered at baseline, three, six and twelve months.	From baseline, compared to the control groups, video-communication significantly reduced feelings of loneliness at one week ($\beta = -1.21$, $P = .002$) and three months ($\beta = -2.84$, $P = .003$), and depressive status at three months. From baseline scores, changes in emotional social support scores at one week and three months and appraisal support scores at one week and three months were found to be significantly higher in the experimental groups compared to the changes in the control groups.	None
				van der Heide et al. (2012)	Participants: n=130 older adults with no prior experience of video communication, mean age=73.2. Living in the community, acrossing homecare, The Netherlands.	Pre-post intervention study (non-randomised): CareTV - video/voice network package, (24/7 communication facility with a nurse practitioner). CareTV applications are (1) Alarm Service (2) Care Service, (3) Good morning/good evening service, (4) Wellfare and housing, and (5) Family Contact.	Loneliness scale (Jong Gierveld & Kamphuis 1985). Clients' experiences were evaluated in open questions in the survey.	Subjects in the experimental group had significantly higher mean emotional and appraisal social support scores at one week ($\beta = .61$, $p < .001$) and three months ($\beta = .68$, $P < .001$), after baseline than those in the control group.	
								Average feeling of loneliness at group level decreased significantly ($p < 0.001$) from 5.97 (sd 2.77) to 4.02 (sd 3.91) between the start and end of the study on a scale from 0 to 11. Social loneliness (5-items) as well as emotional loneliness (6-items) showed significant decreases.	

Included review	Type of review and study numbers	Review focus / aim	AMSTAR2 rating	Primary studies included in our review	Population and setting	Study design, intervention, comparison	Measure of loneliness/ social isolation	Outcomes	Quality tool used in review (rating)
Chen and Schulz (2016)	Systematic review 25 studies	Explore the effects of ICT interventions on reduced social isolation of older people	Critically Low	Cattan <i>et al.</i> (2011)	Participants: n=40 vulnerable, isolated, and/or lonely, age range = mid 50s – early 90s. Mostly living at home, UK.	In-depth interviews; Telephone befriending intervention, <i>Call In Time</i> . Telephone befriending.	Qualitative.	Reduction of social isolation, loneliness, depression, and anxiety; improved state of mind, contentment with life, confidence level, and physical health (less pain).	Effective Public Health Practice Project (EPHPP): N/A
				Savolainen (2008)	Participants: n=8 older adults with frailty, age range: mid 50s – early 90s. Living at home, Sweden	In-depth interviews; Videoconferencing in the ACTION project (ICT to support frail older adults and their family carer). Videoconferencing with family or professional carers.	Qualitative.	Seven out of the eight participants reported a positive impact on loneliness.	EPHPP: N/A
				Torp <i>et al.</i> (2008)	Participants: n=19 carers of people with stroke or dementia. Mean age 73 (57-85), M11, F8. Living at home, Norway.	Pilot cohort study without control group (mixed methods); ICT- and web-based ACTION service. Received broadband connected PC, 9 hours training over 3 weeks; connection to online discussion with peers; videophone for peer communication and contact with health personnel. Peer support facilitated by qualified nurses.	Social contacts measured by the Family and Friendship Contacts Scale. Social support was measured with a 20-item scale developed by Russell <i>et al.</i> (1980). Focus Groups conducted.	Positive and significant change in scores with regard to contact with family and friends ($P = 0.036$) and a sense of social support from other persons ($P = 0.010$). Carers explained that the frequent contact they had via the videophone and discussion forum in-between the monthly meetings was important to build social networks and friendships.	EPHPP: Weak
				Tsai <i>et al.</i> (2010) (Duplicate - see description in Bennett (2015))					EPHPP: Strong
Included review	Type of review and study numbers	Review focus / aim	AMSTAR2 rating	Studies included in our review	Population and setting	Study design, intervention, comparison	Measure of loneliness/ social isolation	Outcomes	Quality tool used in review (rating)
Khosravi <i>et al.</i> (2016)	Systematic review 34 studies	Identify ICTs that are designed to help seniors reduce social isolation and loneliness and assess the effectiveness of this technologies in supporting seniors wellbeing.	Critically Low	Balentine <i>et al.</i> 2010. (Duplicate - see description in Benetto-Montegut (2018)) Hill <i>et al.</i> (2006)	Participants: n=100 women with various chronic illnesses. Intervention: 22 weeks. Participation in an online, asynchronous chat room and health teaching units. WeChat (2005) platform used to deliver the intervention and was available 24/7, allowing women to participate at any convenient time. Access to Koffee Kiatchi, an asynchronous chat room in which they exchanged feelings, expressed concerns, provided support, and shared life experiences. Access to each other and research team via email. Peer support and expert facilitated chat room: 'Health Roundtable'. Details of comparison arm not provided.	RCT: Online self-help support group (Women to Women Program). 22 weeks. Participation in an online, asynchronous chat room and health teaching units. WeChat (2005) platform used to deliver the intervention and was available 24/7, allowing women to participate at any convenient time. Access to Koffee Kiatchi, an asynchronous chat room in which they exchanged feelings, expressed concerns, provided support, and shared life experiences. Access to each other and research team via email. Peer support and expert facilitated chat room: 'Health Roundtable'. Details of comparison arm not provided.	Social support: 15 item Personal Resource Questionnaire (PRQ) 2000 Measure of loneliness: UCLA Loneliness Scale	Statistically significant effects on social support ($p=0.38$) but not on loneliness ($p=206$).	None
				O'Connor <i>et al.</i> (2014)	Participants: n=71 dementia caregivers, age range=54-70. Living at home, USA	Pre-post intervention: 3D virtual environment. Participants interacted using avatars in a virtual chat room (Avatar Social World Life platform) in an 8-week support group.	UCLA Loneliness Scale	Lower levels of depression and loneliness across participants.	None
				Tsai and Tsai, (2011) (Duplicate - see description in Benetto-Montegut (2011)) van der Heide <i>et al.</i> (2012) (Duplicate - see description in Bennett (2015))					None None

Included review	Type of review and study numbers	Review focus / aim	AMSTAR2 rating	Studies included in our review	Population and setting	Study design, intervention, comparison	Measure of loneliness/ social isolation	Outcomes	Quality tool used in review (rating)
Morris <i>et al.</i> (2014)	Systematic Review 18 studies	To conduct a systematic review of studies that assessed the effectiveness of smart technologies in improving or maintaining the social connectedness of older adults who live at home.	Low	Barrera <i>et al.</i> (2002)	Participants: n=160 people with type 2 diabetes. Mean age 59.3 (9.4). M/F: 85. Living at home, USA	4 Group RCT: 1) Information only; 2) Personal Coach and Social Support; 3) Social Support only; 4) Personal Coach and Social Support. Internet-based support group - peer-directed (professionally supported) forum. Real time live chat discussions. Research staff monitored the forum and introduced topics for discussion.	Interpersonal Support Evaluation List (ISEL).	After 3m, individuals who participated in Internet-based social support interventions significantly increased their perceived availability of social support, relative to participants who only had computer access to information about diabetes. Effects bound for general perceptions of support as well as with a measure of support that was designed specifically for individuals who used computer-mediated social support interventions. Intervention effects $f = .27$ for the ISEL items.	Downs and Black checklist Not reported for individual studies. Range 13-22 (median 18.5). Box score = 27.
				Bond <i>et al.</i> (2010)	Participants: n=62 people with diabetes. Mean age Intervention: 66.2 (5.7). Control: 68.2 (6.2). M/F: 42. Living at home, USA	RCT: Web-based interactive intervention. Monitored self-management by nurse; weekly online educational discussion group for peer support; peer email and instant messaging. Interaction between the study nurse and the intervention participants by synchronous (instant messaging and chat) and asynchronous communication (email and a bulletin board). Study nurse, social worker and psychologist moderated sessions. Control received standard diabetes care.	Diabetes-related social support was assessed using the Diabetes Social Support Scale.	Participants who received the Web intervention improved their depression, quality of life, self-efficacy, and social support ratings, compared with the control group ($p=0.001$).	
				Dew <i>et al.</i> (2004)	Participants: n=64 heart transplant recipients and 60 care givers. Age: No data Gender: 46% female Living at home, USA	Matched controlled cohort study; multifaceted web-based intervention. Stress and medical regimen management workshops; monitored discussion groups; access to electronic communication with health professionals. Comparison groups of 40 patients and their caregivers who received only usual care were similarly assessed	Sub-scale reported on Quality of Life (QoL) reflecting social functioning (ability to interact with others and engage in social activities).	QoL in social functioning improved significantly.	
				Guatfson <i>et al.</i> (2005)	Participants: n=229 women with breast cancer. Mean age 51.6 (11.8). Living at home, USA	Cohort study; Integrated web-based programme. Participants taught how to use computer, internet and CHES system (home based, to improve QoL, including participating in a discussion group and Ask an Expert service. Web-based programme provided the resources matched to participants to make weekly phone calls to participants. Comparison group taken from separate recently completed study.	Six-item social support scale for assessing women's perception of emotional and instrumental support.	The discussion group was the most extensively used service (79.65% participants used, with average time of 66.4 minutes over 4 months). Perceived social support ($p<0.000$). Of those who used a peer advocate 77.3% felt somewhat or very much better equipped to deal with their breast cancer and that the peer advocate helped them cope (some, what or very much so) with their breast cancer.	
				Hill <i>et al.</i> (2006), (Duplicate - see description in Khasrawi <i>et al.</i> (2016)) Torp <i>et al.</i> (2008), (Duplicate - see description in Chen & Schulz (2016))					
				Wehnert <i>et al.</i> (2008)	Participants: n=233 women with various chronic illnesses. Mean age 51.8 (2.17). Living at home, USA	RCT: Web-based 22 week intervention programme (Women to Women Program). Peer-led virtual support group, and self-study web skills. Online, asynchronous, peer-led support group (Koffee Klatch) and health teaching units, prepared by the research team, supplemented by asynchronous, expert-facilitated discussions (Health Roundtable). Comparison group received regular care.	Loneliness: UCLA Loneliness Scale. Social support: Personal Resource Questionnaire 2000.	Participants who had access to the online discussion boards showed a significant increase in self-efficacy ($P = .04$) and social support ($P = .03$) compared to the control groups who had no intervention.	
				Wehnert <i>et al.</i> (2011)	Participants: n=309 women with various chronic illnesses. Mean age 55.5. Intervention: 56.1 (7.7), control 55 (9.1). Living at home, USA	RCT: Web-based 11-week intervention (Women to Women Program). Peer-led virtual support group (asynchronous forum, Sharing Circle to exchange feelings and life experiences, give and receive support, discuss issues related to the self-study health teaching units, and share health information) and (b) a series of self-study health teaching units focused on Web skills and the five skills of self-management. Monitored by advance practice nurse. Two comparator arms: Less-intense intervention group (self-study health units), and regular care.	Loneliness: UCLA Loneliness Scale. Social support: Personal Resource Questionnaire 2000.	Self-esteem ($P = .018$), acceptance ($P = .001$), depression ($P = .010$), stress ($P = .005$) and loneliness ($P = .001$) were significantly lower in the intervention group compared to the control group. No statistically significant differences were seen between the two groups in social support ($P = .097$).	

Table 3. AMSTAR-2 ratings for included systematic reviews (displayed as risk of bias).

Review	1. PICO components	2. Protocol	3. Study design explanation	4. Comprehensive search strategy	5. Duplicate study selection	6. Duplicate data extraction	7. Details of excluded studies	8. Description of included studies	9a. Risk of Bias (RoB) assessment (RCTs)	9b. RoB assessment (NRSIs)	10. Funding sources	14. Heterogeneity	15. Publication bias	16. Reports conflicts of interest	Overall rating of quality
(Beneito-Montagut <i>et al.</i> , 2018)	+	-	-	?	+	+	?	+	?	-	-	-	?	-	-
(Bennett, 2015)	+	-	-	+	-	-	+	+	?	?	-	+	?	-	-
(Chen and Schulz, 2016)	+	-	+	?	-	+	+	+	?	?	-	+	?	+	-
(Khosravi <i>et al.</i> , 2016)	+	-	+	?	+	-	?	?	-	-	-	-	?	-	-
(Morris <i>et al.</i> , 2014)	+	-	-	?	+	+	?	+	?	?	-	+	?	-	-

+ = low risk of bias (equivalent to high confidence); ? = moderate or unclear (equivalent to moderate confidence); - = high risk of bias (equivalent to low or critically low confidence); Note chart shows only those items relevant to all included reviews.

also found that the video phone was important for building and maintaining relationships. It is important to note that all interventions included ongoing support to use the technology. See Table 1 for contextual details of reviews and studies on videoconferencing.

Telephone befriending to reduce social isolation. Two reviews included a total of one qualitative and one quantitative study reporting on forms of telephone contact, one of which was a study of telephone befriending. Cattan *et al.*⁴⁹ reported on the *Call in Time* intervention, with qualitative findings from 40 participants. Telephone calls were made to older adults by volunteers, with a project co-ordinator managing the process. Findings included reduced feelings of social isolation, loneliness, depression and anxiety; improved state of mind, contentment with life, confidence level, and physical health (less pain). This study built on an earlier evaluation report that presented data used for the QCA⁶²; this evaluation report was not directly included in any of the reviews, but quantitative data presented within this report suggested that participants had lower wellbeing and social support after the intervention, albeit with a number of caveats.

The only other included study to incorporate telephone contact was Gustafson *et al.*⁵⁰, from the Morris *et al.*³⁷ review, where one element of the intervention was to match up participants with peer advocates, who engaged in weekly phone calls. This was not a telephone befriending service, as the peer advocate had

a different role to that of a befriender. Findings showed that, of those who used a peer advocate 77.3% felt somewhat or very much connected with their peer advocate, and 81.6% felt that the peer advocate helped them cope (somewhat or very much so) with their breast cancer. Perceived social support increased significantly over the four months, but the intervention included more elements than just telephone support (computer and internet training, discussion group, 'ask an expert' service and written guidelines).

Online discussion groups/forums to reduce social isolation and/or loneliness, or to improve/maintain social connectedness. Two reviews contained eight quantitative studies and one mixed-methods (questionnaires and focus groups) on online discussion groups and forums. The studies included synchronous and asynchronous communication: real-time chat discussions, instant peer-messaging, email contact with professionals, and discussion boards. Interventions were designed to support women with chronic illness or breast cancer^{50,54,57,60}, people with diabetes or heart transplant recipients⁵¹⁻⁵³, and caregivers of people with dementia or stroke survivors^{55,56}. The qualitative evidence suggested that discussion groups helped older adults to build social networks and friendships and to feel more familiar with people through regular connections^{55,56}. The quantitative evidence showed mixed results with regard to loneliness and social isolation. The majority of studies showed increases in social support, but only two showed reductions in loneliness, with four studies not measuring loneliness at all. The asynchronous

chat room ‘Koffee Klatch’ in Hill *et al.*⁵⁴ primary study provided a forum for women with chronic illnesses to share their feelings, concerns, life experiences and provide support to each other over 22 weeks, resulting in significant improvement in social support, but not in loneliness, compared to the control group. The Sharing Circle in Weinert *et al.*⁵⁷ provided the same opportunities, with the addition of discussion of self-study units and internet-based health information. This study saw statistically significant improvements in loneliness, but not in social support, compared to the control group. See [Table 1](#) for details of reviews and studies on online discussion groups/forums.

Supported use of social networking sites for mitigating social isolation and loneliness. Two reviews included the same study on social networking sites⁵⁸. The authors of this qualitative study found that the utilisation of a bespoke social networking site had the potential to reduce loneliness in older adults, as there were positive impacts on temporal loneliness (especially in the evening) and on connectedness. Review authors suggested that older adults were more interested in a smaller number of strong relationships mediated through the internet, than they were in a larger network with weak ties. They report that perceived value could have been an issue for older adults, which may have been more obvious through supported social networking service interventions such as that reported by Ballentyne *et al.*⁵⁸.

Multi-tool interventions (PC, training, internet use, messaging, chat groups) to reduce loneliness and/or social isolation, or increase social connectedness. Three reviews included a total of four quantitative studies on multi-tool interventions. van der Heide *et al.*⁵⁹ report on the Care TV package for people receiving home care in The Netherlands. This video and voice network allowed clients to communicate round-the-clock with a nurse practitioner. They received a ‘Good Morning/Goodnight’ call and could use the video facility to call family members. Average feelings of loneliness decreased substantially, with social and emotional loneliness showing pronounced decreases. The three other studies reported on web-based discussion groups in the Women-to-Women programme, with mixed results regarding levels of loneliness and social support. Weinert *et al.*⁵⁷ reported on an RCT of a web-based discussion groups, with a peer-led online support group and self-study units supported by an Advance Practice Nurse. Improvements were found in loneliness, but there was no significant difference in social support between the intervention and comparison groups, following the 11-week intervention. Weinert *et al.*⁶⁰ found significant increases in both loneliness and social support, compared to the control group, over the 22-week intervention. Hill *et al.*⁵⁴, found statistically significant effect on both social support and loneliness after 22 weeks. See [Table 1](#) for contextual details of reviews and studies on multi-tool interventions.

Intervention Component Analysis and Qualitative Comparative Analysis

QCA and ICA were undertaken to help us further identify the processes and mechanisms that were common across the interventions described in [Table 2](#) and the narrative synthesis. To undertake QCA, we first conducted ICA to understand the

nature of the interventions. We inductively coded the nature of intervention features (i.e. components) and used trialists’ informally reported experiences of implementing the interventions (e.g. author reflections reported in introduction and/or discussion sections) to understand the importance and underlying mechanisms of particular features²⁹.

Theory selection and setting up the QCA. A fundamental element of QCA is the selection of an appropriate theory to base the analysis on, and to help identify suitable evidence to extract as part of the ICA (described in the methods section).

Our QCA built on the earlier descriptive and narrative synthesis, and addressed the question: ‘Do the characteristics of social interactions and relationships stated in the fund of sociability theory explain differences between remotely delivered interventions found to be effective compared to those found to be ineffective?’ To gain familiarity with the studies, and attempt to gain ‘deep case knowledge’, we started by reading and re-reading the studies.

Selection of studies (cases)^v for the QCA. We focussed on studies that met our criteria for the QCA including that they (a) presented quantitative results, (b) were remotely delivered, (c) focussed on older adults, and (d) actively sought to strengthen social relationships or prevent/offset loneliness. From the 18 primary studies described above, 12 met these criteria.

Developing a data table. QCA is based on set-theory with sets differentiated as belonging to a successful and unsuccessful set on the basis of their outcome. The outcome can be based on an objective measure or subjective or quality measure⁶³, and on a single measure or a composite indicator⁶⁴. The allocation of studies into a successful set and unsuccessful set can follow different strategies. Firstly, success may be defined through the observation of clinically or statistically significant change in the outcome (for example [65](#)). A second approach is for the researcher to set thresholds for determining success. A third strategy is ensure (approximately) equal representation of un/successful cases by ranking cases according to their effectiveness and allocating studies into un/successful outcome sets. A fourth strategy is to use a more qualitative approach where additional characteristics besides the outcome value are considered to ensure a representation of studies in the un/successful outcome set⁶⁶. In a systematic review, stratifying studies by their characteristics to ensure a diversity in study size or study design among successful studies may ensure a more informative solution is produced.

To generate our outcome sets, and group interventions as being ‘successful’ or ‘unsuccessful’, we calculated an estimated effect size for each study. Effect sizes are used differently within QCA as opposed to meta-analysis; i.e. as a guide to allocating

^v Terminology is used when conducting QCA that is distinct from other research approaches. This includes the use of ‘cases’ to denote studies; we use both terms interchangeably as appropriate.

studies to successful (most effective) or unsuccessful (least effective) outcome sets, rather than to provide a pooled estimate of effect with precision. Most effect sizes were based on measures of social support, which we regarded as a measure of a particular aspect of social isolation. The exceptions were Schwindenhammer⁴⁶ and O'Connor, *et al.*⁵⁵ where a measure of loneliness was the only suitable outcome available. However, we did attempt to express the effect sizes in a common rubric where possible, e.g. prioritising post-test measures for studies that involved randomisation of participants or clusters (five studies), and change measures where these data were not available. For those studies with a comparator group (eight studies), effect sizes were calculated in the standard way see Thomas *et al.*, 2017⁶⁷; for those studies that employed a pre- and post- evaluation design an effect size was estimated based on changes in the pre- and post- individual scores divided by the standard deviation at pre-test^{68,69}; in some cases this involved using mean differences as proxy information and other approximations.

Using the effect size for indicative purposes, we grouped interventions into those that were 'successful' (studies with effect sizes over 0.5), 'partially successful' (studies with effect sizes between 0.2 and 0.5) and 'not successful' (studies with effect sizes under 0.2 or suggested negative impacts) based on thresholds suggested by Cohen⁷⁰ for interpreting effect sizes. However, combining the different study designs, and particularly those with and without a comparison group, using the same approach could lead us to overstate the effectiveness of studies without a comparison group. To mitigate this possibility we also present the results of a sensitivity analysis, where we imposed an additional 'penalty' on studies without a control group – studies with effect sizes of 0.5 and over were deemed to provide partial evidence of success (0.66); studies with effect sizes between 0.2 and 0.5 were deemed to provide weak evidence of being 'not successful' studies (0.33); while studies with values lower than 0.2 were deemed to provide strong evidence of being 'not successful' (0). This is akin to adding in additional 'qualitative' information – in this case on study design – to distinguish studies as belonging to a successful and unsuccessful outcome set⁶⁶. We also examined the potential impact of omitting these four studies, although this is not a preferred option given that QCA models typically need 10 or more cases as a minimum.

To create our data table, a coding scheme was developed to determine whether the conditions reflecting the fund of sociability processes were actually present in the studies (see 15). The results of this coding and the data table are presented below in Table 4.

Truth table. As we had a limited set of cases for the number of conditions, our analytical strategy involved first creating a 'truth table' based on six conditions, and then producing a reduced truth table containing four conditions and minimised solution³³. A 'truth table' sorts cases according to the configuration of conditions they exhibit. Although we noted that both 'availability' and 'control' were conditions generally only observed in successful intervention studies, they did not appear to be as critical to outcome success as the other conditions, appearing in fewer studies. Our reduced truth table thus contained four

conditions (intimacy, interaction, support and nurturance) with five of a possible 16 configurations represented (Table 5). Two configurations are observed as triggering a successful outcome; in one, supported by five studies, all four conditions are present; in the second, supported by two studies, three of four conditions are present. On the right side of the table is a column marked consistency; this indicates the strength to which studies that belong to the condition set are also a subset of the outcome set. A value of 1 indicates perfect consistency; all cases in the configuration are strong members of the condition set and the successful outcome set; and there is strong evidence that these intervention characteristics trigger successful outcomes. A value of 0 indicates perfect inconsistency and there is no evidence that these intervention characteristics trigger successful outcomes. Values in between indicate some degree of ambiguity, which was expected given that we used a "fuzzy-set coding scheme" which allowed studies to be partial members of sets (using a value of 0.85 to denote membership).

Boolean minimisation and formation of a solution. We applied Boolean minimisation to obtain the simplest expression of those conditions (intervention processes) that were associated with triggering a successful intervention. We developed a complex solution based on the observed data only, and found that those interventions that ensured the following processes took place were those in the successful outcome set:

INTIMACY and INTERACTION and SUPPORT

Within QCA, information from unobserved configurations (logical remainders) can be used to simplify the solution and check the quality of the solution. We incorporated these logical remainders to develop two further solutions^{vi}, although incorporating logical remainders in this model did not help to simplify the solution above. Our model and details of its fit are presented below (Table 6). The high consistency value for the solution suggests that when this configuration of conditions is observed in an intervention, it is generally sufficient to trigger a successful intervention (i.e. a substantial change in social support). The coverage statistic suggests that the model broadly accounts for the successful interventions observed.

Sensitivity and additional technical quality checks. Using an alternative measure of effect size that incorporates a 'penalty' reflecting the greater uncertainty around pre-post studies, we re-ran the analysis described above. The truth table (Table 7) with this alternative outcome showed one configuration of successful studies. This suggested that studies which incorporated all three processes observed earlier, as well as nurturance, were those that triggered a successful outcome (using a slightly lower consistency value of 0.825, which is still well within recommended thresholds³³). Coverage was slightly lower for this solution, although the solution still provided a comprehensive explanation of why some interventions were successful (Table 8). The replication of the same three core conditions provides a degree of triangulation that our main solution

^{vi} Known as a parsimonious and intermediate solution.

Table 4. Data table for Qualitative Comparative Analysis.

Study	Effect Size Estimate	Effect Size Set	Effect Size Set (sensitivity analysis)	Intimacy	Interaction	Control	Nurturance	Support	Availability
Barrera	0.530	1	1	1	0.66	0.66	1	1	1
Bond	0.634	1	1	1	1	0.33	1	0.66	1
Gustafson	0.619	1	1	1	1	0.33	1	0.66	0.66
Tsai 2010 ⁹	0.036	0	0	0.33	0	0	0	0.33	0
Tsai 2011 ¹⁰	0.051	0	0	0.33	0	0	0	0.33	0
Weinert 2008 ¹¹	0.314	0.66	0.66	1	1	0.66	1	1	0.66
Weinert 2011	0.131	0	0	1	1	0.33	1	0	0.66
Schwindenhammer ¹²	-0.098	0	0	0.33	0	0	0	0.33	0
Dew ¹³	0.287	0.66	0.33	0.66	0.66	0.33	0	1	0.66
Torp ¹⁴	0.552	1	0.66	1	1	1	1	1	1
Cattan	-0.687	0	0	0.66	0.33	0	0	0	0
O'Connor ¹⁵	0.375	0.66	0.33	1	1	0.33	0.33	0.66	0.33

See also notes in methodology for further explanation.

⁹ Effect size based on post-test measurement and total social support at three months.

¹⁰ Effect size based on post-test measurement.

¹¹ SD estimated from Weinert 2011, equal sample sizes assumed.

¹² Effect size based on post-test measurement.

¹³ Note – effect size based on pre-post results for heart transplant recipients who received the intervention.

¹⁴ SD estimated from baseline value.

¹⁵ Mean and SD estimated from chart, error bars assumed to be based on SD (estimate of 12).

Table 5. Reduced truth table.

Configuration (1=Present; 0=Absent)	Intimacy	Interaction	Support	Nurturance	Outcome	Number of studies	Consistency	PRI (Proportional Reduction in Inconsistency)
A	1	1	1	1	1	5	0.921	0.907
B	1	1	1	0	1	2	1	1
C	1	1	0	1	0	1	0.502	0.405
D	0	0	0	0	0	3	0	0
E	1	0	0	0	0	1	0	0

identified in Table 6 provides a robust account; the inclusion of nurturance as an additional condition below is not contradictory, but suggestive that as a condition it may help to distinguish a smaller pool of studies as successful.

We also undertook quality checks to understand whether our solution, or the assumptions we made in its derivation, could also predict unsuccessful outcomes, and found little evidence that this was a possibility. We also explored whether focussing only on the 10 studies that measured social support would change our interpretation, and again found little evidence that this would influence the model. Similarly, focussing only on studies that had a comparison group showed a similar pattern descriptively^{vii}.

^{vii} However, running a model based on only eight studies with four conditions would not be appropriate.

Table 6. Solution.

	Solution Consistency	PRI (Proportional Reduction in Inconsistency)	Solution coverage	Studies
INTIMACY*INTERACTION* SUPPORT	0.936	0.921	0.829	(Barrera <i>et al.</i> , 2002, Bond <i>et al.</i> , 2010, Dew <i>et al.</i> , 2004, Gustafson <i>et al.</i> , 2005, O'Connor <i>et al.</i> , 2014, Torp <i>et al.</i> , 2008, Weinert <i>et al.</i> , 2008)

Table 7. Truth table – sensitivity analysis with alternative outcome.

Configuration (1=Present; 0=Absent)	Intimacy	Interaction	Support	Nurturance	Outcome	Number of studies	Consistency	PRI (Proportional Reduction in Inconsistency)
A	1	1	1	1	1	5	0.842	0.794
B	1	1	1	0	0	1	0.502	0.405
C	1	1	0	1	0	2	0.5	0
D	0	0	0	0	0	3	0	0
E	1	0	0	0	0	1	0	0

Table 8. Solution – sensitivity analysis with alternative outcome.

	Solution Consistency	PRI (Proportional Reduction in Inconsistency)	Solution coverage	Studies
INTIMACY*INTERACTION* NURTURANCE*SUPPORT	0.842	0.794	0.729	(Barrera <i>et al.</i> , 2002, Bond <i>et al.</i> , 2010, Gustafson <i>et al.</i> , 2005, Torp <i>et al.</i> , 2008, Weinert <i>et al.</i> , 2008)

Interpretation of the solution. The successful outcome set contained those interventions that: (i) supported participants to form ‘intimate’ relationships and express their feelings freely without self-consciousness between peers; (ii) ensured that there were shared characteristics between participants and their peers (beyond a single experience, and beyond geography alone); and (iii) included some form of pastoral care or support (e.g. light-touch oversight of a discussion forum by professionals or opportunities for participants to contact professionals for advice). This configuration explained the majority of the successful outcomes we observed.

Taken together, albeit with some caveats, these characteristics can form a set of design principles for future interventions that are delivered remotely which aim to increase support available to older adults and offset the risks of social isolation

and loneliness. The interventions that were not in the successful outcome set did not provide evidence that all three processes had been part of the interventions, and some indicated that processes to the contrary had taken place.

Summary and discussion

Summary of findings

In this rapid review of reviews, narrative synthesis showed that supported video-communication interventions are regarded positively by older adults and can have positive effects on loneliness and social support. However, the quantitative evidence remains uncertain and, although they were placed in the effective set of studies in QCA analysis, uncertainty about effectiveness is a shared conclusion in other similar reviews⁷¹. Evidence about online discussion groups and forums also demonstrated mixed results, with increases in social support,

but less evidence for improvements in loneliness. Telephone befriending has not been widely researched, but qualitative evidence suggested this intervention model may be helpful in addressing loneliness and social isolation, although quantitative evidence did not show this. The evidence for social networking sites was weak. Multi-tool interventions showed decreases in loneliness, but not always increases in social support. Clearly, these interventions vary greatly, so it is difficult to isolate the effective elements. Similarly, conceptualisations of loneliness and social isolation vary, making comparisons and conclusions challenging.

Using QCA, we looked beyond specific models of intervention to explore which intervention processes are aligned with being in an effective intervention set. We have shown that the following processes are enabled in effective interventions including (i) supporting development of intimate relationships; (ii) supporting interactions through ensuring participants share experiences/characteristics; (iii) supporting participants through pastoral guidance.

Discussion

Gaps in the evidence. Despite our extensive searches, we found only one study of telephone befriending⁴⁹ included within a single systematic review. This was also the one of the few studies that made use of volunteers. There was no information provided about the training and support provided to the volunteers, as the focus was on the experience of older adults receiving the service. Similarly, we found little information about training and support provided to staff members supporting other forms of intervention. Information, communication, support, moderation and mediation was provided to older adults by research staff and health professionals (nurses, psychologists, advance nurse practitioners) within the primary studies, but there was little detail about *how* staff (paid and volunteer) were trained or supported to provide these. The evidence identified in the QCA finds that successful interventions are effective because they are able to enhance complex psychosocial processes and abilities, highlighting that staff may need specialist training in delivering interventions successfully. In addition, support and training is likely to be important for managing the wellbeing of those delivering the intervention. Guidelines published elsewhere suggest volunteers or staff members should receive high-quality training and regular supervision to be competent⁷², yet the call for NHS Volunteer Responders to make telephone calls to isolated older adults did not include any offers of training or support⁵.

Most of the studies included in this review involved some form of new technology, with just two involving an intervention delivered through (traditional) telephone. No study examined an intervention delivered through a smartphone. Similarly, our inclusion criteria could have theoretically allowed other forms of remotely delivered interventions to be included, such as letter writing, although no such study was identified. These forms of interventions could be purposively considered in future reviews, with a recent intervention involving cross-generational letter writing suggestive of positive impacts for older and younger people alike⁷³. There may be scope in the future for inter or

cross-generational interventions that can help to provide both befriending, and technological support, while maintaining the principles outlined earlier.

We found few studies reporting on low intensity psychosocial interventions, which could be due to our focus on loneliness and/or social isolation as outcomes of interest. In the broader literature, whilst some studies have demonstrated positive impacts on depression, wellbeing and general mental health of delivering therapies through remote means⁷⁴, several of these interventions may not specifically address loneliness and are not targeted at older adults⁷⁵.

Since this overview was published, two further systematic reviews have been published in the area examining the role of digital technologies⁷⁶; both have suggested that the evidence on the effectiveness of certain forms of remotely delivered interventions is inconclusive. These inconclusive findings may reflect issues with the intervention approach and its implementation, but may also reflect the reality that evidence in this area is challenging to synthesise and characterised by heterogeneity in study design (with a preponderance of weak or flawed designs) and heterogeneity in outcomes that makes drawing conclusions challenging. For example, some of the interventions included in recent meta-analyses and in our own review were characterised as exhibiting negligible intervention impacts when considering differences in post-test outcome scores alone, although some of these studies also reported significant differences in the change between post-test and baseline, indicating that alternative analytical frameworks such as a meta-analysis of change scores should also be considered in future. In addition, the extent to which studies may impact on social isolation but not loneliness is unknown but is worth further investigation. Furthermore, given that the majority of studies in this area do not implement robust RCT designs, reviews that place stricter inclusion criteria on the study design may only capture a narrow slice of the evidence base. As reflected in the protocol, our own review set out to examine effectiveness in the anticipation of extant meta-analyses in the field, although at the time of searching (April 2020), no eligible review had undertaken meta-analyses.

Empowering and supporting older adults involved in remote interventions. Overall, the results here suggest that older adults can be empowered to support each other through online discussions and forums. In the narrative synthesis we found reviews containing several studies with peer support, provided through synchronous and asynchronous messaging, chat rooms and discussion forums. This challenges the assumption that older adults must always be on the receiving end of an intervention to address social isolation and loneliness. When we moved to study-level synthesis, we also found that studies that enabled older adults to feel that their contributions could improve the outcomes of others (i.e. improved levels of self-worth^{50–52,56,60}) tended to be successful interventions^{viii}. As the mobilisation of

^{viii} A condition reflecting self-worth was not used in the final QCA models because of the small number of studies.

thousands of volunteers takes place to support older adults who are currently shielding in the COVID-19 pandemic, recognising that older adults can be both providers and recipients of support simultaneously is likely to be an important principle to adhere to in the design of activities.

Strengths and limitations. The strengths of this rapid systematic review of reviews include the transparent and robust approach to searches, data extraction, review quality appraisal and analysis, ensured through pre-publication of a protocol on the EPPI-Centre website. Despite the rapid nature of this review process, we have conducted the review according to systematic review methodology⁷⁷. In this case, the rapid element of the review was primarily reflected in the decision to exclude reviews focussed on caregivers from the synthesis; other stages were conducted according to standard systematic reviewing practice. A further strength was the diversity of synthesis approaches conducted, including QCA and ICA.

Searching for systematic reviews means that we may have missed some more recent primary studies in this area, but it ensured that our review was achievable within the four-week timeframe required for a timely response during the COVID-19 crisis. We applied the AMSTAR2 quality appraisal tool to the included reviews, although the reviews included in the synthesis were found to have a low quality rating. In addition, we did not conduct any quality assessment of the primary studies that we looked at in more detail. Some of these had been assessed by the review authors, but many had not. There were very few identifiably robust primary studies that met our inclusion criteria. Only one primary study was identified by review authors as 'strong', with others rated as 'weak' or with no quality appraisal at all. The poor, or lack of, quality rating for many of the included studies means that findings should be considered with caution. In addition, few of the studies considered potential adverse impacts of the interventions. However, this is the case for many reviews in this research area and is not unique to our rapid review.

Owing to the rapid nature of this review, we focussed on reviews addressing interventions to mitigate loneliness or social isolation on the general older adult population. This meant we excluded reviews identified through the searches focussing exclusively on caregivers that may have provided additional insights. Other limitations included our treatment of primary studies in the QCA, where neither the precision of the effect size, study design, or quality were included in the model or the allocation into different outcome sets in our main model. Studies with weaker designs, and effect sizes derived from these, were treated in the same way as those with more robust designs in our main model. Although this is not uncommon in QCA practice, further synthesis could be conducted focussing on only those studies with a more robust design in future.

Further research and conclusions. Loneliness and social isolation are extremely complex phenomena⁸, and require a deep understanding and deliberative treatment that was beyond the remit of this rapid review. The risk of running unsuccessful interventions may be higher than many triallists appear to recognise,

and a failure to ensure that the processes identified as important in effective interventions are incorporated into intervention design may have adverse impacts for older adults, for example in heightening their feelings of alienation⁷⁸. Our findings do not lead us to recommend one particular mode of delivering befriending, social support, or low-intensity psychosocial interventions over another (e.g. videoconferencing, telephone calls, chat rooms or forums), and all may be of benefit, but our findings do suggest that the principles highlighted from the QCA should be incorporated into the delivery of an intervention. Our original intention had been to examine the effectiveness of these approaches, although due to the heterogeneity in study design and the absence of existing meta-analyses in the literature at the time of searching, we were unable to do this directly. Instead in our QCA analyses, we identified studies and qualitatively allocated these into successful and unsuccessful groups based on their effect size and explored common characteristics of successful studies; we consider this to be a prudent way of mediating a need for a rapid evidence to inform policy with the need to implement robust and transparent methods to synthesise this evidence. As discussed, this literature has been developed further since the present review was completed^{71,76,79,80}, and further progression in this area is being tracked through living maps of synthesised evidence⁸¹.

We were surprised by the identification of only one systematic review including a telephone befriending intervention. Given the UK Government's interest in encouraging volunteers to make phone calls to physically isolating and shielding older adults, under the 'stay at home' guidance, a systematic review of telephone befriending interventions is needed, to identify evidence to inform policy in this area. A review by Sharma *et al.*⁸² suggested that a large portion of such interventions may be found in grey literature. In the current context of the COVID-19 pandemic, a number of befriending interventions are being delivered by a variety of organisations, and there is scope to incorporate learning from these in future systematic reviews in this area.

As the training and support components of the technology-mediated interventions were unreported in the reviews and studies that we synthesised, there is a need to search for these elsewhere. Evaluations of existing telephone befriending and psychosocial support services, often found in the grey literature, could act as a starting point. Additional valuable information could be obtained through contact with voluntary sector and NHS organisations delivering befriending, peer support and low-intensity psychosocial interventions. A review of these training and support components could add valuable insight for policy-makers and service providers to ensure that volunteers are well trained, empowered and supported in delivering interventions adhering to the principles outlined earlier. Although we believe all of the intervention modes in scope here have the capacity to include the processes found to lead to more successful interventions (supporting the development of intimate relationships; supporting interactions through ensuring participants share experiences/characteristics; provide pastoral guidance), a more encompassing piece of research is needed in order to

identify which mode is most effective, or has the greatest potential, for changing outcomes.

Data availability

Underlying data

All underlying data as published on the Open Science Framework

Open Science Framework (OSF): Rapid systematic review of systematic reviews: what befriending, social support and low intensity psychosocial interventions, delivered remotely, are effective in reducing social isolation and loneliness among older adults? How do they work? <https://doi.org/10.17605/OSF.IO/VS2UX²¹>

This project contains the following underlying data:

- Data and example evidence for studies included in QCA and conditions included in truth tables.docx
- Data extracted on reviews and primary studies for narrative synthesis.docx
- Data used for QCA.csv

Extended data

Open Science Framework (OSF): Rapid systematic review of systematic reviews: what befriending, social support and low intensity psychosocial interventions, delivered remotely, are effective in reducing social isolation and loneliness among older

adults? How do they work? <https://doi.org/10.17605/OSF.IO/VS2UX²¹>

This project contains the following extended data:

- Search history - medline example.docx
- Further details of exclusion criteria.docx

Reporting guidelines

Open Science Framework (OSF): PRIMSA checklist for “Rapid systematic review of systematic reviews: what befriending, social support and low intensity psychosocial interventions, delivered remotely, are effective in reducing social isolation and loneliness among older adults? How do they work?”. <https://doi.org/10.17605/OSF.IO/VS2UX²¹>

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

Acknowledgements

This paper represents independent research commissioned and funded by the National Institute for Health Research Policy Research Programme. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, the Department of Health and Social Care, or its partners.

References

1. PHE: **Guidance on Social Distancing for Everyone in the UK**. London: Public Health England. 2020; accessed April 22nd 2020.
[Reference Source](#)
2. Mueller AL, McNamara MS, Sinclair DA: **Why does COVID-19 disproportionately affect older people?** *Aging (Albany NY)*. 2020; **12**(10): 9959.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. Pyle E, Evans D: **Loneliness-What characteristics and circumstances are associated with feeling lonely**. Newport: Office for National Statistics, 2018.
[Reference Source](#)
4. Steptoe A, Shankar A, Demakakos P, et al.: **Social isolation, loneliness, and all-cause mortality in older men and women**. *Proc Natl Acad Sci U S A*. 2013; **110**(15): 5797–801.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. NHS: **‘Your NHS Needs You’ – NHS call for volunteer army**. London: National Health Service. 2020; accessed May 12th 2020.
[Reference Source](#)
6. Victor C, Grenade L, Boldy D: **Measuring loneliness in later life: a comparison of differing measures**. *Rev Clin Gerontol*. 2005; **15**(1): 63–70.
[Publisher Full Text](#)
7. Age UK: **Loneliness and isolation—understanding the difference and why it matters**. London: UK. 2018. Accessed: March 14th 2021.
[Reference Source](#)
8. Hayanga B, Kneale D, Phoenix A: **Understanding the friendship networks of older Black and Minority Ethnic people living in the United Kingdom**. *Ageing Soc*. 2020.
[Publisher Full Text](#)
9. Smith KJ, Victor C: **Typologies of loneliness, living alone and social isolation, and their associations with physical and mental health**. *Ageing Soc*. 2019; **39**(8): 1709–30.
[Publisher Full Text](#)
10. Valtorta NK, Kanaan M, Gilbody S, et al.: **Loneliness, social isolation and social relationships: what are we measuring? A novel framework for classifying and comparing tools**. *BMJ Open*. 2016; **6**(4): e010799.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Cattan M, White M, Bond J, et al.: **Preventing social isolation and loneliness among older people: a systematic review of health promotion interventions**. *Ageing Soc*. 2005; **25**(1): 41–67.
[Publisher Full Text](#)
12. Victor C, Mansfield L, Kay T, et al.: **An overview of reviews: the effectiveness of interventions to address loneliness at all stages of the life-course**. London, UK: What Works Centre for Wellbeing. 2018.
[Reference Source](#)
13. Jarvis MA, Padmanabhanunni A, Balakrishna Y, et al.: **The effectiveness of interventions addressing loneliness in older persons: An umbrella review**. *Int J Nurs Sci*. 2020; **12**: 100177.
[Publisher Full Text](#)
14. Veronese N, Galvano D, D’Antiga F, et al.: **Interventions for reducing loneliness: An umbrella review of intervention studies**. *Health Soc Care Community*. 2020;
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Boulton E, Kneale D, Stansfield C, et al.: **Rapid review of reviews: what remotely delivered interventions can reduce social isolation and loneliness among older adults?** London & Manchester, UK: EPPI-Centre & NIHR Older People and Frailty PRU. 2020.
[Reference Source](#)
16. Hunt H, Pollock A, Campbell P, et al.: **An introduction to overviews of reviews: planning a relevant research question and objective for an overview**. *Syst Rev*. 2018; **7**(1): 39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Smith V, Devane D, Begley CM, et al.: **Methodology in conducting a systematic review of systematic reviews of healthcare interventions**. *BMC Med Res Methodol*. 2011; **11**(1): 15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

18. Caird J, Sutcliffe K, Kwan I, *et al.*: **Mediating policy-relevant evidence at speed: are systematic reviews of systematic reviews a useful approach?** *Evidence & Policy: A Journal of Research, Debate and Practice*. 2015; **11**(1): 81–97. [Publisher Full Text](#)
19. Chipps J, Jarvis MA, Ramlall S: **The effectiveness of e-Interventions on reducing social isolation in older persons: A systematic review of systematic reviews.** *J Telemed Telecare*. 2017; **23**(10): 817–27. [PubMed Abstract](#) | [Publisher Full Text](#)
20. Moher D, Liberati A, Tetzlaff J, *et al.*: **Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement.** *Ann Intern Med*. 2009; **151**(4): 264–69, W64. [PubMed Abstract](#) | [Publisher Full Text](#)
21. Kneale D, Boulton E: **Rapid systematic review of systematic reviews: what befriending, social support and low intensity psychosocial interventions, delivered remotely, are effective in reducing social isolation and loneliness among older adults? How do they work?** 2020. <http://www.doi.org/10.17605/OSF.IO/VS2UX>
22. Boulton E, Kneale D, Stansfield C, *et al.*: **What are effective befriending, social support, and low intensity psychosocial interventions delivered remotely to reduce social isolation and loneliness among older adults and how do they work? A rapid review of systematic reviews (protocol).** London: EPPI-Centre. 2020. [Reference Source](#)
23. Hayanga B: **The effectiveness and suitability of interventions for social isolation and loneliness for older people from minoritised ethnic groups living in the UK.** London: University College London. 2021.
24. Cornwell EY, Waite LJ: **Measuring social isolation among older adults using multiple indicators that include randomised or non-randomised studies of healthcare interventions, or both.** *BMJ*. 2017; **358**: j4008. 2009; **64** Suppl 1(Suppl 1): i38–i46. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. CRD: **Database of Abstracts of Reviews of Effects (DARE): Quality-assessed Reviews.** Centre for Reviews and Dissemination, University of York. 2014; accessed April 20th 2020. [Reference Source](#)
26. Thomas J, Graziosi S, Brunton J, *et al.*: **EPPI-Reviewer: advanced software for systematic reviews, maps and evidence synthesis.** London: UCL Social Research Institute, University College London 2020.
27. Shea BJ, Reeves BC, Wells G, *et al.*: **AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both.** *BMJ*. 2017; **358**: j4008. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Snijlsteit B, Oliver S, Vojtkova M: **Narrative approaches to systematic review and synthesis of evidence for international development policy and practice.** *J Dev Effect*. 2012; **4**(3): 409–29. [Publisher Full Text](#)
29. Sutcliffe K, Thomas J, Stokes G, *et al.*: **Intervention Component Analysis (ICA): a pragmatic approach for identifying the critical features of complex interventions.** *Syst Rev*. 2015; **4**(1): 140. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Schneider CQ, Wagemann C: **Set-theoretic methods for the social sciences: A guide to qualitative comparative analysis.** Cambridge University Press. 2012. [Reference Source](#)
31. Ragin CC: **Redesigning social inquiry: Fuzzy sets and beyond.** Wiley Online Library. 2008. [Reference Source](#)
32. Thomas J, O'Mara-Eves A, Brunton G: **Using qualitative comparative analysis (QCA) in systematic reviews of complex interventions: a worked example.** *Syst Rev*. 2014; **3**(1): 67. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Kneale D, Sutcliffe K, Thomas J: **Critical Appraisal of Reviews Using Qualitative Comparative Analyses (CARU-QCA): a tool to critically appraise systematic reviews that use qualitative comparative analysis.** Cochrane Colloquium. Santiago, Chile. 2019. [Reference Source](#)
34. Weiss RS: **The fund of sociability.** *Trans-action*. 1969; **6**(9): 36–43. [Reference Source](#)
35. Bennett N: **The Impact of Video-Communication on Older Adults' Psychological Well-Being: A Mixed Methods Study.** University of Essex. 2015. [Reference Source](#)
36. Chen YRR, Schulz PJ: **The Effect of Information Communication Technology Interventions on Reducing Social Isolation in the Elderly: A Systematic Review.** *J Med Internet Res*. 2016; **18**(1): e18. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Morris ME, Adair B, Ozanne E, *et al.*: **Smart technologies to enhance social connectedness in older people who live at home.** *Australas J Ageing*. 2014; **33**(3): 142–52. [PubMed Abstract](#) | [Publisher Full Text](#)
38. Khosravi P, Rezvani A, Wiewiara A: **The impact of technology on older adults' social isolation.** *Comput Human Behav*. 2016; **63**: 594–603. [Publisher Full Text](#)
39. Benito-Montagut R, Cassián-Yde N, Begueria A: **What do we know about the relationship between internet-mediated interaction and social isolation and loneliness in later life?** *Qual Ageing Older Adults*. 2018; **19**(1): 14–30. [Publisher Full Text](#)
40. Corry M, Neenan K, Brabyn S, *et al.*: **Telephone interventions, delivered by healthcare professionals, for providing education and psychosocial support for informal caregivers of adults with diagnosed illnesses.** *Cochrane Database Syst Rev*. 2019; **5**(5): CD012533. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Elvish R, Lever SJ, Johnstone J, *et al.*: **Psychological interventions for carers of people with dementia: A systematic review of quantitative and qualitative evidence.** *Couns Psychother Res*. 2013; **13**(2): 106–25. [Publisher Full Text](#)
42. Hopwood J, Walker N, McDonagh L, *et al.*: **Internet-Based Interventions Aimed at Supporting Family Caregivers of People With Dementia: Systematic Review.** *J Med Internet Res*. 2018; **20**(6): e216. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. Lins S, Hayder-Beichel D, Rücker G, *et al.*: **Efficacy and experiences of telephone counselling for informal carers of people with dementia.** *Cochrane Database Syst Rev*. 2014; **2014**(9): CD009126. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
44. Savolainen L, Hanson E, Magnusson L, *et al.*: **An Internet-based videoconferencing system for supporting frail elderly people and their carers.** *J Telemed Telecare*. 2008; **14**(2): 79–82. [PubMed Abstract](#) | [Publisher Full Text](#)
45. Tsai HH, Tsai YF: **Older nursing home residents' experiences with videoconferencing to communicate with family members.** *J Clin Nurs*. 2010; **19**(11–12): 1538–43. [PubMed Abstract](#) | [Publisher Full Text](#)
46. Schwindenhammer TM: **Videoconferencing intervention for depressive symptoms and loneliness in nursing home elders.** Illinois State University, 2014.
47. Tsai HH, Tsai YF: **Changes in depressive symptoms, social support, and loneliness over 1 year after a minimum 3-month videoconferencing program for older nursing home residents.** *J Med Internet Res*. 2011; **13**(4): e93. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. Tsai HH, Tsai YF, Wang HH, *et al.*: **Videoconference program enhances social support, loneliness, and depressive status of elderly nursing home residents.** *Ageing Ment Health*. 2010; **14**(8): 947–54. [PubMed Abstract](#) | [Publisher Full Text](#)
49. Cattan M, Kime N, Bagnall AM: **The use of telephone befriending in low level support for socially isolated older people— an evaluation.** *Health Soc Care Community*. 2011; **19**(2): 198–206. [PubMed Abstract](#) | [Publisher Full Text](#)
50. Gustafson DH, McTavish FM, Stengle W, *et al.*: **Use and impact of eHealth system by low-income women with breast cancer.** *J Health Commun*. 2005; **10**(Suppl 1): 195–218. [PubMed Abstract](#) | [Publisher Full Text](#)
51. Barrera M Jr, Glasgow RE, McKay HG, *et al.*: **Do Internet-based support interventions change perceptions of social support?: An experimental trial of approaches for supporting diabetes self-management.** *Am J Community Psychol*. 2002; **30**(5): 637–54. [PubMed Abstract](#) | [Publisher Full Text](#)
52. Bond GE, Burr RL, Wolf FM, *et al.*: **The effects of a web-based intervention on psychosocial well-being among adults aged 60 and older with diabetes.** *Diabetes Educ*. 2010; **36**(3): 446–56. [PubMed Abstract](#) | [Publisher Full Text](#)
53. Dew MA, Goycoolea JM, Harris RC, *et al.*: **An internet-based intervention to improve psychosocial outcomes in heart transplant recipients and family caregivers: development and evaluation.** *J Heart Lung Transplant*. 2004; **23**(6): 745–58. [PubMed Abstract](#) | [Publisher Full Text](#)
54. Hill W, Weinert C, Cudney S: **Influence of a computer intervention on the psychological status of chronically ill rural women: preliminary results.** *Nurs Res*. 2006; **55**(1): 34–42. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
55. O'Connor MF, Arizmendi BJ, Kaszniak AW: **Virtually supportive: a feasibility pilot study of an online support group for dementia caregivers in a 3D virtual environment.** *J Aging Stud*. 2014; **30**: 87–93. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. Torp S, Hanson E, Hauge S, *et al.*: **A pilot study of how information and communication technology may contribute to health promotion among elderly spousal carers in Norway.** *Health Soc Care Community*. 2008; **16**(1): 75–85. [PubMed Abstract](#) | [Publisher Full Text](#)
57. Weiner C, Cudney S, Comstock B, *et al.*: **Computer intervention impact on psychosocial adaptation of rural women with chronic conditions.** *Nurs Res*. 2011; **60**(2): 82–91. [PubMed Abstract](#) | [Publisher Full Text](#)
58. Ballantyne A, Trenwith L, Zubrinich S, *et al.*: **I feel less lonely: what older people say about participating in a social networking website.** *Qual Ageing Older Adults*. 2010; **11**(3): 25. [Publisher Full Text](#)
59. van der Heide LA, Willems CG, Spreeuwenberg MD, *et al.*: **Implementation**

- of CareTV in care for the elderly: the effects on feelings of loneliness and safety and future challenges. *Technol Disabil.* 2012; **24**(4): 283–91.
[Publisher Full Text](#)
60. Weinert C, Cudney S, Hill WG: **Rural women, technology, and self-management of chronic illness.** *Can J Nurs Res.* 2008; **40**(3): 114–34.
[PubMed Abstract](#) | [Free Full Text](#)
 61. Demiris G, Oliver DRP, Hensel B, *et al.*: **Use of videophones for distant caregiving: an enriching experience for families and residents in long-term care.** *J Gerontol Nurs.* 2008; **34**(7): 50–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
 62. Cattan M, Kime N, Bagnall A: **Low-level support for socially isolated older people: An evaluation of telephone befriending.** London: Help the Aged, 2008.
[Reference Source](#)
 63. Berg-Schlösser D, De Meur G, Rihoux B, *et al.*: **Qualitative Comparative Analysis (QCA) as an Approach.** In: Rihoux B, Ragin CC: *Configurational comparative methods Qualitative comparative analysis (QCA) and related techniques.* Thousand Oaks, CA, USA: Sage Publishing 2009: 19–32.
[Reference Source](#)
 64. Harris K, Kneale D, Lasserson TJ, *et al.*: **School-based self-management interventions for asthma in children and adolescents: a mixed methods systematic review.** *Cochrane Database Syst Rev.* 2019; **1**(1): CD011651.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 65. Candy B, King M, Jones L, *et al.*: **Using qualitative evidence on patients' views to help understand variation in effectiveness of complex interventions: a qualitative comparative analysis.** *Trials.* 2013; **14**(1): 179.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 66. Kahwati LC, Lewis MA, Kane H, *et al.*: **Best practices in the Veterans Health Administration's MOVE! Weight management program.** *Am J Prev Med.* 2011; **41**(5): 457–64.
[PubMed Abstract](#) | [Publisher Full Text](#)
 67. Thomas J, O'Mara-Eves A, Kneale D, *et al.*: **Synthesis Methods for Combining and Configuring Quantitative Data.** In: Gough D, Oliver S, Thomas J, eds. *An Introduction to Systematic Reviews.* London: Sage 2017.
[Reference Source](#)
 68. York RO: **Statistics for human service evaluation.** SAGE Publications 2016.
[Reference Source](#)
 69. Masi CM, Chen HY, Hawkey LC, *et al.*: **A meta-analysis of interventions to reduce loneliness.** *Pers Soc Psychol Rev.* 2011; **15**(3): 219–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 70. Cohen J: **Statistical power analysis for the behavioral sciences.** Academic press 2013.
[Reference Source](#)
 71. Noone C, McSharry J, Smalle M, *et al.*: **Video calls for reducing social isolation and loneliness in older people: a rapid review.** *Cochrane Database Syst Rev.* 2020; **5**(5): CD013632.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 72. NICE: **Depression in adults: recognition and management (Clinical Guideline 90).** National Institute for Health & Care Excellence: London: NICE, 2009.
[Reference Source](#)
 73. Binnie G: **Loneliness and the letter: Co-developing cross-generational letter writing with higher education students and older people.** *Research for All.* 2019; **3**(1): 59–73.
[Publisher Full Text](#)
 74. Kaonga NN, Morgan J: **Common themes and emerging trends for the use of technology to support mental health and psychosocial well-being in limited resource settings: A review of the literature.** *Psychiatry Res.* 2019; **281**: 112594.
[PubMed Abstract](#) | [Publisher Full Text](#)
 75. Burholt V, Windle G, Gott M, *et al.*: **Technology-Mediated Communication in Familial Relationships: Moderated-Mediation Models of Isolation and Loneliness.** *Gerontologist.* 2020; **60**(7): 1202–1212.
[PubMed Abstract](#) | [Publisher Full Text](#)
 76. Shah SGS, Nogueras D, van Woerden HC, *et al.*: **Are digital technology interventions effective to reduce loneliness in older adults? A systematic review and meta-analysis.** *medRxiv.* 2020.
[Publisher Full Text](#)
 77. Gough D, Oliver S, Thomas J: **An Introduction to Systematic Reviews.** London: Sage, 2017.
[Reference Source](#)
 78. Wong A, Chau AK, Fang Y, *et al.*: **Illuminating the psychological experience of elderly loneliness from a societal perspective: A qualitative study of alienation between older people and society.** *Int J Environ Res Public Health.* 2017; **14**(7): 824.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 79. Gorenko JA, Moran C, Flynn M, *et al.*: **Social Isolation and Psychological Distress Among Older Adults Related to COVID-19: A Narrative Review of Remotely-Delivered Interventions and Recommendations.** *J Appl Gerontol.* 2021; **40**(1): 3–13.
[PubMed Abstract](#) | [Publisher Full Text](#)
 80. Dube L, Paquet C, Cisneros-Franco M, *et al.*: **Knowledge Synthesis for Mechanistic and Targeted In-Person and Digital Social-Connection Intervention for Wellness and Resilience in Older Adults in Pandemic Context and Beyond.** McGill University. 2021.
[Reference Source](#)
 81. IPPO: **Global Evidence Networks.** 2021.
[Reference Source](#)
 82. Sharma D, Blair L, Clune SJ: **Reducing Later Life Loneliness: A Systematic Literature Review of Loneliness Interventions.** *ICHAA 2018.* 2018.
[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status: ? ?

Version 2

Reviewer Report 02 July 2021

<https://doi.org/10.5256/f1000research.55641.r84065>

© 2021 Shah S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

? **Syed Ghulam Sarwar Shah**

¹ NIHR Oxford Biomedical Research Centre, Oxford University Hospitals NHS Foundation Trust, Oxford, UK

² Radcliffe Department of Medicine, University of Oxford, Level 6, West Wing, John Radcliffe Hospital, Oxford, UK

Thanks to the authors for considering reviewers' comments and submitting a revision version of the manuscript, which is greatly improved. However, I have a few minor comments as follows:

1. The title is too long and it does not read smoothly. The authors may consider revising it so it reads better and is exciting.
2. Definition of older adults: It is given in the introduction section (para 1) as well as in the Methods section. This duplication can be removed by removing it from the methods section because it is again given in the population of interest in the inclusion criteria.
3. The definitions of 'loneliness and social isolation' are also repeated. These terms are defined in the introduction section (para 3) and again in the 'outcomes' in the methods section. Duplication of this information may be avoided.
4. Update of a reference: Ref # 76 Shah *et al.* published as a preprint in medRxiv has now been accepted by the Journal of Medical Internet Research; Hence, this reference may be updated as follows: Shah SGS, Nogueras D, van Woerden HC, Kiparoglou V. Evaluation of the Effectiveness of Digital Technology Interventions to Reduce Loneliness in Older Adults: A Systematic Review and Meta-Analysis. Journal of Medical Internet Research. 19/04/2021:24712 (forthcoming/in press)
<https://preprints.jmir.org/preprint/24712/accepted> (and Shah *et al.*, 2021¹).

References

1. Shah SGS, Nogueras D, van Woerden HC, Kiparoglou V: Evaluation of the Effectiveness of Digital Technology Interventions to Reduce Loneliness in Older Adults: Systematic Review and Meta-

analysis. *J Med Internet Res.* 2021; **23** (6): e24712 [PubMed Abstract](#) | [Publisher Full Text](#)

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Public Health, Loneliness, Digital technology for health, Health inequalities, Systematic reviews and meta-analysis, Quantitative research, Patients' access to electronic health/medical records.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Version 1

Reviewer Report 25 January 2021

<https://doi.org/10.5256/f1000research.29908.r75713>

© 2021 Chipps J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Jennifer Anne Chipps

School of Nursing, University of the Western Cape, Cape Town, South Africa

The authors did use the PRIMSA reporting guidelines. I do have a few concerns:

1. There are already a few umbrella reviews - these are not commented on and as to what his paper adds.
2. Which methodology for umbrella reviews were used?
3. More detail needs to be provided on the criteria of the primary papers extracted from the systematic reviews. E.g. What was the inclusion and exclusion criteria, and how was the quality of the primary papers assessed?
4. In methodology the 'rapid' process needs to be explained – e.g. what time period.
5. Did the study include all residents in community and residential homes?
6. To address the question of effectiveness - only RCTS or QE studies should be included – there are qualitative papers in the synthesis which makes this confusing.
7. Social isolation and loneliness are separate concepts – need to be defined – this also related to the broad heterogeneous measures included.

8. What are diverse population of older persons?

The ICA and QCA is interesting and a new addition to these umbrella reviews

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Yes

Are sufficient details of the methods and analysis provided to allow replication by others?

Partly

Is the statistical analysis and its interpretation appropriate?

Yes

Are the conclusions drawn adequately supported by the results presented in the review?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Mental Health and systematic reviews

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 22 Mar 2021

Dylan Kneale, University College London, London, UK

Response to Reviewers:

We thank the Reviewer for her constructive comments, which has given us the opportunity to improve the paper. We have addressed the comments and have revised the manuscript accordingly. Our point-by-point response is included below and we hope that we have satisfied the concerns of the reviewer.

Response to Professor Jennifer Chipps:

1. There are already a few umbrella reviews - these are not commented on and as to what this paper adds.

Thank you for this comment, enabling us to highlight what our paper adds. We have stated in the Introduction that a 'number of evidence reviews have highlighted the diverse range of interventions to alleviate loneliness amongst older adults in a variety of settings'. We included references for Cattan et al., 2005 and Victor et al., 2018 here. We have now added Jarvis et al., 2020, which had not been identified in our searches in April 2020, and Veronese et al., 2020, which had not been published until after this research was conducted. In addition, we noticed an error in the referencing of one umbrella review (Chipps et al) which

was introduced at the wrong place in the document. At the end of the introduction, we have made it clearer that our paper adds the explicit consideration of particular interventions (befriending, social support and low intensity psychosocial interventions) to address loneliness and social isolation, together with the innovative use of Intervention Component Analysis and Qualitative Comparative Analysis.

'This rapid review examines evidence specifically on whether befriending, social support, and low intensity psychosocial interventions delivered remotely can reduce social isolation or loneliness among older adults.'

'(iii) Generate new understandings on how interventions work and which core components and processes are associated with successful interventions, using the innovative methods of Intervention Component Analysis and Qualitative Comparative Analysis.'

2. Which methodology for umbrella reviews was used?

We have added the following text to the first paragraph of the Methods section:

Although broad frameworks for conducting overviews exist (Smith, Devane et al. 2011), specific guidance that can be used in a directive way to carry out overviews is lacking. Nevertheless, we followed elements of practice recommended by Caird and colleagues (2015) in balancing some of the challenges of conducting reviews of reviews with the need to produce policy-relevant evidence at speed in the context of the COVID-19 pandemic; the present review was developed over a four week period in April-May 2020. We also examined how existing reviews of reviews in the area, and particularly a review conducted by Chipps and colleagues (2017), navigated the challenge around differences between the scope of a systematic review and the scope of the review of the reviews.

3. More detail needs to be provided on the criteria of the primary papers extracted from the systematic reviews. E.g. What was the inclusion and exclusion criteria, and how was the quality of the primary papers assessed?

Thank you for pointing out that this should be included. We have added the following text to the 'Study selection and data extraction' and 'Critical appraisal' sections:

'Once eligible reviews had been identified, primary papers were extracted if they met the criteria in line with the review inclusion and exclusion criteria stated above. That is:

Population: older adults, located in the community or residential care, socially isolated or lonely, or at risk of social isolation or loneliness.

Intervention: interventions that sought to reduce levels of social isolation or loneliness, through strengthening individuals' social contacts and social relationships. Interventions were delivered on a one-to-one basis (e.g. befriending), or as remote group-based interventions (e.g. discussion groups).

Comparator/control: studies with most forms of control group (randomised and non-randomised) and those without a control group (pre-post designs).

Outcomes: measures of social isolation or loneliness, including bespoke and proxy measures, such as social connectedness.'

'The quality of the primary studies was reported where it had been assessed by the review authors. Not all reviews included a quality assessment of their included studies.'

4. In methodology the 'rapid' process needs to be explained – e.g. what time period.

We have included the time period in the first paragraph of the Methods section: 'the present review was developed over a four week period in April-May 2020.'

5. Did the study include all residents in community and residential homes?

We have added the following text into the Search strategy section:

'1) Population: older and middle-aged populations aged 50+ years in the community and in residential settings.

6. To address the question of effectiveness - only RCTS or QE studies should be included – there are qualitative papers in the synthesis which makes this confusing.

Thank you for highlighting this issue, which we have addressed through:

- editing the title of the manuscript to better reflect our research;

The original title of the manuscript reflected the Protocol for the review which was to look at if, and how, interventions worked. However, at the time of searching (April 2020), no eligible review had undertaken meta-analyses, and we have clarified this and modified the title to reflect the content of the manuscript.

- adding a note on the study design in the Methods section:

'As we were expecting some heterogeneity in the question being addressed by reviews, and expected this to be reflected in the design of primary studies included within reviews, we did not specify that source systematic reviews had to be confined to a particular study design. In line with previous reviews in the field, we expected studies measuring quantitative outcomes to be composed of single-group pre-post studies, non-randomised comparison studies, and randomised comparison group studies.'

7. Social isolation and loneliness are separate concepts – need to be defined – this also related to the broad heterogenous measures included.

We have defined social isolation and loneliness, as separate concepts, in the introduction and have added text to highlight this in the outcomes section of the 'Inclusion and exclusion criteria' section:

We conceptualise loneliness as an emotional response by individuals when there is a 'deficit between their desired and actual quality and quantity of social engagement and relationships ^{5, p64}. Social isolation reflects the number of social contacts that people have (UK 2018), and people who are socially isolated tend to have social networks of low density

that are not maintained through frequent engagement⁶. Both loneliness and social isolation are conceptually distinct from living alone, the latter having limited utility as a proxy for either social isolation or loneliness⁷. However, we recognise that defining social isolation and loneliness is challenging, particular as researchers have used terms involving social relationships, including social isolation, loosely (Valtorta, Kanaan et al. 2016). Furthermore, while we recognise social isolation and loneliness as distinct concepts, here we explore both simultaneously as the COVID-19 pandemic and measures adopted to mitigate its spread have exacerbated both isolation and loneliness.

8. What are diverse population of older persons?

Thank you for highlighting this lack of clarity. We have removed the word 'diverse' from the 'Inclusion and exclusion criteria' section, as it was unnecessary.

Competing Interests: We have no competing interests to declare

Reviewer Report 25 January 2021

<https://doi.org/10.5256/f1000research.29908.r76527>

© 2021 Shah S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Syed Ghulam Sarwar Shah

¹ NIHR Oxford Biomedical Research Centre, Oxford University Hospitals NHS Foundation Trust, Oxford, UK

² Radcliffe Department of Medicine, University of Oxford, Level 6, West Wing, John Radcliffe Hospital, Oxford, UK

Many thanks for inviting me to review the manuscript entitled "Rapid systematic review of systematic reviews: what befriending, social support and low intensity psychosocial interventions, delivered remotely, are effective in reducing social isolation and loneliness among older adults? How do they work?"

Loneliness is a major public health issue that has become critical now due to increased social isolation and lockdowns in the current COVID-19 pandemic.

Therefore, this manuscript reporting a rapid review of reviews on remote interventions and their effectiveness in tackling loneliness in older people is timely.

I have the following comments on the manuscript.

Abstract:

In systematic reviews, reporting of the sources of literature, selection criteria and publication

period are important but these are missing in the abstract. The authors need to add this information in the methods section of the abstract. If however there were no strict criteria such as restrictions on publication dates, then these need to be mentioned. The population of interest was 'older adults' and their age is reported as 50 years and above. This means no upper age limit therefore some participants might be very old such as above 70 years of age and they could be called as 'elderly'. The two terms i.e. older and elderly, are interpreted differently; therefore, the authors might like to change the term 'older adults' to 'adults aged 50 years and above' to avoid the confusion. The Results section should also provide a summary of data extracted about characteristics of studies, population and interventions. Determination of the effectiveness of the interventions was one of the objectives and authors have reported a mixed quantitative evidence of effectiveness in the results section but nothing about the effectiveness is reported in the conclusion. The authors should include concluding remarks on the effectiveness in the conclusion in the abstract.

Introduction:

In the introduction section, the authors mention older adults over 70 years while in the abstract they report 50 years plus. The authors need to be consistent in reporting the term older people and years of age covered. The authors might like to describe the term 'older adults' including age in years included in the review in the methods section.

The authors have mentioned that 'During the height of the COVID-19 pandemic these interventions were of limited...'. This suggests that the peak intensity of the COVID-19 pandemic is over, which it is not because there are second and third waves of COVID in different countries. For example, the UK is currently going through a very serious second wave of COVID-19 and strict lockdown is in place across the country. Therefore, the authors need to change the above statement such as 'during the height of the first wave of COVID-19 in 2020...'.

Methods:

Different researchers were involved in the screening and shortlisting of articles but no information on the interrater reliability/agreement is reported. Reporting of a relevant statistic such as the Kappa static will be helpful.

Results:

The results section includes some paragraphs such as the '*Theory selection and setting up the QCA*', '*Developing a data table*' and '*Truth table*'. These paragraphs describe the theory and methodological procedures; hence, these should be reported in the methods section.

Discussion:

The authors might like to refer to some recent meta-analyses on the effectiveness of interventions to tackle loneliness, which show that digital interventions are not effective in adults aged 70 years and above (Shah *et al.*, 2020¹) and especially video calls (Noone *et al.*, 2020²).

Conclusion:

The conclusion section is more about the inclusion of the processes and principles suggested in the QCA. The authors need to include their overall inference about the effectiveness of different types interventions reviewed in the population of interest.

References

1. Shah S, Nogueras D, van Woerden H, Kiparoglou V: Are digital technology interventions effective to reduce loneliness in older adults? A systematic review and meta-analysis. *medRxiv*. 2020.

[Publisher Full Text](#)

2. Noone C, McSharry J, Smalle M, Burns A, et al.: Video calls for reducing social isolation and loneliness in older people: a rapid review. *Cochrane Database of Systematic Reviews*. 2020. [Publisher Full Text](#)

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Yes

Are sufficient details of the methods and analysis provided to allow replication by others?

Partly

Is the statistical analysis and its interpretation appropriate?

Partly

Are the conclusions drawn adequately supported by the results presented in the review?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Public Health, Loneliness, Digital technology for health, Health inequalities, Systematic reviews and meta-analysis, Quantitative research, Patients' access to electronic health/medical records.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 22 Mar 2021

Dylan Kneale, University College London, London, UK

Response to Reviewers.

We thank the Reviewer for his constructive comments, which has given us the opportunity to improve the paper. We have addressed the comments and have revised the manuscript accordingly. Our point-by-point response is included below and we hope that we have satisfied the concerns of the reviewer.

Response to Dr Shah:

○ **Abstract:**

In systematic reviews, reporting of the sources of literature, selection criteria and publication period are important but these are missing in the abstract. The authors need to add this information in the methods section of the abstract. If however there were no strict criteria such as restrictions on publication dates,

then these need to be mentioned. The population of interest was 'older adults' and their age is reported as 50 years and above. This means no upper age limit therefore some participants might be very old such as above 70 years of age and they could be called as 'elderly'. The two terms i.e. older and elderly, are interpreted differently; therefore, the authors might like to change the term 'older adults' to 'adults aged 50 years and above' to avoid the confusion. The Results section should also provide a summary of data extracted about characteristics of studies, population and interventions. Determination of the effectiveness of the interventions was one of the objectives and authors have reported a mixed quantitative evidence of effectiveness in the results section but nothing about the effectiveness is reported in the conclusion. The authors should include concluding remarks on the effectiveness in the conclusion in the abstract.

- **Introduction:**

In the introduction section, the authors mention older adults over 70 years while in the abstract they report 50 years plus. The authors need to be consistent in reporting the term older people and years of age covered. The authors might like to describe the term 'older adults' including age in years included in the review in the methods section.

The authors have mentioned that 'During the height of the COVID-19 pandemic these interventions were of limited...'. This suggests that the peak intensity of the COVID-19 pandemic is over, which it is not because there are second and third waves of COVID in different countries. For example, the UK is currently going through a very serious second wave of COVID-19 and strict lockdown is in place across the country. Therefore, the authors need to change the above statement such as 'during the height of the first wave of COVID-19 in 2020...'.

Thank you for your comments and helpful suggested revisions to the abstract. We have included some further detail about the methods employed, within the confines of the word limit. We have added the fields which the 11 databases covered ('from the fields of health, social care, psychology and social science). We have also added the selection was guided by our PICOS criteria.

Thank you for pointing out the inconsistencies with regard to the age groups covered in the manuscript. The review included research studies with populations aged 50 years and above, whereas the UK Government guidance aimed at protecting older adults during the first wave of the pandemic was targeted at those aged 70 years and above. Thank you for helping us to clarify the time period with your wording about the first wave. We have changed the wording of the introduction to:

'During the height of the first wave of the COVID-19 pandemic in 2020, millions of people aged 70 years and over were advised to avoid social contact with those outside their household. with older age identified as a risk factor for poorer COVID-19 prognosis (Mueller, McNamara et al. 2020).'

We have also changed the wording in the abstract, although the word limit has restricted what we could add here:

‘During the COVID-19 pandemic ‘social distancing’ has highlighted the need to minimise loneliness and isolation among older adults (aged 50+).’

With regard to the use of the term ‘older adults’, we feel that this term does encompass the full range of ages from 50 years and upwards. The term is commonplace in our research groups and we would prefer to keep it as it is, as many people in their 50s and 60s experience poor health and life transitions at an earlier age, meaning that they live with age-related conditions from 50+. As recommended, we have ensured that the term ‘older adults’ is defined in the introduction and at the beginning of our methods section, so that the reader is clear about the population in our paper, and is used consistently throughout. As such, we have changed all instances of ‘older people’ to older adults’.

Introduction: ‘Here we use a broad definition of ‘older’ adult, defined as those aged 50+, which captures those in middle age who may be nearing or experiencing age-related transitions, such as retirement or unpaid caring, or living with age-related long term conditions.’

Methods: ‘For the purposes of this review, we define ‘older adults’ as those aged 50 years and above.’

- **Methods:**
Different researchers were involved in the screening and shortlisting of articles but no information on the interrater reliability/agreement is reported. Reporting of a relevant statistic such as the Kappa static will be helpful.

We have now reported the level of agreement (93%) in the manuscript.

- **Results:**
The results section includes some paragraphs such as the ‘Theory selection and setting up the QCA’, ‘Developing a data table’ and ‘Truth table’. These paragraphs describe the theory and methodological procedures; hence, these should be reported in the methods section.

The theory has now been moved to the methods section. However, as the data table and truth table sections also include results, and provide a description of how to interpret the data presented, we think that these sections should remain in the results. As QCA is a relatively new method applied to systematic reviews, we believe that having this explanation alongside the results will provide clarity for the reader.

- **Discussion:**
The authors might like to refer to some recent meta-analyses on the effectiveness of interventions to tackle loneliness, which show that digital interventions are not effective in adults aged 70 years and above (Shah *et al.*, 2020¹) and especially video calls (Noone *et al.*, 2020²).

Thank you for this suggestion, which has enabled us to improve the Discussion section. We have added the following paragraph at the end of the 'Gaps in the evidence' section:

'Since this overview was published, two further systematic reviews have been published in the area examining the role of digital technologies (Noone, McSharry et al. 2020, Shah, Nogueras et al. 2020); both have suggested that the evidence on the effectiveness of certain forms of remotely delivered interventions is inconclusive. These inconclusive findings may reflect issues with the intervention approach and its implementation, but may also reflect the reality that evidence in this area is challenging to synthesise and characterised by heterogeneity in study design (with a preponderance of weak or flawed designs) and heterogeneity in outcomes that makes drawing conclusions challenging. For example, some of the interventions included in recent meta-analyses and in our own review were characterised as exhibiting negligible intervention impacts when considering post-test outcome scores alone, although significant changes between post-test and baseline scores suggest that a meta-analysis of change scores should also be considered in future. Furthermore, given that the majority of studies in this area do not implement robust RCT designs, reviews that place stricter inclusion criteria on the study design may only capture a narrow slice of the evidence base. As reflected in the protocol, our own review set out to examine effectiveness in the anticipation of extant meta-analyses in the field, although at the time of searching (April 2020), no eligible review had undertaken meta-analyses.'

- **Conclusion:**

The conclusion section is more about the inclusion of the processes and principles suggested in the QCA. The authors need to include their overall inference about the effectiveness of different types interventions reviewed in the population of interest.

Thank you for highlighting this. We have added the following text to the end of the first paragraph of the 'Further research and conclusions' section:

'Our original intention had been to examine the effectiveness of these approaches, although due to the heterogeneity in study design and the absence of existing meta-analyses in the literature at the time of searching, we were unable to do this directly. Instead in our QCA analyses, we identified studies and qualitatively allocated these into successful and unsuccessful groups based on their effect size and explored common characteristics of successful studies; we consider this to be a prudent way of mediating a need for a rapid evidence to inform policy with the need to implement robust and transparent methods to synthesise this evidence. As discussed, this literature has been developed further since the present review was completed (Gorenko, Moran et al. 2020, Noone, McSharry et al. 2020, Shah, Nogueras et al. 2020, Dubé, Paquet et al. 2021), and further progression in this area is being tracked through living maps of synthesised evidence (IPPO 2021).'

Competing Interests: We have no competing interests to declare

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research