REVIEW

Review of the nutritional composition, medicinal, phytochemical and pharmacological properties of *Citrus reticulata* Blanco (Rutaceae) [version 1; peer review: 1 approved, 1 not approved]

Collen Musara¹, Elizabeth Bosede Aladejana¹, Silas Mufambi Mudyiwa²

¹Medicinal Plants and Economic Development (MPED) Research Centre, Department of Botany, University of Fort Hare, Alice, Eastern Cape, 5700, South Africa

²Department of Livestock and Pasture Science, University of Fort Hare, Alice, Eastern Cape, 5700, South Africa

V1 First published: 02 Dec 2020, 9:1387
<https://doi.org/10.12688/f1000research.27208.1>
 Latest published: 02 Dec 2020, 9:1387
<https://doi.org/10.12688/f1000research.27208.1>

Abstract

Citrus reticulata Blanco is a moderately-sized fruit tree widely used as herbal medicine worldwide. The nutritional composition, medicinal uses, phytochemistry and pharmacological properties of *C. reticulata* were critically reviewed in the current study. The literature linked to *C. reticulata* properties was obtained from multiple internet sources including Elsevier, Google Scholar, SciFinder, Web of Science, Pubmed, BMC, Science Direct, and Scopus. Ethnopharmacological research identified antioxidants such as vitamin C, carotenoids and phenolic compounds, also a source of sugars, organic acids, amino acids, pectins, minerals and volatile organic compounds as components of *C. reticulata*. As a medicinal plant, *C. reticulata* is used for the treatment of dyspepsia, gastro-intestinal distension, cough with profuse phlegm, hiccup and vomiting. The crude extracts of *C. reticulata* fruits have depicted anti-inflammatory, anticholesterolemic, analgesic, antiasthmatic, antiscorbutic, antiseptic, antitussive, carminative, expectorant, stomachic. With more people becoming nutrition-conscious, there has been an increase in the demand for the use of citrus fruits and their by-products as traditional medicines for conventional healthcare in developing countries.

Keywords

Citrus reticulata, medicinal, nutrition, pharmacological, phytochemistry

Open Peer Review

Approval Status

	1	2
version 1
02 Dec 2020	view	view

1. **Uma Rani Sinniah** , Universiti Putra Malaysia, Serdang, Malaysia

2. **Maan B. Rokaya** , The Czech Academy of Sciences, Průhonice, Czech Republic

Any reports and responses or comments on the article can be found at the end of the article.

This article is included in the **Agriculture, Food and Nutrition** gateway.

Corresponding author: Elizabeth Bosede Aladejana (ecaladejana@ufh.ac.za)

Author roles: **Musara C:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Aladejana EB:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Mudiyiwa SM:** Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Musara C *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Musara C, Aladejana EB and Mudiyiwa SM. **Review of the nutritional composition, medicinal, phytochemical and pharmacological properties of *Citrus reticulata* Blanco (Rutaceae) [version 1; peer review: 1 approved, 1 not approved]** F1000Research 2020, 9:1387 <https://doi.org/10.12688/f1000research.27208.1>

First published: 02 Dec 2020, 9:1387 <https://doi.org/10.12688/f1000research.27208.1>

Introduction

Citrus reticulata Blanco is a large species belonging to the family Rutaceae, with various varieties and hybrids¹. It includes popular citrus types such as Satsumas, Clementines, Tangerines² and the Mediterranean mandarin^{3,4}. Tangerine is a group of orange-coloured citrus fruits consisting of mandarin hybrids⁵, although the term tangerine is used interchangeably with mandarin.

Mandarins, like other citrus species, are indigenous to the subtropical and tropical zones of Asia, particularly China and Cochin-China^{6,7}. Some researchers have reported that mandarins, alongside other citrus species, evolved in a region including Vietnam, South China, India and Japan^{8,9}. They are now widely cultivated around the world in the warm temperate and tropical areas^{8,10–12}. Mandarins account for 22–25 per cent of world citrus production among the commercially cultivated citrus species^{9,13,14}. The major citrus growing regions of the world and their estimated output is shown in Figure 1.

Mandarins, including both monoembryonic and polyembryonic cultivars and many interspecific hybrids, are the most diversified category of citrus fruits^{15,16}. Nevertheless, a remarkable similarity has been documented between mandarin cultivars at molecular and isoenzymatic levels^{17–19}. There are 36 mandarin species, according to Tanaka²⁰, while Swingle recognized only three species, one of them being *C. reticulata* consisted of 34 species of Tanaka's system²¹. Chromosome studies have shown that the genus has a stable chromosome number of $2n=18$,

except for a few polyploids, with a small number of chromosome markers in the conventional karyotype²². The medicinal uses, phytochemistry and pharmacological properties of *C. reticulata* were examined in the current report.

Botanical description

The mandarin plant is a spiny, evergreen, bushy shrub growing 2–8 m tall²³ with most varieties averaging 7.5 m¹³. The tree has a dense top with slender branches bearing dark green, lance-shaped leaves with a prominent midrib. Petioles are narrowly winged or slightly margined. This tree bears white scented flowers followed by oval to flattened, sweet-fleshed, golden fruits¹³. The mandarin fruit resembles other oranges, is smaller and more oblate than oranges, round in shape, orange in colour, sweet in taste, with a thin, loose, easy-to-peel skin and can be easily damaged by cold. The fruit is up to 8 cm in diameter with easily separable segments².

Flowering is induced through low-temperature stress or soil water deficit stress^{24,25}. Under subtropical climates, flowering is an annual event occurring during spring^{26–28}. In tropical areas, flowering is a continuous event, mostly determined by moisture availability from sufficient rain or water supply^{27–29} while in temperate regions it occurs with the onset of winter³⁰.

Methods and justification of the study

The literature search was performed from March 2020 to June 2020. A mixed-method review approach which involved combining quantitative and qualitative research was used to

Figure 1. World main citrus growing zones and their annual productions. This figure has been reproduced from Mahato *et al.* under the terms of the [Creative Commons Attribution 4.0 International license \(CC-BY-4.0\)](https://creativecommons.org/licenses/by/4.0/)⁹.

compile the review. Information on nutritional composition, medicinal uses, phytochemical and pharmacological properties of *C. reticulata* was gathered from textbooks, theses and online research articles from databases such as Elsevier, Google Scholar, Scopus, Science Direct, SciFinder, PubMed, BMC, and Web of Science. These data sources were chosen based on the topic covered and the main search key terms included “taxonomy, botany, distribution, nutritional composition, ethnobotanical uses, biological and chemical properties” in relation to *C. reticulata*. Search terms were set to be in the title, keywords and abstract. To avoid too much filtering of literature, the terms were searched individually. Focusing, on its multipurpose roles, *C. reticulata* production and utilization can be a catalyst for the development of rural households and community livelihoods. It is therefore imperative to document its nutritional composition, medicinal use and pharmacological properties. With more people becoming nutrition-conscious, demand for

citrus fruits and their by-products has grown even in developing countries.

Nutritional composition

Mandarin is a rich source of vitamins C and A, proteins, dietary fibres and essential minerals such as calcium, potassium, phosphorus and magnesium. Also, they contain minute quantities of vitamins B1, B2, B3, B5, B6, B9 and E^{8,13}. On average, 100 g of mandarin orange consists of 85% water (85.2 g), 13% carbohydrates (13.34 g), 0.81 g protein, 0.38 g dietary fibre and 0.31 g fat^{8,13}. Table 1 shows the nutritional composition of raw mandarins.

The sugars, acids, carotenoids, polyphenols, limonoids and vitamins in *C. reticulata* determine the flavour of the fruit. The vitamins, fibre and health-boosting plant compounds like flavonoids provide many health benefits to humans who eat

Table 1. Nutritional value of raw mandarins. The table has been reproduced with permission from Liu *et al.*⁸.

Nutrient	Content in 100g	Nutrient	Content in 100g
Energy	53 kcal/222 kJ	Vitamin A	34 µg/ 681I.U.
Proteins	0.81 g	Vitamin E (alpha-tocopherol)	0.20 mg
Carbohydrates	13.34 g	Vitamin C	26.7 mg
Sugars	10.58 g	Vitamin B1 (Thiamin)	0.058 mg
Ash	0.38 g	Vitamin B2 (Riboflavin)	0.036 mg
Dietary fibre	1.8 g	Vitamin B3 (Vitamin PP, Niacin)	0.376 mg
Total Lipids (Fats)	0.31 g	Vitamin B6	0.078 mg
Total saturated fatty acids	0.039 g	Folate (Vitamin B9)	16 µg
Total monosaturated fatty acids	0.060 g	Choline	10.2 mg
Total polysaturated fatty acids	0.065 g	Pantothenic acid (Vitamin B5)	0.216 mg
Omega 3 (n-3) fatty acids	0.018 g	beta-Cryptoxanthin	407 µg
Omega 6 (n-6) fatty acids	0.047 g	alpha-Carotene	101 µg
Calcium	37 mg	beta-Carotene	155 µg
Iron	0.15 mg	Betaine	0.1 mg
Magnesium	12 mg	Cholesterol	0 mg
Phosphorus	20 mg	Phytosterols	-
Potassium	166 mg	Xanthophylls	138 µg
Sodium	2 mg		
Zinc	0.07 mg		
Copper	0.042 mg		
Manganese	0.039 mg		
Selenium	0.1 µg		

the fruits and related by-products. For example, the vitamin B complex helps prevent infections, helps promote cell health, energy levels, proper nerve functions, hormone and cholesterol production, and cardiovascular health. In particular, mandarin fruits are rich in β -cryptoxanthine, xanthophyll with pro-vitamin A activity³¹.

Mandarins are well-accepted by consumers owing to their pleasant flavours and abundant phytochemicals. With more people becoming nutrition-conscious, there has been a growing demand for citrus fruits like mandarins, and their by-products¹³.

Chemical constituents

The chemical constituents of *C. reticulata* peel, juice and fruit are shown in Table 2. The peel has high magnesium and carotenoid content³². Methyl-N-methyl anthranilate, a natural antinociceptive compound, has been isolated from mandarin leaves³³. Secondary metabolites such as terpenoids, flavonoids and phenolics compounds act as deterrents to insects and microbial attack³⁴.

Essence oil and aroma

Mandarin is a source of essential oils which are characterized by a fresh-juice fragrance that is widely used in citrus juice products as a natural flavoring agent^{8,35}. The essential oils contain volatile compounds, mainly aldehydes, limonene, ketones, esters, alcohols, terpenes, β -myrcene, 3-carene and α -pinene which provides the distinctive aromas and tastes of citrus fruits^{1,36,37}. Limonene, preceded by γ -terpinene, p-cymene,

α -pinene and myrcene, is the most abundant compound in mandarin essential oil^{1,38–43}.

The essential oils are greatly utilized as fragrance materials in beverages, foods, medical formulations, perfumery, toiletries and other cosmetic products⁴⁴. To some extent, they can also be used as traditional medicine^{2,9,45}. The Chinese use the dried peel of the fruit in the regulation of ch'I (energy/vitality) and to enhance digestion. The leaves and juvenile twigs are a source of essential oil called 'petitgrain oil'²³.

Medicinal uses

The edible part of the raw mandarin fruit possesses antioxidants such as vitamin C, carotenoids and phenolic compounds. The fruit is also a rich source of amino acids, sugars, organic acids, amino acids, pectins, minerals and volatile organic compounds^{46–51}. These constituents are essential for the proper functioning of the body by protecting it against chronic diseases and providing basic nutrition⁵². The dietary fibre and phenolic compounds in mandarins are useful in the formulation of functional foods³². Mandarin fruit also contains coumarins, for instance, bergapten which sensitizes the skin to sunlight⁵³.

The fruit has been reported to possess laxative, aphrodisiac, antiemetic, astringent and tonic properties^{54,55} while the fruit peel regulates skin moisture, softens hard and rough skin and cleanses oily skin⁵⁶. Traditionally, it is also used as a stomachic and carminative^{56–58}. Both the pericarp and endocarp are anticholesterolemic, analgesic, antiseptic, antiasthmatic,

Table 2. Chemical constituents of *C. reticulata*.

Part	Constituents	References
Peel	Fat, protein, ash, magnesium, carotenoids, dietary fibre and polyphenols	32
Juice	Beta-cryptoxanthin xanthophyll esters (zeaxanthin and lutein)	59
	Carotenoids, such as beta-cryptoxanthin.	60,61
Seed	Limonoids, including obacunone 17 beta-D-glucopyranoside, nomilinic acid 17 beta-D-glucopyranoside, limonin, nomilin, and a limonoid glucoside mixture.	62
Fruit	Alkaloids, flavonoids, tannins, phenols and saponins	34
Hexane leaves extract	Alkaloid, carbohydrate, cardiac glycosides and terpenoid	63
Methanol leaves extract	Carbohydrate, cardiac Glycosides	63
Ethanol leaves extract	Alkaloid, carbohydrate, cardiac glycosides and protein	63
Ethanol extract of the aerial part	Carbohydrates, phytosterols, flavonoids, leucoanthocyanins, and tannins	64
Methanol extract of the air-dried fruit peels	n-hexacosanoic acid, reticulataursenoside, citrusterylarachidate and citruslanosteroside	56
Ethanol extracts of the fruit peel	Saponins, tannins, cardiac glycosides, carbohydrates, reducing sugars, compound reducing sugars, terpenoids, flavonoids and sterols.	65
	Carbohydrates, amino acids, flavonoids, steroids, tannins and phenolic derivatives.	66

anti-inflammatory, antiscorbutic, antitussive, carminative, expectorant and stomachic^{57,67}. Therefore, they are used in the treatment and management of dyspepsia, gastro-intestinal distension, cough with profuse phlegm, hiccup and vomiting^{58,67}. The unripened green exocarp is used in the treatment of chest pains and hypochondrium, gastro-intestinal distension, swelling of the liver and spleen and cirrhosis of the liver. The seed is analgesic and carminative, thus used in the treatment of hernia, lumbago, mastitis and pain or swellings of the testes⁶⁷.

Pharmacological properties of *C. reticulata*

Antimicrobial properties

The ethanolic extract of mandarin fruit shell, a traditional herbal medicine used for gastric ulcer treatments in China, showed activity against five clinical strains of *Helicobacter pylori* at the minimum inhibitory concentration (MIC) close to 60 µg/mL⁶⁸. The essential oil of this plant has shown antimicrobial activity with the zone of inhibition varying from 9.16 to 27.63 mm against *Escherichia coli*, *Listeria innocua*, Methicillin-Resistant *Staphylococcus aureus*, *S. aureus* and *Candida albicans*⁴³. In a comparative study, the peel ethanol extract of *C. reticulata* inhibited the growth of all the Gram-positive bacteria tested, with the highest zone of inhibition of 20.33 ± 1.527 mm against *Bacillus* spp. However, the juice extract showed more activity against the Gram-negative bacteria with a maximum zone of inhibition of 11.33 ± 1.154 mm against *Klebsiella pneumonia*⁶⁹. Zainab *et al.*⁷⁰ also reported that the peel extract of *C. reticulata* exhibited a high zone of inhibition against *S. aureus* (28 mm) while *E. coli*, *S. typhi* and *P. aeruginosa* showed resistance to the peel extracts. The presence of flavanones in the peel of *C. reticulata* could be responsible for the efficacy of the peel extract than that of the juice⁷¹. Yashaswini and Arvind⁷² carried out a study to determine the antibacterial potential of *C. reticulata* var. Kinnow peel extracts against pathogenic strains of *S. aureus*, *E. coli*, *P. aeruginosa* and *K. pneumonia*. The acetone extract inhibited the growth of *K. pneumonia* and *E. coli* with a MIC value of 68.75 µg/mL and maximum zone of inhibition of 7.93 mm and 7.75 mm against *K. pneumonia* and *E. coli* respectively.

Sultana *et al.*⁵⁷ have reported that the volatile oil of *C. reticulata* peel possesses antimicrobial activities against *Escherichia coli*, *Staphylococcus aureus*, *Aspergillus flavus*, *Aspergillus niger*, *Aspergillus fumigatus*, and *Candida albicans*. Thus, suggested that the volatile oil could be useful for the treatment of skin disorders and the therapy can be incorporated into the cosmetic formulation. Based on the findings of this review, the essential oil, juice and peel extracts of *C. reticulata* may possess beneficial antibacterial agents that can be exploited in controlling unwanted bacterial infections. The peel oils of mandarins exhibit toxic insecticidal and antibacterial properties⁸.

Anticancer activity

Kang *et al.*⁷³ have reported that the methanol extracts (100 g/mL) of *C. reticulata* peel showed increased apoptosis on SNU-C4, human colon cancer cells through Bax-related *caspase-3* activation, thus, suggested the use of *C. reticulata* on colon cancer patients. In an *in vitro* study, two flavone glucosides isolated from the mandarin fruit peel showed differentiation-inducing

activity in mouse myeloid leukaemia cells (M1), and the cells exhibited phagocytic activity⁷⁴. In addition, hexane and dichloromethane bark extracts of *C. reticulata* assayed against human lung adenocarcinoma cell line A549, human breast adenocarcinoma cell line MCF7, human Caucasian prostate adenocarcinoma cell line PC3, and one normal human prostate cell line PNT2 revealed that the extracts possess good apoptosis-inducing activity against the human cancer cell lines. Thus, the authors concluded that the hexane or dichloromethane extract of the bark of *C. reticulata* is a good crude drug treatment against lung, breast and prostate cancer. However, further *in vitro* and *in vivo* testing would be required before any recommendation of this drug can be given⁷⁵.

The anticancer potential of *Citrus medica* (2 morphotypes), *C. sinensis*, *C. maxima*, *C. limon* and *C. reticulata* peels were investigated using *in vitro* assays and *in vivo* cancer models⁷⁶. The finding depicted that both the extracts and EOs of *C. reticulata* peels had significant activity against Dalton's Lymphoma Ascites (DLA) cell line in an MTT assay. The peel oil showed 91.9% and 100% cell death at 25 and 50 µg/mL, respectively, while the water extract showed 49.8% cell death at 5 µg/mL and 100% cell death at 25 and 50 µg/mL respectively. The *in vivo* study revealed that mice pre-treated with *C. reticulata* peel extract were significantly (50%) protected from DLA compared to post-treated mice (33%) without any obvious toxic symptoms. The volatiles (essential oils, limonoids) and non-volatiles (mainly polymethoxy flavones) in *Citrus* peels have been recognized as their bioactive/anticancer constituents^{76,77}.

The antiproliferative activity of limonoids extracted from *C. reticulata* was evaluated against a series of human cancer cell lines⁶². Limonoids exhibited significant growth inhibitory effects at high concentration of 100 µg/mL against human breast cancer cell lines (MCF-7). However, it could not inhibit leukaemia (HL-60), ovary (SKOV-3), cervix (HeLa), stomach (NCI-SNU-1) and liver (Hep G2) cancer cells lines⁶².

Neuropharmacological activities

Gbaj *et al.*⁷⁸ evaluated the anxiolytic potentials of methanol and aqueous peels extracts of *C. reticulata* in Libya using an elevated plus-maze. The result revealed that the peel extracts exhibited significant anxiolytic activity. In addition, the anxiolytic effect of naringin has been confirmed in 6-8 weeks old mice weighing 30 to 35 g⁷⁹.

Antigenotoxicity effects

Hassan *et al.*⁶⁴ investigate the protective effect of the ethanol extract of the aerial part of *C. reticulata* cultivated in Saudi Arabia against genotoxicity induced by benzo(a)pyrene (BaP) in mice using the comet assay. In the mice treated with BaP, there was a significant increase in the DNA fragmentation in the liver tissues of male mice and an increased rate of DNA damage in mice blood cells. However, the liver and blood cells of the mice treated with ethanol extract demonstrated significant protection by inhibiting the rate of DNA damage. It was concluded that the aerial part of *C. reticulata* could be useful to reduce the genotoxicity induced by hazardous chemical agents⁶⁴. The presence of flavonoid compounds and various secondary

metabolites could be responsible for the protective effect, pharmacological and therapeutic properties of *C. reticulata*^{80–82}.

Antioxidant effects

Boudries *et al.*⁴³ investigated the antioxidant activities of *C. reticulata*, *C. reticulata* cultivar Wilking and *C. clementine* from Algeria using 1,1-diphenyl-2-picrylhydrazil (DPPH) and reducing power. The essential oil (EO) of *C. reticulata* exhibited the strongest DPPH free radical-scavenging activity in a dose-dependent manner, followed by clementine and wilking EOs. Also, in a concentration-dependent manner, the EO of *C. reticulata* showed the greatest reducing power followed by wilking and clementine EOs. According to Junior *et al.*⁸³, the antioxidant nature of the citrus essential oils in terms of free radical scavenging may be due to the antioxidant activity of limonene, which was the main constituent of the oil. The peel of *C. reticulata* was evaluated for antioxidant activity, the results displayed prominent, concentration-dependent free-radical scavenging activity on stable DPPH free radicals and reactive hydroxyl radicals⁸⁴.

Also, the fruit peel of *C. reticulata*, *Zingiber officinale* and *Sesamum indicum* were investigated for their antioxidant activities using the DPPH radical scavenging technique. The findings revealed that *Z. officinale* had the highest antioxidant activity followed by *C. reticulata*, and *S. indicum*. The antioxidant activity of these plants could be attributed to a wide variety of constituents, such as the flavonoid content which are considered as major biological antioxidants⁸⁵.

Cardiovascular effects

Rincon *et al.*³² suggested the use of tangerine peel in reducing the risk of cardiovascular diseases and some associated with lipid oxidation.

Hepatoprotective activity

The protective effect of the essential oils of *C. reticulata* on isoniazid induced hepatotoxicity in Wistar rats was investigated⁸⁶. About 50 gm/kg, p.o. of isoniazid was administered for 30 days in order to induced liver damage in the rats. A total of 200 mg/kg, p.o. of the essential oil was administered daily for 30 days, while the standard group received Liv52⁸⁶. The result revealed a significantly elevated level of ALT, AST, bilirubin and a decreased total protein content in the rats treated with only isoniazid as compared to the group that do not received isoniazid. However, a significant reduction in all the biochemical parameters was observed in the rats treated with the essential oil and Liv52⁸⁶.

CYP450 effects

In an *in vitro* study, the effect of tangeretin (a flavonoid isolated from tangerine juice) on hydroxylation of midazolam, a CYP3A4 probe was evaluated using human liver microsomes and recombinant CYP3A4. The finding revealed that tangeretin is a potent and regioselective stimulator of midazolam 1'-hydroxylation and complementary DNA-expressed CYP3A4⁸⁷. However, further studies are required as the authors have indicated that tangerine juice might not have a clinical effect on CYP3A-mediated drug metabolism in humans.

Antihypercholesterolemic effects

Omer *et al.*⁶⁵ investigated the antihypercholesterolemic potential of the crude ethanolic extracts of *C. reticulata* fruit peel in an *in vivo* study. The findings revealed that daily administration of 250 mg/kg and 500 mg/kg doses of the extracts to the albino rats for four weeks produced a reduction in serum low-density lipoprotein-cholesterol, total cholesterol and triglycerides levels. Also, a significant elevation in serum high-density lipoprotein-cholesterol was observed, thus, indicating their cardioprotective effects and potential as therapeutic antihypercholesterolemic agents⁶⁵. Hence, the efficacy of *C. reticulata* peels extracts on the observed lipid profile parameters might be attributed to the presence of polymethoxylated flavones which occur in the fruit peels.

Anti-ageing potential

According to a study by Apraj and Pandita⁶⁶, both hot and cold alcoholic extracts of *C. reticulata* exhibited strong anti-collagenase and anti-elastase activity, indicating its anti-ageing ability. However, further study is required to determine whether the extracts can be incorporated into skincare products as anti-wrinkle agents.

Toxicity

The oral administration of *C. reticulata* extracts up to 200 mg/kg has been reported to be safe⁷⁸. Also, Li *et al.*⁶⁸ reported that a single oral dose of 16 g/kg of naringin does not produce acute oral toxicity in rats.

Other uses

Worldwide, the mandarin fruit is eaten as fresh produce⁸⁸. It is peeled and eaten plain, used in salads, desserts and main dishes or cooked in puddings, cakes and confectionery. The peel, pulp and seeds are often discarded as waste or they can be processed into animal feed⁸⁸.

Many by-products, including pectin, dried pulp, molasses, marmalades, candied peel, peel seasoning, purees, beverage bases, alcohol, bland syrup, citric acid, seed oil and flavonoids can be obtained from mandarin fruits^{4,36,89}. The dried peel or rind has a sweet-spicy flavour often used as a flavouring in cakes⁴ or as a spice for cooking, baking, drinks or candy⁹⁰. Rind powder extract, a rich source of phenolic compounds having free radical scavenging activity can be used as an anti-oxidant in meat products⁸⁸.

Conclusion

C. reticulata is an important plant which contains some compounds and nutritional values that are of great health importance. The fruit is rich in antioxidants and phenolic compounds, sugars, organic acids, amino acids, pectins, minerals and volatile organic compounds. These substances are essential for the proper functioning of the body by protecting it against chronic diseases, provides basic nutrition and useful in the formulation of functional foods. Based on research carried out, the fruits and peels of *C. reticulata* have been reported to possess neuropharmacological, hepatoprotective, anticancer, antimicrobial, antigenotoxicity, antioxidant activities, antihypercholesterolemic and cardiovascular effects. The usage

of *C. reticulata* was shown to be broad, ranging from dietary applications to the treatment of terminal medical conditions, thus, it is imperative to carry out more research on the toxicity of this plant. Since the peels contain bioactive constituents of pharmacological importance, further investigations should be conducted to investigate whether it could be boiled and

consumed orally, establish dosage ranges for safe consumption and evaluate target-organ toxicity.

Data availability

Underlying data

No data are associated with this article

References

- Sawamura M, Thi Minh Tu N, Onishi Y, et al.: **Characteristic odor components of *Citrus reticulata* Blanco (ponkan) cold-pressed oil.** *Biosci Biotechnol Biochem.* 2004; **68**(8): 1690–1697.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Boughendjioua H, Boughendjioua Z: **Chemical composition and biological activity of essential oil mandarin (*Citrus reticulata*) cultivated in Algeria.** *Int J Pharm Sci Res.* 2017; **44**(1): 179–184.
[Reference Source](#)
- Ortiz JM: **Botany: Taxonomy, morphology and physiology of fruits, leaves and flowers.** In: Dugo G and Di Giacomo A. *Citrus: The genus Citrus*. New York: Taylor and Francis; 2002; 16–35.
[Reference Source](#)
- Dharmawan J: **Characterization of volatile compounds in selected citrus fruits from Asia.** PhD Thesis, National University of Singapore. Singapore. 2009.
[Reference Source](#)
- Webber HJ: **Cultivated varieties of citrus.** In: Webber H. J and L. D. Batchelor. *The citrus industry, Volume I: History, botany and breeding*. Berkeley: University of California Press; 1948; 475–668.
- Tolkowsky S: **Hesperides: A history of the culture and use of citrus fruits.** London: John Bale, Sons & Curnow Ltd; 1938.
[Reference Source](#)
- Gmitter FG, Hu X: **The possible role of Yunnan, China, in the origin of contemporary 78 Citrus species (Rutaceae).** *Econ Bot.* 1990; **44**: 267–277.
[Publisher Full Text](#)
- Liu YQ, Heying E, Tanumihardjo SA: **History, global distribution, and nutritional importance of citrus fruits.** *Compr Rev Food Sci Food Saf.* 2012; **11**(6): 530–545.
[Publisher Full Text](#)
- Mahato N, Sinha M, Sharma K, et al.: **Modern Extraction and Purification Techniques for Obtaining High Purity Food-Grade Bioactive Compounds and Value-Added Co-Products from Citrus Wastes.** *Foods.* 2019; **8**(11): 523.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cronje PJ, Barry GH, Huysamer M: **Postharvest rind breakdown of 'Nules Clementine' mandarin is influenced by ethylene application, storage temperature and storage duration.** *Postharvest Biol Tec.* 2011; **60**(3): 192–201.
[Publisher Full Text](#)
- Zubair M, Komal R, Nasir R, et al.: **Antimicrobial potential of various extract and fractions of leaves of *Solanum nigrum*.** *Int J Phytomed.* 2011; **3**(1): 63–67.
[Reference Source](#)
- Zubair M, Balal RM, Aqueel MA, et al.: **Nutritional assessment of Kinnow Mandarin fruit (*Citrus reticulata* Blanco), infected by few sucking insect-pests of citrus.** *Pakistan Journal of Nutrition.* 2015; **14**(8): 487–491.
[Publisher Full Text](#)
- Putnik P, Barba FJ, Lorenzo JM, et al.: **An integrated approach to Mandarin processing: Food safety and nutritional quality, consumer preference, and nutrient bioaccessibility.** *Compr Rev Food Sci Food Saf.* 2017; **16**(6): 1345–1358.
[Publisher Full Text](#)
- Usman M, Fatima B: **Mandarin (*Citrus reticulata* Blanco) breeding.** In: Al-Khayri J, Jain S and Johnson D. *Advances in plant breeding strategies: Fruits*. Cham: Springer; 2018.
[Publisher Full Text](#)
- Hodgson RW: **Horticultural varieties of citrus.** In: Reuther W, Webber HJ, and Batchelor LD. *The Citrus industry*, rev. University of California Press; 1967.
[Reference Source](#)
- Barrett HC, Rhodes AM: **A numerical taxonomic study of affinity relationships in cultivated citrus and its close relatives.** *Syst Bot.* 1976; **1**(2): 105–136.
[Publisher Full Text](#)
- Esen A, Scora RW: **Amylase polymorphism in citrus and some related genera.** *Am J Bot.* 1977; **64**(3): 305–309.
[Publisher Full Text](#)
- Machado MA, Coletta-Filho HD, Targón MLPN, et al.: **Genetic relationship of Mediterranean mandarins (*Citrus deliciosa* Tenore) using RAPD markers.** *Euphytica.* 1996; **92**: 321–326.
[Publisher Full Text](#)
- Coletta-Filho HD, Machado MA, Targón MLPN, et al.: **Analysis of the genetic diversity among mandarins (*Citrus* spp.) using RAPD markers.** *Euphytica.* 1998; **102**: 133–139.
[Publisher Full Text](#)
- Tanaka T: **Fundamental discussion of Citrus classification.** *Stud Citrol.* 1977; **14**: 1–6.
- Swingle WT, Reece PC: **The botany of citrus and its wild relatives.** In: W. Reuther, H. J Webber and L. D Batchelor. *The Citrus Industry 1. History, distribution, botany, and varieties*. Berkeley: University of California; 1967; 190–430.
- Guerra M, Pedrosa A, Silva AEB, et al.: **Chromosome number and secondary constriction variation in 51 accessions of a Citrus germplasm bank.** *Braz J Genet.* 1997; **20**(3): 489–496.
[Publisher Full Text](#)
- Tropical Plants Database, Ken Fern.** tropical.theferns.info. 2020.
[Reference Source](#)
- Nishikawa F: **Regulation of floral induction in citrus.** *J Japan Soc Hort Sci.* 2013; **82**(4): 283–292.
[Publisher Full Text](#)
- Jhade RK, Huchche AD, Dwivedi SK: **Phenology of flowering in citrus: Nagpur mandarin (*Citrus reticulata* Blanco) perspective.** *International Journal of Chemical Studies.* 2018; **6**(2): 1511–1517.
[Reference Source](#)
- Krajewski A, Rabe E: **Citrus flowering: A critical review.** *J Hort Sci.* 1995; **70**: 357–374.
- Iglesias DJ, Cercos M, Colmenero-Flores JM, et al.: **Physiology of citrus fruiting.** *Braz J Plant Physiol.* 2007; **19**(4): 333–362.
[Publisher Full Text](#)
- Stander J: **The reproductive phenology of citrus. II: Citrus floral ontogeny.** *Technology.* 2015; 65–69.
[Reference Source](#)
- Schneider H: **Anatomy of citrus.** In: W. Reuther. *The citrus industry*. Berkeley: University of California. 1968; 1–85.
- Micheloud NG, Castro DC, Buyatti MA, et al.: **Factors affecting phenology of different Citrus varieties under the temperate climate conditions of Santa Fe, Argentina.** *Rev Bras Frutic Jaboticabal.* 2018; **40**(1).
[Publisher Full Text](#)
- Chen Q, Wang D, Tan C, et al.: **Profiling of flavonoid and antioxidant activity of fruit tissues from 27 Chinese local citrus cultivars.** *Plants (Basel).* 2020; **9**(2): 196.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rincon AM, Vasquez AM, Vasquezand FC: **[Chemical composition and bioactive compounds of flour of orange (*Citrus sinensis*), tangerine (*Citrus reticulata*) and grapefruit (*Citrus paradisi*) peels cultivated in Venezuela].** *Arch Latinoam Nutr.* 2005; **55**(3): 305–310.
[PubMed Abstract](#)
- Correa E, Quinones W, Echeverri F: **Methyl-N-methylantranilate, a pungent compound from *Citrus reticulata*. Blanco leaves.** *Pharm Biol.* 2016; **54**(4): 569–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Okwu DE: **Citrus fruits: A rich source of phytochemicals and their roles in human health.** *Int J Chem Sci.* 2008; **6**(2): 451–471.
[Reference Source](#)
- Shaw PE: **Essential oils.** In: S. Nagy, P. E. Shaw and M. K.Veldhuis. *Citrus science and technology, Volume I: Nutrition, anatomy, chemical composition and bioregulation*. Westport: AVI Publishing Co; 1977; 427–478.
- Braddock RJ: **Handbook of citrus by-products and processing technology.** New York: Wiley; 1999.
[Reference Source](#)
- Tao N, Liu Y, Zhang J, et al.: **Chemical composition of essential oil from the**

- peel of Satsuma mandarin. *Afr J Biotechnol*. 2008; 7(9): 1261–64.
[Reference Source](#)
38. Baalouamer A, Meklai BY, Frisse D, et al.: **The chemical composition of some cold-pressed citrus oils produced in Algeria.** *J Essent Oil Res*. 1992; 4(3): 251–258.
[Publisher Full Text](#)
 39. Verzera A, Trozzi A, Mondello L, et al.: **Uruguayan essential oil. Part X. Composition of the oil of *Citrus Clementine* Hort.** *Flavour Fragr J*. 1998; 13(3): 189–195.
[Publisher Full Text](#)
 40. Lota ML, Rocca-Serra D, Torri F, et al.: **Chemical variability of peel and leaf essential oils of 15 species of mandarins.** *Biochem Syst Ecol*. 2017; 29(1): 77–104.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. Choi HS: **Volatile constituents of satsuma mandarins growing in Korea.** *Flavour Fragr J*. 2004; 19(5): 406–412.
[Publisher Full Text](#)
 42. Viuda-Martos M, Ruiz-Navajas Y, Fernandez-Lopez J, et al.: **Chemical composition of mandarin (*C. reticulata* L.), grapefruit (*C. paradisi* L.), lemon (*C. limon* L.) and orange (*C. sinensis* L.) essential oils.** *Journal of Essential Oil Bearing Plants*. 2009; 12(2): 236–243.
[Publisher Full Text](#)
 43. Boudries S, Loupassaki Y, Ladjal Ettoumi S, et al.: **Chemical profile, antimicrobial and antioxidant activities of *Citrus reticulata*. and *Citrus clementina* (L.) essential oils.** *Intern Food Res J*. 2017; 24(4): 1782–1792.
[Reference Source](#)
 44. Baser KHC, Demirci F: **Chemistry of essential oils.** In: R.G.Berger. *Flavours and fragrances: Chemistry, bioprocessing and sustainability*. Berlin: Springer-Verlag; 2007; 43–86.
 45. Imbesi A, De Pasquale A: ***Citrus* species and their essential oils in traditional medicine.** In: Dugo G and DiandDi Giacomo A. *Citrus: The genus Citrus*. New York: Taylor and Francis; 2002; 577–601.
 46. Ye XQ, Chen JC, Liu DH, Jiang P, et al.: **Identification of bioactive composition and antioxidant activity in young mandarin fruits.** *Food Chem*. 2011; 124(4): 1561–6.
[Publisher Full Text](#)
 47. Matsumoto H, Ikoma Y: **Effect of different postharvest temperatures on the accumulation of sugars, organic acids, and amino acids in the juice sacs of Satsuma mandarin (*Citrus unshiu*. Marc) fruit.** *J Agric Food Chem*. 2012; 60(39): 9900–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 48. Zhang YM, Sun YJ, Xi WP, et al.: **Phenolic compositions and antioxidant capacities of Chinese wild mandarin (*Citrus reticulata*. Blanco) fruits.** *Food Chem*. 2014; 145: 674–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
 49. Rodrigo MJ, Cilla A, Barbera R, et al.: **Carotenoid bioaccessibility in pulp and fresh juice from carotenoid-rich sweet oranges and mandarins.** *Food Funct*. 2015; 6(6): 1950–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 50. Antoine S, Paillé O, Gibon Y, et al.: **Short- and long-term effects of carbohydrate limitation on sugar and organic acid accumulation during mandarin fruit growth.** *J Sci Food Agric*. 2016; 96(11): 3906–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
 51. Chen DL, Huang YR, Liang HL, et al.: **Column chromatographic extraction for quickly separating the volatiles, flavonoids, and pectin from tangerine peel.** *Sep Sci Technol*. 2016; 51(3): 485–93.
[Publisher Full Text](#)
 52. Al-Snafi AE: **Nutritional value and pharmacological importance of citrus species grown in Iraq.** *IOSR J Pharm*. 2016; 6(8): 76–108.
[Publisher Full Text](#)
 53. Bown D: **Encyclopaedia of herbs and their uses.** London: Dorling Kindersley; 1995.
[Reference Source](#)
 54. Chopra RN, Nayar SL, Chopra IC: **Glossary of Indian medicinal plants (Including the Supplement).** New Delhi: Council of Scientific and Industrial Research; 1986.
 55. Anonymous: **The wealth of India: A dictionary of Indian raw materials and industrial products.** New Delhi: National Institute of Science Communication and Information Resources; 2000; 65–69.
[Reference Source](#)
 56. Khan MA, Ali M, Alam P: **Phytochemical investigation of the fruit peels of *Citrus reticulata* Blanco.** *Nat Prod Res*. Formerly Natural Product Letters. 2010; 24(7): 610–620.
[PubMed Abstract](#) | [Publisher Full Text](#)
 57. Sultana HS, Ali M, Panda BP: **Influence of volatile constituents of fruit peels of *Citrus reticulata* Blanco on clinically isolated pathogenic microorganisms under in-vitro.** *Asian Pac J Trop Biomed*. 2012; 2(3): S1299–S1302.
[Publisher Full Text](#)
 58. Apraj VD, Pandita NS: **Evaluation of skin anti-aging potential of *Citrus reticulata* Blanco peel.** *Pharmacognosy Res*. 2016; 8(3): 160–168.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 59. Wingerath T, Stahl W, Sies H: **Beta-Cryptoxanthin selectively increases in human chylomicrons upon ingestion of tangerine concentrate rich in beta-cryptoxanthin esters.** *Arch Biochem Biophys*. 1995; 324(2): 385–390.
[PubMed Abstract](#) | [Publisher Full Text](#)
 60. Granado F, Olmedilla B, Blanco I, et al.: **Major fruit and vegetable contributors to the main serum carotenoids in the Spanish diet.** *Eur J Clin Nutr*. 1996; 50(4): 246–250.
[PubMed Abstract](#)
 61. Irwig MS, El Sohemy A, Baylin A, et al.: **Frequent intake of tropical fruits that are rich in beta-cryptoxanthin is associated with higher plasma beta-cryptoxanthin concentrations in Costa Rican adolescents.** *J Nutr*. 2002; 132(10): 3161–3167.
[PubMed Abstract](#) | [Publisher Full Text](#)
 62. Tian Q, Miller EG, Ahmad H, et al.: **Differential inhibition of human cancer cell proliferation by citrus limonoids.** *Nutr Cancer*. 2001; 40(2): 180–184.
[PubMed Abstract](#) | [Publisher Full Text](#)
 63. Balamurugan P, Rajkumar AR, Prasad MP: **Comparative phytochemical analysis of Rutaceae family (*Citrus* Species) extracts.** *Int J Sci*. 2014; 148–150.
[Reference Source](#)
 64. Hassan AZ, Ahmed KM, Abu-Gabal NS, et al.: **Phytochemical and genotoxicity studies of *Citrus reticulata* aerial part in mice.** *Egyptian Pharmaceutical Journal*. 2017; 16(2): 87.
[Publisher Full Text](#)
 65. Omer SS, Elsidig IM, Mohammed AE, et al.: **Phytochemical screening, antioxidant activity and lipid profile effects of *Citrus reticulata* fruit peel, *Zingiber officinale*. Rhizome and *Sesamum indicum* seed extracts.** *Intern J Medical Health Biomed Bioeng Pharm Eng*. 2015; 9: 797–804.
[Reference Source](#)
 66. Apraj VD, Pandita NS: **Pharmacognostic and phytochemical evaluation of *Citrus reticulata* Blanco peel.** *Int J Pharmacogn Phytochem Res*. 2014; 6(26): 328–331.
[Publisher Full Text](#)
 67. Yeung HC: **Handbook of Chinese herbs and formulas.** Los Angeles: Institute of Chinese Medicine; 1985; 1.
[Reference Source](#)
 68. Li Y, Xu C, Zhang Q, et al.: **In vitro anti-*Helicobacter pylori*. action of 30 Chinese herbal medicines used to treat ulcer diseases.** *J Ethnopharmacol*. 2005; 98(3): 329–333.
[PubMed Abstract](#) | [Publisher Full Text](#)
 69. Shakyia A, Luitel P, Kumari R, et al.: **Comparative study of antibacterial activity of juice and peel extract of citrus fruits.** *Tribhuvan University J Microbiol*. 2019; 7(6): 82–88.
[Publisher Full Text](#)
 70. Zainab AGC, Kadhim NK: **Antimicrobial activity of different aqueous lemon extracts.** *J Appl Pharm Sci*. 2013; 3(6): 74–78.
[Reference Source](#)
 71. Levaj B, Dragovic V, Bursac D, et al.: **Determination of flavonoids in pulp and peel of mandarin fruits.** *Agric Conspec Sci*. 2009; 74(3): 221–225.
[Reference Source](#)
 72. Yashaswini Y, Arvind: **Antimicrobial Properties of Orange (*Citrus reticulata* var. Kinnow). Peel Extracts against Pathogenic Bacteria.** *Int J Curr Microbiol App Sci*. 2018; 7(3): 737–746.
[Publisher Full Text](#)
 73. Kang SA, Park HJ, Kim MJ, et al.: ***Citri reticulatae* Viridereticulatae Viride pericarpium extract induced apoptosis in SNU-C4, human colon cancer cells.** *J Ethnopharmacol*. 2005; 97(2): 231–235.
[PubMed Abstract](#) | [Publisher Full Text](#)
 74. Sugiyama K, Umehara M, Kuroyanagi A, et al.: **Studies on the differentiation inducers of myeloid leukemic cells from citrus species.** *Chem Pharm Bull (Tokyo)*. 1993; 41(4): 714–719.
[PubMed Abstract](#) | [Publisher Full Text](#)
 75. PLANTMEDS: **Potential antimalarial and anticancer lead compound discovery from Cameroonian medicinal plants.** 2018.
[Reference Source](#)
 76. Nair A, Kurup SRR, Nair AS, et al.: ***Citrus* peels prevent cancer.** *Phytomed*. 2018; 50: 231–237.
[Publisher Full Text](#)
 77. Koolaji N, Shammugasamy B, Schindeler A, et al.: **Citrus peel flavonoids as potential cancer prevention agents.** *Curr Dev Nutr*. 2020; 4(5): nzaa025.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 78. Gbaj MA, Sadawe IA, Meiqal NM, et al.: **Evaluation of neuropharmacological activities of methanolic and aqueous extracts of *Citrus reticulata* (Rutaceae) fruit peels.** *Am J Biomed Sci Res*. 2019; 2(4): 131–135.
[Publisher Full Text](#)
 79. Fernandez SP, Nguyen M, Yow TT, et al.: **The flavonoid glycosides, myricitrin, gossypin and naringin exert anxiolytic action in mice.** *Neurochem Res*. 2009; 34(10): 1867–1875.
[PubMed Abstract](#) | [Publisher Full Text](#)
 80. Borokini TI, Omotayo FO: **Phytochemical and ethnobotanical study of some selected medicinal plants from Nigeria.** *J Med Plant Res*. 2012; 6(7): 1106–1118.
[Publisher Full Text](#)
 81. Boligon AA, Sagrillo MR, Machado LF, et al.: **Protective effects of extracts**

- and flavonoids isolated from *Scutiabuxifolia* Reissek against chromosome damage in human lymphocytes exposed to hydrogen peroxide. *Molecules*. 2012; **17**(5): 5757–5769.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
82. El-Rafie HM, Mohammed RS, Hamed MA, *et al.*: **Phytochemical and biological studies of total ethanol and petroleum ether extracts of *Terminalia bentzoe*. (L.) leaves.** *Int J Pharmacogn Phytochem Res*. 2016; **8**: 592–603.
[Reference Source](#)
 83. Junior EF, Souza P, Nascimento J, *et al.*: **Antinociceptive and antiinflammatory properties of the ethanolic extract of *Pouteria ramiflora*. roots.** *Latin Amer J Pharm*. 2009; **28**(6): 812–818.
[Reference Source](#)
 84. Tumbas VT, Četkovic GS, Dilas SM, *et al.*: **Antioxidant activity of mandarin (*Citrus reticulata*) peel.** *Acta Periodica Technologica*. 2010; **41**: 195–203.
[Publisher Full Text](#)
 85. Bravo L: **Polyphenols: chemistry, dietary sources, metabolism, and nutritional significance.** *Nutr Rev*. 1998; **56**(11): 317–333.
[PubMed Abstract](#) | [Publisher Full Text](#)
 86. Kangralkar VA, Gavimath CV, Venkatesh V, *et al.*: **Protective effect of essential oils of *Citrus reticulata* on isoniazid induced hepatotoxicity in wistar rats.** *Intern J Pharm Appl*. 2010; **1**(2): 59–61.
 87. Backman JT, Maenpaa J, Belle DJ, *et al.*: **Lack of correlation between *in vitro* and *in vivo* studies on the effects of tangeretin and tangerine juice on midazolam hydroxylation.** *Clin Pharmacol Ther*. 2000; **67**(4): 382–390.
[PubMed Abstract](#) | [Publisher Full Text](#)
 88. Rafiq R, Kaul SA, Sofi N, *et al.*: **Nayik Citrus peel as a source of functional ingredient: A review.** *J Saudi Soc Agric Sci*. 2018; **17**: 351–358.
 89. Braddock RJ: **By-products of citrus fruit.** *Food Technol*. 1995; **49**(9): 74–77.
[Reference Source](#)
 90. Morton JF: **Mandarin orange.** In: *Fruits of warm climates*. Miami; 1987.
[Reference Source](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 13 January 2022

<https://doi.org/10.5256/f1000research.30063.r100915>

© 2022 Rokaya M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Maan B. Rokaya

Institute of Botany, The Czech Academy of Sciences, Průhonice, Czech Republic

Manuscript is relevant for indexing, but it needs major changes. Please see the comments below:

Abstract

- Authors have not mentioned why this review is necessary. Is it because this species is important as medicine, food or any other things? Please write aims.
- In the abstract, mention how many compounds were found until now. How many diseases are treated? What pharmacological studies are carried out?
- What is the conclusion of your review, which is quite not clear in abstract?

Introduction

- I suggest authors to start with importance of plant species as medicine and food. Then mention about *Citrus reticulata* which is used as food and medicine. Mention some figures about uses, production and trade. Then in separate section mention about varieties in brief (too much detail is not required). Introduction should be followed by aims of your study. I guess it will be wise to see some good review papers if you want to write up to the indexable standard.
- Botanical description is not very necessary but if you want to keep, make it brief.
- Justification of the study should be at the end of Introduction not as separate section.

Method

- Literature review was from March 2020-June 2020. Rather than this, mention how many literatures were considered and mention the oldest and newest literature. Again, please see some review papers to write your text in indexable way.

Nutritional composition

- Were these components from the same species or from same species but with different varieties? Need to mention this very clearly.

- Table 1 is redundant.

Chemical constituents, oil and aroma

- Not in detail and needs much more elaboration. From which plant parts were they extracted? Mention in detail. I know it is mentioned in Table 2 but it should be written in detail. Please categorize compounds into different groups so that it will be easy for readers to follow.
- In the beginning mention how many compounds have been isolated so far.
- Give chemical structures of all the compounds.
- Antioxidant property is considered as property with no pharmacological relevance (Gafner, 2018). So, discuss according to this.

Pharmacological properties

- In this section, only information about extracts and results are mentioned, which is not enough. It is also necessary to write about what methods were used. The doses, type of experimental animal/cell/bacteria, controls (positive/negative if any) are required. Please write results in detail but remember that it should be concise. Also make a table to show all these. Please follow some good review papers on specific plant species published elsewhere and to help improve your manuscript.
- Conclusions should directly link to your findings. Write about its importance and future prospective. What are the study gaps and what more is needed?

Overall, this manuscript should undergo Major revision.

Is the topic of the review discussed comprehensively in the context of the current literature?

Yes

Are all factual statements correct and adequately supported by citations?

Yes

Is the review written in accessible language?

Yes

Are the conclusions drawn appropriate in the context of the current research literature?

Partly

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

<https://doi.org/10.5256/f1000research.30063.r75652>

© 2021 Sinniah U. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Uma Rani Sinniah

Department of Crop Science, Faculty of Agriculture, Universiti Putra Malaysia, Serdang, Malaysia

The article is well written for the purpose intended. All the subject matter namely nutritional composition, medicinal, phytochemical and pharmacological properties as highlighted in the title of this article has been adequately reviewed and cited.

Secondary metabolites in plants are of major interest and are often taught in plant biochemistry courses. The content of secondary metabolites differ in different plant species and the use of such plant phytochemical for various purposes such as antibacterial, antimicrobial, anti cancer, anti-inflammatory etc, also differ among the plant species. This paper provides a comprehensive outlook and will be a good specimen to demonstrate the types of secondary metabolites present and its bioactive compounds in relation to pharmacological activities. The article can be a one stop reference on the properties of *Citrus reticulata* up to 2020.

Just a few minor typos are present, which are highlighted [here](#).

Is the topic of the review discussed comprehensively in the context of the current literature?

Yes

Are all factual statements correct and adequately supported by citations?

Yes

Is the review written in accessible language?

Yes

Are the conclusions drawn appropriate in the context of the current research literature?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Plant science on agronomy, biochemistry and phytochemistry

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 26 Jan 2021

Elizabeth Bosede Aladejana

Thank you for taking the time to evaluate our article. All the grammatical errors would be

corrected. Thank you.

Competing Interests: None

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research