Check for updates

CASE REPORT

REVISED Case Report: Bilateral mandibular buccal bifurcation cysts [version 2; peer review: 2 approved, 1 approved with reservations]

Ashwag Aloyouny¹, Hamad Albagieh², Soad Mansour¹, Fahmy Mobarak³

¹Basic Dental Science Department, College of Dentistry, Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia

²Oral Medicine and Diagnostic Science Department, College of Dentistry, King Saud University, Riyadh, Saudi Arabia

³Surgery Department, College of Dentistry, Cairo University, Cairo, Egypt

v2 First published: 23 Dec 2020, 9:1502
<https://doi.org/10.12688/f1000research.28000.1>
 Second version: 21 Jan 2021, 9:1502
<https://doi.org/10.12688/f1000research.28000.2>
 Latest published: 19 Feb 2021, 9:1502
<https://doi.org/10.12688/f1000research.28000.3>

Abstract

Buccal bifurcation cyst (BBC) is a rare inflammatory odontogenic cyst, which commonly affects children in the first decade of life. We report a case of a seven-year-old healthy boy with bilateral BBC, which involved unerupted incomplete permanent mandibular first molars. A review of the literature in English language revealed few similar cases. We reviewed 16 manuscripts of bilateral mandibular BBC, reporting a total of 20 cases since 1970. The clinical features of bilateral mandibular BBC summarized here could assist specialists with an accurate diagnosis and provide patients with optimal management.

Keywords

Buccal Bifurcation Cysts, Bilateral mandibular cysts, Children, Oral Cyst

Open Peer Review

Approval Status

	1	2	3
version 3			
(revision)		view	
19 Feb 2021			
version 2			
(revision)	view	view	view
21 Jan 2021			
version 1			
23 Dec 2020	view		

1. **Denise Tostes Oliveira** , University of São Paulo, Bauru, Brazil
2. **Konstantinos I. Tosios** , National and Kapodistrian University of Athens, Athens, Greece
3. **Shadia Abdel-Hameed Elsayed** , Al Azhar University, Cairo, Egypt
Taibah University, Medina, Saudi Arabia

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Ashwag Aloyouny (aloyouas-j@hotmail.com)

Author roles: **Aloyouny A:** Conceptualization, Funding Acquisition, Investigation, Resources, Supervision, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Albagieh H:** Conceptualization, Supervision, Visualization, Writing – Original Draft Preparation; **Mansour S:** Investigation, Supervision, Validation, Writing – Review & Editing; **Mobarak F:** Conceptualization, Investigation, Supervision, Visualization

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2021 Aloyouny A *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Aloyouny A, Albagieh H, Mansour S and Mobarak F. **Case Report: Bilateral mandibular buccal bifurcation cysts [version 2; peer review: 2 approved, 1 approved with reservations]** F1000Research 2021, 9:1502 <https://doi.org/10.12688/f1000research.28000.2>

First published: 23 Dec 2020, 9:1502 <https://doi.org/10.12688/f1000research.28000.1>

REVISED Amendments from Version 1

Two sentences added to the article:

1) It is worth mentioning that the etiopathogenesis of BBC is the same of paradental cyst, according to World Health Organization Classification of Head and Neck Tumours (2017).

2) The diagnosis of BBC should be established based on the clinical, radiographical and microscopic characteristics.

Any further responses from the reviewers can be found at the end of the article

Introduction

Buccal bifurcation cyst (BBC) is a rare inflammatory odontogenic cyst. The first case of BBC was reported in 1983 by Stoneman and Worth¹. Children between the age of 4 and 14 years are most commonly affected. In addition, BBC occurs more in the mandible and most likely involves the permanent mandibular first molar. Typically, BBC causes many undesirable oral manifestations such as buccal swelling at the affected area, delayed tooth eruption or partially tooth eruption associated with deep periodontal pockets. In some cases, pain and infection associated with pus could be present².

Radiographically, BBC is illustrated as a well-defined radiolucent lesion surrounded by sclerotic rim. The lesion either involves the roots of a partially erupted tooth or surrounds an unerupted tooth; usually the permanent mandibular first molar³. Histopathologically, the BBC cyst wall is lined by a non-keratinized stratified squamous epithelium with inflammatory lymphocytes infiltrate. Surgical excisional procedure is performed for complete removal of the cysts with the involved teeth to reduce the risk of recurrence of the epithelial cysts⁴.

Here, we report a case of a seven-year-old healthy boy with a chief complaint of painful, slowly growing lower jaw swellings on both right and left sides consistent with bilateral BBC, which involved unerupted incomplete permanent mandibular first molars. A review of the literature revealed few similar cases. We report this case to add to an additional case of bilateral mandibular BBC to the literature.

Case presentation

A seven-year-old healthy boy presented to the Oral and Maxillofacial Surgery Clinic with a chief complaint of painful, slowly growing lower jaw swellings on both right and left sides. The patient had no past medical, surgical, or dental history. His guardian reported no known drug and food allergies. The guardian also denied having family history of genetic related diseases or syndromes. The swellings caused esthetic disfiguring of the patient's face, which led to social exclusion.

On examination, extraoral, bilateral asymmetric swellings of the lower part of the face with no lymph node involvement were noted. Intraorally, painful bilateral, hard, bony mandibular swellings covered with normal color mucosa, extending from the mesial aspect of the mandibular second primary molars on both sides including the retromolar areas, was observed. On palpation, the affected site revealed expansion of the buccal cortical plates on both sides. Both mandibular permanent first molars were not clinically erupted.

Diagnostic radiographs were taken. A panoramic radiograph illustrated bilateral, well-defined radiolucencies surrounded by sclerotic margins and including the unerupted incomplete permanent mandibular first molars (36 and 46, according to the FDI World Dental Federation Notation). The panoramic image showing right radiolucency measured around 1 cm in its greatest dimension, and involved the unerupted permanent mandibular first molar (46, according to the FDI World Dental Federation Notation), and not involving the inferior mandibular cortical bone and the adjacent areas. On the other hand, the left radiolucency measured around 3cm in its greatest dimension, and involved the unerupted permanent mandibular first molar (36, according to the FDI World Dental Federation Notation), the inferior mandibular cortical bone and the adjacent areas. Moreover, the left radiolucent cyst distally displaced the permanent mandibular second molar tooth bud (37, according to the FDI World Dental Federation Notation) (Figure 1). Provisional diagnosis of the bilateral lesions suggested bilateral BBC, dentigerous cyst, and paradental cyst.

The guardian was informed about the patient's condition, possible management and complications. Then, a consent form was signed before starting the complete surgical removal of the lesions under general anesthesia. An incision was made, then a buccal mucoperiosteal flap was raised in the mandible at the mandibular permanent first molar area on both right and left sides to expose the bone. The buccal plate was expanded and thinned; especially in the left side. An access to the cyst area was made by performing an osteotomy on both sides. The whole cyst lining was removed with the attached mandibular first molars. Then, the cavity was enucleated and curetted thoroughly (Figure 2), to reduce the risk of recurrence.

The specimens were sent to the histopathology laboratory in two containers. The histopathologic analysis of the multiple serial sections of the two specimens revealed multiple pieces of cyst wall lined by small strip of non-keratinized stratified squamous epithelium with mixed inflammatory reaction and surrounding edematous granulation tissue formed of numerous proliferating capillary type vascular spaces. Also, the specimens

Figure 1. A seven-year-old healthy boy with mixed dentition and bilateral BBC which involved unerupted incomplete permanent mandibular first molars.

Figure 2. (a and b) Clinical intraoral photographs show bilateral surgical removal of the bilateral BBC with the involved permanent teeth.

under the microscope showed bland looking spindle-shaped fibroblasts and degenerated bony spicules, which was consistent with cystic lesions (Figure 3). Based on the clinical examination, radiographic interpretation, and histopathologic analysis, the final diagnosis was bilateral BBC.

Six months later, the clinical and radiographic examination of both surgery sites showed signs of gradual healing and resolution and increased bone density in the areas. One year later, no clinical evidence of lesion recurrence, and radiographic imaging illustrated that the affected areas were completely filled up with bone.

Discussion

Bilateral BBC is an uncommon inflammatory odontogenic cyst. It usually arises from the buccal side of the mandibular permanent first molar in children. The pathophysiology of BBC is still undetermined. It is claimed that the inflammation is caused by the deep periodontal pocket caused by tilted tooth. Other authors claim that the reason behind the cyst formation could be because of a defect in the tooth eruption, which may lead to inflammation, proliferation of the epithelial cells and cyst formation¹⁻⁵. It is also postulated that enamel projections at the area covered by reduced enamel epithelium between the cemento-enamel junction and the furcation of the tooth could lead to cyst formation⁶. It is worth mentioning that the etio-pathogenesis of BBC is the same of paradental cyst, according to World Health Organization Classification of Head and Neck Tumours (2017).

Clinically, bilateral BBC has few challenging similarities with dentigerous cyst, and paradental cyst. For example, all previously mentioned cysts are commonly associated with a tooth. However, paradental cysts and dentigerous cysts usually occur in adults in the mandibular third molar region^{7,8}. Dentigerous cyst is classified as a developmental cyst that may also occur in the mandibular first molar areas and sometimes interfere with the process of normal tooth eruption in children. On the other hand, paradental cyst is considered as an inflammatory cyst, which commonly occurs in the lateral surface of a vital tooth usually the first and second mandibular molars of primary teeth.

Figure 3. High-power histology shows the cyst wall with chronic inflammatory lymphocytes and two to four cell layers thick of proliferating lining of nonkeratinized stratified squamous epithelium.

Radiographic interpretation is essential to assist recognize and differentiate between different cystic lesions. Normally cystic lesions have a well-defined, unilocular radiolucency with a sclerotic margin; however, cysts sometimes present with different features. For instance, paradental cysts present on the lateral crown of an incompletely erupted tooth, while dentigerous cysts usually surround the crown of an unerupted tooth. Every rule has an exception, it is challenging to differentiate between the circumferential dentigerous cyst and the BBC radiographically as both illustrated as radiolucent cysts involving a completely unerupted tooth. Histopathologically, the cyst wall under the microscope shows chronic inflammatory lymphocytes and two to four cell layers thick of proliferating lining of nonkeratinized stratified squamous epithelium.

The diagnosis of BBC should be established based on the clinical, radiographical and microscopic characteristics. The clinical examination, radiographic interpretation and histopathologic analysis of both lesions displaying cystic origin

Table 1. Data of reported cases with Bilateral mandibular buccal bifurcation cysts since 1970.

Author	Year	Patient age	Patient gender	Number of bilateral cases	Associated tooth	Degree of tooth eruption	Management	Follow up
Stanback ⁹	1970	9 years	M	1	Bilateral mandibular permanent first molars/vital	Not erupted	Marsupialization	2 years
Swerdlhoff ¹⁰	1980	7 years	Not reported	1	Not reported	Not reported	Enucleation	6 months
Vedtofte and Praetorius ¹¹	1989	13 years	F	2	Case1: Bilateral mandibular permanent second molars/vital	Fully erupted	Enucleation	1–6 years
		Not reported	Not reported		Case2: Bilateral mandibular permanent first molars/vital	Not reported	Enucleation	1–6 years
Packota <i>et al.</i> ¹²	1990	8 years	Not reported	1	Bilateral mandibular permanent first molars/vital	Partially erupted	Enucleation	6 months
Bohay <i>et al.</i> ¹³	1992	Not reported	Not reported	1	Bilateral mandibular permanent first molars/vital	Not reported	Enucleation	8 months
Martinez-Conde <i>et al.</i> ¹⁴	1995	11 years	M	1	Bilateral mandibular permanent second molars/vital	Partially erupted	Enucleation/tooth extraction	0
David <i>et al.</i> ⁵	1998	8 years	M	3	Case 1: Bilateral mandibular permanent first molars/vital	Fully erupted	Regress without procedure	1–2 years
		9 years	M		Case2: Bilateral mandibular permanent first molars/vital	Not erupted	9 months follow up then enucleation right side/no procedure left side	1–2 years
		7 years	M		Case3: Bilateral mandibular permanent first molars/vital	Fully erupted	Daily irrigation with saline and hydrogen peroxide	1–2 years
Shohat I <i>et al.</i> ¹⁵	2003	13 years	M	2	Case1: Bilateral mandibular permanent second molars/vital	Fully erupted	Enucleation/tooth extraction	2 years
		8 years	M		Case2: Bilateral mandibular permanent first molars/vital	Fully erupted	Enucleation/tooth extraction	2 years
Gallego ¹⁶	2007	8 years	M	1	Bilateral mandibular permanent first molars/vital	Fully erupted	Enucleation left side/no procedure right side	1 year
Corona-Rodriguez <i>et al.</i> ¹⁷	2011	7 years	M	1	Bilateral mandibular permanent first molars/vital	Erupted	No procedure right side/enucleation left side	6 months
Ramos <i>et al.</i> ¹⁸	2012	9 years	M	1	Bilateral mandibular permanent first molars/vital	Erupted	Enucleation	1 year
Borgonovo <i>et al.</i> ¹⁹	2012	8 years	M	1	Bilateral mandibular permanent first molars/vital	Not erupted	Enucleation	1 year
Boffano P <i>et al.</i> ²⁰	2012	9 years	M	1	Bilateral mandibular permanent first molars/vital	Not reported	Enucleation	6 months
Issler A <i>et al.</i> ²¹	2013	8 years	F	1	Bilateral mandibular permanent first molars/vital	Not erupted	Enucleation	18 months
Bautista <i>et al.</i> ²²	2019	7 years	F	1	Bilateral mandibular permanent first molars/vital	Left first molar was not erupted	Enucleation/bone graft	0
						Right first molar was partially erupted		
Present case	2020	7 years	M	1	Bilateral mandibular permanent first molars/vital	Not erupted	Enucleation/tooth extraction	1 year

M: Male; F: female

with bilateral buccal expansion involving both mandibular first molars in a seven-year-old child associated with bilateral BBC was presented here. This reported case had similar clinical, radiographical and histopathological features of bilateral mandibular BBC. It is worth mentioning that the case presented here is very rare because it describes bilateral mandibular BBC. To the best of our knowledge, the bilateral characteristic of BBC is very rare; there are only few cases have been reported in the English-language literature. In this article, we reviewed 16 manuscripts of bilateral mandibular BBC, reporting a total of 20 cases in the period between 1970 to 2019. Furthermore, these cases occurred in different ages starting from 4 years to 13 years with predilection to male patients (Table 1).

Through the past years different approaches have been performed to manage the BBC. In some cases, the way of management was only by following up the lesion with no interventions; some lesions showed different degrees of regression and others needed intervention^{5-8,17}. On the other hand, other cases underwent surgical enucleation of the cysts either with the involved tooth or maintaining the involved tooth. In the current case, we chose to perform a surgical procedure under general anesthesia to enucleate both bilateral mandibular BBCs with the involved teeth to reduce the risk of recurrence and to fulfill the patient's parent's desire.

Conclusion

BBC typically affects children in the first decade of life. BBC occurs in the buccal area of the mandibular first molar. Only a

few bilateral mandibular BBC cases were reported in the literature. Although bilateral mandibular BBC is uncommon, the diagnosis would be less challenging if it is established by the correlation of the clinical examination, radiographic interpretation and histopathological analysis. Moreover, the clinical features of bilateral mandibular BBC summarized in this review could assist specialists to an accurate diagnosis and provide patients with optimal management.

Patient perspective

The parents mentioned that the surgery has an excellent impact on their son's life. Also, the outcomes met their expectations as the facial swellings disappeared.

Consent

Written informed consent was obtained from the patient's father for publication of this case report and accompanying images.

Data availability

All data underlying the results are available as part of the article and no additional source data are required.

Acknowledgements

This research was funded by the Deanship of Scientific Research at Princess Nourah bint Abdulrahman University through the Fast-track Research Funding Program.

References

1. Stoneman DW, Worth HM: **The mandibular infected buccal cyst--molar area.** *Dent Radiogr Photogr.* 1983; **56**(1): 1-14.
[PubMed Abstract](#)
2. Pompura JR, Sándor GK, Stoneman DW: **The buccal bifurcation cyst: a prospective study of treatment outcomes in 44 sites.** *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 1997; **83**(2): 215-221.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. White S, Pharoah M: **Oral Radiology: Principles and Interpretation.** 7th ed. 2013; 696.
[Reference Source](#)
4. Neville BW, Damm DD, Allen CW, et al.: **Oral and Maxillofacial Pathology.** 3rd Edition. 2008; 984.
[Reference Source](#)
5. David LA, Sándor GK, Stoneman DW: **The buccal bifurcation cyst: in non-surgical treatment an option?** *J Can Dent Assoc.* 1998; **64**(10): 712-717.
[PubMed Abstract](#)
6. Craig GT: **The paradental cyst. A specific inflammatory odontogenic cyst.** *Br Dent J.* 1976; **141**(1): 9-14.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Phillipsen HP, Reichart PA, Ogawa I, et al.: **The inflammatory paradental cyst: a critical review of 342 cases from a literature survey, including 17 new cases from the author's files.** *J oral Pathol Med.* 2004; **33**(3): 147-155.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Jones AV, Craig GT, Franklin CD: **Range and demographics of odontogenic cysts diagnosed in a UK population over a 30-year period.** *J oral Pathol Med.* 2006; **35**(8): 500-507.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Stanback JS: **The management of bilateral cysts of the mandible.** *Oral Surg Oral Med Oral Pathol.* 1970; **30**(5): 587-91.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Swerdloff M, Alexander SA, Ceen RF, et al.: **Bilateral mandibular dentigerous cysts in a seven-year-old child.** *J Pedod.* 1980; **5**(1): 77-84.
[PubMed Abstract](#)
11. Vedtofte P, Praetorius F: **The inflammatory paradental cyst.** *Oral Surg Oral Med Oral Pathol.* 1989; **68**(2): 182-188.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Packota GV, Hall JM, Lanigan DT, et al.: **Paradental cysts on mandibular first molars in children: report of five cases.** *Dentomaxillofac Radiol.* 1990; **19**(3): 126-132.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Bohay RN, Weinberg S, Thorner PS: **The paradental cyst of the mandibular permanent first molar: report of a bilateral case.** *ASDC J Dent Child.* 1992; **59**(5): 361-365.
[PubMed Abstract](#)
14. Martinez-Conde R, Aguirre JM, Pindborg JJ: **Paradental cyst of the second molar: Report of a bilateral case.** *J Oral Maxillofac Surg.* 1995; **53**(10): 1212-1214.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Shohat I, Buchner A, Taicher S: **Mandibular buccal bifurcation cyst: enucleation without extraction.** *Int J Oral Maxillofac Surg.* 2003; **32**(6): 610-613.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Gallego L, Baladrón J, Junquera L: **Bilateral mandibular infected buccal cyst: a new image.** *J Periodontol.* 2007; **78**(8): 1650-1654.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Corona-Rodriguez J, Torres-Labardini R, Velasco-Tizcareño M, et al.: **Bilateral buccal bifurcation cyst: case report and literature review.** *J oral Maxillofac Surg.* 2011; **69**(6): 1694-1696.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Ramos LMA, Vargas PA, Coletta RD, et al.: **Bilateral Buccal Bifurcation Cyst: Case Report and Literature Review.** *Head Neck Pathol.* 2012; **6**(4): 455-459.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Borgonovo AE, Reo P, Grossi GB, et al.: **Paradental cyst of the first molar:**

Report of a rare case with bilateral presentation and review of the literature. *J Indian Soc Pedod Prev Dent.* 2012; **30**(4): 343–348.
[PubMed Abstract](#) | [Publisher Full Text](#)

20. Boffano P, Gallesio C, Roccia F, *et al.*: **Bilateral buccal bifurcation cyst.** *J Craniofac Surg.* 2012; **23**(6): e643-e645.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Issler A, Bornert F, Clauss F, *et al.*: **Mandibular buccal bifurcation cyst**

treatment: Report of two cases and literature review. *Med Buccale Chir Buccale.* 2013; **19**(2): 77–84.
[Publisher Full Text](#)

22. Bautista CRG, Milhan NVM, Anka MDVEA, *et al.*: **Bilateral mandibular buccal bifurcation cyst: a case report emphasizing the role of imaging examination in the diagnosis.** *Autops Case Rep.* 2019; **9**(2): e2018073.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 11 February 2021

<https://doi.org/10.5256/f1000research.54072.r79130>

© 2021 Elsayed S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Shadia Abdel-Hameed Elsayed

¹ Faculty of Dental Medicine for Girls, Al Azhar University, Cairo, Egypt

² College of Dentistry, Taibah University, Medina, Saudi Arabia

Thank you for this interesting, well written case study.

- The table of cases from past literature is very informative.
- Some points need to be explained.
- Please add the prevalence rate of such unusual cases to the introduction with citation.
- To evaluate the lesion extent, was CT or CBCT performed prior to surgery? If so, please include a photograph of representing cut.
- Please also include an extraoral clinical photograph.
- Add details on the prescribed drugs and instructions for postoperative care.
- Please address previous citations regarding surgical management of this unusual cyst with complete tooth removal in the discussion.

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Partly

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Partly

Is the case presented with sufficient detail to be useful for other practitioners?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Oral & Maxillofacial Surgery

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 15 Feb 2021

Ashwag Aloyouny, College of Dentistry Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia

- Please add the prevalence rate of such unusual cases to the introduction with citation (Done)
- To evaluate the lesion extent, was CT or CBCT performed prior to surgery? If so, please include a photograph of representing cut.
No CT or CBCT were performed.
- Please also include an extraoral clinical photograph.
No extraoral photograph was taken
- Add details on the prescribed drugs and instructions for postoperative care. (Done)

Competing Interests: No competing interests

Reviewer Report 02 February 2021

<https://doi.org/10.5256/f1000research.54072.r77129>

© 2021 Tosios K. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Konstantinos I. Tosios

Department of Oral Medicine and Pathology, School of Dentistry, National and Kapodistrian University of Athens, Athens, Greece

This is a case of bilateral buccal bifurcation cysts that is unusual in that (a) it caused an extensive esthetic disfiguration leading to “social exclusion” and (b) necessitated extraction of two unerupted permanent molar teeth. However, as there are no clinical photographs, intra-oral or extra-oral, the first feature cannot be documented. Even more, the authors do not contemplate on

the reasons that dictated extractions: does the risk of recurrence (how common is it?) balance the loss of two permanent first molars? A more thorough discussion is necessary.
More specific comments:

Introduction

- The Introduction contains information (i.e. "Children between the age of 4 and 14 years are most commonly affected.") not supported by relevant references.

Clinical Presentation

- The swellings are described as "painful" by the patient's guardian. It would be interesting to know whether those swellings were, also, painful on palpation and/or showed pus excretion, or whether the authors noticed any orifice on the mucosa through which they could probe the cyst.
- The teeth are described as "unerupted incomplete permanent mandibular first molars" (Case presentation, paragraph 3). Do they mean partially/incompletely erupted, as they were clinically not evident but radiographically not surrounded by bone?
- The histopathologic description should be reorganized: "multiple pieces of cyst wall" are profoundly consistent with "cystic lesions", while it is the presence "of non-keratinized stratified squamous epithelium", not "bland looking spindle-shaped fibroblasts and degenerated bony spicules" that make the findings "consistent with cystic lesions". Fig. 3 is not of proper quality.

Discussion

- The Discussion contains information not supported by relevant references.
- The main aim of Discussion is for the authors to discuss their findings and offer (if possible) new insights. For instance, the authors correctly state that the inflammation caused by a deep periodontal pocket, or a defect in tooth eruption are possible reasons for the development of . Which of those factors possibly caused the lesions in their cases? How do they explain the bilaterality of the lesion?
- The authors should be very careful in what they state. For instance, in the 2nd paragraph they write:
  - "Clinically, bilateral BBC has few challenging similarities with dentigerous cyst, and paradental cyst." Only bilateral BCC has those similarities, not the usual unilateral variant?
  - "all previously mentioned cysts are commonly associated with a tooth". This is common for most cysts, inflammatory or developmental!
  - "paradental cysts and dentigerous cysts usually occur in adults in the mandibular third molar region" and later "paradental cyst... which commonly occurs in the lateral surface of a vital tooth usually the first and second mandibular molars of primary teeth." Paradental cyst usually develops distally to the third permanent molar.
- "Radiographic interpretation is essential to assist recognize and differentiate between different cystic lesions". Please correct syntax.
- "on the lateral crown". Wrong expression.

- “Every rule has an exception, it is challenging to differentiate between the circumferential dentigerous cyst and the BBC radiographically as both illustrated as radiolucent cysts involving a completely unerupted tooth”. The authors correctly include this variety of dentigerous cyst in the d.d. of the present case (why don’t they state it in the case presentation), but they should give more information not only for the sake of completeness of their work, but also for “tutorial” reasons. The “Every rule has an exception” does not fit in that case, while the whole phrase needs reconsideration.
- “Histopathologically, the cyst wall under the microscope shows chronic inflammatory lymphocytes and two to four cell layers thick of proliferating lining of nonkeratinized stratified squamous epithelium”. Repetition of well known information. It would more interesting to know if histopathologic examination can differentiate the cysts included in the d.d.
- “The diagnosis of BBC should be established based on the clinical, radiographical and microscopic characteristics. The clinical examination, radiographic interpretation and histopathologic analysis of both lesions displaying cystic origin with bilateral buccal expansion involving both mandibular first molars in a seven-year-old child associated with bilateral BBC was presented here. This reported case had similar clinical, radiographical and histopathological features of bilateral mandibular BBC. It is worth mentioning that the case presented here is very rare because it describes bilateral mandibular BBC. To the best of our knowledge, the bilateral characteristic of BBC is very rare; there are only few cases have been reported in the English-language literature.” This paragraph says the same thing many times. Please reconsider.

Is the background of the case’s history and progression described in sufficient detail?

Partly

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Partly

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

No

Is the case presented with sufficient detail to be useful for other practitioners?

No

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Oral Pathology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 22 January 2021

<https://doi.org/10.5256/f1000research.54072.r77849>

© 2021 Oliveira D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Denise Tostes Oliveira

Department of Surgery, Stomatology, Pathology and Radiology, Area of Pathology, Bauru School of Dentistry, University of São Paulo, Bauru, Brazil

There are no suggestions. The manuscript is ok.

Is the background of the case's history and progression described in sufficient detail?

No

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

No

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

No

Is the case presented with sufficient detail to be useful for other practitioners?

No

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Oral pathology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 18 January 2021

<https://doi.org/10.5256/f1000research.30969.r77339>

© 2021 Oliveira D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Denise Tostes Oliveira

Department of Surgery, Stomatology, Pathology and Radiology, Area of Pathology, Bauru School of Dentistry, University of São Paulo, Bauru, Brazil

This is an interesting and well-documented case report of bilateral buccal bifurcation cysts (BBC). It is important to include in the manuscript that the etiopathogenesis of BBC is the same of paradental cyst, according to World Health Organization Classification of Head and Neck Tumours (2017). In addition, the diagnosis should be established based in the clinical/radiographic and microscopic characteristics.

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Yes

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Yes

Is the case presented with sufficient detail to be useful for other practitioners?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Oral pathology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research