CASE REPORT

Case Report: Acute kidney failure leading to permanent haemodialysis due to hyperoxaluria following one-anastomosis gastric bypass-related rapid weight loss. [version 1; peer review: 1 approved, 1 approved with reservations]

Angelo Miranda¹, Andrea Rosato¹, Andrea Costanzi¹, Lucia Pisano², Sara Colzani², Sara Auricchio², Giulio Mari¹ , Pietro Achilli³ , Dario Maggioni¹

¹General Surgery Departement, Desio Hospital, Desio, Italy, 20843, Italy

²Nephrological Unit, Desio Hospital, Desio, Italy, 20843, Italy

³Univerity of Milan, Residency in general Surgery, Milano, Italy, 20100, Italy

V1 First published: 26 Feb 2020, 9:155
<https://doi.org/10.12688/f1000research.22109.1>
 Latest published: 29 May 2020, 9:155
<https://doi.org/10.12688/f1000research.22109.2>

Abstract

The one-anastomosis laparoscopic gastric bypass (OAGB) has been proven to provide good weight loss, comorbidity improvement, and quality of life with follow-up longer than five years. Although capable of improving many obesity-related diseases, OAGB is associated with post-operative medical complications mainly related to the induced malabsorption. A 52-year-old man affected by nephrotic syndrome due to a focal segmental glomerulosclerosis underwent OAGB uneventfully. At three months post-surgery, the patient had lost 40kg, reaching a BMI of 32. The patient was admitted to the nephrology unit for acute kidney injury with only mild improvement in renal function (Scr 9 mg/dl); proteinuria was still elevated (4g/24h), with microhaematuria. A renal biopsy was performed: oxalate deposits were demonstrated inside tubules, associated with acute and chronic tubular and interstitial damage and glomerulosclerosis (21/33 glomeruli). Urinary oxalate levels were found to be elevated (72mg/24h, range 13-40), providing the diagnosis of acute kidney injury due to hyperoxaluria, potentially associated to OAGB. No recovery in renal function was observed and the patient remained dialysis dependent. Early and rapid excessive weight loss in patients affected by chronic kidney insufficiency could be associated with the worsening of renal function. Increased calcium oxalate levels associated with OAGB-related malabsorption could be a key factor in kidney injury.

Keywords

OAGB, kidney failure, Calcium oxalate, weight loss

Open Peer Review

Approval Status

	1	2
version 2 (revision) 29 May 2020		 view
version 1 26 Feb 2020	 view	 view

1. **Matteo Uccelli**, Policlinico San Marco,
Zingonia-Osio Sotto, Italy

2. **Mohammad Kermansaravi** , Iran
University of Medical Sciences, Tehran, Iran

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Giulio Mari (giul_mari@yahoo.it)

Author roles: **Miranda A:** Conceptualization; **Rosato A:** Conceptualization; **Costanzi A:** Data Curation; **Pisano L:** Data Curation; **Colzani S:** Methodology; **Auricchio S:** Data Curation; **Mari G:** Supervision; **Achilli P:** Supervision; **Maggioni D:** Funding Acquisition

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Miranda A *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Miranda A, Rosato A, Costanzi A *et al.* **Case Report: Acute kidney failure leading to permanent haemodialysis due to hyperoxaluria following one-anastomosis gastric bypass-related rapid weight loss. [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2020, 9:155 <https://doi.org/10.12688/f1000research.22109.1>

First published: 26 Feb 2020, 9:155 <https://doi.org/10.12688/f1000research.22109.1>

Introduction

Bariatric surgery is known to be the most effective means of inducing durable weight loss in obese populations¹⁻³. In particular, bariatric malabsorptive procedures are effective metabolic therapies capable of improving many obesity-related diseases^{4,5}.

Among bariatric procedures claiming to be effective in providing a metabolic improvement in obese patients, the one-anastomosis laparoscopic gastric bypass (OAGB) has been proven to provide good weight loss, comorbidity improvement, and quality of life even with follow-up longer than five years⁶. Despite the positive reported results in terms of surgical outcomes and effectiveness in reducing several obesity-related diseases, OAGB is associated with post-operative medical complications, mainly related to the induced malabsorption⁷. Focusing on renal function, bariatric surgery is capable of being a protective factor against renal failure^{8,9}; however, the association between induced malabsorption and particular renal syndromes can significantly alter the metabolism of numerous serum metabolites, leading to direct kidney damage¹⁰. When reported, such alterations of the metabolic framework tend to appear a relatively long time after surgery and are described as a late complication of weight loss¹¹. However, rapid and early weight loss could aggravate and accelerate such a process.

Here, we report a rare case of acute renal failure occurring three months after laparoscopic OAGB caused by hyperoxaluria associated with rapid and early excessive weight loss.

Case presentation

A white Italian 52-year-old man employed at an insurance company underwent OAGB in February 2019 for stage III morbid obesity (135kg, BMI 45).

Past medical history was characterized by pathological obesity, type 2 diabetes, obstructive sleep apnoea syndrome and nephrotic syndrome due to focal segmental glomerulosclerosis. In 2011, at the time of the initial kidney biopsy, his weight was 110kg, his blood pressure and kidney function were normal, and proteinuria was 9g/24h. He was then treated with steroids and after five months cyclosporin was added, leading to complete remission of the nephrotic syndrome. He was also treated with renin-angiotensin-system blockers. In 2015, his weight was 125kg and his serum creatinine (SCr) and proteinuria began to rise, until 2018 when his weight was 134kg, SCr was 1.6mg/dl and proteinuria 4.8g/24h, without nephrotic syndrome. The patient had tried to follow a hypocaloric and hypoproteic diet with no success. The worsening of proteinuria was interpreted as secondary to obesity; in the meantime, steroids and cyclosporin were both contraindicated.

The patient was then referred to a surgeon for bariatric surgery. In February 2019, the surgical procedure was carried out uneventfully under general anaesthesia using the laparoscopic technique. The gastric sleeve was fashioned with a 60 mm stapler (six charges) and Seamguard. The operating surgeon was highly experienced in laparoscopic bariatric surgery, having performed more than five hundred procedures. The patient was discharged on day three after surgery after the methylene

blue test did not reveal any signs of staple line leak. Follow-up was done monthly for the first six months after surgery with outpatient visits. The patient's SCr was unvaried. In March 2019, the patient's body weight was 114.5 kg; his sCr was 2.12 mg/dl in March and 2.23 mg/dl in April 2019. In May 2019, his body weight was 95 Kg. His water and food intake were very poor; he had stopped taking vitamin and citrate supplements and his SCr had raised up to 16 mg/dl. The excess weight loss was 64% at two months; at three months after surgery, the patient had lost 40kg, reaching a BMI of 32.

The patient was admitted to our nephrology unit for acute kidney injury. His blood pressure was 140/80 mmHg, heart rate was 77 bpm and urine output was about 2000 ml/day. On admission, SCr was 16.6 mg/dl, urea was 235 mg/dl, sodium and potassium levels were normal despite metabolic acidosis (HCO_3^- 16 mmol/l), proteinuria was 3 g/day and mild microhaematuria was detected using a urine dipstick. Renal ultrasound was normal. The patient was given fluids (2 litres of polysaline solution intravenously for 10 days and an oral intake of water of 1.5 litres) with only mild improvement in renal function (SCr 8mg/dl). A new renal biopsy was performed; the puncture was ultrasound-guided using 16g × 20 mm Bard Monopty needles and two specimens were obtained for histologic examination. Oxalate deposits were demonstrated inside tubules, associated with acute and chronic tubular and interstitial damage and glomerulosclerosis (21/33 glomeruli). Urinary oxalate level was then found to be elevated (72mg/24h, range 13–40), providing the diagnosis of acute kidney injury due to hyperoxaluria potentially associated to OAGB.

In July 2019, renal replacement therapy was started. Online hemodiafiltration was administered through a central venous catheter and the treatment was performed three times a week for four hours without ultrafiltration. The patient was also treated with oral sodium bicarbonate (NaHCO_3 , 2 g/day), potassium and magnesium citrate (2.56 g/day), calcium carbonate (2.5g/day) and sevelamer (2.4 g/day). Renal function was assessed monthly by creatinine and urea clearance tests, but no recovery in renal function was observed and the patient remained dialysis dependent (creatinine clearance 11 ml/min, urea clearance 4 ml/min in January 2020). The patient's body weight is now steady at 84Kg.

Discussion

Kidney failure can complicate the post-operative course after bariatric surgery. Its aetiology can be multifactorial and complex in obese populations. Pre-existing factors such as diabetes, hypertension, and chronic kidney disease (CKD) can play a role in worsening the renal insult during the early post-operative period^{12,13}. In addition to dehydration, which is the main cause of acute kidney failure after bariatric surgery, the alteration of several metabolic assets induced by a rapid and consistent weight loss can become a key factor in triggering kidney failure¹⁴. One metabolite potentially involved in the acute onset of kidney failure but rarely associated to it, is calcium oxalate¹⁵.

The reasons for the onset of hyperoxaluria are not yet completely understood. There are various hypotheses: one of the

main reasons seems to be the malabsorption of lipids since, in this condition, the calcium found inside the intestinal lumen tends to bind to fatty acids instead of forming insoluble precipitates of calcium oxalate; the oxalate thus remains in a soluble state and can be reabsorbed by the ileal walls¹⁶. Another possible explanation is the increase in bile secretion into the colon due to the lack of intestinal reabsorption of bile salts, with consequent increase in the permeability of the colic walls due to a decrease in the function of the epithelial barrier. Finally, another hypothesis is the decrease in colonization of the colon by oxalate-metabolizing bacteria such as *Oxalobacter formigenes*, able to metabolize oxalate¹⁷.

The association between excessively rapid weight loss and renal damage due to calcium oxalate is not widely documented in the literature and therefore remains a hypothesis based on a valid rationale¹⁸.

Bariatric surgery is able to reduce the most dangerous risk factors for end stage renal disease such as hypertension and diabetes, but post-operative complications in CKD patients are slightly higher than the general population and surgeons must be aware of oxalate nephropathy, because it's a rare but often irreversible cause of acute kidney injury^{19,20}.

Patient with pre-existing chronic kidney disease may be considered to be higher risk for secondary oxalate nephropathy but this has not yet been established. Monitoring the 24-hour urinary oxalate excretion rate might be a useful tool to prevent oxalate nephropathy in high risk patients. For such patients, a careful selection of the right bariatric procedure (malabsorptive vs restrictive) should therefore be performed²¹.

Conclusions

The early and rapid excessive weight loss in patients affected by chronic kidney insufficiency could be associated with the worsening of renal function. Increased calcium oxalate levels associated with OAGB-related malabsorption could be a key factor in kidney injury.

Consent

Written informed consent for the publication of their clinical details and clinical images and video was obtained from the patient.

Data availability

Underlying data

All data underlying the results are available as part of the article and no additional source data are required.

Extended data

Zenodo: Acute kidney failure leading to permanent hemodialysis due to hyperoxaluria following OAGB related rapid weight loss. Case report. <https://doi.org/10.5281/zenodo.3609350>²²

This project contains the following extended data:

- Surgical procedure video (in MP4 file format)
- Creatinine levels pre-surgery (in JPG format)
- Creatinine levels post-surgery (in JPG format)

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

References

1. Schauer PR, Kashyap SR, Wolski K, *et al.*: **Bariatric surgery versus intensive medical therapy in obese patients with diabetes.** *N Engl J Med.* 2012; **366**(17): 1567–76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
2. Mingrone G, Panunzi S, De Gaetano A, *et al.*: **Bariatric surgery versus conventional medical therapy for type 2 diabetes.** *N Engl J Med.* 2012; **366**(17): 1577–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Dixon JB, O'Brien PE, Playfair J, *et al.*: **Adjustable gastric banding and conventional therapy for type 2 diabetes: a randomized controlled trial.** *JAMA.* 2008; **299**(3): 316–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Sugerman HJ, Wolfe LG, Sica DA, *et al.*: **Diabetes and hypertension in severe obesity and effects of gastric bypass-induced weight loss.** *Ann Surg.* 2003; **237**(6): 751–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Batsis JA, Romero-Corral A, Collazo-Clavell ML, *et al.*: **Effect of weight loss on predicted cardiovascular risk: change in cardiac risk after bariatric surgery.** *Obesity (Silver Spring).* 2007; **15**(3): 772–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Neuberger M, Blanchet MC, Gignoux B, *et al.*: **Long-Term Outcomes After One-Anastomosis Gastric Bypass (OAGB) in Morbidly Obese Patients.** *Obes Surg.* 2019.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Bhandari M, Nautiyal HK, Kosta S, *et al.*: **Comparison of one-anastomosis gastric bypass and Roux-en-Y gastric bypass for treatment of obesity: a 5-year study.** *Surg Obes Relat Dis.* 2019; **15**(12): 2038–2044.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Neff KJ, Baud G, Raverdy V, *et al.*: **Renal Function and Remission of Hypertension After Bariatric Surgery: a 5-Year Prospective Cohort Study.** *Obes Surg.* 2017; **27**(3): 613–619.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Alkandari A, Ashrafian H, Sathyapalan T, *et al.*: **Bariatric Surgery Modulates Urinary Levels of MicroRNAs Involved in the Regulation of Renal Function.** *Front Endocrinol (Lausanne).* 2019; **10**: 319.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Gero D, Raptis DA, Vleeschouwers W, *et al.*: **Defining Global Benchmarks in Bariatric Surgery: A Retrospective Multicenter Analysis of Minimally Invasive Roux-en-Y Gastric Bypass and Sleeve Gastrectomy.** *Ann Surg.* 2019; **270**(5): 859–867.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Thakar CV, Kharat V, Blanck S, *et al.*: **Acute kidney injury after gastric bypass surgery.** *Clin J Am Soc Nephrol.* 2007; **2**(3): 426–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Nor Hanipah Z, Panchai S, Augustin T, *et al.*: **Impact of Early Postbariatric Surgery Acute Kidney Injury on Long-Term Renal Function.** *Obes Surg.* 2018; **28**(11): 3580–3585.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Currie A, Chetwood A, Ahmed AR: **Bariatric surgery and renal function.** *Obes Surg.* 2011; **21**(4): 528–39.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Weingarten TN, Gurrieri C, McCaffrey JM, *et al.*: **Acute kidney injury following**

- bariatric surgery. *Obes Surg.* 2013; **23**(1): 64–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Liu H, Ye T, Yang X, *et al.*: **H19 promote calcium oxalate nephrocalcinosis-induced renal tubular epithelial cell injury via a ceRNA pathway.** *EBioMedicine.* 2019; **50**: 366–378.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Nelson WK, Houghton SG, Milliner DS, *et al.*: **Enteric hyperoxaluria, nephrolithiasis, and oxalate nephropathy: potentially serious and unappreciated complications of Roux-en-Y gastric bypass.** *Surg Obes Relat Dis.* 2005; **1**(5): 481–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Patel BN, Passman CM, Fernandez A, *et al.*: **Prevalence of hyperoxaluria after bariatric surgery.** *J Urol.* 2009; **181**(1): 161–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Nasr SH, D'Agati VD, Said SM, *et al.*: **Oxalate nephropathy complicating Roux-en-Y Gastric Bypass: an underrecognized cause of irreversible renal failure.** *Clin J Am Soc Nephrol.* 2008; **3**(6): 1676–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Chang AR, Grams ME, Navaneethan SD: **Bariatric Surgery and Kidney-Related Outcomes.** *Kidney Int Rep.* 2017; **2**(2): 261–270.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Chang AR, Chen Y, Still C, *et al.*: **Bariatric surgery is associated with improvement in kidney outcomes.** *Kidney Int.* 2016; **90**(1): 164–171.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Lumlertgul N, Siribamrungwong M, Jaber BL, *et al.*: **Secondary Oxalate Nephropathy: A Systematic Review.** *Kidney Int Rep.* 2018; **3**(6): 1363–1372.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Achilli P: **Acute kidney failure leading to permanent hemodialysis due to hyperoxaluria following OAGB related rapid weight loss. Case report.** 2020. <http://www.doi.org/10.5281/zenodo.3609350>

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 18 May 2020

<https://doi.org/10.5256/f1000research.24383.r63316>

© 2020 Kermansaravi M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Mohammad Kermansaravi

Minimally Invasive Surgery Research Center, Iran University of Medical Sciences, Tehran, Iran

It's a well written manuscript, but I have some suggestions:

1. In abstract and introduction, it's better write " one-anastomosis gastric bypass (OAGB)" instead of " one-anastomosis laparoscopic gastric bypass (OAGB)"
2. In case presentation, you mentioned that the patient underwent sleeve gastrectomy! Sleeve or OAGB? Please clarify the surgical technique.
3. There is not any strong evidence about rapid weight loss related nephropathy. How do you result in? Please clarify.
4. How do you justify the relation between nephropathy and OAGB? Rapid weight loss if imaginary has a relation with rapid weight loss, it could happen in any type of weight loss therapies.

Is the background of the case's history and progression described in sufficient detail?

Partly

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Partly

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Partly

Is the case presented with sufficient detail to be useful for other practitioners?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Bariatric surgery, metabolic surgery, dvanced laparoscopic surgery

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 26 May 2020

Giulio Mari, Desio Hospital, Desio, Italy

We corrected the OAGB definition as indicated.

We better described the OAGB procedure in the case presentation.

As requested by the reviewer we clarified the possible correlation between OAGB, rapid weight loss and kidney failure due to calcium oxalate.

The case report that we report describes a kidney failure following OAGB and rapid weight loss. The only change in the renal function across surgery was the calcium oxalate levels increase. We therefore make an hypothesis based on a pathophysiological rationale that will need further analysis to be proven.

Competing Interests: We disclose any competing interests

Reviewer Report 28 February 2020

<https://doi.org/10.5256/f1000research.24383.r60620>

© 2020 Uccelli M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Matteo Uccelli

Department of Surgery, S.I.C.OB. (Italian Society of Bariatric Surgery) Referral Center for Bariatric Surgery, San Donato Group, Policlinico San Marco, Zingonia-Osio Sotto, Italy

- The case report described here presents a useful suggestion in deciding which bariatric procedure should be chosen in nephropathic patients.
- The patient description is well explained and the discussion is well written as it establishes a sort of guide line that deserves to be taken into consideration.
- Kidney function in obese patients is a main issue and a main concern. Analyzing which kind of pathological disorder lays underneath kidney insufficiency should be done before

choosing the type of bariatric procedure.

References

1. Magalhães DSC, Pedro JMP, Souteiro PEB, Neves JS, et al.: Analyzing the Impact of Bariatric Surgery in Kidney Function: a 2-Year Observational Study. *Obes Surg.* **29** (1): 197-206 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Yes

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Yes

Is the case presented with sufficient detail to be useful for other practitioners?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: bariatric surgery

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research