Check for updates

RESEARCH ARTICLE

REVISED The benefit of cinnamon (*Cinnamomum burmannii*) in lowering total cholesterol levels after consumption of high-fat containing foods in white mice (*Mus musculus*) models [version 2; peer review: 2 approved]

Previously titled: Benefit of cinnamon (*Cinnamomum burmannii*) in lowering total cholesterol level after consumption of high-fat containing foods in white mice (*Mus musculus*) models

Annisa Pulungan¹, Yunita Sari Pane ²

¹Faculty of Medicine, Universitas Sumatera Utara, Medan, North Sumatera, Indonesia

²Department of Pharmacology, Faculty of Medicine, Universitas Sumatera Utara, Medan, North Sumatera, Indonesia

V2 First published: 06 Mar 2020, 9:168
<https://doi.org/10.12688/f1000research.22311.1>

Latest published: 29 May 2020, 9:168
<https://doi.org/10.12688/f1000research.22311.2>

Abstract

Background: Hypercholesterolemia is a condition where cholesterol levels in the body exceed the normal range. If the condition is longer, it can cause metabolic and cardiovascular diseases. The therapy of synthetic drugs has side effects that can be fatal (rhabdomyolysis). Needed to find natural remedies with minimal side effects. There are many nutritional components contained in cinnamon, such as cinnamaldehyde. The cinnamaldehyde, a substance that is thought to affect cholesterol levels. The study aims to determine the efficacy of *Cinnamomum burmannii* in lowering total cholesterol levels of mice (*Mus musculus*) given high-fat feed.

Methods: This is an experimental study with a pre-post control study design. The groupings were performed by a simple random sampling method. The male mice were divided into five groups (n=6/group): 1) negative control (aquadest); 2) positive control of high-fat containing food (HFC; quail yolk); 3) HFC + cinnamon extract (CE; dose 2mg/20g body weight (BW); 4) HFC + CE (dose 4mg/20gBW); 5) HFC + CE (dose 8mg/20gBW). The study was conducted for 28 days. Consumption of quail yolk as HFC to increasing cholesterol in mice. The intervention of CE started on day 15 and ended on day 28. Measurement of total cholesterol and BW of mice was performed on days 0, 14 and 28.

Results: The comparison of total cholesterol levels in the K1 group (120.3 ± 5.53 mg/dl) to K2 (107.3 ± 3.61 mg/dl), K3 (106.8 ± 4.57 mg/dl) and K4

(106.7 ± 0.51 mg/dl) showed decreased significantly different (p = 0.001) in groups consuming CE. However, there was not a significant

Open Peer Review

Approval Status

	1	2
version 2		
(revision)
29 May 2020
version 1
06 Mar 2020

1. **Tahereh Farkhondeh** , Birjand University of Medical Sciences, Birjand, Iran

2. **Mourad Akdad** , Faculty of Science and Technology Errachidia (FSTE), Errachidia, Morocco

Any reports and responses or comments on the article can be found at the end of the article.

change between groups in mice BW ($p = 0.419$).

Conclusions: The cinnamon (*Cinnamomum burmannii*) proved can be lowering of total cholesterol levels for 14 days in mice compared without given cinnamon after consumption of high-fat containing foods.

Keywords

hypercholesterolemia, cinnamon, high-fat feed, quail yolk, body weight of mice

This article is included in the **Agriculture, Food and Nutrition** gateway.

Corresponding author: Yunita Sari Pane (yunita@usu.ac.id)

Author roles: Pulungan A: Conceptualization, Formal Analysis, Investigation, Writing – Original Draft Preparation; Pane YS: Conceptualization, Formal Analysis, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Pulungan A and Pane YS. This is an open access article distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Pulungan A and Pane YS. **The benefit of cinnamon (*Cinnamomum burmannii*) in lowering total cholesterol levels after consumption of high-fat containing foods in white mice (*Mus musculus*) models [version 2; peer review: 2 approved]** F1000Research 2020, 9:168 <https://doi.org/10.12688/f1000research.22311.2>

First published: 06 Mar 2020, 9:168 <https://doi.org/10.12688/f1000research.22311.1>

REVISED Amendments from Version 1

In version 2 of the abstract manuscript, we revised in background by added more information about the needed to treat hyperlipidemia to prevented metabolic and CV diseases, and the importance of found natural remedies that might potentially be antihyperlipidemia agents with minimal side effects. The aims of study had confirmed.

The added information in methods about the used of male mice as research subjects to avoided bias caused by hormonal differences between male and female mice. The used of quail yolk as High Fat (HFC) food to increase cholesterol levels in mice. The quail yolk consumed was based on information from literature and preliminary studies. Details of the results with the addition of the mean value $\pm$ SD in the cholesterol level of each groups, but detailed information was not disclosed in the mice's body weight results because the results did not significantly difference. We have compiled new words to conclude this research for the better as part of the conclusion. We stated that the study was conducted for 28 days but we gave *Cinnamomum Burmani* for 14 days.

Any further responses from the reviewers can be found at the end of the article

Introduction

Hypercholesterolemia is a state where the cholesterol level in the body exceeds the normal range. Hypercholesterolemia can increase the risk of atherosclerosis, coronary artery disease, pancreatitis, diabetes mellitus, liver disease and renal disease, etc¹.

Data from the Indonesian Ministry of Health in 2019 reported that coronary artery disease to be the first killer, 26.4% of all deaths in Indonesia². Out of the 17 million the premature death (under the age of 70) due to non communicable disease, most cases are in low-and middle income countries, and 37% are caused coronary heart disease and 6.7 million of these were triggered by stroke (WHO, 2015)³.

Many side effects caused by the condition of hypercholesterolemia need good care besides improving lifestyle by reducing the intake of high fat foods, also taking drugs, such as statin drugs. Ezad *et al.* (2018) reported that the use of statin drugs can cause side effects such as myopathy, increase levels of liver enzymes, increase the risk of impairment or memory loss, and can even be fatal caused rhabdomyolysis. Therefore, it is important to find natural remedies that are efficacious in reducing cholesterol levels with minimal side effects⁴. In Indonesia, the development of traditional medicine is performed by exploring herbs known to be beneficial for health, and this information is passed from generation to generation. One such herb is *gambier* (*Uncaria gambir* Roxb.)⁵, which has antioxidant effects and lowers blood glucose when treating type 2 diabetes mellitus (T2DM). There are also *bangun-bangun* leaves (*Coleus amboinicus*), which are believed to be effective in relieving pain when distilled as ethanol and water extract⁶. Traditional medicine that comes from plants is commonly used by many people in Indonesia. One of these herbs is cinnamon (*Cinnamomum burmannii*). It has the efficacy to lower blood glucose levels. It is potential as an antidiabetic because it contains high levels of cinnamaldehyde. Cinnamon is often eaten

in daily food. Cinnamon is a native plant in Indonesia and can be found abundant in Central of Java, (Karanganyar), West Sumatra (Padang), Jambi (Kerinci), etc⁷.

The consuming high fat feed can increase cholesterol in the blood. Lipid levels of yolks of quail are higher than chicken egg yolks⁸. Preliminary studies have shown success in creating hypercholesterolemia mice models by consuming high-fat feed using quail yolk. Mice were used in this study, because they have biology similar to humans, and therefore can be a model for human hyperlipidemia⁹⁻¹¹. This study was conducted to determine the efficacy of *Cinnamomum burmannii* in lowering total cholesterol levels of mice (*Mus musculus*) given high-fat feed.

Methods

This is an experimental study in an animal model with a pre-post control study design. This study was conducted at the Pharmacology Laboratory of the Faculty of Medicine, Universitas Sumatera Utara, Indonesia.

Ethics

This study was approved by the Health Research Ethical Committee of Universitas Sumatera Utara (No:55/TGL/KEPK FK USU-RSUP HAM/2019).

Mice were given high-fat containing food (HFC) to create hyperlipidemic mouse model. The intervention was carried out by giving different CE doses to find the dose that reduces the total cholesterol level of the mice. Cholesterol total levels were obtained from the blood by cutting the mice's tail. To ensure the experimental animals remained in a comfortable condition, the mice were given anesthesia procedures before cutting off the tail^{12,13}.

Experimental animals

The animals were purchased from the Department Biology of Mathematics and Scientific Faculty USU. In total, 30 male white mice (*Mus musculus*), Swiss Webster strain, 10 to 12 weeks old, and weighing 25 – 40 g were used. Before conducting the study, the mice were adapted to their cages (plastic (30 × 20 × 10 cm) and covered with fine wire mesh; base of the cages was covered with rice husks as thick as 0.5 – 1 cm and replaced every day during the study) for 2 weeks before the experiment started. They had 12 hours of daylight (6:00 A.M. – 6:00 P.M.) and 12 hours of dark (6:00 P.M. – 6:00 A.M.). The mice were fed with standard feed (CP 551) from PT Charoen Pokphand-Indonesia and water was given *ad libitum*. Room temperature and humidity were kept at normal ranges. They were weighed once a week to avoid stress.

Allocation and treatment groups

After the adaptation period, the mice were randomly divided into five groups, with each group consisting of 6 mice.

The sample size was calculated according to Federer's formula¹⁴:

$$(t-1)(n-1) > 15$$

t = the number of groups

n = the number of samples

All 30 mice were given a number (1–30) using a SPIDOL marker pen, then randomized by putting the numbers in an envelope and dividing them into 5 groups according to the numbers are taken from the envelope⁹.

Groups were as follows: 1) K0, negative control group/placebo, no treatment or high-fat food (only given aquadest 0.5 ml); 2) K1, positive control group, diet of HFC (quail yolk); 3) K2, HFC + cinnamon extract (CE) dose 2mg/20g body weight (BW); 4) K3, HFC + CE dose 4mg/20gBW; 5) K4, HFC + CE dose 8mg/20gBW.

The study was conducted for 28 days.

Cinnamon extract. CE was given to the mice every morning at 8:00 am. Animals were kept during research in the mice laboratory in the Pharmacology laboratory. Before being given CE, it was dissolved with aquadest. The extract was given to the mice orally using feeding tubes. CE interventions began on the 15th day until the 28th day.

Cinnamon extract (Cinnamomum burmannii) was obtained as Herbilogy Cinnamon Extract Powder® (PT.Phytochemindo Reksa, Bogor, Indonesia; batch no. 033CT).

High-fat containing food. The quail yolk was used to induce hypercholesterolemia, because the concentration of the lipid in quail yolk is higher than chicken egg yolk⁸. A dosage of 0.5ml/day was given until day 28. The administration of quail yolk was done orally using feeding tubes as per the administration of CE.

Determination of dose of cinnamon. Based on research by Vanessa *et al.* in 2013, who saw a decrease in total blood cholesterol levels in white rats (*Rattus norvegicus*) by administering instant cinnamon powder drink (*Cinnamomum burmannii* BI.) at a dose of 14.4 mg and 43.2 mg for 7 days. In their research, it was found that a dose of 14.4 mg can reduce cholesterol levels in rats. Therefore we used a dose of 14.4 mg, because it was effective and efficient¹⁵. We converted its dose to mice and increased it with variation doses of 2, 4 and 8 mg. The value converted from 200g rat to 20g mice is 0.14.

Rat dose = 14.4 mg

Conversion to mice = $0.14 \times 14.4 \text{ mg} = 2.01 \text{ mg}$

~2 mg/20 gBW mice

Laboratory analysis

Measurement of total cholesterol and BW of mice was performed on days 0, 14 and 28 (every two weeks). Anesthesia ketamine/

xylazine 0.1 ml was injected intraperitoneal before the mice's tail cutting off^{12,13}.

Mice tails were cut 2 – 5mm to draw blood, which was then assessed for total cholesterol using the digital *autocheck*® cholesterol measurement tool. Mice were weighed using a digital scale.

Statistical analysis

Data were analyzed using SPSS 24. The number average of sample data presented as mean $\pm$ SD. The one-way ANOVA statistical analysis to indicate the effects of treatments for all group. If the result is significant ($p < 0.05$), then bootstrapping was performed using *post-hoc* Bonferroni for differentiating between each group.

Results

Figure 1 shows that groups K0 (negative controls), K1 (positive controls), K2 (CE 2 mg/20gBW), K3 (CE 4 mg/20gBW), and K4 (CE 8 mg/20gBW), had a decrease in total cholesterol levels. There is a significant difference in total cholesterol among the groups ($p=0.001$ between groups). A post-hoc Bonferroni test was performed to see the difference in total cholesterol averages within each group.

It was seen that there was a difference in average total cholesterol levels between K0 ($107.5 \pm 1.87 \text{ mg/dl}$) vs K1 ($120.3 \pm 5.53 \text{ mg/dl}$), ($p = 0.001$). This shows that in the K1 group (positive control) given quail yolk succeeded in increasing cholesterol in mice with a significant difference compared to the K0 group (negative control), which was only given aquadest. Besides, differences in total cholesterol levels in the K1 group ($120.3 \pm 5.53 \text{ mg/dl}$) compared with K2 ($107.3 \pm 3.61 \text{ mg/dl}$), K3 ($106.8 \pm 4.57 \text{ mg/dl}$) and K4 ($106.7 \pm 0.51 \text{ mg/dl}$) groups showed that a significant decrease in total cholesterol levels ($p = 0.001$). This proves that CE is efficacious in reducing total cholesterol levels, while in groups K2, K3, and K4, which were all given CE, there was not a significant difference between total cholesterol levels ($p > 0.05$).

Figure 2 shows that there were increases in BW in all groups. The largest increase of BW was in group K2, which was the positive control group who were given HFC quail yolk at 0.5 ml/20gBW. The smallest increase in BW was found in group K4, which was the group provided with HFC quail yolk and CE with the dose of 8 mg/20gBW (highest dose in this study). One-way ANOVA showed that there was no significant difference in BW between groups ($p = 0.419$), which could be inferred that the action of giving CE gave no effect increasing BW in mice.

Figure 1. Total cholesterol per treatment group before (day 14) and after (day 28) treatment. K0, negative control group, no treatment or high-fat food (aquadest); K1, positive control group, diet of high-fat containing food (HFC; quail yolk); K2, HFC + cinnamon extract (CE) dose 2mg/20g body weight (BW); K3, HFC + CE dose 4mg/20gBW; K4, HFC + CE dose 8mg/20gBW.

Figure 2. Total body weight per treatment group before (in day 14) and after (in day 28) treatment. K0, negative control group, no treatment or high-fat food (aquadest); K1, positive control group, diet of high-fat containing food (HFC; quail yolk); K2, HFC + cinnamon extract (CE) dose 2mg/20g body weight (BW); K3, HFC + CE dose 4mg/20gBW; K4, HFC + CE dose 8mg/20gBW. *Mean delta = (average BW at the end of the study) - (average BW at the beginning of the study).

Discussion

The aim of study to investigate cholesterol levels can be lowering in mice given variation of doses CE after consumption of quail yolk for a high-fat diet. Mice was used in this study because they have a similar biology to human, and can therefore be a model for human hyperlipidemia⁹⁻¹¹. In this study, it was proven that consumption of 0.5ml quail yolk for 28 days increased the total cholesterol level between the negative and positive control groups (*post-hoc* Bonferroni K0 vs K1; $p=0.001$).

There was a significant difference in the total decrease in cholesterol after treatment among all groups in this research.

However, this is not consistent with Vanessa *et al.* (2013) who stated that there was a decrease, even though the difference is not statistically significant. Their studies were only done for 14 days, with an increase of 3x the initial dose (14.4 mg to 43.2 mg), while our study was carried out 28 days, leading to differing results¹⁵.

Cholesterol is formed by the action of HMG-CoA reductase enzyme (3-hydroxy-3-methylglutaryl-CoA). If hypercholesterolemia is left without implementing proper diet or treatment, it can cause occlusion in a blood vessel. Hypercholesterolemia treated to medicines, such as statin could be given¹⁶. The use of

statins will competitively block HMG-CoA reductase and efficiently reduce serum LDL cholesterol. But, treatment using statin can cause rhabdomyolysis⁴. Thus, it needed research on traditional or herbal medicine, e.g. CE, that needs developed because CE is predicted to have minimal adverse effects.

Cinnamon (*Cinnamomum burmannii*) has cinnamaldehyde as it's biggest compound. Cinnamaldehyde, a phenolic component abundantly found in *Cinnamomum*⁷. Bandara *et al.* (2011) stated that cinnamon had the ability to be an antioxidant, antiviral, antifungal, antimicrobe, antitumor, and can lower cholesterol and blood pressure¹⁷. Cinnamon is believed to have a direct role in lipid metabolism to prevent hypercholesterolemia and hypertriglyceridemia, as well as in preventing free fatty acids with its strong lipolytic activity¹⁸. In the present study we believe that the increase of cholesterol levels due to quail yolk could be decreased with cinnamon maybe caused by its ability to block HMG-CoA reductase enzyme and suppress lipid peroxidation through increased antioxidant enzyme activity¹⁹. Abeysekera *et al.* (2017) reported cinnamaldehyde that highest compound in bark extracts of Ceylon cinnamon possess moderate cholesterol esterase and cholesterol micellization inhibition and bile acid binding in vitro. It could lower cholesterol levels because it contributed to bile acid synthesis²⁰.

Our study showed that there was no difference in BW of the mice between groups ($p = 0.419$). This is not consistent with Vafa *et al.* (2012) stated that consuming 3 grams of cinnamon for 8 weeks could decrease some biochemical and anthropometric variables compared to previous states significantly, i.e. a decrease of 1.19% body weight, 1.54% body mass index and 1.36% body fat. They conducted clinical study in T2DM patients who were given cinnamon supplements (*Cinnamomum zeylanicum*). In addition to lowering BW variables, it could also decrease fasting blood glucose level by about 9.2%, and 6.12% of HbA1c and 15.38% of triglyceride levels²¹. In contrary, Alsoodeeri *et al.* (2020) reported weight gain in rats

(Albino rats) which are given *Cinnamomum cassia* after consuming high fat compared to body weight at the beginning until end of the study in each groups¹⁸.

Leaf and Antonio (2017) stated that the higher the food intake (with Fat Mass/Fat Free Mass), the higher the increase in BW²². In the present study, in addition to standard food intake, giving hypercholesterol and interventional food should increase BW, because the mice had increased energy intake. The amount of food intake will affect the amount of energy intake, which will be saved as fat and impact on the mice's BWs, since energy intake is inversely proportional to physical activity. A high-fat diet group has low sensitivity to leptin, which results in increased appetite and food intake, thus increases the BW¹⁶. However, this was not the case in our study.

Conclusion

In the present study showed cinnamon (*Cinnamomum burmannii*) extracts proved can be lowering of total cholesterol level for 14 days after consuming high-fat containing foods in mice models compared to does not given cinnamon.

Data availability

Underlying data

Figshare: data Analysis-cholesterol and bodyweight of mice.docx, <https://doi.org/10.6084/m9.figshare.11901174.v23>

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

Acknowledgments

The author gives appreciation and thanks to the Pharmacology Laboratory of the Faculty of Medicine, Universitas Sumatera Utara for allowing the author to use the facility for collecting data in this research. In addition, thanks to all lecturers and examiners who had given advice and suggestions to the author, and all other parties who had contributed to this research.

References

1. Stapleton PA, Goodwill AG, James ME, *et al.*: **Hypercholesterolemia and microvascular dysfunction: interventional strategies.** *J Inflamm (Lond)*. 2010; 7: 54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
2. Kemenkes RI: **Hari Jantung Sedunia (World Heart Day): Your Heart is Our Heart Too.** 2019.
[Reference Source](#)
3. WHO (World Health Organization), 2015.
[Reference Source](#)
4. Ezad S, Cheema H, Collins N: **Statin-induced rhabdomyolysis: a complication of a commonly overlooked drug interaction.** *Oxf Med Case Reports*. 2018; 2018(3): omx104.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Pane YS, Ganie RA, Lindarto D, *et al.*: **The Effect of Gambier Extract on The Levels of Malondialdehyde, Superoxide Dismutase, and Blood Glucose in Type 2 Diabetes Mellitus Patients.** *Asian J Pharm Clin Res*. 2018; 11(10): 121–124.
[Publisher Full Text](#)
6. Pane YS, Sufitni S, Lumongga, F, *et al.*: **The Effectiveness of Coleus Amboinicus Leaf Extracts as an Analgetic Activity on Mice Exposed to Acetic acid.** *Asian J Pharm Clin Res*. 2018; 11(9): 301–4.
[Publisher Full Text](#)
7. Plumeriastuti H, Budiastut, Effendi MH, *et al.*: **Identification of bioactive compound of the essential oils of *Cinnamomum burmannii* from several areas in Indonesia by gas chromatography–mass spectrometry method for antidiabetic potential.** *Natl J Physiol Pharm Pharmacol*. 2019; 9(4): 279–283.
[Publisher Full Text](#)
8. Ukachukwu UG, Ozougwu VEO, Nwankwo NE: **A Comparative Study on the Total Cholesterol, Triacylglycerides and Lipid Concentrations of Quail and Chicken Eggs.** *International Journal of Research in Pharmacy and Biosciences*. 2017; 4(10): 11–16.
[Reference Source](#)
9. Russell WMS, Burch RL: **The Principles of Humane Experimental Technique.** London: Methuen & Co.Ltd. (Reissued: 1992, Universities Federation for Animal Welfare, Herts, England.) 1959.
[Reference Source](#)

10. Fitzpatrick A: **Ethics and animal research.** *J Lab Clin Med.* 2003; **141**(2): 89–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Festing MF: **Principles: the need for better experimental design.** *Trends Pharmacol Sci.* 2003; **24**: 341–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. **Vertebrate Animal Research.** The University of Iowa.
[Reference Source](#)
13. Stewart K, Schroeder VA: **Anesthesia Induction and Maintenance.** *Basic Biology.*
[Reference Source](#)
14. Federer WY: **Experimental Design, Theory and Application.** New York: Mac. Millan. hal. 1963; 544.
[Reference Source](#)
15. Vanessa R, Purwijantiningih LME, Aida Y: **Pemanfaatan Minuman Serbuk Kayu Manis (*Cinnamomum burmanii*.Bl.) Untuk Menurunkan Kadar Kolesterol Total Darah Pada Tikus Putih.** 2014.
[Reference Source](#)
16. Cherg S, Young J, Ma H: **HMG-CoA reductase (3-hydroxy-3-methyl-glutaryl-CoA reductase) (HMGR).** *The Journal of American Science.* 2008; **4**(3): ISSN 1545-1003.
[Reference Source](#)
17. Bandara T, Uluwaduge I, Jansz ER: **Bioactivity of Cinnamon with Special Emphasis on Diabetes Mellitus: A review.** *Int J Food Sci Nutr.* 2012; **63**(3): 380–386.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Alsoodeeri FN, Alqabbani HM, Aldossari NM: **Effects of Cinnamon (*Cinnamomum cassia*) Consumption on Serum Lipid Profiles in Albino Rats.** *J Lipids.* 2020; 2020: 8469830.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Lee JS, Jeon SM, Park EM, *et al.*: **Cinnamate Supplementation Enhances Hepatic Lipid Metabolism and Anti Oxidant Defense Systems In High Cholesterol Fed Rat.** *J Med Food.* 2003; **6**(3): 183–191.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. Abeysekera WPKM, Arachchige SPG, Ratnasooriya WD: **Bark Extracts of Ceylon Cinnamon Possess Antilipidemic Activities and Bind Bile Acids *In Vitro*.** *Evid Based Complement Alternat Med.* 2017; 2017: 7347219.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Vafa M, Mohammadi F, Shidfar F, *et al.*: **Effects of cinnamon consumption on glycemic status, lipid profile and body composition in type 2 diabetic patients.** *Int J Prev Med.* 2012; **3**(8): 531–6.
[PubMed Abstract](#) | [Free Full Text](#)
22. Leaf A, Antonio J: **The Effects of Overfeeding on Body Composition: The Role of Macronutrient Composition - A Narrative Review.** *Int J Exerc Sci.* 2017; **10**(8): 1275–1296.
[PubMed Abstract](#) | [Free Full Text](#)
23. Pulungan A, Pane YS: **data Analysis-cholesterol and bodyweight of mice.docx.** *figshare, Dataset.* 2020.
<http://www.doi.org/10.6084/m9.figshare.11901174.v2>

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 12 June 2020

<https://doi.org/10.5256/f1000research.26537.r64105>

© 2020 Farkhondeh T. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Tahereh Farkhondeh

Cardiovascular Diseases Research Center, Birjand University of Medical Sciences, Birjand, Iran

It is acceptable.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: metabolic disorder

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 14 April 2020

<https://doi.org/10.5256/f1000research.24610.r61729>

© 2020 Akdad M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Mourad Akdad

Department of Biology, Faculty of Science and Technology Errachidia (FSTE), Errachidia, Morocco

This study aimed to evaluate the effect of cinnamon extract (C. E) on cholesterol levels in mice with hypercholesterolemia. Authors demonstrated that C. E is useful to decrease cholesterol level, so its a promising plant to explore in the treatment of hypercholesterolemia and the related illness (hyperlipidemia, diabetes, atherosclerosis...).

1. Cinnamon has been shown to reduce total cholesterol, low-density lipoprotein and triglyceride levels, as well as increasing high-density lipoprotein levels. » precise the model of the study.
2. In introduction section, you must conclude by the aim of your research, this phrase "Mice were used in this study because of they have a similar biology to humans, and can therefore be a model for human hyperlipidemia⁸⁻¹⁰. must be include in discussion.
3. Avoid repetition such as: "This study was conducted at the Pharmacology Laboratory of the Faculty of Medicine, Universitas Sumatera Utara, Indonesia. » « The experiments took place at the Laboratory of Pharmacology Universitas Sumatera Utara (USU). »
4. What is the type of extract (aqueuous, ethanolic.....), and the part of plant by which industry do extraction.
5. Try to explain why you don't have a dose-dependent effect in your experiment.
6. « Figure 1 shows that groups K0 (negative controls), K1 (positive controls), K2 (CE 2 mg/20gBW), K3 (CE 4 mg/20gBW), and K4 (CE 8 mg/20gBW), had a decrease in total cholesterol levels. » why all groups have a decrease of Total Cholesterol, you don't precise the days of this results you must clarify the idea.
7. In the section results, days of measurement must be righted in the description to clarify the idea.
8. The smallest increase in BW was found in group K5 », K5 ??, you would like to say K4?
9. In the present study we believe that the increase of cholesterol levels due to quail egg yolk could be decreased with cinnamon by its ability to block HMG-CoA reductase enzyme and suppress lipid peroxidation through increased antioxidant enzyme activity¹⁸. » the mechanism involved in the antihypercholesterolemic effect must be declared with precaution, 'may be'.
10. Our study showed that there was no difference in BW of the mice between groups ($p = 0.419$). This is not consistent with Vafa *et al.* (2012)¹ who stated that consuming 3 grams of cinnamon for 8 weeks could decrease some biochemical and anthropometric variables compared to previous states significantly, i.e. a decrease of 1.19% body weight, 1.54% body mass index and 1.36% body fat. In addition to lowering BW variables, it could also decrease fasting blood glucose level by about 9.2%, and 6.12% of HbA1c and 15.38% of triglyceride levels²⁰. » model used by the study cited must be declared.
11. In the present study, a high-fat containing food used in combination with cinnamon extract (*Cinnamomum burmannii*) for 28 days could decrease total cholesterol levels in mice. » this conclusion must be reformulated, high-fat containing food was used only to have the model for studying the effect of CE on cholesterol level, so the conclusion must be related only to CE.

References

1. Vafa M, Mohammadi F, Shidfar F, Sormaghi MS, et al.: Effects of cinnamon consumption on glycemic status, lipid profile and body composition in type 2 diabetic patients. *Int J Prev Med.* 2012; **3** (8): 531-6 [PubMed Abstract](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Ethnopharmacology, medicinal plants, diabetes, hypertension

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 02 April 2020

<https://doi.org/10.5256/f1000research.24610.r61018>

© 2020 Farkhondeh T. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Tahereh Farkhondeh

Cardiovascular Diseases Research Center, Birjand University of Medical Sciences, Birjand, Iran

The present study investigated the effect of cinnamon extract on body weight and cholesterol levels of mice-fed HFD. It is a good idea.

Some revisions should be done:

Abstract

1. In the method section several point should be cleared:
 1. Type of cinnamon extract.
 2. Female or male mice.

Introduction

1. The aim of study should be better mentioned. it was not clear the question about the effect of plant on the lipid profile. In the past paragraph, the author described the lipid lowering effect of extract. Did this evidence extracted from traditional medicine reports?

Method

1. Did you used xylazine with ketamine for analgesic effect It should be mentioned. Animal ethics did not accept this anesthesia procedure.
2. Please more about cholesterol measurement (technique, instrument, kit).

Conclusion

1. If you tested the quail egg yolks as a high-fat diet that induces hypercholesterolemia. It is necessary in the abstract and introduction in the aim section.
2. In the first paragraph, you should about your findings without analytic indicator such as p-value.
3. You did not measure the changes in TG or FFA therefore, you should not concluded about in the conclusion section.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: metabolic disorder

I confirm that I have read this submission and believe that I have an appropriate level of

expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 06 Apr 2020

yunita sari pane, universitas sumatera utara, Indonesia

RESPONSE to review's **Tahereh Farkhondeh**

Cardiovascular Diseases Research Center, Birjand University of Medical Sciences, Birjand, Iran

The present study investigated the effect of cinnamon extract on body weight and cholesterol levels of mice-fed HFD. It is a good idea.

Some revisions should be done:

Abstract

1. In the method section several points should be cleared:

1. Type of cinnamon extract.

Answer:

Type of cinnamon extract used in this study had been explained in part of METHODS (page 3 manuscript in F1000research).

We used *Cinnamomum burmannii* in powder extract.

2. Female or male mice.

Answer:

The sample animal were male mice. This is one of the inclusion criteria of this study. Its mean to homogenized (make smaller of the differences in each samples) for prevent BIAS. As we known, there are differences between male and female will increasing number of lipid and dysfunction metabolic, including of hyperlipidemia. It related to level of hormones.

Introduction

1. The aim of study should be better mentioned.

Answer:

Thank you for your comment, we would confirm our statement on the aim of this study.

The aim of the study to determine *cinnamomum burmannii* effective in reducing total cholesterol levels of mice (*Mus musculus*) given high fat feed.

It was not clear the question about the effect of plant on the lipid profile.

Answer:

We had explained about the effect of plant on the lipid profile related to few references. Please see in part of DISCUSSION (page 5 manuscript in F1000research).

Cinnamon is believed to have a direct role in lipid metabolism and preventing hypercholesterolemia and hypertriglyceridemia, as well as in preventing free fatty acids with its strong lipolytic activity¹⁷.

In the present study we believe that the increase of cholesterol levels due to quail egg yolk could be decreased with cinnamon by its ability to block HMG-CoA reductase enzyme and

suppress lipid peroxidation through increased antioxidant enzyme activity¹⁸. In research by Pai *et al.* (2013), it was stated that cinnamaldehyde could lower cholesterol and triglyceride levels by the action of some enzymes secreted in certain amounts, which might contribute to bile acid synthesis¹⁹.

(This study needed continuous some research in the future to measure number of cinnamaldehyde in Cinnamon Extract Powder[®] per 1 mg CE used. Then we'll do new research to prove how CE mechanism as anti hypercholesterolemia, Is it blocked by HMG-CoA reductase enzyme and suppress lipid peroxidation through increased antioxidant enzyme activity or dysfunction of bile acid synthesis or both mechanism) ?

In the past paragraph, the author described the lipid lowering effect of extract. Did this evidence extracted from traditional medicine reports?

Answer:

Yes, the explain about "this has been suggested as a substance.....triglyceride levels" better moves to part of DISCUSSION, because we took from one of reference. We will add inform about the quail egg yolks as a high-fat diet that induces hypercholesterolemia in part of ABSTRACT.

We can not add inform in the AIM of study. Because, if we added in the Aims of study, it was not consequence to the title of study. If we changed title "Benefit of cinnamon (*Cinnamomum burmannii*) in lowering total cholesterol level after consumption of the quail egg yolk as a high-fat diet that induces hypercholesterolemia". It will make the title very long and not in accordance to the rules of journals F1000Research.

Method

1. Did you used xylazine with ketamine for analgesic effect It should be mentioned. Animal ethics did not accept this anesthesia procedure.

Answer:

Yes, we did. We corrected uncomplete in methods ketamin with xylazine. We choosed of **Injectable Anesthetic Agents, we used** Ketamine/xylazine* We follow to GUIDELINES - PREPARATION OF KETAMINE/XYLAZINE COCKTAIL FOR MICE. Dose Ketamine 80-100 mg/kg intra peritoneal (IP) and Xylazine 10-12.5 mg/kg IP.(reference 11. Vertebrate Animal Research. The University of Iowa).

The administration of effective anaesthetic agents to avoid impact/effects of the experimental procedures when we did cut off mice tail to collect blood sample with anesthesia to make minimize pain and prevent distress caused of pain in mice.

1. Please more about cholesterol measurement (technique, instrument, kit).

ANSWER:

In Laboratory analysis study showed how was technique to take blood from tail
*correction in methods study cut of mice tail.

"Mice tails were cut 2 – 5 mm to draw blood, which was then assessed for total cholesterol using the digital autocheck[®](merck of digital cholesterol measured tool). We used stick

cholesterol to measured total cholesterol of mice.

Regarded to follow **National Centre for the Replacement, Refinement & Reduction (NC3Rs) of Animals in Research** to measured (page 4 manuscript in F1000research).

Conclusion

1. If you tested the quail egg yolks as a high-fat diet that induces hypercholesterolemia. It is necessary in the abstract and introduction in the aim section.

Answer:

Yes you right. We had tried to add in abstract before, but it was more than 250 words, it was not allow with Journal F1000research rules. But we will try again to put in your comment in abstract.

1. In the first paragraph, you should about your findings without analytic indicator such as p-value.

Answer:

Im sorry, I can't catch what you mean in this statement. Would you like to describe more detail?

1. You did not measure the changes in TG or FFA therefore, you should not concluded about in the conclusion section.

Answer:

Yes, thank you for your comment. We are agree with you.

Conclusion:

The cinnamon extract (*Cinnamomum burmannii*) proved it can be decrease of total cholesterol levels of for 14 days in mice consuming high-fat containing compared to group without given CE.

Competing Interests: No competing interests were disclosed

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research