REVIEW

Oral lichen planus: a novel staging and algorithmic approach and all that is essential to know [version 1; peer review: 4 approved]

Eran Shavit ¹⁻³, Hagen Klieb^{4,5}, Neil H. Shear^{1,6,7}

¹Division of Dermatology, Department of Medicine, Sunnybrook Health Sciences Centre, University of Toronto, Toronto, ON, Canada

²Dermatology Unit, Barzilai University Medical Center, Ashkelon, Israel

³Faculty of Health Sciences, Ben-Gurion University of the Negev, Beer-Sheba, Israel

⁴Department of Dentistry, Sunnybrook Health Sciences Centre, University of Toronto, Toronto, ON, Canada

⁵Department of Anatomic Pathology, Sunnybrook Health Sciences Centre, University of Toronto, Toronto, ON, Canada

⁶Department of Medicine (Dermatology, Clinical Pharmacology, and Toxicology), Sunnybrook Health Sciences Centre, University of Toronto, Toronto, Ontario, Canada

⁷Department of Pharmacology, Sunnybrook Health Sciences Centre, University of Toronto, Toronto, ON, Canada

V1 First published: 24 Mar 2020, 9(F1000 Faculty Rev):206
<https://doi.org/10.12688/f1000research.18713.1>

Latest published: 24 Mar 2020, 9(F1000 Faculty Rev):206
<https://doi.org/10.12688/f1000research.18713.1>

Abstract

Lichen planus (LP) is a chronic autoimmune disease. Oral lesions may occur in isolation or in combination with other affected mucocutaneous sites. Oral LP (OLP) may present as one of the various manifestations of LP or may present as a disease sui generis with a broad range of severity. Despite this disease targeting the oral mucosa, its great impact on quality of life is underrated. In this article, we provide an updated review of the pathophysiology and epidemiology of OLP and offer guidance for its diagnosis and management. We also propose an algorithmic approach to the clinical forms of OLP and a novel staging system to facilitate management decisions.

Keywords

lichen planus, oral lichen planus, erosive oral lichen planus, atrophic oral lichen planus

Open Peer Review

Approval Status

	1	2	3	4
version 1 24 Mar 2020				

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **José López-López**, University of Barcelona, Barcelona, Spain
2. **Maria Jose Garcia-Pola Vallejo**, Universidad de Oviedo, Oviedo, Spain
3. **Andrés Blanco-Carrión**, University of Santiago de Compostela, Santiago de Compostela, Spain

---

4. Kobkan Thongprasom, Chulalongkorn

University, Bangkok, Thailand

Any comments on the article can be found at the end of the article.

Corresponding author: Eran Shavit (erans29@gmail.com)

Author roles: Shavit E: Writing – Original Draft Preparation, Writing – Review & Editing; Klieb H: Validation, Writing – Review & Editing; Shear NH: Conceptualization, Supervision, Validation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Shavit E *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Shavit E, Klieb H and Shear NH. **Oral lichen planus: a novel staging and algorithmic approach and all that is essential to know [version 1; peer review: 4 approved]** F1000Research 2020, 9(F1000 Faculty Rev):206 <https://doi.org/10.12688/f1000research.18713.1>

First published: 24 Mar 2020, 9(F1000 Faculty Rev):206 <https://doi.org/10.12688/f1000research.18713.1>

Introduction

Lichen planus (LP) is a chronic inflammatory disease that affects the skin, hair, nails, and mucous membrane including the oral cavity. It usually presents with recurrences and periods of clinical exacerbations and remissions; however, it may be a transient disorder¹. Oral LP (OLP) is a common disease that afflicts 1–2% of the population, most commonly middle-aged females, and is persistent². A chronic course is typical, and spontaneous resolution is uncommon^{3,4}. OLP may take a progressing course and therefore require lifetime surveillance. Although LP is managed medically by dermatologists, OLP is mostly encountered and managed by oral surgeons. The aim of this review was to summarize the essentials required to understand OLP to facilitate its diagnosis and management and to encourage a multidisciplinary approach.

Our literature search was performed using PubMed with keywords such as “oral lichen planus” and/or “oral lichen planus” with other keywords to extend the search such as “etiology”, “epidemiology”, “diagnosis”, “management”, and “therapy”. Excluded from our review were articles that did not specifically adhere to OLP, such as oral lichenoid reactions, or articles that include other nonspecific stomatitis indicated in the article title, and articles published in languages other than English. Generally, no restrictions were made on any article type, but clinical trials and reviews were our primary focus. Additionally, some individual articles retrieved manually from the reference list of the relevant papers were also included.

The aim of this article is to review the etiopathogenesis, diagnosis, and treatment of OLP and to propose a novel algorithm and staging approaches to facilitate the work of clinicians who may encounter them in their practice.

Background on OLP

OLP is a common disease encountered worldwide, with a female preponderance^{2–4}. The overall incidence of OLP is uncertain but is estimated to be between 0.5 and 1.5%⁵. The involvement of the oral mucosa in systemic LP patients is between 70 and 77%⁶. The disease commonly presents in the older population, between the fifth and sixth decade of life. This disease is uncommonly seen in younger patients. This was supported in a recent study investigating oral mucosal lesions in teenagers: only 0.13% out of more than 6,000 patients actually suffered from OLP as a cause of mucosal lesion⁷. The disease is influenced by local aggravating factors including tobacco smoking, dry mouth, mechanical irritants, and bacterial plaque. Despite the latter-mentioned aggravating factor, patients should be reassured that OLP is not an infectious disease.

The etiology of OLP is still unknown. Also, the pathogenesis is not entirely clear, but it is thought to arise from an immune response presumably involving CD4⁺ and CD8⁺ T lymphocytes producing cytokines, interleukin-2, and tumor necrosis factor within the oral epithelium that induce a chronic inflammatory response and keratinocyte apoptosis⁸. Hence, there is no surprise that this disease has also been associated with other illnesses that share an immunological basis⁹. OLP is seen in families,

which implies that genetic predisposition exists for this disease^{10,11}. Indeed, genetic polymorphisms of various human leukocyte antigen (HLA) markers have been suggested; gene polymorphisms of different HLA markers, as well as inflammatory cytokines and chemokines, have been associated with the presence of LP. Although the cause of these polymorphisms is not clear, their occurrence supports the autoantigen hypothesis¹².

Interestingly, the association with other autoimmune diseases, not restricted to OLP, has been recently rationalized with the discovery of ‘Survivin’, a multifunctional protein that is part of the inhibitor of apoptosis family and is functionally important for cell division, apoptosis, and, possibly, microRNA biogenesis. This protein has been found to be involved in various autoimmune diseases, hence shedding more light on their pathogenesis¹³.

The great impact OLP has on its patients is so tremendous that one study examining 50 patients with OLP compared with controls showed a greater degree of sleep problems in such a way that their sleep deprivation has led to psychiatric disorders such as depression and anxiety¹⁴. Another study showed that the longer the duration of subjective symptoms, the poorer the quality of life and the higher the level of perceived stress^{15,16}. These findings were further backed up in a few recent reviews, including one from the past year, that all indicate that the development of OLP and especially the exacerbation of this disease is linked with stress, anxiety, depression, and eventually impairment of the patient’s quality of life^{17–19}. The rationale behind this was proved with elevated salivary and/or urinary cortisol levels that corresponded to increased anxiety and depressive states¹⁹.

Comorbid risks with OLP

LP with oral involvement has been linked to other systemic diseases with different levels of evidence and various results. The strongest association exists with hepatitis C virus (HCV)²⁰. Although HCV has been strongly linked to systemic LP, not many studies have focused solely on OLP²¹. Also, the coexistence of HCV infection and OLP may be relevant in some geographic regions such as southeast Asia and southern Europe²² but less relevant in other regions such as central Europe, as it was exhibited in another recent study²³. LP has also been said to be affected by some other viruses such as those of the herpes family, but, in contrast to the latter, this was not substantiated with OLP²⁴. However, a much more significant association has been determined recently between human papilloma virus (HPV) and OLP; this association, as with HCV, also varies with different geographic populations. This association suggests a causal role in the malignant progression of OLP, which is of great significance, since OLP is benign in nature, but with this association malignant transformation is a potential risk, although it may not apply in all cases²⁵. Other associations that are mostly related to LP and not OLP in the strict sense include diabetes mellitus, hypertension, hypercholesterolemia, and rarely thymoma^{9,21,22,26}. Good’s syndrome is another example of a disease that has an association with OLP but an even

rarer occurrence when thymoma is combined with hypogammaglobulinemia²⁶.

Autoimmune diseases were present in 7% of OLP patients in one study but were not statistically significant when compared to a control group²².

Only a few studies showed a true association between OLP and other systemic diseases, of which only a few were controlled studies with a high level of evidence²⁷ (see Table 1 for a summary of studies associated with systemic diseases). However, equal or more studies have depicted results that disprove these associations, and indeed most of these associations are based on cases series and retrospective studies with relatively small samples studied^{28–32}.

Differential diagnosis and prognosis

The diagnosis of OLP begins with a thorough history and physical examination. Patients may or may not have symptoms. Although there is no consensus regarding the subtypes of OLP, commonly few variants present, including reticular, papular, plaque-like, atrophic, and ulcerative (erosive). The reticular form of OLP is typically asymptomatic and characterized by lacy or striated keratosis. Patients may be aware of an altered oral texture (usually described as a rough sensation similar to sandpaper). On occasion, there will be irritation with hot and acidic foods or beverages. There can be chronic and lingering oral burning discomfort with erythematous and ulcerative lesions. Although experienced dentists and dermatologists can often diagnose OLP clinically, a biopsy may be necessary to exclude immunobullous disorders (e.g. mucous membrane pemphigoid and pemphigus vulgaris) if there is extensive ulceration and mucosal desquamation. Biopsy is

essential and must also be considered if there is concern regarding dysplastic or neoplastic transformation³², for example, a region of persistent and progressive erythema, induration, or tissue friability. Moreover, persistent symptomatic or asymptomatic red and white lesions in the oral cavity should never be ignored; these oral lesions must be biopsied. Patients complaining of dysphagia or odynophagia should be referred to a gastroenterologist to exclude esophageal LP with endoscopy.

OLP must be distinguished from other lichenoid lesions, including lichenoid drug reactions, lichenoid amalgam reactions, and graft-versus-host disease. Signs and symptoms of these disorders mimic OLP, and biopsy will be of limited value since the histopathological features are similar to OLP, but direct immunofluorescence is of importance to confirm the diagnosis. Nevertheless, a thorough medical history will usually facilitate diagnosis³³. Although currently there is no blood test to determine the diagnosis of OLP, blood tests may be required to rule out the presence of associated comorbidities such as diabetes mellitus, dyslipidemia, thyroid diseases, and chronic liver diseases.

Histopathology of OLP

The histological features of OLP include epithelial hyperkeratosis (orthokeratosis or parakeratosis), liquefactive degeneration of the basement membrane, and classically a band-like lymphocytic infiltrate in the superficial lamina propria comprising colloid or Civatte bodies, which are degenerated keratinocytes in the interface of the epithelium and mucosa. A saw-tooth pattern of the epithelial ridges is more commonly expected in the dermis of LP rather than in the mucosa of OLP^{33–35}. Histopathological evaluation is essential to precisely determine the diagnosis of OLP versus other diagnoses but more often to

Table 1. Summary of previous studies supporting the association of OLP with other diseases since 2000^{2,9,25,28,36,37}.

Author/Journal/year	PubMed ID	Country	Sample size	Systemic disease	Study design and methods	Outcome
Ma J, <i>PLoS One</i> . 2016 ²⁵	27571417	China	835 cases and 734 controls	HPV	Meta-analysis	HPV 16 and 18 showed strong association with OLP
Lauritano <i>et al. Head Face Med</i> . 2016 ⁹	27113338	Italy	87 OLP patients	HTN, DM, HCV thyroiditis	Retrospective chart review	Only one patient developed a malignant transformation (1.2%)
Barbosa <i>et al., Int J Dermatol</i> . 2015 ³⁶	25534406	Brazil	37 OLP patients	HTN, DM Less HCV, anxiety	Case series	No significant association was observed between the variables studied and clinical form or Sx
Gümrü B. <i>Med Oral Patol Oral Cir Bucal</i> . 2013 ²⁸	23524413	Turkey	370 OLP patients	HTN, DM, anxiety & depression	Retrospective chart review	Multiple sites in the majority of patients No malignant transformation
Bermejo-Fenoll <i>et al. Oral Pathol Med</i> . 2010 ³⁷	20456611	Spain	550 OLP patients	HTN, rheumatic diseases, GI disorders, anxiety & depression	A retrospective descriptive study	Five patients (0.9%) developed SCC
Eisen. <i>J Am Acad Dermatol</i> . 2002 ²	11807431	USA	723 OLP patients	HCV	Retrospective study	Oral SCC developed in six patients(0.8%)

DM, diabetes mellitus; HCV, hepatitis C virus; HPV, human papilloma virus; HTN, hypertension; GI, gastrointestinal; SCC, squamous cell carcinoma; Sx, symptoms.

rule out the possibility of neoplastic transformation that is included in the differential diagnosis but also may be merely a complication of longstanding OLP.

Current classification of OLP

There are a few variants of OLP, including atrophic, erosive, bullous plaque-like, and reticular forms. However, these variants are often intermingled. The reticular form presents with characteristic lacy white striae with a diffuse and bilateral distribution in the oral mucosa commonly affecting the buccal mucosa, ventral tongue, and gingiva. The atrophic and erosive forms present as red patches and ulceration, respectively. Plaque-form LP resembles leukoplakia but has a multifocal distribution. The bullous variant is rare, with bullae or vesicles that can resemble other oral immunobullous disorders (see [Figure 1](#) for various types of OLP).

Currently, there is no consensus regarding the classification of OLP, and in various reports the disease is characterized based on the different clinical forms and/or whether the disease is symptomatic or not. We believe that this gap should be better addressed in order to facilitate the work of clinicians. For the sake of simplicity, one approach is to categorize the OLP based on the degree of involvement as mild, moderate, or severe. However, since there are many subjective complaints, i.e. the mild cases may be described as severe in certain individuals, these complaints may and should dictate management. Another option is to divide the disease according to whether or not symptoms are available, namely symptomatic or asymptomatic. But this is not an adequate approach that will differentiate the different variants of this disease. For instance, whereas the reticular variant may be typically asymptomatic, the other variants may manifest with variable burning discomfort that is exacerbated by certain acidic and tart foods and beverages as well as with toothpastes and mouthwashes. There can be severe lingering pain affecting the patient's quality of life but little or no impact when the disease is asymptomatic in one study³⁸. For the clinical approach to various forms of OLP, see [Figure 2](#).

Forming an algorithmic approach is essential, since it is not always possible to generalize and to decide on management based solely on the clinical morphology. Nevertheless, we sought to suggest an algorithmic approach to facilitate the diagnosis of OLP (see [Figure 2](#)). It is essential to emphasize that our novel algorithm is only a suggested one and should be tested before widely accepted. Also, we would like to suggest a novel classification system for this disease in which it is separated into three stages—I, II, and III—that will correspond to, but better depict, three levels of severity—mild, moderate, and severe, respectively (see [Table 2](#)). Determining the level of involvement is pivotal for establishing the management strategy. Although it is a disease localized to one region, it should be considered as a systemic variant of LP. Moreover, a high prevalence of esophageal involvement was found in patients with systemic LP³⁹. In another recent study, 62% of the sample studied had esophageal involvement, of which 90% of patients with esophageal disease had oral involvement. However, that cohort was composed of a small sample⁴⁰.

Complications of OLP

We believe that OLP is not only associated with other systemic diseases (as mentioned previously) but also a systemic disease by itself that presents regionally. Naturally, it is impacted by other diseases that may shorten the lifespan of patients. But other complications do exist, including infection and malignant transformation. Despite the wide use of topical corticosteroids to manage patients with OLP, studies have failed to show a statistically significant increase in the risk for oral candidiasis; however, clinicians should be aware of it and treat it when it appears rather than prophylactically⁴¹.

The risk of malignant transformation is uncertain, but, although it exists, it is most probably much lower than once previously thought. Nevertheless, it was enough to include OLP in one of the oral malignant disorders⁴². In one systematic review, the overall rate of OLP patients who eventually suffered malignant transformation to squamous cell carcinoma (SCC) was

Figure 1. Images of patients with desquamative gingivitis, a leukoplakia like oral lichen planus, and reticular type oral lichen planus. We confirm that we have obtained written consent to use these images from the patients included in this figure.

Oral lichen planus – proposed diagnostic algorithm

Abbreviations: Bx, biopsy; F/U, follow up; OLP, oral lichen planus; R/O, rule out; Rx, treatment
 *e.g. white spongy nevus

Oral lichen planus – proposed diagnostic algorithm (cont'd- 1)

Abbreviations: Bx, biopsy; OLP, oral lichen planus; R/O, rule out; Rx, treatment

Oral Lichen planus – proposed diagnostic algorithm (cont'd-2)

Abbreviations: Bx, biopsy; DIF, direct immunofluorescence; OLP, oral lichen planus; R/O, rule out

Figure 2. Oral lichen planus – proposed diagnostic algorithm. Bx, biopsy; DIF, direct immunofluorescence; F/U, follow up; OLP, oral lichen planus; R/O, rule out; Rx, treatment; *e.g. white spongy nevus

Table 2. A new classification of OLP.

Stages of disease	Description	Management plan	Recommended discipline
I: mild	No symptoms	No pharmacotherapy indicated May use nonpharmaceutical substances	Dentist
II: moderate	Mild-to-moderate symptoms (e.g. sensitivity with spicy foods, mild lingering pain)	Low-potency corticosteroids/alternative TCIs, treat PRN	Preferably oral maxillofacial surgeon specialist and/or dermatologist
III: severe	Severe or recalcitrant symptoms/systemic manifestations and/or diffuse involvement of the entire oral cavity	First-line high-potency corticosteroids, systemic treatment (see text)	Dermatologist or another immunotherapy specialist May involve consultants as well such as ENT specialists

ENT, ear, nose, and throat; OLP, oral lichen planus; PRN, *pro re nata* (when necessary); TCI, topical calcineurin inhibitor.

1%⁴³. Another more recent study showed that patients with OLP were 4.8 times more likely to have oral SCC than the matched referents⁴⁴. In another Finnish population survey involving more than 13,000 LP patients, the risks of cancers of the oral mucosa, esophagus, and larynx were significantly elevated. The standardized incidence ratio for cancer of the tongue was

12.4 (95% confidence interval [CI] 9.45–16.0) and for cancer of the oral cavity was 7.97 (95% CI 6.79–9.24)⁴⁵. Two other recent systematic reviews have also showed a low risk of potential malignant transformation: one with an overall malignant transformation rate to SCC of 1.4%⁴⁶ and another with a combined malignant transformation rate to SCC

of 1.14%⁴⁷. However, both studies have identified specific risk factors for malignant transformation to include tongue localization, atrophic-erosive lesions, tobacco smoking, and alcohol consumption^{46,47}.

Management

There is no consensus despite scarce reports about an algorithmic approach towards the management of OLP⁴⁸. In general, management should be directed towards symptoms. No therapy is advocated in the absence of symptoms. Irritating foods, beverages, and oral hygiene products (e.g. minty toothpastes) should be avoided. Optimum oral hygiene and regular dental cleanings are helpful for minimizing plaque and gingival inflammation with the potential to exacerbate this condition. Pharmacotherapy is indicated, however, when symptoms are severe, lingering, or interfering with daily functions (e.g. toothbrushing, eating). The rationale behind the therapy is based on further insight into the pathogenesis of OLP as an immune-mediated disease related to T-lymphocyte immunological dysfunction, and within the implicated cytokines, such as TNF-alpha, IFN-gamma, TNF-alpha, IL-6, and IL-8, which has paved the way for the utilization of immunosuppressant therapies. Therefore, commonly, topical corticosteroids are utilized initially for symptomatic OLP, and, if topical therapy fails, systemic therapy is considered.

However, owing to the chronic nature of this disease, a complete cure is very difficult to achieve. There is no uniform approach to treatment, with treatment varying from one individual to another. Nevertheless, in recalcitrant cases, calcineurin inhibitors (such as tacrolimus and pimecrolimus) have been introduced as a second line of therapy. These medications should be used by experts with caution, preferably for short-term treatments, owing to the risk of developing oral SCC after its usage^{49,50}.

Only scarce clinical trials are available that examined the effect of systemic medications such as methotrexate, systemic pulse therapy of corticosteroids, alitretinoin, and, more recently, apremilast^{51,52}. Studies on systemic therapies in the form of energy-based devices such as photodynamic therapy or laser therapy, or these in combination with corticosteroids, were also published^{53–56}. However, many more reports advocate the

utilization of topical medications, especially corticosteroids either alone or in combination in various chemical preparations⁵⁷. More than a dozen studies have investigated the use of topical calcineurin inhibitors (pimecrolimus or tacrolimus) either alone or compared to corticosteroids^{58,59}. Also, other topical components such as cyclosporine have been examined^{60–62}. Fewer attempts are made with various other substances such as *Aloe vera* gel or other “natural” substances or topical preparations such as tazarotene with various results in anecdotal reports^{63–65}. What is the reason for more reports of various topical preparations versus systemic therapies? One explanation is that clinicians and patients are settling in the use of topical medications for OLP and reserving systemic medications for when the former fails. Another explanation is since there is no gold standard therapy nor consensus or guidelines on how to treat these patients, the management decision is being made on a case-by-case basis.

Conclusions

OLP is a common disease with an overwhelmingly negative impact on the patient's quality of life. The diagnosis is mostly based on clinical grounds, but it seems that histological and immunofluorescence confirmation is necessary owing to many reasons we have mentioned. Despite being localized only to the oral cavity, in our opinion, OLP should be recategorized as a systemic disease because of the major impact it has and should not be considered as merely one of the manifestations of LP. We do not know whether there is an immunological relationship among LP, OLP, etc., or if indeed this will help us define a therapeutic approach that aligns with the ideal target at each stage. Nevertheless, there is still no uniform approach towards the management of OLP and, except for the widely used topical corticosteroids, the therapy varies greatly between clinicians and medical institutions. This might stem from the lack of uniform classification. We have suggested an easier new classification system and algorithmic approach to the different variants of the disease. Our suggested algorithm should be critically appraised and tested in the future. However, we hope that this will facilitate the work of clinicians. Having said that, not only newer classifications but also more novel therapeutic options are required to provide more hope for OLP patients. Overall, we suggest adopting a multidisciplinary holistic approach and recommend close monitoring for these patients.

References

- Asch S, Goldenberg G: **Systemic treatment of cutaneous lichen planus: an update**. *Cutis*. 2011; **87**(3): 129–34.
[PubMed Abstract](#)
- Eisen D: **The clinical features, malignant potential, and systemic associations of oral lichen planus: a study of 723 patients**. *J Am Acad Dermatol*. 2002; **46**(2): 207–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Axéll T, Rundquist L: **Oral lichen planus—a demographic study**. *Community Dent Oral Epidemiol*. 1987; **15**(1): 52–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Dissemond J: **Oral lichen planus: an overview**. *J Dermatolog Treat*. 2004; **15**(3): 136–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Di Stasio D, Guida A, Salerno C, et al.: **Oral lichen planus: a narrative review**. *Front Biosci (Elite Ed)*. 2014; **6**: 370–6.
[PubMed Abstract](#)
- Payeras MR, Cherubini K, Figueiredo MA, et al.: **Oral lichen planus: Focus on etiopathogenesis**. *Arch Oral Biol*. 2013; **58**(9): 1057–69.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Amadori F, Bardellini E, Conti G, et al.: **Oral mucosal lesions in teenagers: a cross-sectional study**. *Ital J Pediatr*. 2017; **43**(1): 50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**

8. Nogueira PA, Carneiro S, Ramos-E-Silva M: **Oral lichen planus: an update on its pathogenesis.** *Int J Dermatol.* 2015; **54**(9): 1005–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Lauritano D, Arrica M, Lucchese A, *et al.*: **Oral lichen planus clinical characteristics in Italian patients: a retrospective analysis.** *Head Face Med.* 2016; **12**: 128.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Bermejo-Fenoll A, López-Jornet P: **Familial oral lichen planus: presentation of six families.** *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2006; **102**(2): e12–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Lu SL, Qi XM, Dong G, *et al.*: **Clinical characteristics and analysis of familial oral lichen planus in eight Chinese families.** *Exp Ther Med.* 2016; **12**(4): 2281–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Gorouhi F, Davari P, Fazel N: **Cutaneous and mucosal lichen planus: a comprehensive review of clinical subtypes, risk factors, diagnosis, and prognosis.** *ScientificWorldJournal.* 2014; **2014**: 742826.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. **F** Gravina G, Wasén C, García-Bonete MJ, *et al.*: **Survivin in autoimmune diseases.** *Autoimmun Rev.* 2017; **16**(8): 845–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
14. Adamo D, Ruoppo E, Leuci S, *et al.*: **Sleep disturbances, anxiety and depression in patients with oral lichen planus: a case-control study.** *J Eur Acad Dermatol Venereol.* 2015; **29**(2): 291–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. **F** Radwan-Oczko M, Zwyrtek E, Owczarek JE, *et al.*: **Psychopathological profile and quality of life of patients with oral lichen planus.** *J Appl Oral Sci.* 2018; **26**: e20170146.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
16. Lundqvist EN, Wahlén YB, Bergdahl M, *et al.*: **Psychological health in patients with genital and oral erosive lichen planus.** *J Eur Acad Dermatol Venereol.* 2006; **20**(6): 661–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. **F** Cerqueira JDM, Moura JR, Arsati F, *et al.*: **Psychological disorders and oral lichen planus: A systematic review.** *J Invest Clin Dent.* 2018; **9**(4): e12363.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
18. **F** Hasan S, Ahmed S, Kiran R, *et al.*: **Oral lichen planus and associated comorbidities: An approach to holistic health.** *J Family Med Prim Care.* 2019; **8**(11): 3504–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
19. **F** Manczyk B, Gólda J, Biniak A, *et al.*: **Evaluation of depression, anxiety and stress levels in patients with oral lichen planus.** *J Oral Sci.* 2019; **61**(3): 391–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
20. Baccaglini L, Thongprasom K, Carrozzo M, *et al.*: **Urban legends series: lichen planus.** *Oral Dis.* 2013; **19**(2): 128–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Tovuș S, Parlătescu I, Gheorghe C, *et al.*: **Oral lichen planus: a retrospective study of 633 patients from Bucharest, Romania.** *Med Oral Patol Oral Cir Bucal.* 2013; **18**(2): e201–e206.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. López-Jornet P, Parra-Pérez F, Pons-Fuster A: **Association of autoimmune diseases with oral lichen planus: a cross-sectional, clinical study.** *J Eur Acad Dermatol Venereol.* 2014; **28**(7): 895–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Remmerbach TW, Liese J, Krause S, *et al.*: **No association of oral lichen planus and hepatitis C virus infection in central Germany.** *Clin Oral Investig.* 2016; **20**(1): 193–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Lodi G, Scully C, Carrozzo M, *et al.*: **Current controversies in oral lichen planus: report of an international consensus meeting. Part 1. Viral infections and etiopathogenesis.** *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2005; **100**(1): 40–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Ma J, Zhang J, Zhang Y, *et al.*: **The Magnitude of the Association between Human Papillomavirus and Oral Lichen Planus: A Meta-Analysis.** *PLoS One.* 2016; **11**(8): e0161339.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Motegi SI, Uchiyama A, Yamada K, *et al.*: **Lichen planus complicated with thymoma: Report of three Japanese cases and review of the published work.** *J Dermatol.* 2015; **42**(11): 1072–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. **F** Cassol-Spanemberg J, Rodríguez-de Rivera-Campillo ME, Otero-Rey EM, *et al.*: **Oral lichen planus and its relationship with systemic diseases. A review of evidence.** *J Clin Exp Dent.* 2018; **10**(9): e938–e944.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
28. Gümrü B: **A retrospective study of 370 patients with oral lichen planus in Turkey.** *Med Oral Patol Oral Cir Bucal.* 2013; **18**(3): e427–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Xue JL, Fan MW, Wang SZ, *et al.*: **A clinical study of 674 patients with oral lichen planus in China.** *J Oral Pathol Med.* 2005; **34**(8): 467–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
30. Ingafou M, Leao JC, Porter SR, *et al.*: **Oral lichen planus: a retrospective study of 690 British patients.** *Oral Dis.* 2006; **12**(5): 463–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Vučićević Boras V, Savage NW, Brailo V, *et al.*: **The significance of oral and systemic factors in Australian and Croatian patients with oral lichen planus.** *Acta Dermatovenereol Croat.* 2014; **22**(2): 97–102.
[PubMed Abstract](#)
32. Shen ZY, Liu W, Zhu LK, *et al.*: **A retrospective clinicopathological study on oral lichen planus and malignant transformation: analysis of 518 cases.** *Med Oral Patol Oral Cir Bucal.* 2012; **17**(6): e943–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. **F** Yamanaka Y, Yamashita M, Innocentini LMA, *et al.*: **Direct Immunofluorescence as a Helpful Tool for the Differential Diagnosis of Oral Lichen Planus and Oral Lichenoid Lesions.** *Am J Dermatopathol.* 2018; **40**(7): 491–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
34. Setterfield JF, Black MM, Challacombe SJ: **The management of oral lichen planus.** *Clin Exp Dermatol.* 2000; **25**(3): 176–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Cheng YS, Gould A, Kurago Z, *et al.*: **Diagnosis of oral lichen planus: A position paper of the American Academy of Oral and Maxillofacial Pathology.** *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2016; **122**(3): 332–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Barbosa NG, Silveira EJ, Lima EN, *et al.*: **Factors associated with clinical characteristics and symptoms in a case series of oral lichen planus.** *Int J Dermatol.* 2015; **54**(1): e1–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Bermejo-Fenoll A, Sánchez-Siles M, López-Jornet P, *et al.*: **A retrospective clinicopathological study of 550 patients with oral lichen planus in south-eastern Spain.** *J Oral Pathol Med.* 2010; **39**(6): 491–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Karbach J, Al-Nawas B, Moergel M, *et al.*: **Oral health-related quality of life of patients with oral lichen planus, oral leukoplakia, or oral squamous cell carcinoma.** *J Oral Maxillofac Surg.* 2014; **72**(8): 1517–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Quispel R, van Boxel OS, Schipper ME, *et al.*: **High prevalence of esophageal involvement in lichen planus: a study using magnification chromoendoscopy.** *Endoscopy.* 2009; **41**(3): 187–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
40. Kern JS, Technau-Hafsi K, Schwacha H, *et al.*: **Esophageal involvement is frequent in lichen planus: study in 32 patients with suggestion of clinicopathologic diagnostic criteria and therapeutic implications.** *Eur J Gastroenterol Hepatol.* 2016; **28**(12): 1374–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Marable, Bowers LM, Stout TL, *et al.*: **Oral candidiasis following steroid therapy for oral lichen planus.** *Oral Dis.* 2016; **22**(2): 140–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
42. Warnakulasuriya S, Johnson NW, van der Waal I: **Nomenclature and classification of potentially malignant disorders of the oral mucosa.** *J Oral Pathol Med.* 2007; **36**(10): 575–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Fitzpatrick SG, Hirsch SA, Gordon SC: **The malignant transformation of oral lichen planus and oral lichenoid lesions: a systematic review.** *J Am Dent Assoc.* 2014; **145**(1): 45–56.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Lianos V, Torgerson RR, Ramos-Rodríguez AJ, *et al.*: **Incidence of squamous cell carcinoma in oral lichen planus: a 25-year population-based study.** *Int J Dermatol.* 2019; **58**(3): 296–301.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
45. **F** Halonen P, Jakobsson M, Heikinheimo O, *et al.*: **Cancer risk of Lichen planus: A cohort study of 13,100 women in Finland.** *Int J Cancer.* 2018; **142**(1): 18–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
46. **F** Giuliani M, Troiano G, Cordaro M, *et al.*: **Rate of malignant transformation of oral lichen planus: A systematic review.** *Oral Dis.* 2019; **25**(3): 693–709.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
47. **F** González-Moles MÁ, Ruiz-Ávila I, González-Ruiz L, *et al.*: **Malignant transformation risk of oral lichen planus: A systematic review and comprehensive meta-analysis.** *Oral Oncol.* 2019; **96**: 121–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
48. Lodi G, Scully C, Carrozzo M, *et al.*: **Current controversies in oral lichen planus: report of an international consensus meeting. Part 2. Clinical management and malignant transformation.** *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2005; **100**(2): 164–78.
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Morita M, Asoda S, Tsunoda K, *et al.*: **The onset risk of carcinoma in patients continuing tacrolimus topical treatment for oral lichen planus: a case report.** *Odontology.* 2017; **105**(2): 262–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. **F** Thongprasom K: **Challenges in using topical calcineurin inhibitors as a treatment for recalcitrant oral lichen planus.** *Br J Dermatol.* 2019; **181**(6): 1120.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

51. Lajevardi V, Ghodsi SZ, Hallaji Z, *et al.*: **Treatment of erosive oral lichen planus with methotrexate.** *J Dtsch Dermatol Ges.* 2016; **14**(3): 286–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
52. Bettencourt M: **Oral Lichen Planus Treated With Apremilast.** *J Drugs Dermatol.* 2016; **15**(8): 1026–8.
[PubMed Abstract](#)
53. Kunz M, Urošević-Maiwald M, Goldinger SM, *et al.*: **Efficacy and safety of oral alitretinoin in severe oral lichen planus—results of a prospective pilot study.** *J Eur Acad Dermatol Venereol.* 2016; **30**(2): 293–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
54. **F** Mostafa D, Moussa E, Alnouaem M: **Evaluation of photodynamic therapy in treatment of oral erosive lichen planus in comparison with topically applied corticosteroids.** *Photodiagnosis Photodyn Ther.* 2017; **19**: 56–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
55. **F** Mirza S, Rehman N, Alrahliah A, *et al.*: **Efficacy of photodynamic therapy or low level laser therapy against steroid therapy in the treatment of erosive-atrophic oral lichen planus.** *Photodiagnosis Photodyn Ther.* 2018; **21**: 404–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
56. Huang Z, Wang Y, Liang Q, *et al.*: **The application of a carbon dioxide laser in the treatment of superficial oral mucosal lesions.** *J Craniofac Surg.* 2015; **26**(3): e277–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
57. Dillenburg CS, Martins MA, Munerato MC, *et al.*: **Efficacy of laser phototherapy in comparison to topical clobetasol for the treatment of oral lichen planus: a randomized controlled trial.** *J Biomed Opt.* 2014; **19**(6): 068002.
[PubMed Abstract](#) | [Publisher Full Text](#)
58. **F** Hambly JL, Haywood A, Hattingh L, *et al.*: **Comparison between self-formulation and compounded-formulation dexamethasone mouth rinse for oral lichen planus: a pilot, randomized, cross-over trial.** *J Invest Clin Dent.* 2017; **8**(3).
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
59. Arduino PG, Carbone M, Della Ferrera F, *et al.*: **Pimecrolimus vs. tacrolimus for the topical treatment of unresponsive oral erosive lichen planus: a 8 week randomized double-blind controlled study.** *J Eur Acad Dermatol Venereol.* 2014; **28**(4): 475–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
60. Sonthalia S, Singal A: **Comparative efficacy of tacrolimus 0.1% ointment and clobetasol propionate 0.05% ointment in oral lichen planus: a randomized double-blind trial.** *Int J Dermatol.* 2012; **51**(11): 1371–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
61. Thongprasom K, Chaimusig M, Korkij W, *et al.*: **A randomized-controlled trial to compare topical cyclosporin with triamcinolone acetonide for the treatment of oral lichen planus.** *J Oral Pathol Med.* 2007; **36**(3): 142–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
62. Conrotto D, Carbone M, Carrozzo M, *et al.*: **Ciclosporin vs. clobetasol in the topical management of atrophic and erosive oral lichen planus: a double-blind, randomized controlled trial.** *Br J Dermatol.* 2006; **154**(1): 139–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
63. Mansourian A, Momen-Heravi F, Saheb-Jamee M, *et al.*: **Comparison of aloe vera mouthwash with triamcinolone acetonide 0.1% on oral lichen planus: a randomized double-blinded clinical trial.** *Am J Med Sci.* 2011; **342**(6): 447–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
64. Petrucci M, De Benedittis M, Grassi R, *et al.*: **Oral lichen planus: a preliminary clinical study on treatment with tazarotene.** *Oral Dis.* 2002; **8**(6): 291–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
65. **F** Hashem AS, Issrani R, Elsayed TEE, *et al.*: **Topical hyaluronic acid in the management of oral lichen planus: A comparative study.** *J Investig Clin Dent.* 2019; **10**(2): e12385.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

- 1. Kobkan Thongprasom**
Oral Medicine Department, Faculty of Dentistry, Chulalongkorn University, Bangkok, Thailand
Competing Interests: No competing interests were disclosed.
- 2. Andrés Blanco-Carrión**
Stomatology Department, School of Dentistry, University of Santiago de Compostela, Santiago de Compostela, Spain
Competing Interests: No competing interests were disclosed.
- 3. Maria Jose Garcia-Pola Vallejo**
Departamento de Cirugía y Especialidades Médico-Quirúrgicas, Facultad de Medicina, Universidad de Oviedo, Oviedo, Spain
Competing Interests: No competing interests were disclosed.
- 4. José López-López**
Oral Health and Masticatory System Group (Bellvitge Biomedical Research Institute) IDIBELL, University of Barcelona, Barcelona, Spain
Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research