RESEARCH ARTICLE

Determinants of safe delivery utilization among Indonesian women in eastern part of Indonesia [version 1; peer review: 2 approved]

Ferry Efendi ^{1,2}, Susy Katikana Sebayang^{3,4}, Erni Astutik ^{4,5},
Setho Hadisyatmana ^{1,2}, Eka Mishbahatul Mar'ah Has ¹, Heri Kuswanto⁶

¹Faculty of Nursing, Universitas Airlangga, Surabaya, Indonesia

²School of Nursing & Midwifery, La Trobe University, Melbourne, Australia

³Department of Biostatistics and Population Studies, Faculty of Public Health, Universitas Airlangga, Banyuwangi Campus, Banyuwangi, Indonesia

⁴Research Group for Health and Wellbeing of Women and Children, Faculty of Public Health, Universitas Airlangga, Banyuwangi, Indonesia

⁵Department of Epidemiology, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

⁶Department of Statistics, Institut Teknologi Sepuluh Nopember (ITS), Surabaya, Indonesia

V1 First published: 05 May 2020, 9:332
<https://doi.org/10.12688/f1000research.23324.1>

Latest published: 23 Sep 2020, 9:332
<https://doi.org/10.12688/f1000research.23324.2>

Abstract

Background: Improving maternal health and reducing maternal mortality are part of the United Nations global Sustainable Development Goals for 2030. Ensuring every woman's right to safe delivery is critical for reducing the maternal mortality rate, especially in Indonesia. Our study aimed to identify determinants of safe delivery utilization among women in the eastern Indonesia.

Methods: This study was cross-sectional and used data from the 2017 Indonesian Demographic and Health Survey (IDHS). A total of 2,162 women who had their last child in the five years preceding the survey and lived in the eastern part of Indonesia were selected as the respondents. Chi-squared test and binary logistic regression were used to understand the determinants of safe delivery.

Results: Higher child rank and interval ≤ 2 years (OR: 0.30, 95% CI: 0.19-0.47), unwanted pregnancy at time of becoming pregnant (OR: 1.48, 95% CI: 1.05-2.08), richest wealth quintile (OR: 5.59, 95% CI: 3.37-9.30), more than four antenatal care visits (OR: 3.62, 95% CI: 2.73-4.79), rural residence (OR: 0.49, 95% CI: 0.36-0.66), good composite labor force participation (OR: 1.47, 95% CI: 1.15-1.89), and a good attitude towards domestic violence (OR: 1.33, 95% CI: 1.04-1.69) were found to be significantly associated with facility-based delivery. Higher child rank and interval ≤ 2 years (OR: 0.49, 95% CI: 0.29-0.83), husband/partner having completed secondary or higher education (OR: 2.18, 95% CI: 1.48-3.22), husband/partner having a non-agricultural occupation (OR: 1.35, 95% CI: 1.00-1.81), being in the

Open Peer Review

Approval Status

	1	2
version 2		
(revision)		
23 Sep 2020	view	
version 1		
05 May 2020		
	view	view

1. **Ryan Michael Flores Oducado** , West Visayas State University, Iloilo City, Philippines

2. **Asmaa Salah Eldin Mohamed Saleh** , Beni Suef University, Beni Suef, Egypt

Any reports and responses or comments on the article can be found at the end of the article.

richest wealth quintile (OR: 15.69, 95% CI: 5.53-44.50), and three other factors were found to be significantly associated with skilled assistance delivery.

Conclusions: Safe delivery and facility-based delivery among women in the eastern part of Indonesia were determined by several individual and household factors. An open innovation and partnership process that engages the full range of stakeholders should be developed based on local needs.

Keywords

facility-based delivery, safe delivery, skilled birth delivery.

Corresponding author: Ferry Efendi (ferry-e@fkip.unair.ac.id)

Author roles: **Efendi F:** Conceptualization, Formal Analysis, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Sebayang SK:** Conceptualization, Formal Analysis, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Astutik E:** Conceptualization, Data Curation, Formal Analysis, Methodology, Validation, Writing – Review & Editing; **Hadisuyatmana S:** Conceptualization, Methodology, Writing – Review & Editing; **Has EMM:** Conceptualization, Methodology, Writing – Review & Editing; **Kuswanto H:** Conceptualization, Methodology, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This study was funded by Universitas Airlangga, Surabaya, Indonesia through “Hibah Riset Mandat” (Mandate Research Grant) grant number [335/UN3.14/LT/2019].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2020 Efendi F *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Efendi F, Sebayang SK, Astutik E *et al.* **Determinants of safe delivery utilization among Indonesian women in eastern part of Indonesia [version 1; peer review: 2 approved]** F1000Research 2020, 9:332 <https://doi.org/10.12688/f1000research.23324.1>

First published: 05 May 2020, 9:332 <https://doi.org/10.12688/f1000research.23324.1>

Introduction

Maternal morbidity and mortality is a global health concern (World Health Organization, 2017). Every day in 2017, around 830 mothers died due to pregnancy and childbirth (World Health Organization, 2018). The United Nations Sustainable Development Goals set a target to reduce maternal deaths to 70 per 100,000 live births by 2030 (Nations, 2015). In Indonesia, the maternal mortality rate is still high, at 305 per 100,000 live births (BPS, 2015b). A higher rate was found in the eastern part of Indonesia, namely Nusa Tenggara, Maluku, and Papua Island, than in the other islands (BPS, 2015a). One of the major causes of maternal mortality is bleeding, which is followed by eclampsia (Tejayanti *et al.*, 2012). Safe delivery as the critical policy of making motherhood safer requires skilled health professionals and facility-based delivery across the provinces of Indonesia (Efendi *et al.*, 2019; Kementerian Kesehatan RI, 2014). As an archipelago country, institutional delivery and skilled assistant delivery are still a challenge because of the geographical situation (Belton *et al.*, 2014; Ministry of Health [MoH], 2012). To increase safe delivery for Indonesian mothers, the government has set a goal to reach 85% of institutional deliveries in 2019 (Kementerian Kesehatan RI, 2017). Even though the government has not yet set a goal for skilled attendant delivery specifically in this document, it should be assumed that the government demands the highest standard of health attainment.

The 2017 IDHS found that there is a gap in coverage of institutional delivery and skilled assistant delivery between western provinces and eastern provinces of Indonesia. Eastern provinces of Indonesia, including Bali, Nusa Tenggara Island, Sulawesi Island, Maluku, and Papua Island, have not reached 70% coverage of safe delivery in either institutional delivery or skilled assistant delivery, as depicted in Figure 1 (BPS *et al.*, 2018).

Studies that examine safe delivery have been conducted in some countries. A study conducted in Ethiopia found that residence, religion, educational level, age at first pregnancy, parity, and antenatal care (ANC) attendance have a significant association with safe delivery service utilization (Abera *et al.*, 2011). Another study conducted in Tanzania reported that in

addition to socio-demographic factors, women's empowerment status contributed to the decision to give birth with a health professional (Shimamoto & Gipson, 2015). In a similar vein, studies in 13 sub-Saharan African countries found that living conditions and women's autonomy are key factors of maternal healthcare utilization (Iacoella & Tirivayi, 2019). In Indonesia, a study about facility-based childbirth found that educational level, place of residence, working status, involvement in decision-making, economic status, and ANC visits are significantly associated with health facility delivery among women (Efendi *et al.*, 2019). Furthermore, the gap in age and education between a woman and her husband/partner, women's self-esteem, age at first marriage, and age at pregnancy were found have a high association with institutional delivery among Indonesian women (Kurniati *et al.*, 2018).

The gap in coverage of safe delivery, including institutional delivery and skilled assistant delivery, in western and eastern provinces of Indonesia should be resolved to attain the Sustainable Development Goals by 2030. Therefore, this study aimed to determine safe delivery utilization among Indonesian women in the eastern part of Indonesia.

Methods

Ethical statement

IDHS ethical clearance was obtained from the Indonesian Ministry of Health (MoH). For this study, permission to use the data was obtained from Inner City Fund (ICF) International. This study used existing IDHS data and re-analysis was done under the original consent provided by the participants. Thus, no further consent was obtained from the participants.

Data source

This was an analytical cross-sectional study that used data from the 2017 IDHS. The 2017 IDHS was conducted in 34 provinces in Indonesia from July to December 2017 by the Central Statistics Agency (BPS), National Population and Family Planning Board (BKKBN), and the Ministry of Health with technical help from ICF. The Individual Recode (IR) dataset was downloaded from www.dhsprogram.com after completing registration.

Figure 1. Institutional delivery and skilled assistant delivery by western and eastern provinces of Indonesia.

Sample size and sampling

In the 2017 IDHS, a total of 49,627 women were finished the survey. Two-stage cluster sampling was used to select the respondents. Interviews were performed as privately as possible with a detailed manual as reported by ICF (ICF Macro, 2020).

The inclusion criteria for this study were women aged 15–49 years who had their last child in the five years preceding the survey and lived in the eastern provinces of Indonesia. For the purpose of analysis, we divided Indonesia into two greater parts, western and eastern, based on the geographical location. The eastern provinces included Bali, West Nusa Tenggara, East Nusa Tenggara, North Sulawesi, Central Sulawesi, South Sulawesi, Southeast Sulawesi, Gorontalo, West Sulawesi, Maluku, North Maluku, West Papua and Papua. In total, survey data from 2,162 women meeting the criteria were accessed for this study's analysis.

Variables

The dependent variables in this study were facility-based delivery and assisted delivery. Facility-based delivery was divided into two categories: health facility and non-health facility. Health facility delivery or institutional delivery is delivery that is carried out at a health facility, including public health centers, clinics or maternity homes, and hospitals. The assisted delivery variable was also divided into two categories: health professional and non-health professional. Health professional or skilled assistant delivery is defined as birth delivered with the assistance of skilled providers such as general practitioners, obstetricians, midwives, and skilled nurses.

There were several independent variables in this study. Age difference between man and woman was divided into four categories: woman older than husband, 0–4 years younger, 5–7 years younger, and >7 years younger. Birth rank and interval was divided into five categories: second or third child with interval >2 years; first birth, second or third child with interval ≤2 years; fourth or higher child with interval >2 years; and fourth or higher child with interval ≤2 years. Planning status of births, women who had a birth or several births in the five years prior to their interview were asked whether the pregnancy had been wanted at the time it occurred (wanted then) or whether it had been wanted but had occurred sooner than wanted (wanted later), or whether the woman had wanted no further children at the time (unwanted). Husband/partner's education attainment was divided into three categories: incomplete primary education/none, complete primary or some secondary, and completed secondary or higher. Husband/partner's occupation was divided into two categories: agricultural and non-agricultural. Wealth quintile was categorized as poorest, poorer, middle, richer, and richest (Rutstein & Johnson, 2004). Number of household members was divided into two categories: households that have less than four members and households with four or more members. Number of ANC visits was categorized as less than four times and four times or more. Covered by health insurance was divided into two categories: yes and no. Residence was categorized as urban and rural.

Women's empowerment variables, including composite labor force participation, attitude towards domestic violence (wife-beating), decision-making power, and women's knowledge level, were divided into three categories: poor, moderate, and good. Further details on how these variables were assessed can be found in study as conducted by Sebayang, *et al.* (2019).

Data analysis

The determinants of safe delivery were analyzed using a Chi-square test and binary logistic regression. Both analyses were performed in Stata version 14. The variables were significant at a p-value of 0.05, and the strength of the association was assessed using odds ratio (OR) with a 95% confidence interval (CI).

Results

Among the women who were included in this study, 71.6% used a health facility and 86.2% were assisted by a health professional at their last birth. The majority of the respondents are 0–4 years younger than their husband (41.2%), from the poorest wealth quintile (41.8%), have four or more members in the household (87.6%), are covered by health insurance (64.3%), and live in a rural residence (66.7%). Concerning the husband/partner's education and occupation, 47.3% have completed secondary or higher education and more than half work in an agricultural occupation (54.1%). For almost half the respondents, their last child was a second or third child with an interval more than two years (44.9%). The majority of respondents had more than four ANC visits (88.4) and their pregnancy was wanted when they became pregnant (82.1%). In terms of women's empowerment, most respondents have good composite labor force participation (35.7%), a moderate attitude towards domestic violence (34.8%), poor decision-making power (35.3%), and a poor level of knowledge (34.6%). Details about the descriptive characteristics of the respondents are shown in Table 1.

In the bivariate analysis, most of the variables showed a significant association with a p-value of 0.05 with both outcomes: facility-based delivery and assisted delivery. For the facility-based delivery outcome, three variables have a p-value of more than 0.05 (pregnancy was wanted at time of becoming pregnant, number of household members, decision-making power), while for the assisted delivery outcome, four variables were not significant (age difference between man and woman, pregnancy was wanted at time of becoming pregnant, number of household members, decision-making power). Details about the bivariate analysis are shown in Table 2 and Table 3.

In the binary logistic regression analysis, delivery at a health facility was associated with several variables. Women who lived in a rural residence [AOR=0.49; 95% CI=0.36-0.66] were less likely to deliver in a health facility. A similar result was found for women whose last child was a fourth or higher child with an interval of two years or under [AOR=0.30; 95% CI=0.19-0.47]. Women from the richest wealth quintile family and those who had four or more ANC visits were five [AOR=5.59; 95% CI=3.37-9.30] and three times more likely to deliver in a health facility [AOR=3.62; 95% CI=2.73-4.79], respectively.

Table 1. Characteristics of the respondents regarding determinants of safe delivery utilization among Indonesian women in eastern part of Indonesia (n=2,162).

Variable	n	%
Facility-based delivery		
Non-health facility	615	28.4
Health facility	1,547	71.6
Assisted delivery		
Non-health professional	298	13.8
Health professional	1,864	86.2
Age difference between man and woman		
Woman older than man	450	20.8
0–4 years	890	41.2
5–7 years	434	20.1
>7 years	388	17.9
Birth rank and interval		
Second or third child, interval >2 years	970	44.9
First birth	603	27.9
Second or third child, interval ≤2 years	131	6.0
Fourth or higher child, interval >2 years	384	17.8
Fourth or higher child, interval ≤2 years	74	3.4
Planning status of births		
Then	1,774	82.1
Later	246	11.4
No more	142	6.5
Husband/partner's education attainment		
Incomplete primary education/none	299	13.8
Complete primary or some secondary	841	38.9
Completed secondary or higher	1,022	47.3
Husband/partner's occupation		
Agricultural	1,169	54.1
Non-agricultural	993	45.9
Wealth quintile		
Poorest	903	41.8
Poorer	454	21.0
Middle	315	14.6
Richer	253	11.7
Richest	237	11.0
Number of household members		
<4 members	268	12.4
≥4 members	1,894	87.6
Number of antenatal care visits		
<4 visits	251	11.6
≥4 visits	1,911	88.4

Variable	n	%
Covered by health insurance		
No	772	35.7
Yes	1,390	64.3
Residence		
Urban	719	33.3
Rural	1,443	66.7
Composite labor force participation		
Poor	692	32.0
Moderate	698	32.3
Good	772	35.7
Attitude toward domestic violence		
Poor	674	31.2
Moderate	753	34.8
Good	735	34.0
Decision-making power		
Poor	764	35.3
Moderate	703	32.5
Good	695	32.1
Women's knowledge level		
Poor	748	34.6
Moderate	714	33.0
Good	700	32.4

Women who have good composite labor force participation [AOR=1.47; 95% CI=1.15-1.89] and a moderate attitude towards domestic violence [AOR=1.38; 95% CI=1.10-1.73] were more likely to deliver in a health facility. Women whose pregnancy was unwanted when they became pregnant [AOR=1.48 95% CI=1.05-2.08] were also more likely to deliver in a health facility. Details about the binary logistic regression analysis with a facility-based delivery outcome are shown in [Table 4](#).

According to the assisted delivery outcome, women whose last child was a fourth or higher child with an interval of two years or under [AOR=0.49; 95% CI=0.29-0.83] were less likely to deliver with a health professional. Women whose husband completed secondary or higher education were two times [AOR=2.18; 95% CI=1.48-3.22] more likely to deliver with a health professional. Likewise, women who had four or more ANC visits and were from the richest wealth quintile were three [AOR=3.83; 95% CI=2.77-5.30] and 15 times [AOR=15.69; 95% CI= 5.53-44.50] more likely to be helped by a health professional, respectively.

Women whose husband worked in a non-agricultural occupation [AOR=1.35; 95% CI=1.00-1.81] were more likely to deliver with a health professional. A similar result was

Table 2. Bivariate analysis of women's characteristics and facility-based delivery outcome.

Variable	Non-health facility		Health facility		X²
	N	%	N	%	
Age difference between man and woman					
Woman older than man	117	26.1	333	73.9	17.23**
0–4 years	262	29.4	628	70.6	
5–7 years	108	24.8	326	75.2	
>7 years	127	32.8	261	67.2	
Birth rank and interval					
Second or third child, interval >2 years	241	24.9	729	75.1	200.57***
First birth	124	20.6	479	79.4	
Second or third child, interval ≤2 years	48	36.8	83	63.2	
Fourth or higher child, interval >2 years	156	40.6	228	59.4	
Fourth or higher child, interval ≤2 years	45	60.2	29	39.8	
Planning status of births					
Then	513	28.9	1261	71.1	3.51
Later	68	27.8	178	72.2	
No more	34	23.9	108	76.1	
Husband/partner's education attainment					
Incomplete primary education/none	143	47.9	156	52.1	233.59***
Complete primary or some secondary	277	32.9	564	67.1	
Completed secondary or higher	194	19.0	828	81.0	
Husband/partner's occupation					
Agricultural	423	36.2	746	63.8	162.94***
Non-agricultural	191	19.2	802	80.8	
Wealth quintile					
Poorest	423	46.8	480	53.2	607.20***
Poorer	105	23.2	349	76.8	
Middle	45	14.3	270	85.7	
Richer	29	11.3	224	88.7	
Richest	13	5.6	224	94.4	
Number of household members					
<4 members	67	25.2	201	74.8	3.40
≥4 members	547	28.9	1,347	71.1	
Number of antenatal care visits					
<4 visits	153	61.6	98	38.4	328.54***
≥4 visits	461	24.1	1,450	75.9	
Covered by health insurance					
No	242	31.3	530	68.7	10.16*
Yes	373	26.8	1,017	73.2	
Residence					
Urban	88	12.2	631	87.8	296.56***
Rural	527	36.5	916	63.5	

Variable	Non-health facility		Health facility		X ²
	N	%	N	%	
Composite labor force participation					
Poor	248	35.8	444	64.2	123.78***
Moderate	221	31.7	477	68.3	
Good	145	18.8	627	81.2	
Attitude toward domestic violence					
Poor	237	35.1	437	64.9	46.49***
Moderate	194	25.8	559	74.2	
Good	184	25.0	551	75.0	
Decision making power					
Poor	218	28.5	546	71.5	0.64
Moderate	195	27.7	508	72.3	
Good	202	29.0	493	71.0	
Women's knowledge level					
Poor	290	38.8	458	61.2	145.48***
Moderate	187	26.2	527	73.8	
Good	137	19.6	563	80.4	

*p<0.05; **p<0.01; ***p<0.001

Table 3. Bivariate analysis of women's characteristics and assisted delivery outcome.

Variable	Non-health professional		Health facility		X²
	N	%	N	%	
Age difference between man and woman					
Woman older than man	56	12.4	394	87.6	7.55
0–4 years	125	14.0	765	86.0	
5–7 years	54	12.5	380	87.5	
>7 years	63	16.3	325	83.7	
Birth rank and interval					
Second or third child, interval >2 years	113	11.6	857	88.4	136.16***
First birth	53	8.8	550	91.2	
Second or third child, interval ≤2 years	22	16.6	109	83.4	
Fourth or higher child, interval >2 years	87	22.7	297	77.3	
Fourth or higher child, interval ≤2 years	24	31.9	50	68.1	
Planning status of births					
Then	254	14.3	1,520	85.7	5.43
Later	29	11.8	217	88.2	
No more	15	10.5	127	89.5	
Husband/partner's education attainment					
Incomplete primary education/none	106	35.4	193	64.6	367.17***
Complete primary or some secondary	130	15.5	711	84.5	
Completed secondary or higher	61	6.0	961	94.0	
Husband/partner's occupation					

Variable	Non-health professional		Health facility		X ²
	N	%	N	%	
Agricultural	229	19.6	940	80.4	155.85***
Non-agricultural	69	6.9	924	93.1	
Wealth quintile					
Poorest	243	26.9	660	73.1	
Poorer	39	8.6	415	91.4	509.25***
Middle	11	3.4	304	96.6	
Richer	3	1.3	250	98.7	
Richest	2	0.7	235	99.3	
Number of household members					
<4 members	30	11.3	238	88.7	3.24
≥4 members	267	14.1	1,627	85.9	
Number of antenatal care visits					
<4 visits	102	40.5	149	59.5	364.74***
≥4 visits	197	10.3	1,714	89.7	
Covered by health insurance					
No	121	15.7	651	84.3	8.03*
Yes	177	12.7	1,213	87.3	
Residence					
Urban	38	5.3	681	94.7	139.61***
Rural	260	18.0	1,183	82.0	
Composite labor force participation					
Poor	125	18.0	567	82.0	
Moderate	119	17.0	579	83.0	95.35***
Good	56	7.2	716	92.8	
Attitude toward domestic violence					
Poor	111	16.5	563	83.5	20.25**
Moderate	106	14.1	647	85.9	
Good	80	10.9	655	89.1	
Decision-making power					
Poor	119	15.6	645	84.4	7.34
Moderate	92	13.1	611	86.9	
Good	86	12.4	609	87.6	
Women's knowledge level					
Poor	180	24.1	568	75.9	231.52***
Moderate	74	10.4	640	89.6	
Good	43	6.1	657	93.9	

*p<0.05; **p<0.01; ***p<0.001

Table 4. Binary logistic regression analysis with a facility-based delivery outcome.

Variable	AOR	CI	
		Lower	Upper
Age difference between man and woman			
Woman older than man	Ref		
0–4 years	0.88	0.69	1.11
5–7 years	1.14	0.88	1.46
>7 years	0.96	0.73	1.26
Birth rank and interval			
Second or third child, interval >2 years	Ref		
First birth	1.37**	1.09	1.71
Second or third child, interval ≤2 years	0.60**	0.43	0.84
Fourth or higher child, interval >2 years	0.66***	0.52	0.82
Fourth or higher child, interval ≤2 years	0.30***	0.19	0.47
Planning status of births			
Then	Ref		
Later	1.01	0.77	1.32
No more	1.48*	1.05	2.08
Husband/partner's education attainment			
Incomplete primary education/ none	Ref		
Complete primary or some secondary	1.23	0.94	1.60
Completed secondary or higher	1.26	0.94	1.69
Husband/partner's occupation			
Agricultural	Ref		
Non-agricultural	1.17	0.93	1.47
Wealth quintile			
Poorest	Ref		
Poorer	1.94***	1.53	2.47
Middle	2.85***	2.06	3.94
Richer	3.23***	2.25	4.64
Richest	5.59***	3.37	9.30
Number of household members			
<4 members	Ref		
≥4 members	1.02	0.75	1.39
Number of antenatal care visits			
<4 visits	Ref		
≥4 visits	3.62***	2.73	4.79

Variable	AOR	CI	
		Lower	Upper
Covered by health insurance			
No	Ref		
Yes	1.15	0.95	1.38
Residence			
Urban	Ref		
Rural	0.49***	0.36	0.66
Composite labor force participation			
Poor	Ref		
Moderate	1.14	0.93	1.40
Good	1.47**	1.15	1.89
Attitude toward domestic violence			
Poor	Ref		
Moderate	1.38**	1.10	1.73
Good	1.33*	1.04	1.69
Decision-making power			
Poor	Ref		
Moderate	0.85	0.69	1.06
Good	0.84	0.66	1.06
Women's knowledge level			
Poor	Ref		
Moderate	1.13	0.92	1.40
Good	1.17	0.93	1.47

*p<0.05; **p<0.01; ***p<0.001

AOR, adjusted odds ratio; CI, confidence interval.

found for women with good composite labor force participation [AOR=1.58; 95% CI=1.11–2.26] and a good level of knowledge [AOR=1.76; 95% CI=1.25–2.46] (Table 5).

Discussion

Delivery was regarded as safe when it was attended by a skilled birth attendant and took place in a health facility. This study found that several variables have a significant association with facility-based delivery and assisted delivery. Women from the richest wealth quintile were more likely to have a delivery in a health facility than those from the poorest wealth quintile. This finding is consistent with that of a previous study in Indonesia. The wealth index of the household would contribute to the access to health care services, including institutional delivery (Caulfield *et al.*, 2016; Do *et al.*, 2015; Efendi *et al.*, 2019; Roro *et al.*, 2014). Women from the richest wealth

Table 5. Binary logistic regression analysis with assisted delivery.

Variable	AOR	CI	
		Lower	Upper
Age difference between man and woman			
Woman older than man	Ref		
0-4 years	0.89	0.65	1.22
5-7 years	0.99	0.71	1.36
>7 years	1.01	0.72	1.42
Birth rank and interval			
Second or third child, interval >2 years	Ref		
First birth	1.28	0.93	1.76
Second or third child, interval ≤2 years	0.80	0.48	1.33
Fourth or higher child, interval >2 years	0.73	0.53	1.00
Fourth or higher child, interval ≤2 years	0.49**	0.29	0.83
Planning status of births			
Then	Ref		
Later	1.12	0.74	1.68
No more	1.58	0.96	2.59
Husband/partner's education attainment			
Incomplete primary education/ none	Ref		
Complete primary or some secondary	1.90***	1.42	2.55
Completed secondary or higher	2.18***	1.48	3.22
Husband/partner's occupation			
Agricultural	Ref		
Non-agricultural	1.35*	1.00	1.81
Wealth quintile			
Poorest	Ref		
Poorer	2.33***	1.70	3.18
Middle	5.14***	3.19	8.28
Richer	10.84***	5.34	22.01
Richest	15.69***	5.53	44.50
Number of household members			
<4 members	Ref		
≥4 members	0.87	0.56	1.35
Number of antenatal care visits			
<4 visits	Ref		
≥4 visits	3.83***	2.77	5.30

Variable	AOR	CI	
		Lower	Upper
Covered by health insurance			
No	Ref		
Yes	1.19	0.94	1.51
Residence			
Urban	Ref		
Rural	0.84	0.54	1.30
Composite labor force participation			
Poor	Ref		
Moderate	0.99	0.76	1.29
Good	1.58*	1.11	2.26
Attitude toward domestic violence			
Poor	Ref		
Moderate	1.24	0.94	1.63
Good	1.30	0.96	1.77
Decision-making power			
Poor	Ref		
Moderate	0.89	0.69	1.14
Good	0.99	0.73	1.34
Women's knowledge level			
Poor	Ref		
Moderate	1.54**	1.13	2.11
Good	1.76**	1.25	2.46

*p<0.05; **p<0.01; ***p<0.001

AOR, adjusted odds ratio; CI, confidence interval.

quintile were more likely to have a delivery with a skilled assistant than those from the poorest wealth quintile. This finding is consistent with that of a previous study conducted in Bangladesh (Muhammed *et al.*, 2017). Women from low-income families may find it difficult to pay for a skilled assistant delivery, so they prefer to give birth without professional assistance (Muhammed *et al.*, 2017). Therefore, the coverage of health insurance must be enhanced so that women in all the wealth quintiles can have equal access to health care services.

A higher child rank and interval of ≤2 years was associated with a lower chance of women having a delivery in a health facility and being assisted by a health professional. This result is similar to those of studies conducted in Ethiopia and Nigeria (Abera *et al.*, 2011; Ononokpono & Odimegwu, 2014). Women with a higher child rank will have more experience with pregnancy and delivery, so they feel that they have the confidence to have a delivery outside a health facility

(Abera *et al.*, 2011). Another argument is that women have limited access to health services due to the burden of their economic situation (Ononokpono & Odimegwu, 2014). The results for delivery with a skilled birth attendant are similar to those of studies conducted in Sudan and Ethiopia (Mustafa & Mukhtar, 2015; Wilunda *et al.*, 2015). Women with a higher birth rank tend to rely on their experience from previous pregnancies, believing they already know about childbirth. Consequently, they choose to give birth without professional assistance (Mustafa & Mukhtar, 2015). Review studies conducted in African countries also highlighted the link between higher parity and lower likelihood of facility-based delivery (Moyer & Mustafa, 2013). Therefore, health education about safe delivery should prioritize mothers with a high child rank by giving them greater access to free health care services.

Women who had more than four ANC visits during their pregnancy were found to be three times more likely to have a safe delivery. This is consistent with the results of studies conducted in Uganda and Ethiopia (Abera *et al.*, 2011; Atusiimire *et al.*, 2019). Furthermore, a population-based study conducted in Bangladesh had a similar result, which emphasized the positive effect of the ANC on utilization of health facility-based delivery (Pervin *et al.*, 2012). The ANC can prevent unsafe delivery because it will provide health education for the mother, giving information and recommending the place of delivery according to the mother's and fetus's condition (Atusiimire *et al.*, 2019). Women who had more than four ANC visits during their pregnancy were found to be more likely to have a skilled assistant delivery. This is consistent with the result of a study that was conducted in Kenya (Gitimu *et al.*, 2015). ANC attendance will influence the decision of the mother to have an assisted delivery because the ANC emphasizes the importance of safe delivery (Gitimu *et al.*, 2015). ANC visits must be optimized for pregnant women so that mothers are more exposed to information about safe delivery. The information that the mother receives will influence the decision on where to deliver the baby. Therefore, a minimum number of ANC visits should be given to all pregnant women so they can monitor the condition of the baby and have more knowledge about safe pregnancy and delivery.

Another finding was that women who wanted no more pregnancies when they became pregnant were more likely to give birth in a health facility. This finding is consistent with that of a study conducted in Egypt (Marston & Cleland, 2003). However, it is inconsistent with the results of a study conducted in Bangladesh, which showed that women who have an unintended pregnancy were less likely to visit an ANC service and more likely to have a home delivery (Kamal, 2013). There was no study that explained this issue, as it may be related to the social norms and health system of the country itself. Therefore, this topic should be analyzed further by considering other variables.

Women from a rural residence were found to be less likely to have a delivery in a health facility. This finding is simi-

lar to those of previous studies conducted in Bangladesh and Indonesia (Efendi *et al.*, 2019; Kamal, 2013; Kenea & Jisha, 2017). Living in an urban residence allows easier access to health facilities than living in rural areas. In addition, access to information is easier in urban areas so information about safe delivery can be spread more easily (Kamal, 2013). Therefore, the gap between rural and urban areas should be taken into consideration by the government regarding the issue of maternal and child health.

Women who had good composite labor force participation and a good attitude towards domestic violence were more likely to have a delivery in a health facility and be assisted by a health professional. This is consistent with the result of studies conducted in Ethiopia and Bangladesh (Kamal, 2013; Tiruneh *et al.*, 2017). In Bangladesh, women who were against domestic violence and more independent economically were more likely to have four or more ANC service visits, which may lead the women to have a delivery in a health facility (Kamal, 2013). Women who had good composite labor force participation and a good knowledge level were more likely to have a delivery with a skilled assistant. This is consistent with the result of studies conducted in Senegal and Tanzania (Shimamoto & Gipson, 2015). If women have greater empowerment, in terms of knowledge and economic power, this will lead to improvement in their health. They can choose the best for their health, including choosing to have a safe delivery (Prata *et al.*, 2017). Therefore, gender equality needs to be improved so women can make decisions about their health.

Husband/partner's education attainment was found to be significantly associated with skilled assistant delivery. Women whose husband/partner had completed secondary or higher education were more likely to have a skilled assistant delivery. This is consistent with the results of studies conducted in Kenya and Somalia (Gitimu *et al.*, 2015; Yusuf *et al.*, 2017). Husbands with a higher level of education will have more knowledge about health, including safe delivery (Kifle *et al.*, 2018). As the head of the household, the husband's knowledge will affect the reproductive health decisions (Yusuf *et al.*, 2017). Engagement of the husband in the issue of maternal health should be expanded in all levels of the community.

Women whose husband/partner's occupation was non-agricultural were found to be significantly more likely to have a delivery with professional assistance. This finding is similar to those of studies conducted in Nigeria and Ethiopia (Adewemimo *et al.*, 2014; Fekadu & Regassa, 2015). The husband's occupation will affect the family's income. If the family income increases, the decision to have a skilled assistant delivery will also be affected. In addition, another study conducted in Ethiopia found that women whose husbands work in a non-agricultural occupation tend to use an ANC service, which encourages the decision to give birth with professional assistance (Tsegay *et al.*, 2013). Therefore, health promotion about safe delivery is important for the husband/partner, especially for husbands/partners whose occupation is agricultural.

Limitations and strengths

This study used secondary data from the 2017 IDHS, so the selection of the variables was determined by the availability of the data. Another limitation is that some questions in the survey needed respondents to recall what happened five years preceding the survey, so the information may not be precisely stated. In addition to the limitations, however, this study has strengths. The sample of this study was selected using two-stage cluster sampling, so the data were nationally representative. Therefore, the results can provide recommendations for policymakers to develop effective regulation so the coverage gap of safe delivery between western and eastern provinces in Indonesia can be reduced.

Conclusions

Safe delivery was found to be determined by several factors, which reflected the need for multi-stakeholder intervention in increasing the practice of safe delivery across the country. Programmatic and structured policies that target poor women and those with a low education level and encourage husbands/

partners' participation in this issue may help increase the prevalence of safe delivery in the eastern part of Indonesia. This study gives some recommendations to the policymakers, such as health promotion about safe delivery should be prioritized for women who have a high birth rank. Moreover, not only the women, but also their husband/partner should be involved in health education. The coverage of health insurance and health facilities should be enhanced so that everyone can have equal access to health services. Furthermore, the women's empowerment program should be maximized so that all the women can choose the best for their health.

Data availability

Data used in this study is available online from the Indonesian 2017 Demographic and Health Survey (DHS) website under the 'Individual Recode' section. Access to the dataset requires registration and is granted only for legitimate research purposes. A guide for how to apply for dataset access is available at: <https://dhsprogram.com/data/Access-Instructions.cfm>.

References

- Abera M, Gebremariam A, Belachew T: **Predictors of safe delivery service utilization in arsi zone, South-East Ethiopia**. *Ethiop J Health Sci*. 2011; 21(Suppl 1): 95–106.
[PubMed Abstract](#) | [Free Full Text](#)
- Adewemimo AW, Msuya SE, Olaniyan CT, et al.: **Utilisation of skilled birth attendance in Northern Nigeria: a cross-sectional survey**. *Midwifery*. 2014; 30(1): e7–e13.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Atusiimire LB, Waiswa P, Atuyambe L, et al.: **Determinants of facility based-deliveries among urban slum dwellers of Kampala, Uganda**. *PLoS One*. 2019; 14(4): e0214995.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Belton S, Myers B, Ngana FR: **Maternal deaths in eastern Indonesia: 20 years and still walking: an ethnographic study**. *BMC Pregnancy Childbirth*. 2014; 14: 39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- BPS: **Angka Kematian Ibu Menurut Pulau (per 100.000 kelahiran hidup)**. 2015a.
[Reference Source](#)
- BPS: **Hasil Survei Penduduk Antar Sensus 2015 [Result of the 2015 Intercensal Population Census]**. Jakarta: Badan Pusat Statistik. 2015b.
- BPS, BKKBN, RI KK, ICF: **Indonesia Demographic and Health Survey 2002**. 2018.
- Caulfield T, Onyo P, Byrne A, et al.: **Factors influencing place of delivery for pastoralist women in Kenya: a qualitative study**. *BMC Womens Health*. 2016; 16(1): 52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Do M, Soelaeman R, Hotchkiss DR: **Explaining inequity in the use of institutional delivery services in selected countries**. *Matern Child Health J*. 2015; 19(4): 755–763.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Efendi F, Ni'mah AR, Hadisuyatmana S, et al.: **Determinants of Facility-Based Childbirth in Indonesia**. *ScientificWorldJournal*. 2019; 2019: 1–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fekadu M, Regassa N: **Skilled delivery care service utilization in Ethiopia: analysis of rural-urban differentials based on national demographic and health survey (DHS) data**. *Afr Health Sci*. 2014; 14(4): 974–984.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gitimu A, Herr C, Oruko H, et al.: **Determinants of use of skilled birth attendant at delivery in Makueni, Kenya: a cross sectional study**. *BMC Pregnancy Childbirth*. 2015; 15: 9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Iacoella F, Tirivayi N: **Determinants of maternal healthcare utilization among married adolescents: Evidence from 13 Sub-Saharan African countries**. *Public Health*. 2019; 177: 1–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- ICF Macro: **Demographic and Health Survey Interviewer's Manual**. Rockville: ICF International. 2020.
[Reference Source](#)
- Kamal SM: **Preference for institutional delivery and caesarean sections in Bangladesh**. *J Health Popul Nutr*. 2013; 31(1): 96–109.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kenea D, Jisha H: **Urban-rural disparity and determinants of delivery care utilization in Oromia region, Ethiopia: Community-based cross-sectional study**. *Int J Nurs Pract*. 2017; 23(1).
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kifle MM, Kesete HF, Gaim HT, et al.: **Health facility or home delivery? Factors influencing the choice of delivery place among mothers living in rural communities of Eritrea**. *J Health Popul Nutr*. 2018; 37(1): 22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kurniati A, Chen CM, Efendi F, et al.: **Factors influencing Indonesian women's use of maternal health care services**. *Health Care Women Int*. 2018; 39(9): 1–16.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Marston C, Cleland J: **Do unintended pregnancies carried to term lead to adverse outcomes for mother and child? An assessment in five developing countries**. *Popul Stud (Camb)*. 2003; 57(1): 77–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
- MoH: **Luas Wilayah dan Kondisi Geografis Menjadi Tantangan Besar Pembangunan Kesehatan Provinsi Papua - Sehat Negeriku**. 2012.
[Reference Source](#)
- Moyer CA, Mustafa A: **Drivers and deterrents of facility delivery in sub-Saharan Africa: a systematic review**. *Reprod Health*. 2013; 10: 40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Muhammed G, Kibria A, Ghosh S, et al.: **Factors affecting deliveries attended by skilled birth attendants in Bangladesh**. *Matern Health Neonatol Perinatol*. 2017; 3: 7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mustafa MH, Mukhtar AM: **Factors associated with antenatal and delivery care in Sudan: analysis of the 2010 Sudan household survey**. *BMC Health Serv Res*. 2015; 15: 452.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nations U: **About the Sustainable Development Goals**. 2015.
[Reference Source](#)
- Ononokpono DN, Odimegwu CO: **Determinants of Maternal Health Care Utilization in Nigeria: a multilevel approach**. *Pan Afr Med J*. 2014; 17(Suppl 1): 2.
[PubMed Abstract](#) | [Free Full Text](#)
- Organizations WH: **WHO Safe Childbirth Checklist**. In *WHO*. Geneva: World

Health Organization. 2017.

[Reference Source](#)

Organizations WH: **Maternal mortality**. 2018.

[Reference Source](#)

Pervin J, Moran A, Rahman M, *et al.*: **Association of antenatal care with facility delivery and perinatal survival - a population-based study in Bangladesh**. *BMC Pregnancy Childbirth*. 2012; **12**(1): 111.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Prata N, Tavrow P, Upadhyay U: **Women's empowerment related to pregnancy and childbirth: introduction to special issue**. *BMC Pregnancy Childbirth*. 2017; **17**(Suppl 2): 352.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Kementrian Kesehatan RI: **Rencana Strategis Kementerian Kesehatan 2015-2019**. 2017.

[Reference Source](#)

Kementrian Kesehatan RI: **Situasi Kesehatan Ibu**. In *Buletin Jendela Data & Informasi Kesehatan*. Jakarta: Pusat Data Kemenkes RI. 2014.

[Reference Source](#)

Roro MA, Hassen EM, Lemma AM, *et al.*: **Why do women not deliver in health facilities: a qualitative study of the community perspectives in south central Ethiopia?** *BMC Res Notes*. 2014; **7**(1): 556.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Rutstein SO, Johnson K: **The DHS wealth index**. In *DHS Comparative Reports No. 6*. 2004.

[Reference Source](#)

Sebayang SK, Efendi F, Astutik E: **Women's empowerment and the use of**

antenatal care services: analysis of demographic health surveys in five Southeast Asian countries. *Women Health*. 2019; **59**(10): 1155–1171.

[PubMed Abstract](#) | [Publisher Full Text](#)

Shimamoto K, Gipson JD: **The relationship of women's status and empowerment with skilled birth attendant use in Senegal and Tanzania**. *BMC Pregnancy Childbirth*. 2015; **15**(1): 154.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Tejayanti T, Harimat, Sabatini K, *et al.*: **Disparitas Akses dan Kualitas: Kajian Determinan Kematian Maternal di Lima Region Indonesia**. Jakarta: MoH-UNFPA. 2012.

Tiruneh FN, Chuang KY, Chuang YC: **Women's autonomy and maternal healthcare service utilization in Ethiopia**. *BMC Health Serv Res*. 2017; **17**(1): 718.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Tsegay Y, Gebrehiwot T, Goicolea I, *et al.*: **Determinants of antenatal and delivery care utilization in Tigray region, Ethiopia: a cross-sectional study**. *Int J Equity Health*. 2013; **12**(1): 30.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Wilunda C, Quaglio G, Putoto G, *et al.*: **Determinants of utilisation of antenatal care and skilled birth attendant at delivery in South West Shoa Zone, Ethiopia: a cross sectional study**. *Reprod Health*. 2015; **12**: 74.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Yusuf MS, Kodhiambo M, Muendo F: **Determinants of Access to Skilled Birth Attendants by Women in Galkacyo District, Somalia**. *Asian Journal of Medicine and Health*. 2017; **4**(4): 1–7.

[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 21 August 2020

<https://doi.org/10.5256/f1000research.25747.r66055>

© 2020 Saleh A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Asmaa Salah Eldin Mohamed Saleh

Community Health Nursing, Faculty of Nursing, Beni Suef University, Beni Suef, Egypt

This is a good written article, but there are little things I think to make it perfect if the author follows it.

First: in Introduction: the author does not mention study hypotheses. Explain why the current research is important.

What value does the paper add?

What practical benefits will it provide?

Second: Sample and sampling, it is important to add and mention the procedure of taking sampling for this huge sample size.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: community health nursing, geriatric health, MCH, health promotion, school health, pediatrics, maternity

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 26 May 2020

<https://doi.org/10.5256/f1000research.25747.r63312>

© 2020 Oducado R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Ryan Michael Flores Oducado

College of Nursing, West Visayas State University, Iloilo City, Philippines

This article looks into the determinants of safe delivery in the eastern part of Indonesia. The report addresses an important topic and contributes to the advancement of knowledge on maternal health and in reducing maternal mortality.

I only have a few comments regarding the research article:

1. The authors mentioned that the dependent variables were facility-based delivery and assisted delivery. Will it be more appropriate to consider a) place of delivery and b) proportion of women assisted by skilled birth attendants as dependent variables?
2. It was reported in the article that 71.6% used a health facility and 86.2% were assisted by a health professional at their last birth. I wonder, isn't that health professionals typically assist health facility delivery? What is the reason for the discrepancy?
3. Please clarify women's knowledge level. This variable pertains to women's level of knowledge about what?
4. An interesting finding in this study was that women who wanted no more pregnancies when they became pregnant were more likely to give birth in a health facility. The authors mentioned that it might be related to the social norms and health system of the country itself. Can the authors expound on this? Can the researchers offer their plausible explanation of this result?
5. Can the authors restate the conclusions in the abstract? Can household factors be changed to a more encompassing term reflective of the result? Can the facility-based delivery term be deleted and focus on safe delivery as directed in the title of the article?
6. Can the use of secondary data be mentioned in the abstract?

Thank you very much for the opportunity to review this article.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Public Health and Community Health Nursing

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 04 Sep 2020

Ferry Efendi, Faculty of Nursing, Universitas Airlangga, Surabaya, Indonesia

We thank all reviewers for their generous and positive comments on our manuscript. With regard to their reviews, we have edited the manuscript to respond to their comments. In particular, we have made updates in track changes to indicate changes as suggested. We thank you for the opportunity to submit our revised manuscript.

Reviewer 1

This article looks into the determinants of safe delivery in the eastern part of Indonesia. The report addresses an important topic and contributes to the advancement of knowledge on maternal health and in reducing maternal mortality.

I only have a few comments regarding the research article:

1. The authors mentioned that the dependent variables were facility-based delivery and assisted delivery. Will it be more appropriate to consider a) place of delivery and b) proportion of women assisted by skilled birth attendants as dependent variables?

Thank you for these constructive comments, we have changed the terminology into place of delivery (health facility Vs non-health facility) and type of assistance at delivery (skilled birth attendants Vs unskilled birth attendants) as guided by guide to DHS statistics DHS-7 and

your comments.

2. It was reported in the article that 71.6% used a health facility and 86.2% were assisted by a health professional at their last birth. I wonder, isn't that health professionals typically assist health facility delivery? What is the reason for the discrepancy?

Thank you for raising this issue, all delivery in health facility must be assisted by health professionals. However, not all delivery assisted by health professionals occurred in health facility.

3. Please clarify women's knowledge level. This variable pertains to women's level of knowledge about what?

Thank you very much, we have added the explanation in variables section.

4. An interesting finding in this study was that women who wanted no more pregnancies when they became pregnant were more likely to give birth in a health facility. The authors mentioned that it might be related to the social norms and health system of the country itself. Can the authors expound on this? Can the researchers offer their plausible explanation of this result?

Thank you very much, we have added the explanation in discussion section.

5. Can the authors restate the conclusions in the abstract? Can household factors be changed to a more encompassing term reflective of the result? Can the facility-based delivery term be deleted and focus on safe delivery as directed in the title of the article?

Thank you very much, we have revised the abstract.

6. Can the use of secondary data be mentioned in the abstract?

Thank you very much, we have added in the abstract section.

Competing Interests: None

Author Response 04 Sep 2020

Ferry Efendi, Faculty of Nursing, Universitas Airlangga, Surabaya, Indonesia

We thank all reviewers for their generous and positive comments on our manuscript. With regard to their reviews, we have edited the manuscript to respond to their comments. In particular, we have made updates in track changes to indicate changes as suggested. We thank you for the opportunity to submit our revised manuscript.

Reviewer 2

1. This is a good written article, but there are little things I think to make it perfect if the author follows it.

First: in Introduction: the author does not mention study hypotheses. Explain why the current research is important. What value does the paper add? What practical benefits will it provide?

Thank you very much, we have added on introduction section for the hypothesis and the value. While, we have added the practical benefits on the limitation and strength section.

2. Second: Sample and sampling, it is important to add and mention the procedure of taking sampling for this huge sample size.

Thank you very much, we have added on sample size and sampling section.

Competing Interests: None

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research