REVIEW

Understanding transport processes in lichen, *Azolla*–cyanobacteria, ectomycorrhiza, endomycorrhiza, and rhizobia–legume symbiotic interactions [version 1; peer review: 3 approved]

Rahul Roy¹, Anke Reinders², John M Ward¹, Tami R McDonald³

¹Department of Plant and Microbial Biology, University of Minnesota, Minnesota, USA

²College of Continuing and Professional Studies, University of Minnesota, Minnesota, USA

³Biology Department, St. Catherine University, Minnesota, USA

V1 First published: 23 Jan 2020, 9(F1000 Faculty Rev):39
<https://doi.org/10.12688/f1000research.19740.1>

Latest published: 23 Jan 2020, 9(F1000 Faculty Rev):39
<https://doi.org/10.12688/f1000research.19740.1>

Abstract

Intimate interactions between photosynthetic and non-photosynthetic organisms require the orchestrated transfer of ions and metabolites between species. We review recent progress in identifying and characterizing the transport proteins involved in five mutualistic symbiotic interactions: lichens, *Azolla*–cyanobacteria, ectomycorrhiza, endomycorrhiza, and rhizobia–legumes. This review focuses on transporters for nitrogen and carbon and other solutes exchanged in the interactions. Their predicted functions are evaluated on the basis of their transport mechanism and prevailing transmembrane gradients of H⁺ and transported substrates. The symbiotic interactions are presented in the assumed order from oldest to most recently evolved.

Keywords

transporters, endomycorrhiza, ectomycorrhiza, rhizobia-legume, *Azolla*-cyanobacteria, lichens

Open Peer Review

Approval Status

	1	2	3
version 1 23 Jan 2020			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Stefanie Wienkoop**, University of Vienna, Vienna, Austria
2. **Kendal Hirschi**, Baylor College of Medicine, Houston, USA
3. **Ulrike Mathesius**, The Australian National University, Canberra, Australia

Any comments on the article can be found at the end of the article.

Corresponding author: John M Ward (jward@umn.edu)

Author roles: **Roy R:** Writing – Original Draft Preparation, Writing – Review & Editing; **Reinders A:** Writing – Original Draft Preparation, Writing – Review & Editing; **Ward JM:** Writing – Original Draft Preparation, Writing – Review & Editing; **McDonald TR:** Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Roy R *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Roy R, Reinders A, Ward JM and McDonald TR. **Understanding transport processes in lichen, *Azolla*–cyanobacteria, ectomycorrhiza, endomycorrhiza, and rhizobia–legume symbiotic interactions [version 1; peer review: 3 approved]** F1000Research 2020, 9(F1000 Faculty Rev):39 <https://doi.org/10.12688/f1000research.19740.1>

First published: 23 Jan 2020, 9(F1000 Faculty Rev):39 <https://doi.org/10.12688/f1000research.19740.1>

Introduction

The vast majority of plant species participate in symbiotic relationships with fungi and bacteria. We know that the formation of symbiotic relationships between species occurred before the first eukaryotic cells which themselves are the product of endosymbiosis that resulted in the formation of mitochondria and chloroplasts. Understanding transmembrane transport processes between species is the key to understanding nutritional symbiotic interactions. This is because direct transfer from cell to cell between species is rare, although there are examples such as secretion between *Agrobacterium* and plant cells. In nutritional symbioses, cells of both species are bounded by a cell membrane and transfer of ions and metabolites occurs through membrane-bound transport proteins. Identifying the transporters involved and characterizing their activity can reveal which ions and metabolites are transported between species.

There has always been great interest in understanding symbiotic systems such as mycorrhiza and lichens, but research at the molecular level has often been too difficult. However, this is an appropriate time to revisit such complex systems and identify transport proteins that function in symbiotic interactions. Technical advances, especially the availability of genome sequences, have had an exceptionally large impact on the ability to study symbiotic interactions. Now, transcriptomic and proteomic experiments involving multiple interacting species can be performed and transporters can be assigned to each interacting partner. Once transporters are identified, heterologous expression experiments can be used to study transport activity, fluorescent protein fusions can be used to localize membrane proteins, and mutational analysis can be used to test whether transporter function is important for the interaction.

In the symbiotic associations discussed in this review, the transport of fixed nitrogen (N) and fixed carbon (C) compounds takes center stage. The reason is that all organisms need large amounts of N and C; photosynthetic plants can fix C, but only bacteria (and Archaea) can fix N. Cyanobacteria can fix both N and C but no eukaryotic organism can do the same. In the symbioses discussed here, large amounts of fixed N and C are transported between species. We will discuss transporters for N and C that have been identified and their activity and point out possibilities for transporters that have not yet been identified. We will also discuss selected transporters for other nutrients, ions, metals, and metabolites that are essential to symbiotic interactions.

To identify novel transporters functioning in a symbiosis, it is important to understand that the prevailing negative membrane potential, the transmembrane pH gradient that is more acidic outside, and the gradient of substrate may narrow the types of transporters likely to serve a particular function. For example, owing to the negative membrane potential, anion exporters are likely to be uniporters, such as anion channels. Exporters for cations are likely to be H⁺-coupled antiporters or pumps. Anion uptake transporters are likely to be H⁺-coupled symporters, and cation uptake transporters are likely to be H⁺-coupled

symporters or channels. For uncharged molecules, efflux transporters could be uniporters, H⁺-coupled antiporters, or pumps, whereas uptake transporters are likely to be H⁺-coupled symporters. It is also important to understand the orientation of membranes at the symbiotic interface. For all of the interactions discussed here, in order for ions and metabolites to be transferred from one organism to another, a transmembrane export step followed by an uptake step is required.

Lichens

Lichens are a morphologically and taxonomically diverse assemblage of obligate symbioses involving fungi and photosynthesizers. A typical lichen consists of a tough upper and lower fungal cortex protecting a symbiotic interaction zone consisting of a layer of algal or cyanobacterial cells supported by a network of fungal hyphae (Figure 1). The entirety of this structure is called a thallus. Some lichens harbor two photosynthesizers, often a green alga as the main photobiont and a secondary cyanobacterial photobiont sequestered in wart-like growths called cephalodia. Close to 20% of currently described fungal species form lichens^{1,2}.

Seminal radiolabeling experiments in the 1960s and 1970s established the nature of the carbohydrate transfer from the photosynthesizer to the fungus (3 and references therein). These experiments demonstrated that (1) a single C source is exported from the photobiont, (2) green algae transfer the polyol ribitol (in some cases, erythritol or sorbitol) whereas cyanobacteria transfer glucose, and (3) the transfer appears to be dependent on contact with, or signals from, the fungus, as the photobiont in pure culture does not excrete fixed C even as soon as 24 to 48 hours after isolation. Upon entry into the fungal cells, C is rapidly metabolized into mannitol, possibly through an intermediate such as arabinol. Cyanobacterial photobionts can also fix N into ammonia in specialized cells common to all multicellular cyanobacteria, called heterocysts or heterocytes, and transfer it to the fungus⁴.

Genomes and transcriptomes are enabling a new understanding of the transfer of C and N in lichens. Genomes of several lichenizing fungi have been sequenced⁵⁻¹¹. Although it is not yet possible to consistently reconstitute any mature lichen symbiosis in culture, the transcriptomes of the separated symbionts^{12,13}, of the early stages of symbiotic interaction when cultures of the isolated symbionts are mixed¹⁴, or of mature lichen symbioses in various metabolic states such as desiccation or rehydration¹⁵ allow the identification of potentially important transporters of C and N. These excellent first forays set the stage for molecular dissection of the lichen symbiosis.

A current priority is testing of hypotheses flowing from genome and transcriptome experiments, such as functional characterization of the putative ribitol transporters identified by Yoshino *et al.*¹⁶ in a lichenizing fungus. Interestingly, no exporter/importer pair for any of the symbiotically exchanged nutrients has yet been defined in any lichen system. Also missing from the discussion are sugar exporters in algae and cyanobacteria that could be *sugars will eventually be exported transporters*

Figure 1. Transporters in the lichen symbiosis. Left: cross-section of a lichen thallus with both cyanobacterial and algal photobionts. Left: The cephalodium is a structure that houses the cyanobacterial cells. The cyanobacterial vegetative cells are shown in green, and the heterocysts are shown in white. "Tomentum" refers to the fungal area composed of closely matted mycelia. Right: Detail of symbiotic interface showing interaction of fungal cells with photosynthetic cells. Transporter activities that do not have an identified transporter are indicated with a question mark. Red transporter, C transporter (glucose or polyols such as ribitol); blue transporter, N transporter (ammonium).

(*SWEETs*) or *semiSWEETs*. Likewise, analysis of ammonium transporters (*AMTs*) and amino acid transporters as the conduits of symbiotically exchanged N is also necessary.

Azolla

The small aquatic fern *Azolla filiculoides* (and related species) has a mutualistic relationship with the cyanobacterium *Nostoc azollae* (*Anabaena azollae*) and traditionally has been employed as a biofertilizer in rice paddies. More recently, there has been interest in using *Azolla* for bioremediation as a way to remove ammonium from wastewater or as a protein source¹⁷. *Azolla* has also been implicated in the Arctic *Azolla* Event. Around 50 million years ago, exponential growth of the plant led to a rapid reduction of carbon dioxide (CO_2) levels in the atmosphere, which is thought to have resulted in a cooling of the Earth¹⁸. The genomes of both *A. filiculoides* and *N. azollae* have been sequenced^{19,20}. The leaves of the fern contain cavities that are formed during development by invagination of the epidermis. Cyanobacteria are contained within these gas-filled cavities in close association with trichomes and embedded in a mucilaginous matrix. The structure of the trichomes is similar to that of transfer cells²¹.

In contrast to the legume–rhizobia symbiosis, in *Azolla* the bacterial symbiont is transmitted from one generation to the

next. Whereas the fern can grow without its symbiont, *N. azollae* is unable to exist outside of the plant. This is reflected in the genome of *N. azollae*, which has adapted to the symbiont lifestyle and, compared with related free-living cyanobacteria, appears reduced, and some core genes (for instance, that of phospho-fructokinase) have been lost¹⁹.

Although *N. azollae* is able to perform photosynthesis, it does so only at a low rate and depends largely upon sugar derived from the plant²². Much of the energy that *N. azollae* derives from the sugar it receives from *Azolla* is invested into N fixation. At the stage when N fixation is at its highest level, the bacteria are no longer dividing. The major product of photosynthesis in *Azolla* is sucrose²², but it is not clear in what form C is exported from the plant and taken up into the cyanosymbiont.

In the bacteria, N fixation occurs in specialized cells, called heterocysts or heterocytes, which are more frequent in the symbiont than in related free-living *Nostoc sp.*²³. Fixed N is supplied to *Azolla*, and about 40% of the N is exported from the bacteria and released in the form of ammonium into the leaf cavity²³ (Figure 2).

Even though it has been shown that *Nostoc* receives carbohydrates (at least some of it in the form of sucrose) from its plant

Figure 2. The Azolla–cyanobacteria symbiosis. In *Azolla*, expression of *MOT1*, *AMT2-4*, and an iron transporter related to *AtVIT1* is induced by N-limiting conditions. *MOT1* may be an uptake transporter, *AMT2-4* may transport fixed N from the cyanobacteria into the plant, and the iron transporter may serve as an iron efflux transporter. In the cyanobacteria, a phosphotransferase system (PTS) sugar transporter could function in sugar uptake and AmtB could function in ammonium efflux. Iron and molybdate transporters in the cyanobacteria are hypothetical.

partner²², the necessary transporters of either symbiont have not been identified. The genome of *N. azollae* contains a gene for a phosphoenolpyruvate-dependent sugar phosphotransferase system (PTS). This type of transporter is common in bacteria and catalyzes the uptake and phosphorylation of a range of carbohydrates (for instance, glucose, fructose, or cellobiose)²⁴.

On the plant side, 15 *SWEET* genes are present in the *A. filiculoides* genome, but there is no evidence that any of them is differentially expressed in the presence of the cyanosymbiont²⁵. If *SWEET* proteins are involved in C efflux in the *Azolla* symbiosis, the presence of the symbiont may trigger other post-transcriptional regulation.

N. azollae supplies its plant partner with ammonium, but no specific transporter responsible for this step has been identified.

The genome of *N. azollae* contains an *AmtB* gene, encoding a putative ammonium transporter that could export ammonia into the leaf cavity of *Azolla*. Interestingly, *N. azollae* is lacking other N transporters (for instance, for nitrate or urea)¹⁹. This may be an adaptation to the symbiont lifestyle, preventing *N. azollae* from obtaining N from the plant and making it dependent upon N fixation.

The genome of *A. filiculoides* contains genes for both electrogenic (*AMT1*) and electroneutral (*AMT2*) ammonium transporters¹⁷. Expression of one of the *AMT2* genes, *AfAMT2-4* (Azfi_s0034.g025227.AMT2), is upregulated in the fern when no N is supplied in the growth medium but only in the presence of the cyanobiont²⁵. *AMT2-4* is considered to function in the uptake of ammonium from the leaf cavity into the plant (Figure 2).

The cyanobiont appears to be dependent on *Azolla* for additional nutrients. A molybdate transporter homolog, *AfMOT1* (Azfi_s0167.g054529), and an iron transporter homolog (Azfi_s0018.g014823) are induced in the fern under the same N-limiting conditions as *AMT2-4*²⁵. Both iron and molybdate are required co-factors for nitrogenase, the key enzyme for N fixation in the cyanobacteria. This suggests that, under N-limiting conditions, the plant upregulates Mo and Fe uptake transporters to supply more of these elements to the cyanobiont. *AfMOT1* may function as a molybdate uptake transporter into the plant cells rather than as an exporter to the cyanobiont. The iron transporter Azfi_s0018.g014823 is related to vacuolar Fe/H⁺ antiporter VIT1, so it could function in Fe export from plant cells if it is localized to the plasma membrane.

To fully understand the *Azolla/Nostoc* symbiotic interaction, the key transporters responsible for C efflux from the plant and N uptake into the plant need to be identified. With new technologies such as gene editing, it may be possible to test

whether specific transporters such as *AMT2-4* from *Azolla* are required for the interaction.

Ectomycorrhiza

The ectomycorrhizal symbiosis consists of a plant partner, usually a tree species and a fungal partner, either an ascomycete or a basidiomycete. The fungus initially forms a dense layer of hyphae around the root tips of fine roots, called the mantle. Then hyphae penetrate between the cells of the epidermis and outer cortex of the tree root but without entering the cells. This structure is called the Hartig net. The fungal hyphae and the root cells share an apoplastic space, the symbiotic interface, a structure that is important for symbiosis (Figure 3). The hyphae reach an extended area around the root of the tree, thereby allowing the tree to access nutrients and water beyond the reach of its roots²⁶.

The fungus receives fixed C from its plant partner in the form of hexoses. The tree exports sucrose into the wall space, where

Figure 3. Model for ectomycorrhizal transporters. Transporters drawn with two arrows are H⁺-coupled uptake symporters. Acid invertase, shown in the symbiotic interface, is encoded by the plant. Transporters that are known are named, and transporters that have not been identified are indicated with a question mark. The fungal phosphate uptake transporter (PT) is shown away from the symbiotic interface to emphasize that it is functioning in phosphate uptake from the environment rather than in phosphate export to the plant.

the sucrose is hydrolyzed by plant-derived acid invertase into glucose and fructose. Glucose is taken up preferentially²⁷. In the fungus, glucose is rapidly converted into trehalose and sugar alcohols such as mannitol or arabitol, allowing the fungus to continue to act as a sink for the long-distance transport of sugars²⁷.

In the ectomycorrhizal interaction, the plant benefits from metabolic capabilities of the fungus as well as the increased surface area for absorption from the soil. Interaction with the fungal partner enhances the supply of nutrients, in particular N and P²⁸. The form of N transported to the plant is not clear. Larsen *et al.*²⁹ concluded that amino acids are the main metabolites that the fungus synthesizes and shares with the plant. In their study, they found that expression of amino acid transporters is enriched in both plant and fungus, providing candidate genes to study in the interaction. Ammonium transporters have been identified in both plant and fungus. In poplar, a nitrate transporter in the NRT2 family is induced by the presence of the fungal partner³⁰. TbNrt2, a nitrate uptake transporter from the fungus *Tuber borchii*, is strongly expressed in the Hartig net and mantle and only weakly in free hyphae³¹. There is no evidence for an enrichment of ammonium or inorganic phosphate (Pi) transporters in the ectomycorrhizal association²⁹. Three fungal H⁺-coupled phosphate uptake transporters have been isolated from *Hebeloma*: HcPT1.1, HcPT1.2, and HcPT2^{32–34}. All are likely to function in Pi acquisition by the fungus. HcPT2 is localized at sites of both Pi uptake and Pi release, and RNA interference (RNAi) lines of *Hebeloma* with decreased expression of *HcPT2* form fewer ectomycorrhizae than the control strain³³. However, owing to its predicted transport mechanism and the existing proton gradient with higher proton concentration in the apoplast, it is unlikely that HcPT2 activity is responsible for Pi export to the plant. An additional transporter required for P efflux to the plant has not been identified yet.

It is generally accepted that the form of C taken up by the fungus is hexoses. Several hexose transporters, HXTs or MSTs in the major facilitator superfamily, have been isolated from a range of ectomycorrhizal fungi³⁵. They function as H⁺-coupled uptake transporters^{36,37}. None of them has been shown to be essential for symbiosis, but several are specifically expressed in the root tips or in Hartig net/mantle versus free-living hyphae. In an ectomycorrhizal transcriptome study, sugar transporters were found to be specifically enriched in the transcripts of the fungal partner. No such enrichment occurred in the plant transcriptome²⁹. This may indicate that the fungal partner drives the movement of sugars, at least as far as the transport step is concerned. The plant partner may control the supply of photosynthates through controlling the activity of acid invertase in the apoplast, thereby limiting the amount of hexoses available for the fungal partner to take up.

Although no specific transporters have been shown to be required for ectomycorrhizal symbioses, a lot of progress has been made in genome sequencing and transcriptomic experiments to identify a range of transporters that may be important.

Export of sucrose is likely via SWEET proteins and uptake of hexoses by MSTs in the fungus. It would be very useful to know the forms of N and P that are taken up by the plant, and identifying transporters that are necessary for the interaction will help narrow the list of possibilities.

Endomycorrhiza

Endomycorrhizal or arbuscular mycorrhizal (AM) symbioses are ancient and widespread interactions between obligate biotrophic fungi belonging to the Glomeromycotina and the roots of land plants³⁸. This intimate association enables nutrient exchange, and plants provide up to 20% of the photosynthetically fixed C to the fungi and derive mineral nutrients from the fungal partner^{35,39,40}. Fungal hyphae grow between cells in the root cortex and through the cell wall surrounding plant cells. The fungi do not break through the plasma membrane; they form highly branched structures known as arbuscules in the apoplastic space (Figure 4). The formation of arbuscules is a signal-based process involving significant transcriptional reprogramming and cellular remodeling⁴¹. The arbuscule is surrounded by a plant-derived periarbuscular membrane (PAM), thus being physically separated from the host cytoplasm. Recent studies suggest that the PAM encloses a periarbuscular space (PAS) which contains many hyphal branches, challenging an earlier notion that the PAM trails along each hyphal branch⁴². The arbuscule provides an increased surface area for exchange of nutrients between the two organisms. Membrane transport in AM symbiosis has been studied mainly on the plant side because of the difficulty of working with AM fungi and the complexity of fungal genomes. The AM fungi establish a new C sink in the roots of plants, thus driving the transport of photoassimilates or C reserves from the host plant, preferentially in the form of hexoses, primarily glucose. Studies reveal significant transcriptional upregulation of the members of MST, sucrose transporter (SUT) and SWEET families of transporters in the arbuscule-containing cells of many plant species, although most of the results await genetic and functional validation^{35,40}. Recently, SWEET1b from *Medicago truncatula* was localized to the PAM and demonstrated to transport glucose⁴³. However, loss-of-function *mtsweet1b* mutants show no defect in the plant–fungal symbiotic interaction, indicating the presence of redundant transporters. Interestingly, *sut2* mutants in tomato reveal higher AM fungi colonization rates, suggesting a role for SUT2 in reverse transport of sucrose from the PAS space back to the plant⁴⁴ (Figure 4). Sugar transporters in AM fungi are less well studied. GpMST1 from *Geosiphon pyriformis* and RiMST2 from *Rhizophagus irregularis* transport hexoses and have the highest affinity for glucose^{45,46}. RiMST2 is expressed in arbuscules and intercellular hyphae where it may function in sugar uptake from the plant⁴⁶. Interestingly, active uptake of ¹⁴C-labelled sugars by the extraradical mycelium indicates that sugar transport by the fungus also occurs outside of arbuscules⁴⁷.

Lipids may serve as another source of C transferred from plants to AM fungi. AM fungi depend on the plant host for lipids since they lack genes encoding the cytosolic fatty acid synthase subunits for fatty acid biosynthesis^{48,49}. This

Figure 4. Transporters in endomycorrhizal or arbuscular mycorrhizal (AM) interactions. A plant cell containing an arbuscule is shown. Transporters that are known are named in the illustration. Transporters drawn with two arrows are H^+ -coupled uptake symporters. Transporter activities that do not have an identified transporter are indicated with a question mark. PAS, periarbuscular space.

dependency was revealed by legume mutants defective in lipid biosynthetic genes that have stunted arbuscules and reduced colonization. Other studies using transgenic *Medicago* expressing fatty acid synthesis genes and radioisotopes also demonstrate the transfer of fatty acids from plants to fungi. The ABC-type transporters STR1 and STR2 are required for arbuscule formation in rice and *Medicago*^{50,51} and are predicted to transport beta-monoacylglycerol from plants to fungi^{52,53}. The mechanism of transport or trafficking of lipids at the AM–plant interface is not clear and will require additional research such as using imaging with lipophilic dyes or transport assays using reconstitution of transporter proteins in liposomes.

Owing to the low mobility of phosphate ions in soil, plants use the extensive network of AM fungi to acquire most of their P. Availability of Pi is also a signal for arbuscule maintenance since plant Pi transporter mutants display premature degeneration of arbuscules⁵⁴ and exogenous Pi treatment in rice inhibits fungal colonization and mycorrhizal uptake of Pi⁵⁵. Krajinski *et al.*⁵⁶ revealed that Pi uptake by AM fungi is driven by the transmembrane H^+ gradient. Fungal Pi/ H^+ symporters are responsible for Pi uptake from the soil by the extraradical mycelium, although they are also expressed in the intraradical mycelium, suggesting their role in reabsorption from the PAS. Pi is then trafficked as polyphosphate chains to the intraradical mycelium, where breakdown produces Pi which then gets transported into the PAS via a still-unknown mechanism. AM fungi colonization induces the expression of Pi transporters in rice (*OsPT11*), tomato (*LePT1*), potato (*StPT3*), maize (*ZmPT9*) and *Medicago*

truncatula (*MtPT4*)^{35,40}. Both low- and high-affinity H^+ /Pi symporters of the plant PHT1 family have been implicated in AM fungi symbioses and these cluster in a separate clade from PHT1 transporters in non-AM symbiotic plants. Plant and fungal Pi transporters might also function as transceptors sensing Pi, and further investigation is warranted^{57,58}. Interestingly, high Pi represses *STR1/2* expression, suggesting that plants withhold C under high Pi conditions⁵⁹. For a detailed review of Pi transport in mycorrhiza, see⁶⁰.

N acquisition by AM fungi can provide plant partners with almost one third of their N needs⁶¹. Fungi encode ammonium transporters in the MEP family that could serve as ammonium exporters⁶², and high- and low-affinity MEPs have been reported in *R. irregularis*. AM-inducible *AMTs* are found in all host plants (both dicots and monocots) and have been localized to the PAM⁶³. Since the PAS is acidic, NH_4^+ concentration predominates over NH_3 . Plant *AMTs* are thought to bind NH_4^+ , deprotonate it, and co-transport NH_3 and H^+ (Figure 4). Nitrate might also be transferred since plant nitrate transporters are induced by AM colonization^{35,40}. Nitrate transporters are also induced by high phosphate or low nitrate concentrations, suggesting a complex regulation.

Legume–rhizobia symbiosis

In the rhizobia–legume interaction, the plant responds to signals produced by the bacterium and initiates a developmental program leading to the formation of a plant root structure called the nodule. Within the nodule, the modified rhizobia,

called bacteroids, are housed in intracellular, membrane-bound compartments called symbiosomes. The plant symbiosome membrane (SM), also called the peribacteroid membrane (PBM), surrounds the bacteroids (Figure 5). The plant supplies fixed C and a myriad of other ions, metabolites, and proteins to create a suitable environment for the bacteroids to fix N which is absorbed by the plant. Recent advances in proteomic and transcriptomic analysis have led to the identification of transporters in the PBM.

Transporters involved in the interaction of legumes and rhizobia have been recently reviewed in detail^{64,65}. Despite their central importance for the symbiosis, most of the N and C transporters predicted to be required for the interaction have not been identified.

The PBM contains plasma membrane-type H^+ ATPases that acidify the peribacteroid space and generate a negative membrane potential⁶⁴. To understand transport processes at the PBM, the peribacteroid space should be considered equivalent to the apoplast or cell wall space of a plant cell. Accordingly, solutes are transported from the plant to bacteroids via plant exporter proteins.

It is clear that the form of fixed C taken up by bacteroids is C_4 -dicarboxylates⁶⁶. Bacterial mutants defective in dicarboxylate transporter DctA are unable to fix N. Malate, succinate, and fumarate are transported by DctA^{66,67}. Malate appears to be the most important and only required C_4 -dicarboxylate

supplied to the bacteroids by the plant⁶⁸. It is not surprising that the uptake of dicarboxylates by DctA is coupled to the uptake of at least three H^+ so that the transport reaction is driven by the negative bacterial membrane potential and by the transmembrane pH gradient⁶⁷. The exporter responsible for malate export from the plant across the SM has not yet been identified. There are excellent candidates for dicarboxylate export into the peribacteroid space. Malate is transported by S-type anion channels⁶⁹ in the SLAC family and by aluminum-activated malate transporters (ALMTs)⁷⁰. ALMTs were first identified in wheat and were shown to export malate in response to exposure to aluminum. In *Lotus japonicus*, LjALMT4 transports malate and is expressed in nodules but is localized to the vasculature in nodules⁷¹. Perhaps another member of the ALMT family localizes to the PBM in infected cells as indicated in Figure 5.

The form of N transported from the bacteroid to the plant is thought to be ammonium (either NH_4^+ or NH_3). Although both plants and bacteria encode ammonium transporters in the AMT/MEP/Rh family⁶², the current view is that transporters in the AMT/MEP/Rh family are not involved in either NH_3 release from bacteroids or uptake into infected plant cells. The bacterial AMT *AmtB* is not expressed in bacteroids and therefore N is thought to be exported from bacteroids by diffusion of NH_3 through the bacteroid membrane⁶⁴. The aquaporin homolog NOD26 has been suggested to serve as the ammonium uptake transporter from the peribacteroid space into rhizobia-infected cells. NOD26 is highly expressed, localized to the

Figure 5. Legume–rhizobia transport processes. Plant transporters for malate efflux, iron efflux, and ammonium uptake are shown on the symbiosome membrane. At the plasma membrane of infected plant cells, SST1, MOT1, Nramp1, and PT7 are thought to function in uptake into the plant cells. In the bacteroid membrane, DctA, the bacterial C_4 -dicarboxylate transporter, is shown on the bacteroid membrane. NH_3 diffusion through the bacteroid membrane is indicated by a dashed line.

PBM, and known to transport NH_3 ⁷². Transporters in the AMT1 family were not identified in proteomic studies of nodules^{73,74}. GmSAT1 from soybean was originally considered a candidate for the plant PBM ammonium uptake transporter; however, GmSAT1 was subsequently found to be a transcription factor that affects expression of the yeast *MEP3* ammonium transporter when expressed in yeast⁷⁵.

Since the bacteroid is completely enclosed by the plant SM, it is necessary for the plant to provide all nutrients required by the bacteroid. Recent results demonstrate that uptake into nodule cells is an essential step in providing nutrients to the bacteroid. Molybdenum is required for the nitrogenase enzyme, and in *Medicago* MtMOT1.3⁷⁶ and MtMOT1.2⁷⁷ localize to the plasma membrane in nodule cells and function in molybdate uptake. The molybdate exporter, required to deliver molybdate into the peribacteroid space, has not been identified. Synthesis of nitrogenase in the bacteroid requires a significant amount of sulfur (S) for both S-containing amino acids and Fe-S complexes. Bacteroids have been shown to take up 20-fold times more S than the nodule host cells⁷⁸. The plant SO_4^{2-} transporter SST1 is required for symbiotic N fixation⁷⁹, and *sst1* mutant nodules show less S accumulated in host cells, in the symbiosome space, and in bacteroids⁷⁸. SST1 protein was identified by mass spectroscopy in PBM fractions^{73,74}; however, its localization has not been confirmed by a second method. SST1 belongs to the SULTR family of plant electrogenic H^+ -coupled SO_4^{2-} uptake transporters, so the function of SST1 is most likely S uptake into plant cells in the nodule rather than SO_4^{2-} export to the bacteroids. If SST1 functions in SO_4^{2-} uptake, then its localization is more likely to be on the plasma membrane rather than the PBM. Other plant nutrient uptake transporters localized to the plasma membrane, such as the phosphate transporter GmPT7 from soybean⁸⁰, have been shown to be necessary for optimal N fixation and they also function in uptake into nodule cells rather than nutrient export to the bacteroid (Figure 5).

The biochemical reactions of N fixation that occur in the bacteroid require transition metals. For example, nitrogenase

is the most abundant protein in bacteroids and contains 32 iron atoms and one molybdenum. Cobalt, copper, manganese, nickel, and zinc are also required and must be supplied to the bacteroid. In *Medicago truncatula*, iron transport into rhizobia-infected cells is thought to occur via MtNramp1, which is localized to the plasma membrane in nodule cells⁸¹. The *nramp1-1* mutant has lower nitrogenase activity, indicating that less iron is delivered to the bacteroids in the mutant. MtSEN1 may serve as the iron exporter on the PBM necessary for iron transport into the peribacteroid space⁸². MtSEN1 is in the VIT1/CCC1 family of vacuolar iron exporters that function as H^+ -coupled antiporters⁸³. Subcellular localization of MtSEN1 will be necessary to support this hypothesis. There is a recent review of transporters involved in export of transition metals from the plant and uptake into the bacteroid⁸⁴.

Conclusions

In this review, we discussed five nutritional symbioses and some common features were found; for example, all of the interactions involve the transfer of fixed C and fixed N. When the form of C exported from a photosynthetic partner is sucrose or its hydrolysis products, glucose and fructose, the type of transporter is likely to be a SWEET. The legume–rhizobia interaction is an exception in that C is transferred in the form of the C_4 -dicarboxylate malate. Although the transporter responsible for malate efflux has not been identified, plant anion channels are known to transport malate. The most common form of transported N in these interactions is ammonium. Plants encode both electrogenic (AMT1) and electroneutral (AMT2) ammonium transporters. One common hypothesis is that AMT1s serve as uptake transporters and AMT2s serve as exporters⁶², but the situation seems to be more complicated and interesting; for example, *AMT2-4* from *Azolla* has been suggested to be involved in ammonium uptake into the fern. This is an exciting time to study symbiotic interactions because of advances in genome sequencing, transcriptomics, and gene editing. Identification and analysis of the transporters involved will allow us to better understand how symbiotic organisms share nutrients.

References

- Feuerer T, Hawksworth DL: **Biodiversity of lichens, including a world-wide analysis of checklist data based on Takhtajan's floristic regions.** *Biodivers Conserv.* 2007; **16**(1): 85–98.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lücking R, Rivas Plata E, Chaves J, *et al.*: **How many tropical lichens are there... really.** *Bibliotheca Lichenologica.* 2009; **100**: 399–418.
[Reference Source](#)
- Richardson DHS: **Photosynthesis and Carbohydrate Movement.** In V Ahmadjian, ME Hale, eds *The Lichens.* Academic Press, New York, 1973; 249–288.
[Reference Source](#)
- Rai AN, Rowell P, Stewart WD: **(^{15}N) $_2$ Incorporation and metabolism in the lichen *Peltigera aphthosa* Willd.** *Planta.* 1981; **152**: 544–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Park SY, Choi J, Kim JA, *et al.*: **Draft Genome Sequence of *Cladonia macilentata* KoLR1003786, a Lichen-Forming Fungus Producing Biruloquinone.** *Genome Announc.* 2013; **1**(5): 161.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Park SY, Choi J, Kim JA, *et al.*: **Draft Genome Sequence of Lichen-Forming Fungus *Caloplaca flavorubescens* Strain KoLR1002931.** *Genome Announc.* 2013; **1**(4): 937.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Park SY, Choi J, Lee GW, *et al.*: **Draft Genome Sequence of *Umbilicaria muenhbergii* KoLR1000956, a Lichen-Forming Fungus Amenable to Genetic Manipulation.** *Genome Announc.* 2014; **2**(2): 15272.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Park SY, Choi J, Lee GW, *et al.*: **Draft Genome Sequence of Lichen-Forming Fungus *Cladonia metacorrallifera* Strain KoLR1002260.** *Genome Announc.* 2014; **2**(1): pii: e01065-13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Armstrong EE, Prost S, Ertz D, *et al.*: **Draft Genome Sequence and**

- Annotation of the Lichen-Forming Fungus *Arthonia radiata*.** *Genome Announc.* 2018; 6(14): pii: e00281-18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
10. **F** Wang Y, Yuan X, Chen L, *et al.*: **Draft Genome Sequence of the Lichen-Forming Fungus *Ramalina intermedia* Strain YAF0013.** *Genome Announc.* 2018; 6(23): pii: e00478-18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 11. Armaleo D, Müller O, Lutzoni F, *et al.*: **The lichen symbiosis re-viewed through the genomes of *Cladonia grayi* and its algal partner *Asterochloris glomerata*.** *BMC Genomics.* 2019; 20(1): 605.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 12. Wang YY, Liu B, Zhang XY, *et al.*: **Genome characteristics reveal the impact of lichenization on lichen-forming fungus *Endocarpon pusillum* Hedwig (Verrucariales, Ascomycota).** *BMC Genomics.* 2014; 15: 34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 13. Wang Y, Zhang X, Zhou Q, *et al.*: **Comparative transcriptome analysis of the lichen-forming fungus *Endocarpon pusillum* elucidates its drought adaptation mechanisms.** *Sci China Life Sci.* 2015; 58(1): 89–100.
[PubMed Abstract](#) | [Publisher Full Text](#)
 14. Athukorala SN, Huebner E, Piercey-Normore MD: **Identification and comparison of the 3 early stages of resynthesis for the lichen *Cladonia rangiferina*.** *Can J Microbiol.* 2014; 60(1): 41–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
 15. Junttila S, Laiho A, Gyenesei A, *et al.*: **Whole transcriptome characterization of the effects of dehydration and rehydration on *Cladonia rangiferina*, the grey reindeer lichen.** *BMC Genomics.* 2013; 14: 870.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 16. **F** Yoshino K, Yamamoto K, Hara K, *et al.*: **The conservation of polyol transporter proteins and their involvement in lichenized Ascomycota.** *Fungal Biol.* 2019; 123(4): 318–29.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 17. **F** Brouwer P, Bräutigam A, Buijs VA, *et al.*: **Metabolic Adaptation, a Specialized Leaf Organ Structure and Vascular Responses to Diurnal N₂ Fixation by *Nostoc azollae* Sustain the Astonishing Productivity of *Azolla* Ferns without Nitrogen Fertilizer.** *Front Plant Sci.* 2017; 8: 442.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 18. Speelman EN, Van Kempen MM, Barke J, *et al.*: **The Eocene Arctic *Azolla* bloom: environmental conditions, productivity and carbon drawdown.** *Geobiology.* 2009; 7(2): 155–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
 19. Ran L, Larsson J, Vigil-Stenman T, *et al.*: **Genome erosion in a nitrogen-fixing vertically transmitted endosymbiotic multicellular cyanobacterium.** *PLoS One.* 2010; 5(7): e11486.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 20. **F** Li FW, Brouwer P, Carretero-Paulet L, *et al.*: **Fern genomes elucidate land plant evolution and cyanobacterial symbioses.** *Nat Plants.* 2018; 4(7): 460–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 21. Zheng W, Rang L, Bergman B: **Structural Characteristics of the Cyanobacterium-*Azolla* Symbioses.** In K Pawlowski, ed. *Prokaryotic Symbionts in Plants*. Microbiology Monographs, Springer, Berlin, Heidelberg. 2009; 8: 235–263.
[Publisher Full Text](#)
 22. Kaplan D, Peters GA: **Interaction of carbon metabolism in the *Azolla-Anabaena* symbiosis.** *Symbiosis.* 1998; 6: 53–69.
[Reference Source](#)
 23. Meeks JC: **Physiological Adaptations in Nitrogen-fixing *Nostoc*–Plant Symbiotic Associations.** In K Pawlowski, ed. *Prokaryotic Symbionts in Plants*. Microbiology Monographs, Springer, Berlin, Heidelberg. 2009; 8: 181–205.
[Publisher Full Text](#)
 24. Deutscher J, Francke C, Postma PW: **How phosphotransferase system-related protein phosphorylation regulates carbohydrate metabolism in bacteria.** *Microbiol Mol Biol Rev.* 2006; 70(4): 939–1031.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 25. **F** Eily AN, Pryer KM, Li FW: **A first glimpse at genes important to the *Azolla*–*Nostoc* symbiosis.** *Symbiosis.* 2019; 78(2): 149–62.
[Publisher Full Text](#) | [F1000 Recommendation](#)
 26. Wiemken V, Boller T: **Ectomycorrhiza: gene expression, metabolism and the wood-wide web.** *Curr Opin Plant Biol.* 2002; 5(4): 355–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
 27. Nehls U, Göhringer F, Wittlusk S, *et al.*: **Fungal carbohydrate support in the ectomycorrhizal symbiosis: a review.** *Plant Biol (Stuttg).* 2010; 12(2): 292–301.
[PubMed Abstract](#) | [Publisher Full Text](#)
 28. **F** Nehls U, Plassard C: **Nitrogen and phosphate metabolism in ectomycorrhizas.** *New Phytol.* 2018; 220(4): 1047–58.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 29. Larsen PE, Sreedasyam A, Trivedi G, *et al.*: **Using next generation transcriptome sequencing to predict an ectomycorrhizal metabolome.** *BMC Syst Biol.* 2011; 5: 70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 30. Willmann A, Thomfahde S, Haensch R, *et al.*: **The poplar NRT2 gene family of high affinity nitrate importers: Impact of nitrogen nutrition and ectomycorrhiza formation.** *Environmental and Experimental Botany.* 2014; 108: 79–88.
[Publisher Full Text](#)
 31. Montanini B, Viscomi AR, Bolchi A, *et al.*: **Functional properties and differential mode of regulation of the nitrate transporter from a plant symbiotic ascomycete.** *Biochem J.* 2006; 394(Pt 1): 125–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 32. Garcia K, Haider MZ, Delteil A, *et al.*: **Promoter-dependent expression of the fungal transporter *HcPT1.1* under Pi shortage and its spatial localization in ectomycorrhiza.** *Fungal Genet Biol.* 2013; 58–59: 53–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
 33. **F** Becquer A, Garcia K, Amenc L, *et al.*: **The *Hebeloma cylindrosporum* HcPT2 Pi transporter plays a key role in ectomycorrhizal symbiosis.** *New Phytol.* 2018; 220(4): 1185–99.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 34. **F** Becquer A, Garcia K, Plassard C: **HcPT1.2 participates in Pi acquisition in *Hebeloma cylindrosporum* external hyphae of ectomycorrhizas under high and low phosphate conditions.** *Plant Signal Behav.* 2018; 13(10): e1525997.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 35. Garcia K, Doidy J, Zimmermann SD, *et al.*: **Take a Trip Through the Plant and Fungal Transportome of Mycorrhiza.** *Trends Plant Sci.* 2016; 21(11): 937–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
 36. Polidori E, Ceccaroli P, Saltarelli R, *et al.*: **Hexose uptake in the plant symbiotic ascomycete *Tuber borchii* Vittadini: biochemical features and expression pattern of the transporter TBHT1.** *Fungal Genet Biol.* 2007; 44(3): 187–98.
[PubMed Abstract](#) | [Publisher Full Text](#)
 37. Ceccaroli P, Saltarelli R, Polidori E, *et al.*: **Sugar transporters in the black truffle *Tuber melanosporum*: from gene prediction to functional characterization.** *Fungal Genet Biol.* 2015; 81: 52–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
 38. Lanfranco L, Bonfante P, Genre A: **The Mutualistic Interaction between Plants and Arbuscular Mycorrhizal Fungi.** *Microbiol Spectr.* 2016; 4(6).
[PubMed Abstract](#) | [Publisher Full Text](#)
 39. **F** Guerrero-Galán C, Garcia K, Houdinet G, *et al.*: **HcTOK1 participates in the maintenance of K⁺ homeostasis in the ectomycorrhizal fungus *Hebeloma cylindrosporum*, which is essential for the symbiotic K⁺ nutrition of *Pinus pinaster*.** *Plant Signal Behav.* 2018; 13(6): e1480845.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 40. **F** Lanfranco L, Fiorilli V, Gutjahr C: **Partner communication and role of nutrients in the arbuscular mycorrhizal symbiosis.** *New Phytol.* 2018; 220(4): 1031–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 41. **F** Luginbuehl LH, Oldroyd GED: **Understanding the Arbuscule at the Heart of Endomycorrhizal Symbioses in Plants.** *Curr Biol.* 2017; 27(17): R952–R963.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 42. **F** Ivanov S, Austin J 2nd, Berg RH, *et al.*: **Extensive membrane systems at the host-arbuscular mycorrhizal fungus interface.** *Nat Plants.* 2019; 5(2): 194–203.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 43. **F** An J, Zeng T, Ji C, *et al.*: **A *Medicago truncatula* SWEET transporter implicated in arbuscule maintenance during arbuscular mycorrhizal symbiosis.** *New Phytol.* 2019; 224(1): 396–408.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 44. Bitterlich M, Krügel U, Boldt-Burisch K, *et al.*: **The sucrose transporter SISUT2 from tomato interacts with brassinosteroid functioning and affects arbuscular mycorrhiza formation.** *Plant J.* 2014; 78(5): 877–89.
[PubMed Abstract](#) | [Publisher Full Text](#)
 45. Schüßler A, Martin H, Cohen D, *et al.*: **Characterization of a carbohydrate transporter from symbiotic glomeromycotan fungi.** *Nature.* 2006; 444(7121): 933–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 46. **F** Helber N, Wippel K, Sauer N, *et al.*: **A versatile monosaccharide transporter that operates in the arbuscular mycorrhizal fungus *Glomus* sp is crucial for the symbiotic relationship with plants.** *Plant Cell.* 2011; 23(10): 3812–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 47. Bücking H, Abubaker J, Govindarajulu M, *et al.*: **Root exudates stimulate the uptake and metabolism of organic carbon in germinating spores of *Glomus intraradices*.** *New Phytol.* 2008; 180(3): 684–95.
[PubMed Abstract](#) | [Publisher Full Text](#)
 48. Wever V, Brands M, Dörmann P: **Fatty acid synthesis and lipid metabolism in the obligate biotrophic fungus *Rhizophagus irregularis* during mycorrhization of *Lotus japonicus*.** *Plant J.* 2014; 79(3): 398–412.
[PubMed Abstract](#) | [Publisher Full Text](#)
 49. Tang N, San Clemente H, Roy S, *et al.*: **A Survey of the Gene Repertoire of *Gigaspora rosea* Unravels Conserved Features among Glomeromycota for Obligate Biotrophy.** *Front Microbiol.* 2016; 7: 233.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 50. Zhang Q, Blaylock LA, Harrison MJ: **Two *Medicago truncatula* half-ABC transporters are essential for arbuscule development in arbuscular mycorrhizal symbiosis.** *Plant Cell.* 2010; 22(5): 1483–97.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 51. Gutjahr C, Radovanovic D, Geoffroy J, *et al.*: **The half-size ABC transporters STR1 and STR2 are indispensable for mycorrhizal arbuscule formation in rice.** *Plant J.* 2012; 69(5): 906–20.
[PubMed Abstract](#) | [Publisher Full Text](#)

52. **F** Bravo A, Brands M, Wewer V, *et al.*: **Arbuscular mycorrhiza-specific enzymes FatM and RAM2 fine-tune lipid biosynthesis to promote development of arbuscular mycorrhiza.** *New Phytol.* 2017; **214**(4): 1631–45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
53. **F** Keymer A, Pimprikar P, Wewer V, *et al.*: **Lipid transfer from plants to arbuscular mycorrhiza fungi.** *eLife.* 2017; **6**: pii: e29107.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
54. **F** Javot H, Penmetsa RV, Terzaghi N, *et al.*: **A *Medicago truncatula* phosphate transporter indispensable for the arbuscular mycorrhizal symbiosis.** *Proc Natl Acad Sci U S A.* 2007; **104**(5): 1720–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
55. Kobae Y, Ohmori Y, Saito C, *et al.*: **Phosphate Treatment Strongly Inhibits New Arbuscule Development But Not the Maintenance of Arbuscule in Mycorrhizal Rice Roots.** *Plant Physiol.* 2016; **171**(1): 566–79.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. Krajinski F, Courty PE, Sieh D, *et al.*: **The H⁺-ATPase HA1 of *Medicago truncatula* Is Essential for Phosphate Transport and Plant Growth during Arbuscular Mycorrhizal Symbiosis.** *Plant Cell.* 2014; **26**(4): 1808–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
57. Yang SY, Grönlund M, Jakobsen I, *et al.*: **Nonredundant regulation of rice arbuscular mycorrhizal symbiosis by two members of the phosphate transporter1 gene family.** *Plant Cell.* 2012; **24**(10): 4236–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
58. Xie X, Lin H, Peng X, *et al.*: **Arbuscular Mycorrhizal Symbiosis Requires a Phosphate Transceptor in the *Gigaspora margarita* Fungal Symbiont.** *Mol Plant.* 2016; **9**(12): 1583–608.
[PubMed Abstract](#) | [Publisher Full Text](#)
59. **F** Wang W, Shi J, Xie Q, *et al.*: **Nutrient Exchange and Regulation in Arbuscular Mycorrhizal Symbiosis.** *Mol Plant.* 2017; **10**(9): 1147–58.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
60. **F** Plassard C, Becquer A, Garcia K: **Phosphorus Transport in Mycorrhiza: How Far Are We?** *Trends Plant Sci.* 2019; **24**(9): 794–801.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
61. Govindarajulu M, Pfeffer PE, Jin H, *et al.*: **Nitrogen transfer in the arbuscular mycorrhizal symbiosis.** *Nature.* 2005; **435**(7043): 819–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
62. McDonald TR, Ward JM: **Evolution of Electrogenic Ammonium Transporters (AMTs).** *Front Plant Sci.* 2016; **7**: 352.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
63. Kobae Y, Tamura Y, Takai S, *et al.*: **Localized expression of arbuscular mycorrhiza-inducible ammonium transporters in soybean.** *Plant Cell Physiol.* 2010; **51**(9): 1411–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
64. Udvardi M, Poole PS: **Transport and Metabolism in Legume-Rhizobia Symbioses.** *Annu Rev Plant Biol.* 2013; **64**: 781–805.
[PubMed Abstract](#) | [Publisher Full Text](#)
65. Clarke VC, Loughlin PC, Day DA, *et al.*: **Transport processes of the legume symbiosome membrane.** *Front Plant Sci.* 2014; **5**: 699.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
66. Yurgel SN, Kahn ML: **Dicarboxylate transport by rhizobia.** *FEMS Microbiol Rev.* 2004; **28**(4): 489–501.
[PubMed Abstract](#) | [Publisher Full Text](#)
67. Groeneveld M, Weme RG, Duurkens RH, *et al.*: **Biochemical characterization of the C₄-dicarboxylate transporter DctA from *Bacillus subtilis*.** *J Bacteriol.* 2010; **192**(11): 2900–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
68. **F** Mitsch MJ, diCenzo GC, Cowie A, *et al.*: **Succinate Transport Is Not Essential for Symbiotic Nitrogen Fixation by *Sinorhizobium meliloti* or *Rhizobium leguminosarum*.** *Appl Environ Microbiol.* 2017; **84**(1): pii: e01561-17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
69. Schmidt C, Schroeder JI: **Anion Selectivity of Slow Anion Channels in the Plasma Membrane of Guard Cells (Large Nitrate Permeability).** *Plant Physiol.* 1994; **106**(1): 383–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
70. Piñeros MA, Kochian LV: **A patch-clamp study on the physiology of aluminum toxicity and aluminum tolerance in maize. Identification and characterization of Al³⁺-induced anion channels.** *Plant Physiol.* 2001; **125**(1): 292–305.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
71. Takanashi K, Sasaki T, Kan T, *et al.*: **A Dicarboxylate Transporter, LjALMT4, Mainly Expressed in Nodules of *Lotus japonicus*.** *Mol Plant Microbe Interact.* 2016; **29**(7): 584–92.
[PubMed Abstract](#) | [Publisher Full Text](#)
72. Hwang JH, Ellingson SR, Roberts DM: **Ammonia permeability of the soybean nodulin 26 channel.** *FEBS Lett.* 2010; **584**(20): 4339–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
73. Wienkoop S, Saalbach G: **Proteome analysis. Novel proteins identified at the peribacteroid membrane from *Lotus japonicus* root nodules.** *Plant Physiol.* 2003; **131**(3): 1080–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
74. Clarke VC, Loughlin PC, Gavrin A, *et al.*: **Proteomic analysis of the soybean symbiosome identifies new symbiotic proteins.** *Mol Cell Proteomics.* 2015; **14**(5): 1301–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
75. Marini AM, Springael JY, Frommer WB, *et al.*: **Cross-talk between ammonium transporters in yeast and interference by the soybean SAT1 protein.** *Mol Microbiol.* 2000; **35**(2): 378–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
76. **F** Tejada-Jiménez M, Gil-Díez P, León-Mediavilla J, *et al.*: ***Medicago truncatula* Molybdate Transporter type 1 (MtMOT1.3) is a plasma membrane molybdenum transporter required for nitrogenase activity in root nodules under molybdenum deficiency.** *New Phytol.* 2017; **216**(4): 1223–35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
77. **F** Gil-Díez P, Tejada-Jiménez M, León-Mediavilla J, *et al.*: **MtMOT1.2 is responsible for molybdate supply to *Medicago truncatula* nodules.** *Plant Cell Environ.* 2019; **42**(1): 310–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
78. Schneider S, Schintlmeister A, Becana M, *et al.*: **Sulfate is transported at significant rates through the symbiosome membrane and is crucial for nitrogenase biosynthesis.** *Plant Cell Environ.* 2019; **42**(4): 1180–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
79. Krusell L, Krause K, Ott T, *et al.*: **The Sulfate Transporter SST1 Is Crucial for Symbiotic Nitrogen Fixation in *Lotus japonicus* Root Nodules.** *Plant Cell.* 2005; **17**(5): 1625–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
80. Chen L, Qin L, Zhou L, *et al.*: **A nodule-localized phosphate transporter GmPT7 plays an important role in enhancing symbiotic N₂ fixation and yield in soybean.** *New Phytol.* 2019; **221**(4): 2013–25.
[PubMed Abstract](#) | [Publisher Full Text](#)
81. Tejada-Jiménez M, Castro-Rodríguez R, Kryvoruchko I, *et al.*: ***Medicago truncatula* Natural Resistance-Associated Macrophage Protein1 Is Required for Iron Uptake by Rhizobia-Infected Nodule Cells.** *Plant Physiol.* 2015; **168**(1): 258–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
82. Hakoyama T, Niimi K, Yamamoto T, *et al.*: **The Integral Membrane Protein SEN1 is Required for Symbiotic Nitrogen Fixation in *Lotus japonicus* Nodules.** *Plant Cell Physiol.* 2012; **53**(1): 225–36.
[PubMed Abstract](#) | [Publisher Full Text](#)
83. Kato T, Kumazaki K, Wada M, *et al.*: **Crystal structure of plant vacuolar iron transporter VIT1.** *Nat Plants.* 2019; **5**(3): 308–15.
[PubMed Abstract](#) | [Publisher Full Text](#)
84. Abreu I, Mihelj P, Raimunda D: **Transition metal transporters in rhizobia: Tuning the inorganic micronutrient requirements to different living styles.** *Metallomics.* 2019; **11**(4): 735–55.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Ulrike Mathesius**

Division of Plant Science, Research School of Biology, The Australian National University, Canberra, Australia

Competing Interests: No competing interests were disclosed.

2. **Kendal Hirschi**

Children's Nutrition Research Center, Baylor College of Medicine, Houston, USA

Competing Interests: No competing interests were disclosed.

3. **Stefanie Wienkoop**

Division of Molecular Systems Biology, Department of Ecogenomics and Systems Biology, University of Vienna, Vienna, Austria

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research