Check for updates

STUDY PROTOCOL

Identifying and understanding factors that affect the translation of therapies from the laboratory to patients: a study protocol [version 1; peer review: 1 approved, 1 approved with reservations]

Manoj M. Lal^{1,3}, Joshua Montroy¹, C. Glenn Begley⁴, Tania Bubela⁵, Victoria Hunniford¹, David Ripsman^{1,6}, Neil Wesch¹, Jonathan Kimmelman⁷, Malcolm Macleod⁸, David Moher¹, Alvin Tieu^{1,6}, Lindsey Sikora⁹, Dean A. Fergusson^{1,3}

¹Clinical Epidemiology Program, Ottawa General Hospital Research Institute, Ottawa, Ontario, Canada

²Department of Anesthesiology and Pain Medicine, Ottawa Hospital, Ottawa, Ontario, Canada

³Department of Cellular and Molecular Medicine, University of Ottawa, Ottawa, Ontario, Canada

⁴BioCurate Pty Ltd, Parkville, Victoria, Australia

⁵Faculty of Health Science, Simon Fraser University, Burnaby, British Columbia, Canada

⁶Department of Medicine, University of Ottawa, Ottawa, Ontario, Canada

⁷Biomedical Ethics Unit, McGill University, Montreal, Quebec, Canada

⁸Center for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK

⁹Health Sciences Library, University of Ottawa, Ottawa, Ontario, Canada

V1 First published: 01 Jun 2020, 9:485
<https://doi.org/10.12688/f1000research.23663.1>
 Latest published: 16 Oct 2020, 9:485
<https://doi.org/10.12688/f1000research.23663.2>

Abstract

Background: The process of translating preclinical findings into a clinical setting takes decades. Previous studies have suggested that only 5-10% of the most promising preclinical studies are successfully translated into viable clinical applications. The underlying determinants of this low success rate (e.g. poor experimental design, suboptimal animal models, poor reporting) have not been examined in an empirical manner. Our study aims to determine the contemporary success rate of preclinical-to-clinical translation, and subsequently determine if an association between preclinical study design and translational success/failure exists.

Methods: Established systematic review methodology will be used with regards to the literature search, article screening and study selection process. Preclinical, basic science studies published in high impact basic science journals between 1995 and 2015 will be included. Included studies will focus on publicly available interventions with potential clinical promise. The primary outcome will be successful clinical translation of promising therapies - defined as the conduct of

Open Peer Review

Approval Status

	1	2
version 2		
(revision)
16 Oct 2020
version 1
01 Jun 2020

1. **John Holcomb**, University of Alabama at Birmingham, Birmingham, USA

2. **Jonathan A Lal**, Jacob Institute of Biotechnology and Bioengineering, Sam Higginbottom University of Agriculture, Technology and Sciences, Allahabad, India

at least one Phase II trial (or greater) with a positive finding. A case-control study will then be performed to evaluate the association between elements of preclinical study design and reporting and the likelihood of successful translation.

Discussion: This study will provide a comprehensive analysis of the therapeutic translation from the laboratory bench to the bedside. Importantly, any association between factors of study design and the success of translation will be identified. These findings may inform future research teams attempting preclinical-to-clinical translation. Results will be disseminated to identified knowledge users that fund/support preclinical research.

Keywords

Translational failures, promising therapies, bench-to-bedside research, systematic review, case-control study

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Manoj M. Lalu (mlalu@toh.ca)

Author roles: **Lalu MM:** Conceptualization, Funding Acquisition, Methodology, Resources, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Montroy J:** Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Begley CG:** Methodology, Writing – Review & Editing; **Bubela T:** Methodology, Writing – Review & Editing; **Hunniford V:** Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Ripsman D:** Investigation, Methodology, Writing – Review & Editing; **Wesch N:** Investigation, Methodology, Writing – Review & Editing; **Kimmelman J:** Conceptualization, Methodology, Writing – Review & Editing; **Macleod M:** Methodology, Writing – Review & Editing; **Moher D:** Methodology, Writing – Review & Editing; **Tieu A:** Methodology, Writing – Review & Editing; **Sikora L:** Methodology, Writing – Review & Editing; **Fergusson DA:** Conceptualization, Funding Acquisition, Methodology, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Canadian Institutes of Health Research (Project Grant #365473). CIHR was not involved in the protocol's generation; it does not have any influence on the decision to submit this manuscript.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2020 Lalu MM *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Lalu MM, Montroy J, Begley CG *et al.* **Identifying and understanding factors that affect the translation of therapies from the laboratory to patients: a study protocol [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2020, 9:485 <https://doi.org/10.12688/f1000research.23663.1>

First published: 01 Jun 2020, 9:485 <https://doi.org/10.12688/f1000research.23663.1>

Introduction

Advances in discovery research form the backbone for the development of novel therapeutics. Such translation relies on a “process of turning observations in the laboratory, clinic, and community into interventions that improve the health of individuals and the public”¹. The process is often visualized as a linear model (Figure 1). In reality, however, it comprises a spectrum that masks the complex, iterative and interdisciplinary research and development steps that engage multiple stakeholders^{2,3}. As a result, the successful translation of promising preclinical discovery research into human studies (T1) is rare⁴, and the resource-intensive efforts to evaluate discoveries in sufficient detail to allow them to be available for patients (T2) takes decades rather than years⁵.

While discovery research has inherent value, funding agencies around the world are increasingly focused on improving the success of “bench to bedside” translation (T0-T2), thereby enhancing the potential impact of biomedical research budgets. Certainly, there is great potential for improvement; it is estimated that the US alone invests 28 billion dollars per year on research that cannot be reproduced⁶. Return on investment is poor even in more successful areas. For example, the internal rate of return for health benefits of cardiovascular disease research has been calculated to be 9%⁷. This means that every \$1.00 invested into cardiovascular research provides the equivalent of 9¢ per year in health benefits. Importantly, a clinically translated therapy – one which has obtained regulatory approval – does not mean patients will have access to it. An additional hurdle in the translation into practice (T3) and community (T4) access is whether or not insurance companies and healthcare systems will pay for the therapy. The willingness to pay is largely dependent on a cost-comparative analysis of the potential therapy as well as the needs and resources of the jurisdiction. As this is a critical aspect in the translation spectrum and widespread use of a therapy, it has been thoroughly investigated⁸.

Investigation of translation in the earlier phases of the translational spectrum – from basic research (T0) to clinical trials in humans (T2) – has uncovered striking deficits when moving from one phase to another⁹. These inefficiencies ultimately contribute to a 5–10% rate of successful translation of preclinical “bench” research to approved therapies. This means that approximately 90% of promising discoveries may not directly contribute to improved human health^{5,10,11}. Moreover, translation failures expose clinical research participants to potential harms of investigational products that fail on safety or efficacy grounds^{12,13}. The extent of translational failures diverts scarce funding away from developing interventions that are more likely to benefit patients. Given the recent focus on research waste^{14–17}, it is abundantly clear that bench-to-bedside translation needs solutions to become more efficient.

Three highly cited studies investigating this issue in publicly-available data have been previously published. Contopoulos-Ioannidis *et al.*¹⁸ traced the translation of published, highly promising *in vivo* basic science findings into clinical applications from 1979 to 1983 in six basic science journals. They identified 101 promising therapies and in 2003 (i.e. ~20 years since studies published), 27 entered clinical trials and only 5 were clinically licensed (i.e. only 5% translated into an approved therapeutic). Hackam and Redelmeier¹¹ searched for studies published in seven leading journals between 1980 and 2000 that were cited >500 times and investigated highly promising preclinical agents. Their inclusion criteria resulted in 76 studies of interventions, of which 42 (55%) entered clinical trials and only 8 (10%) were approved for clinical use. These authors attempted multivariable analyses to identify predictors of translation; however, the small sample size “limited power to discern individual predictors of translation.” In the most recent investigation of this issue, Morris *et al.*¹⁹ (2011) performed a pseudo-systematic search to identify studies that had quantified time lags in the development of health interventions. They found time lags of ~20 years for interventions to translate, though this depended on how each study defined if translation occurred. Importantly, they concluded that the “knowledge of time lags is of limited use to those responsible for R&D... who face difficulties in knowing what they should or can do to reduce time lags.” In other words, identifying factors associated with successful translation of preclinical research into clinical trials may be a more practical metric than time lags for decision makers.

In considering this previous research, it is still currently unknown if/how rates of translation from preclinical research to the therapy’s evaluation in clinical trials are associated with study rigor (i.e. internal validity, risk of bias), and external validity (i.e. reproducibility over a range of conditions), among other factors affecting the quality of the preclinical evidence such as construct validity (i.e. the experimental model appropriately represents the clinical condition and setting). Therefore, the aim of this project is to use established knowledge synthesis methods to identify the prevalence of bench-to-bedside translation, map contemporary trends in translation, and identify key modifiable factors associated with successful and unsuccessful bench-to-bedside translation.

Objectives

Objective #1

We will perform a cross-sectional study to estimate the rate of publicly available bench-to-bedside translation. Specifically, we will identify published preclinical¹ research of promising

¹ The term ‘preclinical’ has variable definitions, depending on the context. For the purposes of this study, our group has chosen to focus on preclinical interventional *in vivo* animal experiments, relevant to the betterment of human health and/or treatments for human disease.

Figure 1. The translational research spectrum.

therapies and track how far along the translational spectrum they have progressed.

Regulatory approval of the therapy will be considered successful translation. If approval has not yet been obtained, we will consider a therapy that translated from the preclinical phase (T0) to a Phase II clinical trial (T2) or further, a successful bench-to-bedside translation.

Objective #2

We will next perform a case-control study to identify factors associated with therapies that have been successfully translated from the preclinical study in which it was identified, to a positive clinical trial evaluating treatment efficacy. Specifically, we will consider a therapy to be ‘successfully’ translated if it 1) has obtained regulatory approval, or 2) has a significant, positive finding in a clinical trial evaluating efficacy (presumably a Phase II trial) at the latest stage trial the therapy has advanced to (i.e. the therapy will not be considered successful if it demonstrated efficacy in an earlier trial and subsequently showed null or negative findings in a later efficacy trial).

Methods

Protocol

Any modifications from this protocol will be included in the final report.

Objective #1

The first objective of this study is to estimate the rate of translation from preclinical research in animal models to success in phase II clinical studies of therapeutic efficacy. This will be achieved through a cross-sectional study of preclinical investigations. In order to identify and select relevant studies for inclusion, we will employ methods typically used in systematic reviews²⁰. The objective focuses on two questions. *What is the current rate of preclinical to clinical translation of promising basic science findings? What are the basic characteristics and design features of the promising therapies identified?*

Eligibility criteria. We will identify preclinical studies published in *Science*, *Nature*, and *Nature Medicine*. These journals were selected because they are considered leaders across all domains of preclinical research and are anticipated to publish work that may impact human health. Our selection also reflects the philosophy adopted by the largest previous evaluation of bench-to-bedside research⁵.

Eligible articles will be those published between 1995 and 2015. This timeframe allows a significant time-lag (although not necessarily the full ~20-year time lag described previously). This timeframe will also allow for an evaluation of (in an exploratory manner) whether bench-to-bedside translation has improved in terms of both time-lags and the proportion of therapies advancing to later phases of clinical trials or gaining regulatory approval. Importantly, we anticipate that the journal selection and year range will also provide an adequate sample size to perform the case-control study outlined in Objective #2.

Inclusion criteria

i. Population:

Articles that describe a preclinical, interventional study: any article that includes *in vivo* non-human animal experiments that has not been tested for the same purpose in humans prior to the study.

ii. Intervention and outcome:

A ‘promising therapy’, defined by the following:

- Any therapy introduced to the animal model (i.e. pharmacologic and non-pharmacologic therapies, vaccines, antibodies, etc.), which was still at the developmental stage and did not have a prior application in humans for the specific indication.
- Novel uses of existing therapies (e.g. treatment of a different disease) will also be included.
- The intervention must induce an outcome that positively benefits an indicator of health of the animal (e.g. an immunotherapy that shrinks solid tumor size in the animal model) and/or investigators state that the therapy should be translated clinically based on their findings.

iii. Comparison:

Any comparison to the intervention will be acceptable

Exclusion criteria

- Editorials, commentaries, reviews, news articles, articles that focus on a mechanism of action, pathophysiology, or diagnosis, and articles on agricultural or veterinary applications that would not be feasibly applied to humans.
- Investigators explicitly state that translation should not be attempted based on the study.
- Ex vivo*, *in vitro*, and human clinical trials.

Information sources and search strategy. All included journals are indexed in MEDLINE, therefore this database will be searched from January 1st, 1995, to December 31st, 2015. A validated animal filter will limit results to animal studies²¹. The search strategy will be developed and finalized with the help of an information specialist who has expertise in the design of systematic searches (L. Sikora). The search strategy can be found in *Extended data*²².

Study selection process. The literature search results will be uploaded to Distiller Systematic Review Software (DistillerSR®, Evidence Partners, Ottawa, Canada). DistillerSR is an audit-ready, cloud-based software program that allows for transparent and reproducible work required for an accurate review. Two reviewers will independently screen the titles and abstracts from the search results using the predefined eligibility criteria. A calibration exercise will be performed on the first 50 studies to refine the screening question prior to formally commencing the screening process. Two review authors will assess the eligibility of the full-text articles. Discrepancies between the reviewers will be resolved by discussion or with a third-party

member if consensus cannot be established. Reasons for excluding studies will be recorded.

Data collection process and data items. Standardized forms designed in DistillerSR will be used to extract all study characteristics (see list below). Following a pilot to refine the forms and a calibration exercise to ensure high inter-rater agreement (i.e. above 80%), data will be extracted independently by two reviewers. Disagreements between reviewers will be resolved by discussion or with a third-party member if a consensus cannot be reached. The following data will be collected from eligible preclinical articles, and many of these elements will be incorporated for analyses listed in Objective #2. Data items i–v will be collected from all eligible preclinical articles. Data items vi–x will only be collected from articles selected as either cases or controls in Objective #2:

- i. *Funding:* academic/governmental/charitable vs. biotechnology and/or pharmaceutical companies (defined as reported industry affiliation by an author, financial support, or provision of the therapy being studied)
- ii. *Study characteristics:* study title, first and corresponding authors' name, publication year, journal of publication, country of corresponding author, and the number of authors and affiliated institutions
- iii. *Study population:* animal species, sex, age, presence of comorbid illnesses
- iv. *Type of model:* disease being studied, name of animal model(s) used, number of different models used
- v. *Interventions (promising therapy):* name of intervention, dose-response tested, anticipated application (e.g. preventative, therapeutic, or both)
- vi. *Preclinical outcomes:* primary outcome (or, if none declared, main *in vivo* outcome highlighted by authors), death, adverse events
- vii. *Risk of bias (internal validity):* the Cochrane Risk of Bias tool to evaluate preclinical *in vivo* studies²⁰
- viii. *External validity* (range of experimental conditions tested): more than one animal model used (species or disease model/co-morbid model), inclusion of male and female animals, and multiple treatment doses
- ix. *Completeness of reporting:* an operationalized version of the National Institutes of Health Preclinical Reporting guidelines will be used²³.
- x. *Article metrics:* The H-index of the corresponding author(s) and the number of times the paper has been cited

Identifying clinical translation of promising bench findings. After identifying the set of promising potential therapies, the stage of clinical translation that each therapy has achieved will be identified. In collaboration with an information specialist (L. Sikora), a piloted algorithmic search strategy that identifies

the current state of clinical use will be used. Backward citation analysis (identifying and examining the studies cited in an article) and forward citation analysis (identifying studies that cite an original article or work after it had been published) will be used. The search strategy consists of searching known databases for citation analysis (i.e. Scopus, ISI Web of Science, Google Scholar). Other resources will also be searched including grey literature sources, clinical trial registries, clinical practice guidelines, and conference proceedings. The full draft algorithm is listed in *Extended data*. This approach will be used to determine the furthest stage of clinical testing and whether the agent has received regulatory approval.

Outcomes. The primary outcome of interest will be the successful clinical translation of the identified promising therapies defined as 1) having obtained regulatory approval for the therapy, or 2) the conduct of at least one Phase II trial (or greater) for efficacy, with a statistically significant result favoring the treatment (and no later trials with negative or null results). As per the Federal Drug Agency (FDA) in the United States, the purpose of Phase II trials is “efficacy as well as side effects.”²⁴ A positive finding will be defined as a statistically significant result demonstrating therapeutic superiority compared to placebo/no treatment, or established interventions; or stated non-inferiority compared to currently established interventions in an appropriate designed and conducted clinical study. This primary outcome has been selected as it indicates replication of the findings from the animal models of the preclinical study in the human subjects of the clinical trial.

The secondary outcome is the furthest clinical advances of all promising therapies from Phase I through to regulatory approval. For this outcome we will consider *any* clinical trial that has been performed and/or published. Whether favorable results were obtained for each identified trial (which we will call a “positive” trial) will also be reported. As above, a positive result will be defined by a statistically significant result demonstrating superiority compared to placebo/no treatment, or established interventions; or stated non-inferiority compared to currently established interventions. In determining regulatory approval status, while the dominant regulatory agencies are often regarded as the FDA (US), European Medical Agency (Europe), and Pharmaceuticals and Medical Devices Agency (Japan), approval by any regulatory agency will be accepted. This outcome will be presented as a proportion of studies that advanced to the clinical trial stage. Results will also be stratified by year of publication: 24–20 years ago (to provide a comparison to the landmark article by Contopoulos-Ioannidis *et al.*⁵), 20–10 years ago, and <10 years ago.

Data synthesis and analysis plan. The total number of studies screened, assessed for eligibility, and included in the review (with reasons for exclusion) will be reported according to the PRISMA guidelines for reporting systematic reviews²⁵. For each included study, the characteristics outlined above will be described, with frequencies and proportions reported. For the primary outcome, the time from the preclinical paper publication date, to first positive Phase II trial will be

calculated and presented. Likewise, for the secondary outcomes, the time to furthest advance (Phase I, II, III, IV, or approval), as well time to first positive trial will be calculated using medians and interquartile ranges. Kaplan-Meier curves will also be constructed for all time to event analyses. All analyses will be performed using SAS (Version 9.4).

For all promising therapies, a descriptive analysis of all study characteristics and its research trajectory (i.e. furthest level of human experimentation) will be presented. This will present the cohort of successful and unsuccessful promising therapies published in highly influential journals, as well as the ability to evaluate secular trends in translation.

Risk of bias assessment. Risk of bias will be assessed in duplicate by two independent reviewers as high, low or unclear for six domains of bias identified by the Cochrane Risk of Bias tool²⁰, with the addition of two domains suggested to influence the risk of bias in preclinical studies²⁶. Disagreements will be resolved first by discussion and subsequently by consulting a third-party member, if needed. Graphic representations of risk of bias within and across studies will be conducted using RevMan 5.3 (Cochrane Collaboration, Oxford, United Kingdom).

Objective #2

A case-control study will be performed to evaluate the association between preclinical study characteristics and the incidence of translation. Translated therapies gaining regulatory approval or into a Phase II clinical trial (or later) will be defined as cases; unsuccessfully translated therapies will serve as controls. This study will help identify factors of internal and/or external validity, which are associated with “failed” versus “successful” translation. A case-control design was chosen as the most efficient methodological design due to i.) the rarity of preclinical to clinical translation and ii.) the ability to study multiple factors that may contribute to translational success.

Data source, identification and matching of cases and controls. Upon completion of the cross-sectional study outlined in Objective #1, a comprehensive roster of promising basic science findings published in top-tier journals between 1995 and 2015 will be generated. From this dataset, cases and matched controls will be selected:

Cases: Promising discoveries that became successfully translated will be used as cases in our study. ‘Successfully translated’ will be defined in two ways as mentioned above. In keeping with the primary outcome in Objective #1, successful translation will be defined as having 1) obtained approval from a regulatory agency and/or 2) demonstrated positive findings in a Phase II clinical trial of efficacy - indicating replication of the findings from the preclinical study. As in Objective #1, a positive result will be defined by a statistically significant result demonstrating superiority compared to placebo/no treatment, or established interventions; or stated non-inferiority compared to currently established interventions.

Controls: Highly promising basic science findings which were not successfully translated will serve as controls. Unsuccessful translation will be defined as either a highly promising basic science finding without 1) having obtained regulatory approval, AND 2) any clinical trial (i.e. no trial identified by literature search and/or identified in clinical trial registries such as clinicaltrials.gov) or going to trial and failing for any reason (e.g. failing in safety in Phase I; or efficacy in Phase II). If possible, up to 4 matched controls per case will be selected. This 4:1 ratio represents the most statistically efficient ratio while minimizing bias introduced to the study^{27,28}. Controls will be frequency matched based on year of publication (+/- 5 years for each case, disease of interest, and general area of biomedical research (i.e. cardiology, cancer, immunotherapy, etc.)). These stratifying factors are not predictor variables but have been chosen to ensure the comparability of cases and controls in the sample. Following matching, controls will be randomly selected using a computer-generated algorithm for those cases with >4 matched controls.

Variables. The primary analysis will investigate the association between a set of variables thought to influence translation, and successful translation. It is hypothesized that the following four sets of variables are of high importance:

- i. **Internal validity:** Similarly to clinical studies, preclinical studies that lack methodological rigor usually demonstrate the largest measures of efficacy²⁹⁻³¹. However, this lack of methodological rigor may reduce the likelihood that these studies can be translated. In our study, risk of bias will be assessed by applying the Cochrane Risk of Bias tool²⁰ – modified for suitability to preclinical studies. We will assess six domains identified by this tool (sequence generation, allocation concealment, blinding of personnel, blinding of outcome assessors, incomplete outcome data, and selective outcome reporting), as well as two additional risk of bias domains (conflict of interest and sample size calculation) for a total of eight risk of bias domains. Again, this checklist will be operationalized into a binary outcome; if the majority (i.e. >4) of items on the checklist are met the study will be considered to have high internal validity.
- ii. **External validity:** This will be assessed by the experiment(s) within the preclinical study being performed across a range of conditions: the presence of dose response relationship, more than one animal model (species or method of disease induction) used, male and female animals (or the appropriate sex for sex specific diseases, e.g. prostate or ovarian cancer), and co-morbid animals³². This outcome will be dichotomized, (i.e. if two or more of these four criteria are met, the study will be considered to have high external validity).
- iii. **Funding source:** Evidence suggests that industry involvement may lead to faster translation⁵. This relationship will be examined by determining if the preclinical study involving the promising therapy was industry, government, or philanthropy funded (with

attention also given to whether industry funding was from pharmaceutical or biotechnology companies).

- iv. *Completeness of reporting:* The National Institutes for Health preclinical reporting guidelines will be used to assess completeness of reporting. We have previously operationalized this list into 21 yes or no questions. This outcome will be dichotomized as high or low completeness of reporting when the majority of items (≥ 11) are reported upon or not. As a pre-planned sensitivity analysis, we will also evaluate the preclinical studies as high completeness of reporting when they report on over two thirds of the checklist items (≥ 14).

For the exposure items of validity (internal and external) and reporting the individual items in each checklist will not be weighted.

Exploratory analysis:

- i. Individual components of internal validity: We will assess two selected risk of bias domains (sequence generation and blinding of outcome assessor) individually.
- ii. Individual components of external validity: We will assess all components of external validity individually, rather than as an aggregate outcome.
- iii. *Number of authors and institutions:* We will use the number of co-authors and the number of affiliated institutions on the preclinical paper as an approximation for the degree of collaboration within the preclinical study. With this, we plan to evaluate how the degree of collaboration within a preclinical study affects translation.
- iv. *H-index:* We will use the corresponding author's H-index as an approximation for investigator seniority. With this, we plan to evaluate how the corresponding author's seniority affects preclinical to clinical translation.
- v. *Number of citations:* We will use the number of citations associated with the preclinical paper as an approximation for the degree of dissemination. With this we plan to evaluate how the degree of the preclinical study's dissemination affects translation.

Data analysis and sample size

Sample size. In a preliminary screen of *Science* and *Nature*, 482 promising therapies (184 *Science*; 298 *Nature*) were identified. Application of a conservative estimate suggests that the addition of the third journal (*Nature Medicine*) will provide an additional 150 promising therapies, for a total of 632. In the previous report citing a 5% translation rate¹⁸, 19 of 101 (19%) promising therapies reached the clinical trial stage and demonstrated favorable results. Methodological differences between this study and the previous report may lead to a decrease in the percentage of promising therapies demonstrating favorable results at phase II. A conservative estimate of half the rate from the previous report (9.5%) suggests that roughly 60 of the promising therapies from our sample would demonstrate promising results at phase II or later. Generally, it

is recommended that a regression model contain 10 events per variable^{33,34}. Considering the four variables listed above, we will require 40 cases in order to be adequately powered to perform our proposed analysis.

If an adequate sample size of 40 cases is not reached, we will continue searching for promising preclinical therapies in order to identify additional phase II or later translated therapies. The CAMARADES group has used a machine learning algorithm³⁵ to identify a cohort of over 100,000 reports of *in vivo* research available on PubMed Central. From this we will randomly select reports of *in vivo* preclinical research and assess these for clinical translation. The number of studies randomly selected will depend on the number of additional case required. We will sample iteratively until our predetermined minimal set of 40 cases is obtained.

Analytic plan. A comparison of baseline variables (aside from matching factors) in the cases and controls will be assessed using frequency distributions and univariate descriptive statistics including measures of central tendency and dispersion. Multivariable conditional logistic regression will be performed to determine the adjusted association of our factors to the outcome variable. Odds ratios with accompanying 95% confidence intervals will be calculated. A sensitivity analysis will be performed excluding cases that had a positive clinical trial but had not received regulatory approval. All "analyses will be performed using SAS (version 9.4). Data will be presented as odds-ratios and 95% confidence intervals in forest plots.

Limitations

Our focus on leading journals (*Science*, *Nature* and *Nature Medicine*) may be perceived as a limitation. Given the usually positive findings and higher visibility of studies published therein, this may lead to an over-estimate in the bench-to-bedside translation. Although this selection of journals may represent an over-estimate of translation, we also believe it represents the 'best-in-class' of preclinical research. Alternatively, the reliance upon publicly available data may result in an under-estimate of the bench-to-bedside transition as it is possible that preclinical work from within industry that does not proceed to a clinical trial may not be published. Without examining this omitted data, we may not be presenting a fully accurate assessment of the state of preclinical to clinical trial translation.

Another limitation is that components of internal validity have historically been poorly reported in bench research³⁶, which may not be an accurate reflection of what investigators actually did. A third limitation is that the dichotomization of variables in the case-control could be regarded as an oversimplification of complex concepts. However, our exploratory analyses will allow us to investigate a larger number of factors in more granular detail. Furthermore, our assessment on the external validity and how this may affect translation rests on the assumption that the rationale for initiating a clinical trial is based solely on one preclinical study, rather than several studies from different labs under a range of conditions. In reality, external validity would be best established through multiple preclinical

experiments rather than one with a high level of external validity.

Another important aspect affecting the quality of preclinical evidence is construct validity. Studies have demonstrated that poor construct validity affects reproducibility and downstream clinical translation^{37,38}. Though a checklist to assess construct validity has been developed³², after pilot testing this tool we found that it was infeasible for our study, as it required content expertise for every promising therapeutic to be used appropriately. Thus, another limitation of this study is that we are unable to evaluate the construct validity of the preclinical studies and assess how this aspect affects the translation into clinical trials.

Lastly, it should be acknowledged that the translational spectrum involves many complex steps and various phases. We have chosen to focus on the translation of the *in vivo* animal experiment stage within T0 research; to the stage of therapy evaluation in clinical trials, within T2 research. When we deem a successful translation from preclinical to clinical research, we do not evaluate all stages of preclinical nor clinical research. Furthermore, successful translation to a Phase II clinical trial does not imply that the therapy has been successfully translated to practice or regulatory approval. With this limitation in mind, we will focus on preclinical issues that may affect early translation, rather than potential issues with the clinical trials that subsequently evaluated the therapies in human subjects. We anticipate a future study that will critically analyze the translational clinical trials identified in Objective #1 to address issues associated with those trials.

Dissemination

We will have a one-day, in-person meeting between the investigators and identified knowledge users (Stem Cell Network and BioCanRx, two Government of Canada funded Networks of Centres of Excellence focused on the development of novel therapeutics). Over the one-day meeting we will examine data and results generated by the project. In addition, each knowledge user will speak about their organization's experiences and perspectives on bench-to-bedside translation. The meeting will end with a discussion to identify next steps to be taken by stakeholders to improve translation.

We anticipate the publication of two key papers submitted to appropriate peer-reviews journals. The first will describe key findings of the systematic review and present the results of our case-control study. The second will be a policy piece to describe how our findings may affect stakeholders across the spectrum of bench-to-bedside research. The results of this study will be presented at relevant national and international scientific meetings to promote knowledge transfer.

Amendments

If amendments are required for this protocol, the date of each amendment will be provided with a description for rationale for the change in this section.

Study status

With our systematic search strategy, we have searched the journals *Science*, *Nature*, and *Nature Medicine* and are currently

screening articles to identify promising therapies as per Objective #1 of this study.

Discussion

This study is an innovative project that will use knowledge synthesis methods to identify the incidence of successful translation, map contemporary trends in translation, and identify key modifiable factors associated with successful and unsuccessful bench-to-bedside translation. Additionally, the proposed study will generate the largest dataset to date of promising preclinical therapies and then apply a case-control design to investigate the association of validities with successful or failed translation. This evidence is required to implement a "knowledge-to-action"^{39–41} cycle to improve bench-to-bedside translational research.

Current lags in translation are unknown. Some of our own efforts (and others) at translation suggest that these lags between preclinical findings and first-in-human trials may be shortening^{26,42–45}. This reduction may reflect increased funding towards translational activities, improved research rigor, as well as regulatory changes that have accelerated translation efforts (e.g. FDA's four unique approaches of Fast Track, Breakthrough Therapy, Accelerated Approval, and Priority Review). If this hypothesis is supported by the data, all stakeholders need information about the factors that support improvements in development timelines.

The effect of evolving preclinical reporting guidelines on translation is unknown. In the last decade, funders, journals, and other stakeholders have endorsed reporting guidelines^{36,46} that improve transparency and potentially improve the ability to forecast which interventions may translate successfully. These guidelines have been developed in response to multiple publications, which demonstrated that the majority of basic science findings are irreproducible^{47–49}. Previous studies of bench-to-bedside translation were conducted prior to these guidelines being widely endorsed; thus, a current study is needed to determine if increased awareness and emphasis regarding preclinical reporting and design has improved translation.

Taken together, there is a strong need and justification for the proposed study. This study will provide a contemporary understanding of bench-to-bedside translation, which is required to tailor experimental designs to improve translational efficiency, including the education and uptake of available study design and analysis methods.

Data availability

Underlying data

No data is associated with this article.

Extended data

Open Science Framework: Identifying and understanding factors that affect the translation of therapies from the laboratory to patients, <https://doi.org/10.17605/OSF.IO/WGZ3T>²².

This project contains the following extended data:

- Representative search strategy in PubMed

- Data selection items
- Algorithm to identify clinical translation of promising bench findings

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC0 1.0 Public domain dedication).

References

- National Center for Advancing Translational Sciences: **Translational Science Spectrum**. 2018.
[Reference Source](#)
- Bonter K, Breckenridge Z, Lachance S, *et al.*: **Opportunities and challenges for the cellular immunotherapy sector: a global landscape of clinical trials**. *Regen Med*. 2017; 12(6): 623–636.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bubela T, Bonter K, Lachance S, *et al.*: **More Haste, Less Speed: Could Public-Private Partnerships Advance Cellular Immunotherapies?** *Front Med (Lausanne)*. 2017; 4: 134.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ioannidis JP: **Evolution and translation of research findings: from bench to where?** *PLoS Clin Trials*. 2006; 1(7): e36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Contopoulos-Ioannidis DG, Ntzani E, Ioannidis JP: **Translation of highly promising basic science research into clinical applications**. *Am J Med*. 2003; 114(6): 477–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Freedman LP, Cockburn IM, Simcoe TS: **The Economics of Reproducibility in Preclinical Research**. *PLoS Biol*. 2015; 13(6): e1002165.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Health Economics Research Group, Office of Health Economics & RAND Europe: **Medical research: what's it worth? Estimating the economic benefits from medical research in the UK**. 2008; London: UK Evaluation Forum,
[Reference Source](#)
- Bubela T, *et al.*: **Bringing regenerative medicines to the clinic: the future for regulation and reimbursement**. *Regen Med*. 2015; 10(7): 897–911.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kimmelman J, London AJ: **The structure of clinical translation: efficiency, information, and ethics**. *Hastings Cent Rep*. 2015; 45(2): 27–39.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hay M, Thomas DW, Craighead JL, *et al.*: **Clinical development success rates for investigational drugs**. *Nat Biotechnol*. 2014; 32(1): 40–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hackam DG, Redelmeier DA: **Translation of research evidence from animals to humans**. *JAMA*. 2006; 296(14): 1731–1732.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Yarborough M, Bredenoord A, D'Abramo F, *et al.*: **The bench is closer to the bedside than we think: Uncovering the ethical ties between preclinical researchers in translational neuroscience and patients in clinical trials**. *PLoS Biol*. 2018; 16(6): e2006343.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Feuerstein GZ, Zaleska MM, Krams M, *et al.*: **Missing steps in the STAIR case: a Translational Medicine perspective on the development of NXY-059 for treatment of acute ischemic stroke**. *J Cereb Blood Flow Metab*. 2008; 28(1): 217–219.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chalmers I, Bracken MB, Djulbegovic B, *et al.*: **How to increase value and reduce waste when research priorities are set**. *Lancet*. 2014; 383(9912): 156–165.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Moher D, Glasziou P, Chalmers I, *et al.*: **Increasing value and reducing waste in biomedical research: who's listening?** *Lancet*. 2016; 387(10027): 1573–1586.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ioannidis JP, Greenland S, Hlatky MA, *et al.*: **Increasing value and reducing waste in research design, conduct, and analysis**. *Lancet*. 2014; 383(9912): 166–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Glasziou P, Altman DG, Bossuyt P, *et al.*: **Reducing waste from incomplete or unusable reports of biomedical research**. *Lancet*. 2014; 383(9913): 267–76.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Contopoulos-Ioannidis DG, Ntzani E, Ioannidis JP: **Translation of highly promising basic science research into clinical applications**. *Am J Med*. 2003; 114(6): 477–484.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Morris ZS, Wooding S, Grant J: **The answer is 17 years, what is the question: understanding time lags in translational research**. *J R Soc Med*. 2011; 104(12): 510–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Higgins J, Green S: **Cochrane Handbook for Systematic Reviews of Interventions**. The Cochrane Collaboration, 2015.
[Reference Source](#)
- Hooijmans CR, Tillema A, Leenaars M, *et al.*: **Enhancing search efficiency by means of a search filter for finding all studies on animal experimentation in PubMed**. *Lab Anim*. 2010; 44(3): 170–175.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lalu MM, Hunniford V, Ferguson D, *et al.*: **Identifying and understanding factors that affect the translation of therapies from the laboratory to patients: a study protocol - Extended Data**. 2020.
<http://www.doi.org/10.17605/OSF.IO/WGZ3T>
- Fergusson DA, Wesch NL, Leung GJ, *et al.*: **Assessing the Completeness of Reporting in Preclinical Oncolytic Virus Therapy Studies**. *Mol Ther Oncolytics*. 2019; 14: 179–187.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- U.S. Food and Drug Administration: **The Drug Development Process. Step 3: Clinical Research**. 2019.
[Reference Source](#)
- Moher D, Liberati A, Tetzlaff J, *et al.*: **Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement**. *J Clin Epidemiol*. 2009; 62(10): 1006–1012.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lalu MM, Sullivan KJ, Mei SH, *et al.*: **Evaluating mesenchymal stem cell therapy for sepsis with preclinical meta-analyses prior to initiating a first-in-human trial**. *eLife*. 2016; 5: pii: e17850.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Austin PC: **Statistical criteria for selecting the optimal number of untreated subjects matched to each treated subject when using many-to-one matching on the propensity score**. *Am J Epidemiol*. 2010; 172(9): 1092–1097.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rassen JA, Shelat AA, Myers J, *et al.*: **One-to-many propensity score matching in cohort studies**. *Pharmacoepidemiol Drug Saf*. 2012; 21(Suppl 2): 69–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Crossley NA, Sena E, Goehler J, *et al.*: **Empirical evidence of bias in the design of experimental stroke studies: a metaepidemiologic approach**. *Stroke*. 2008; 39(3): 929–934.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hirst JA, Howick J, Aronson JK, *et al.*: **The need for randomization in animal trials: an overview of systematic reviews**. *PLoS One*. 2014; 9(6): e98856.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Macleod MR, O'Collins T, Horky LL, *et al.*: **Systematic review and metaanalysis of the efficacy of FK506 in experimental stroke**. *J Cereb Blood Flow Metab*. 2005; 25(6): 713–721.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Henderson VC, Kimmelman J, Fergusson D, *et al.*: **Threats to validity in the design and conduct of preclinical efficacy studies: a systematic review of guidelines for in vivo animal experiments**. *PLoS Med*. 2013; 10(7): e1001489.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Concato J, Peduzzi P, Holford TR, *et al.*: **Importance of events per independent variable in proportional hazards analysis. I. Background, goals, and general strategy**. *J Clin Epidemiol*. 1995; 48(12): 1495–501.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Peduzzi P, Concato J, Kemper E, *et al.*: **A simulation study of the number of events per variable in logistic regression analysis**. *J Clin Epidemiol*. 1996; 49(12): 1373–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bannach-Brown A, Przybyla P, Thomas J, *et al.*: **Machine learning algorithms for systematic review: reducing workload in a preclinical review of animal studies and reducing human screening error**. *Syst Rev*. 2019; 8(1): 23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kilkenny C, Browne WJ, Cuthill IC, *et al.*: **Improving bioscience research reporting: the ARRIVE guidelines for reporting animal research**. *PLoS Biol*. 2010; 8(6): e1000412.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Scott S, Kranz JE, Cole J, *et al.*: **Design, power, and interpretation of studies in the standard murine model of ALS**. *Amyotroph Lateral Scler*. 2008; 9(1): 4–15.
[PubMed Abstract](#) | [Publisher Full Text](#)

38. Reynolds JC, Rittenberger JC, Menegazzi JJ: **Drug administration in animal studies of cardiac arrest does not reflect human clinical experience.** *Resuscitation*. 2007; **74**(1): 13–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Canadian Institutes of Health Research: **Guide to Knowledge Translation Planning at CIHR: Integrated and End-of-Grant Approaches.** 2012. (Cat. No. MR4-11/2012E-PDF).
[Reference Source](#)
40. Canadian Institutes of Health Research: **Integrating Ethics and the Knowledge-To-Action Cycle.** 2018.
[Reference Source](#)
41. Straus S, Tetroe J, Graham ID: **Knowledge translation in health care: moving from evidence to practice.** John Wiley & Sons, 2013.
[Reference Source](#)
42. Taljaard M, Ward MR, Kutryk MJ, *et al.*: **Rationale and design of Enhanced Angiogenic Cell Therapy in Acute Myocardial Infarction (ENACT-AMI): the first randomized placebo-controlled trial of enhanced progenitor cell therapy for acute myocardial infarction.** *Am Heart J*. 2010; **159**(3): 354–360.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Granton J, Langleben D, Kutryk MB, *et al.*: **Endothelial NO-Synthase Gene-Enhanced Progenitor Cell Therapy for Pulmonary Arterial Hypertension: The PHACeT Trial.** *Circ Res*. 2015; **117**(7): 645–654.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Eiger BioPharmaceuticals: **A Phase 2, Randomized, Double-Blind, Placebo-Controlled Study of UBenimex in Patients With Pulmonary ARterial HYpertension (WHO Group 1) (LIBERTY).** *ClinicalTrials.gov*. 2018.
[Reference Source](#)
45. Tian W, Jiang X, Tamosiuniene R, *et al.*: **Blocking macrophage leukotriene b₄ prevents endothelial injury and reverses pulmonary hypertension.** *Sci Transl Med*. 2013; **5**(200): 200ra117.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. National Institutes of Health: **Principles and Guidelines for Reporting Preclinical Research.** 2016.
[Reference Source](#)
47. Ioannidis JP, Allison DB, Ball CA, *et al.*: **Repeatability of published microarray gene expression analyses.** *Nat Genet*. 2009; **41**(2): 149–155.
[PubMed Abstract](#) | [Publisher Full Text](#)
48. Prinz F, Schlange T, Asadullah K: **Believe it or not: how much can we rely on published data on potential drug targets?** *Nat Rev Drug Discov*. 2011; **10**(9): 712.
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Begley CG, Ellis LM: **Drug development: Raise standards for preclinical cancer research.** *Nature*. 2012; **483**(7391): 531–533.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 23 September 2020

<https://doi.org/10.5256/f1000research.26110.r69847>

© 2020 Lal J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Jonathan A Lal

Department of Molecular and Cellular Engineering, Jacob Institute of Biotechnology and Bioengineering, Sam Higginbottom University of Agriculture, Technology and Sciences, Allahabad, Uttar Pradesh, India

Reference 4 is too old (from 2006) to claim that translation from preclinical to human trials is rare. This should be updated.

Reference 6 is 5 years old, should work, but if more recent data is available, perhaps replace. Reference 7 is too old to be valid in today's economic development, please update.

For methods it says any changes will be included in the final report. It is to be noted that the peer review is limited to the manuscript itself.

Eligibility criteria limited to 3 prominent journals limits the scope as JAMA and other such high impact factor (preclinical/clinical) journals are left out. At least 5 journals could have been taken into consideration to justify the sample size.

What would be interesting to see is the ratio between successful Phase II trials and those successful Phase II trials which actually reach regulatory approvals.

Furthermore, regulatory approval may not still guarantee widespread adoption. The latter of which could be due to other factors which remain to be explored, including competition. A comment towards this end would be advised.

In limitations it mentions preclinical to Phase II, and eventually analysing the preclinical problems till Phase II. Phase III and/or Phase IV excluded, regulatory touched upon, it seems this should be made clear at the start of the manuscript, that the focus is till Phase II and not bedside directly.

Is the rationale for, and objectives of, the study clearly described?

Yes

Is the study design appropriate for the research question?

Yes

Are sufficient details of the methods provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Not applicable

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 08 Oct 2020

Manoj Lal, Ottawa General Hospital Research Institute, Ottawa, Canada

We thank you for your careful assessment of our protocol and their supportive comments. We have addressed your queries below and within the revised manuscript.

Reference 4 is too old (from 2006) to claim that translation from preclinical to human trials is rare. This should be updated.

Updated citation to: Seyhan, A.A. Lost in translation: the valley of death across preclinical and clinical divide – identification of problems and overcoming obstacles. transl med commun 4, 18 (2019). <https://doi.org/10.1186/s41231-019-0050-7>

Reference 6 is 5 years old, should work, but if more recent data is available, perhaps replace.

Updated citation to: Hutchins BI, Davis MT, Meseroll RA, Santangelo GM (2019) Predicting translational progress in biomedical research. PLoS Biol 17(10): e3000416. <https://doi.org/10.1371/journal.pbio.3000416>

Reference 7 is too old to be valid in today's economic development, please update.

We agree with the reviewer that this is an old reference. We believe this may not have been the strongest argument to present in the introduction, thus we have removed the sentences on return on investments.

For methods it says any changes will be included in the final report. It is to be noted that the peer review is limited to the manuscript itself.

Changed sentence to: *Any deviation from this protocol will be documented in the manuscript that reports the results of this study, upon its completion.*

Eligibility criteria limited to 3 prominent journals limits the scope as JAMA and other such high impact factor (preclinical/clinical) journals are left out. At least 5 journals could have been taken into consideration to justify the sample size.

Our group of investigators considered this very issue at length. Ultimately, there simply is no perfect sampling strategy to address the objectives of the study. We

selected the three journals as they are highly influential and considered 'best-in-class' by many. The number of articles resulting from the search of the three included journals is 20,341. With our available resources, it has taken close to one year to screen all abstracts and full texts in duplicate. Adding more journals would not be feasible with our finite resources. We added the following to the limitation section of the discussion:

Furthermore, our sample is limited to three journals, which may reduce the scope of our work. As automation of meta-research projects gains traction (and tools are validated) broader searches will certainly be possible in the future. However, given that this process currently relies on personnel we are limited to the current strategy.

What would be interesting to see is the ratio between successful Phase II trials and those successful Phase II trials which actually reach regulatory approvals.

We certainly agree with this suggestion. This is something we will report on in the manuscript that summarizes the results of this study. We added the following sentence in the analysis plan of Objective 1:

Additionally, we will calculate the proportion of successful Phase II trials that received regulatory approval.

Furthermore, regulatory approval may not still guarantee widespread adoption. The latter of which could be due to other factors which remain to be explored, including competition. A comment towards this end would be advised.

We also agree with this comment and have addressed this limitation in our discussion:

Furthermore, successful translation to a Phase II clinical trial does not imply that the therapy has been successfully translated to practice or regulatory approval; and regulatory approval may not guarantee widespread adoption. With this limitation in mind, we will focus on preclinical issues that may affect early translation, rather than potential issues with the clinical trials that subsequently evaluated the therapies in human subjects or barriers that affect both regulatory approval and adoption into practice.

In limitations it mentions preclinical to Phase II, and eventually analysing the preclinical problems till Phase II. Phase III and/or Phase IV excluded, regulatory touched upon, it seems this should be made clear at the start of the manuscript, that the focus is till Phase II and not bedside directly.

We would argue that Phase II is considered bedside as the evaluation is being completed on patients. However, we do agree that this is quite different from full clinical adoption and therefore should not be conflated with widespread approval. As such we have modified the following in our manuscript:

Lastly, it should be acknowledged that the translational spectrum involves many complex steps and various phases. Though ultimate clinical translation is the approval and adoption of a therapy, we will focus on a specific stage in the translational spectrum⁴. We have chosen to measure the translation of the in vivo animal experiment stage within T0 research; to the evaluation of the therapies' efficacy in clinical trials, within T2 research.

Competing Interests: None to disclose.

Reviewer Report 17 August 2020

<https://doi.org/10.5256/f1000research.26110.r69397>

© 2020 Holcomb J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

John Holcomb

Division of Acute Care Surgery, Department of Surgery, University of Alabama at Birmingham, Birmingham, AL, USA

The authors have embarked on an important project. They will evaluate the issues related to translation of preclinical data into clinical practice.

I have several comments.

1. The authors should clearly state the narrow focus of their work, largely in the drug development arena. They seem to exclude other research areas such as devices and blood products. Please inform the reader what focus areas are included and excluded.
2. The authors have included only several journals to review. Given the ability to use electronic databases, I would suggest they broaden their search.
3. Lastly, the definition of clinical translation is not completion of phase 2 studies, but rather regulatory approval and adoption into clinical practice. Defining adoption itself is difficult, and should be discussed.

Is the rationale for, and objectives of, the study clearly described?

Yes

Is the study design appropriate for the research question?

Partly

Are sufficient details of the methods provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: injury, resuscitation,

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 08 Oct 2020

Manoj Lal, Ottawa General Hospital Research Institute, Ottawa, Canada

We thank you for your careful assessment of our protocol and their supportive comments. We have addressed your queries below and within the revised manuscript.

The authors have embarked on an important project. They will evaluate the issues related to translation of preclinical data into clinical practice. I have several comments.

1. The authors should clearly state the narrow focus of their work, largely in the drug development arena. They seem to exclude other research areas such as devices and blood products. Please inform the reader what focus areas are included and excluded.

We apologize for the lack of clarity on our inclusion criteria. Our search and strategy for selecting therapies is designed to consider all interventions that have therapeutic benefit. We added blood products and devices in the examples of 'promising therapies' in the inclusion criteria.

A 'promising therapy', defined by the following:

a. Any therapy introduced to the animal model (e.g. pharmacologic and non-pharmacologic therapies, vaccines, antibodies, blood products, implants/devices, etc.), which was still at the developmental stage and did not have a prior application in humans for the specific indication.

1. The authors have included only several journals to review. Given the ability to use electronic databases, I would suggest they broaden their search.

Our group of investigators considered this very issue at length. Ultimately, there simply is no perfect sampling strategy to address the objectives of the study. We selected the three journals as they are highly influential and considered 'best-in-class' by many. The number of articles resulting from the search of the three included journals is 20,341. With our available resources, it has taken close to one year to screen all abstracts and full texts in duplicate. Adding more journals would not be feasible with our finite resources. We added the following to the limitation section of the discussion:

Furthermore, our sample is limited to three journals, which may reduce the scope of our work. As automation of meta-research projects gains traction (and tools are validated) broader searches will certainly be possible in the future. However, given that this process currently relies on personnel we are limited to the current strategy.

1. Lastly, the definition of clinical translation is not completion of phase 2 studies, but

rather regulatory approval and adoption into clinical practice. Defining adoption itself is difficult, and should be discussed.

We agree that full clinical translation is not the completion of a Phase II trial and we also do agree that ultimate clinical translation is the clinical adoption of a therapy. In the current study we are focussed on the translation of preclinical research to first studies demonstrating efficacy in clinical trials. Full clinical translation to approved therapies will also be tracked and we will certainly be reporting on this metric as well.

We have added/modified a sentence in the manuscript to further explain our intention:

Therefore, the aim of this project is to use established knowledge synthesis methods to identify the prevalence of translation from preclinical research (bench) to the first clinical evaluations of efficacy (bedside), map contemporary trends in translation, and identify key modifiable factors associated with successful and unsuccessful bench-to-bedside translation.

We added the following sentence in the analysis plan of Objective 1:

Additionally, we will calculate the proportion of successful Phase II trials that received regulatory approval.

We also added and modified the following in our limitations section:

Lastly, it should be acknowledged that the translational spectrum involves many complex steps and various phases. Though ultimate clinical translation is the approval and adoption of a therapy, we will focus on a specific stage in the translational spectrum⁴. We have chosen to measure the translation of the in vivo animal experiment stage within T0 research to the first evaluation of the therapy's efficacy in clinical trials, within T2 research.

Competing Interests: None to disclose.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research