REVIEW

Drug Therapies for the Management of Sickle Cell Disease

[version 1; peer review: 2 approved]

Parul Rai ¹, Kenneth I. Ataga ²

¹Department of Hematology, St. Jude Children's Research Hospital, Memphis, TN, USA

²Center for Sickle Cell Disease, University of Tennessee Health Science Center, Memphis, TN, USA

v1 First published: 11 Jun 2020, 9(F1000 Faculty Rev):592
<https://doi.org/10.12688/f1000research.22433.1>

Latest published: 11 Jun 2020, 9(F1000 Faculty Rev):592
<https://doi.org/10.12688/f1000research.22433.1>

Abstract

Sickle cell disease (SCD) afflicts millions of people worldwide but is referred to as an orphan disease in the United States. Over the past several decades, there has been an increasing understanding of the pathophysiology of SCD and its complications. While most individuals with SCD in resource-rich countries survive into adulthood, the life expectancy of patients with SCD remains substantially shorter than for the general African-American population. SCD can be cured using hematopoietic stem cell transplantation and possibly gene therapy, but these treatment approaches are not available to most patients, the majority of whom reside in low- and middle-income countries. Until relatively recently, only one drug, hydroxyurea, was approved by the US Food and Drug Administration to ameliorate disease severity. Multiple other drugs (L-glutamine, crizanlizumab, and voxelotor) have recently been approved for the treatment of SCD, with several others at various stages of clinical testing. The availability of multiple agents to treat SCD raises questions related to the choice of appropriate drug therapy, combination of multiple agents, and affordability of recently approved products. The enthusiasm for new drug development provides opportunities to involve patients in low- and middle-income nations in the testing of potentially disease-modifying therapies and has the potential to contribute to capacity building in these environments. Demonstration that these agents, alone or in combination, can prevent or decrease end-organ damage would provide additional evidence for the role of drug therapies in improving outcomes in SCD.

Keywords

Sickle cell disease, Treatment, Novel Drugs, Drug Development, Clinical Trials

Open Peer Review

Approval Status

	1	2
version 1		
11 Jun 2020		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Daniel E. Bauer**, Boston Children's Hospital, Boston, USA
Dana-Farber Cancer Institute, Boston, USA
Harvard Medical School, Boston, USA
2. **Yogen Saunthararajah**, Taussig Cancer Institute, Cleveland Clinic, Cleveland, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Kenneth I. Ataga (kataga@uthsc.edu)

Author roles: Rai P: Investigation, Writing – Original Draft Preparation; Ataga KI: Conceptualization, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: KIA has served on clinical advisory boards and/or served as a consultant for Global Blood Therapeutics, Novartis, Emmaus Life Sciences, Editas Medicine, and Novo Nordisk. PR has no competing interests.

Grant information: This work is supported by an award from the US Food and Drug Administration - R01FD006030 (KIA). *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2020 Rai P and Ataga KI. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Rai P and Ataga KI. **Drug Therapies for the Management of Sickle Cell Disease [version 1; peer review: 2 approved]** F1000Research 2020, 9(F1000 Faculty Rev):592 <https://doi.org/10.12688/f1000research.22433.1>

First published: 11 Jun 2020, 9(F1000 Faculty Rev):592 <https://doi.org/10.12688/f1000research.22433.1>

Introduction

Although referred to as an orphan disease in the United States (US), sickle cell disease (SCD) affects millions of individuals worldwide, with the vast majority residing in sub-Saharan Africa and India¹. SCD is characterized by the presence of sickle hemoglobin (HbS), hemolytic anemia, vaso-occlusive complications, and cumulative end-organ damage. The mortality rate associated with SCD in sub-Saharan Africa remains high, with an estimated 50 to 90% of children dying before the age of 5². However, the majority of children with SCD in resource-rich countries live to adulthood^{3–5}. Despite increased survival to adulthood, individuals with SCD in resource-rich nations continue to have a shorter life expectancy than the general population^{6–9}. There has been substantial progress with the use of allogeneic bone marrow transplantation as a curative therapy in SCD, and increasing evidence supports the curative potential of gene therapy and gene editing^{10,11}. However, as these modalities are not available to the vast majority of patients, most of whom reside in resource-limited countries, the availability of drug therapies that are safe, effective, and affordable remains highly desirable.

This review will focus on approaches to develop drug therapies in SCD, ongoing and recently completed trials, and our perspective on the use of approved drugs.

Pathophysiology

The development of effective therapies for SCD depends on an adequate understanding of its pathophysiology. Although the pathophysiology of SCD is complex and involves multiple pathways, the primary event is due to the polymerization of HbS following deoxygenation¹². The rate and extent of polymer

formation depends on the degree and duration of HbS deoxygenation, presence of fetal hemoglobin (HbF), and the intracellular concentration of HbS. Clinical manifestations of SCD appear to be driven by two major pathophysiological processes: vaso-occlusion with ischemia-reperfusion injury and hemolytic anemia¹³. Vaso-occlusion occurs because of adhesive interactions of leukocytes and sickle RBCs with the endothelium causing microvascular obstruction and subsequent tissue ischemia¹³. These episodes of vascular obstruction are followed by the restoration of blood flow, which promotes further tissue injury by reperfusion. The inflammatory cascade resulting from ischemia-reperfusion is amplified by the activation of CD1d-restricted invariant natural killer T (iNKT) cells¹⁴. The release of free plasma hemoglobin following intravascular hemolysis results in direct scavenging of nitric oxide (NO), as well as the generation of reactive oxygen species, powerful scavengers of NO^{15,16}. NO is usually produced by the endothelium and regulates basal vasodilator tone as well as inhibits the activation of platelets and the coagulation system and the transcriptional expression of nuclear factor κ B (NF κ B)-dependent adhesion molecules, such as vascular cell-adhesion molecule-1, intercellular cell-adhesion molecule-1, and selectins^{17,18}. HbS polymerization as well as its multiple downstream consequences, including endothelial cell injury, endothelial dysfunction, increased oxidant stress, inflammation, and coagulation and platelet activation, are therapeutic targets in SCD (Figure 1). SCD has been dichotomized into two overlapping sub-phenotypes: viscosity-vaso-occlusion (higher hemoglobin levels, possibly increased blood viscosity, and complications such as osteonecrosis, acute chest syndrome, and acute pain crisis) and hemolysis-endothelial dysfunction (increased hemolysis with lower hemoglobin levels and higher levels of hemolytic markers, including reticulocyte

Figure 1. Potential targets of pharmacologic agents in sickle cell disease. HbF, hemoglobin F; NKT, natural killer T-cells; NO, nitric oxide; RBCs, red blood cells. Adapted with permission from Ataga KI and Stocker J. *Br J Haematol*, 2015.

count and serum lactate dehydrogenase, and complications such as leg ulcers, priapism, stroke, and possibly pulmonary hypertension)¹⁹. While somewhat simplistic, this classification is useful to understand the pathobiology of disease complications and may provide guidance on the effects of therapies on disease-related complications. The pathophysiology of SCD has been reviewed more extensively elsewhere^{30–22}.

Drug therapies for sickle cell disease

Despite an improved understanding of the pathophysiology of SCD, available drug treatments remain limited. For many years, hydroxyurea was the only drug available to modify the severity of SCD²³. However, there has been progress in the development of drug therapies with recent approvals of L-glutamine, crizanlizumab, and voxelotor by the US Food and Drug Administration (FDA)^{24–26}. Based on our current understanding of SCD pathophysiology, there are multiple possible approaches to treating the disease and its complications, including inhibition of HbS polymerization and amelioration of multiple downstream sequelae of HbS polymerization.

Inhibition of sickle hemoglobin polymerization

Therapeutic approaches targeting HbS polymerization, including 1) blocking intermolecular contacts in the sickle fiber, 2) induction of HbF synthesis, 3) reduction of intracellular HbS concentration, 4) increase in oxygen affinity, and 5) reduction of the concentration of 2,3-diphosphoglycerate, have been described²⁷. Table 1 lists ongoing studies of anti-sickling agents in SCD.

Induction of fetal hemoglobin. High levels of HbF ameliorate the severity of SCD^{8,28,29}. Hydroxyurea strongly induces HbF³⁰, although the mechanisms by which it does so are still unclear^{30–33}. Hydroxyurea improves erythrocyte deformability, lowers hemolysis, decreases circulating leukocytes and reticulocytes³⁴, reduces adhesion receptor expression^{35–38}, and is an NO donor^{39,40}. In two placebo-controlled, phase III trials, hydroxyurea significantly reduced the frequency of vaso-occlusive crises, acute chest syndrome, blood transfusion requirement, and hospitalizations in adults and children with SCD^{23,41}. Treatment with low, fixed weight-based dosing of hydroxyurea (10 mg/kg daily) also decreased the frequency of SCD-related complications with low toxicity^{42,43}. Hydroxyurea further ameliorates SCD morbidity by decreasing the risk of stroke and chronic kidney disease^{44,45} and may reduce mortality in SCD^{46–49}. Recent studies to optimize hydroxyurea therapy are exploring dose maximization via pharmacokinetics-based dosing^{50–54}. Despite its multiple benefits, a substantial number of patients on hydroxyurea may not obtain an adequate clinical response.

Inhibitors of epigenetic enzymes which repress γ -globin genes, including DNA methyltransferase (DNMT1), histone deacetylases, and lysine-specific demethylase (LSD-1), and γ -globin gene co-activator agonists are being evaluated in SCD. Decitabine (5-aza-2'-deoxycytidine), a DNA hypomethylating agent which depletes DNMT1, increased HbF and total hemoglobin levels at non-cytotoxic doses in patients who

had no responses to hydroxyurea⁵⁵. However, it has a very short plasma half-life and negligible oral bioavailability due to rapid inactivation by cytidine deaminase (CDA)⁵⁵. The combination of escalating doses of oral decitabine and an oral CDA inhibitor, tetrahydrouridine (THU), was well tolerated, and the highest tested dose increased HbF, doubled F-cells by approximately 80% of total RBCs, increased total hemoglobin, and decreased biomarkers of hemolysis, coagulation activation, and inflammation⁵⁶.

Histone deacetylase (HDAC) inhibitors increase HbF levels^{57–63}. Although treatment with intermittent doses of sodium butyrate produced sustained increases in HbF, F-cells, and total hemoglobin⁵⁷, the orally available sodium dimethyl butyrate (HQB-1001) did not significantly increase HbF and appeared to result in more pain crisis compared with placebo⁶³. A phase II study of the global HDAC inhibitor vorinostat was terminated early owing to poor accrual. Selective HDAC1/2 inhibition with ACY-957 increased HbF *in vitro* and represents a promising therapeutic approach with a better safety profile⁶⁴. Based on the success of the LSD-1 inhibitors RN-1 and ORY-3001 in increasing HbF and F-cells in preclinical studies^{65–67}, INCB059872 was investigated (NCT03132324). However, this phase I study was terminated early for business reasons. Pomalidomide, a third-generation immunomodulatory drug, produced modest increases of HbF, augmented erythropoiesis, and preserved bone marrow function following 8 weeks of treatment in transgenic sickle mice⁶⁸. Pomalidomide appeared to increase HbF and total hemoglobin only at the highest dose or with greater than 6 months of exposure⁶⁹. The safety and efficacy of increasing the expression of γ -globin gene co-activators—FOXO3 agonist⁷⁰ (metformin [NCT02983129]), nuclear factor-like 2 (Nrf2) agonist⁷¹ (dimethyl fumarate), and SIRT1⁷²—to augment HbF levels are being explored.

Allosteric modification of sickle hemoglobin to its high-oxygen affinity R-state. Voxelotor, recently approved under the FDA's accelerated approval program, is an orally bioavailable small molecule which binds to α -globin chains of hemoglobin, increases hemoglobin oxygen affinity, and stabilizes the oxyhemoglobin state⁷³. Early studies showed a reduction in markers of hemolysis, with a rapid rise in hemoglobin by day 15⁷⁴. This response was not accompanied by worsening tissue hypoxia, based on serum erythropoietin levels and change in oxygen consumption (VO_2 max), even in participants receiving daily doses of 1,000 mg for 28 days and 900 mg for at least 90 days. Furthermore, voxelotor has been shown to reduce blood viscosity *in vitro*⁷⁵. In a multicenter, phase III study of patients 12 to 65 years of age, randomly assigned to once-daily oral voxelotor doses of 1,500 mg, 900 mg, or placebo, a significantly higher percentage of subjects on 1,500 mg had a hemoglobin response of 1.0 g/dL at 24 weeks compared with placebo⁷⁶. In addition, subjects on the 1,500 mg dose saw significant reductions in indirect bilirubin and reticulocyte counts from baseline. A dose-dependent increase in hemoglobin following treatment with voxelotor was not accompanied by an increase in pain crisis rate. As this trial was not enriched for subjects with frequent pain crises, appropriately designed studies

Table 1. Ongoing clinical trials of anti-sickling agents in sickle cell disease.

Mechanism	Drug	Sponsor	NCT number (study acronym)	Clinical phase/status	Intervention	Number/age	Objective
HbF induction	Nicotinamide vs. THU and decitabine	EpiDestiny, Inc; NIH; NHLBI	NCT04055818	Phase I Recruiting	Oral nicotinamide vs. THU + decitabine for 12 weeks followed by combination for a further 12 weeks	20/≥18 years	Compare effect of oral nicotinamide vs. THU-decitabine and in combination on hemoglobin level at week 12
	Panobinostat (LBH589)	Abdullah Kutlar	NCT01245179	Phase I Active, not recruiting	Oral, for 12 weeks	18/≥18 years	Evaluate safety, HbF, F cells, total hemoglobin, markers of inflammation, QOL measures
	Metformin	Baylor College of Medicine	NCT02981329 (FITMet)	Phase I Recruiting	Hydroxyurea + metformin vs. metformin	56/10–60 years	Change in HbF or total hemoglobin, QOL, RNA sequencing
Allosteric modifier (to the R-state)	Voxelotor (formerly GBT440)	Global blood therapeutics	NCT03943615 (expanded access)	Approved for marketing	Oral	≥12 years	To provide early access to patients without alternative treatment options
			NCT04247594	Phase II Recruiting	Oral, open-label	45/18–60 years	Dose escalation study to evaluate safety and tolerability of doses, 1,500 mg to 3,000 mg daily
			NCT02850406	Phase IIa Recruiting	Part A: single dose Part B: 24 weeks Part C: 48 weeks	125/4–17 years	Pharmacokinetics, change in hemoglobin, effect on hemolysis, TCD velocity, safety
			NCT03573882	Phase III Active, not recruiting	Oral, daily	179/≥12 years	Open-label extension study, adverse events, frequency of SCD-related complications
			NCT04218084 (HOPE kids 2)	Phase III Not yet recruiting	Oral voxelotor vs. placebo	224/2–14 years	Evaluate effect on TCD in children
			NCT04188509	Phase III Enrolling by invitation	Oral, open-label	50/4–18 years	Evaluate safety and tolerability, SCD-related complications
			NCT04000165	Phase I Recruiting	Oral, with two dose escalations after 2 weeks	25/≥18 years	Safety and tolerability, pharmacokinetics, and pharmacodynamics
Allosteric activator of RBC pyruvate kinase-R	AG-348 (Mitapivat sulfate)	NHLBI		Phase I Recruiting	Single and multiple ascending oral doses of FT-4202 vs. placebo	130/12–60 years	Safety, pharmacokinetics, pharmacodynamics
RBC ion transport channels	FT-4202	Forma therapeutics	NCT03815695	Phase I Recruiting	Part A: open-label, dose-escalation study Part B: randomized, placebo-controlled, cross-over study	60/18–55 years	Safety, change in hemoglobin, markers of hemolysis, QOL measures, functional capacity
	SCD-101	Invenux	NCT02380079	Phase Ib Recruiting	Oral, once daily for 1 year	40/≥10 years	Safety, frequency of hospitalizations, duration of hospitalizations, analgesic use, transfusion requirement, QOL measures
	Memantine (NMDAR antagonist)	HaEmek Medical Center, Israel	NCT03247218	Phase IIa/b Recruiting			

HbF, fetal hemoglobin; QOL, quality of life; NHLBI, National Heart, Lung, and Blood Institute; NMDAR, N-methyl-D-aspartate receptor; RBC, red blood cell; SCD, sickle cell disease; TCD, transcranial doppler; THU, tetrahydrodione; VOC, vaso-occlusive crisis.

of patients with frequent pain episodes are needed to determine the effect of voxelotor on reducing such crises. Open-label extension studies to assess the long-term effects of voxelotor are ongoing.

5-hydroxymethyl-2-furfural (5HMF, Aes-103) interacts allosterically with HbS, increasing oxygen affinity and decreasing HbS polymerization and RBC sickling during hypoxia. A phase I trial, with oral doses of up to 4,000 mg, showed no significant side effects⁷⁷, but a phase II trial was terminated early because of unblinding of drug groups (NCT01987908). FT-4202, a novel selective activator of RBC pyruvate kinase, decreases intracellular 2,3-diphosphoglycerate levels, with a resulting increase in hemoglobin–oxygen affinity. It demonstrated a favorable safety profile in healthy individuals⁷⁸ and is being investigated in a phase I study in SCD (NCT03815695).

The oxygen delivery therapeutic agents sanguinate (pegylated bovine carboxyhemoglobin) and MP4CO (pegylated human carboxyhemoglobin) are dual transfer agents which release carbon monoxide when delivering oxygen to hypoxic tissues. They stabilize HbS in its R-state and appear to exhibit anti-inflammatory and anti-apoptotic activity by induction of Nrf2 and heme oxygenase-1^{79,80}. Sanguinate had an acceptable safety profile^{81,82}. A phase II study evaluating its efficacy during acute pain crisis was recently completed (NCT02411708).

Increase red blood cell hydration. The inverse relationship between HbS concentration and delay time suggests that even small decreases in intracellular HbS might be beneficial⁸³. Senicapoc selectively blocks the calcium-activated, potassium efflux (Gardos) channel and improves anemia and hemolysis in SCD^{84,85}. Despite improvements in anemia and hemolysis, a phase III trial showed no significant decrease in the rate of pain crises compared to placebo⁸⁴. Similarly, studies of agents that block the potassium-chloride co-transport channel showed no clinical benefits^{86–88}. SCD101, a botanical drug with an unclear anti-sickling mechanism which might involve dilution of HbS by affecting the RBC membrane, was well tolerated in a phase Ib study, with a decrease in chronic pain and fatigue and improvement in leg ulcers⁸⁹.

Targeting downstream sequelae of sickle hemoglobin polymerization

Antioxidant therapy. Agents that upregulate antioxidant and/or reactive oxygen species scavenging processes have been evaluated in SCD. Glutamine, a conditionally essential amino acid, is a precursor for nicotinamide adenine dinucleotide (NAD) and improves NAD redox potential. In a randomized, multicenter trial of 230 patients with HbSS or HbS β^0 -thalassemia, L-glutamine significantly reduced the number of pain crises, hospitalizations, cumulative hospital days, and frequency of acute chest syndrome compared with placebo⁹⁰. However, the effect size of L-glutamine in reducing pain crises was relatively small, with only a 25% reduction in the median number of pain crises. L-glutamine was well tolerated, although low-grade nausea, non-cardiac chest pain, fatigue, and musculoskeletal pain occurred more frequently than with placebo. Despite some concerns related to the high dropout rate in the trial (36%

L-glutamine arm; 24% placebo arm), L-glutamine was approved by the FDA to reduce the acute complications of SCD in patients 5 years and older²⁴.

In an open-label pilot trial of oral N-acetylcysteine (NAC), treatment with either 1,200 mg or 2,400 mg daily for 6 weeks increased whole blood glutathione levels and decreased erythrocyte outer membrane phosphatidylserine exposure, plasma levels of advanced glycation products, and cell-free hemoglobin in both groups⁹¹. A randomized, placebo-controlled, double-blind trial of NAC at 600 mg twice daily for 6 months did not decrease the rate of SCD-related pain days per patient year, vaso-occlusive crises, hospital admission days, number of admissions, or days with home analgesic use compared with placebo⁹². The safety and efficacy of NAC, administered at a higher dose during pain crisis, is being explored in a phase I/II study (NCT01800526) (Table 2).

Anti-adhesive therapy. Agents targeting adhesion of blood cells to the endothelium have been investigated in SCD. Crizanlizumab, a humanized monoclonal anti-P-selectin antibody, was recently approved for use in patients 16 years and older for the prevention of vaso-occlusive crises²⁵ based on results of a randomized, double-blind, phase II study which evaluated the benefit of 2.5 mg/kg or 5 mg/kg doses versus placebo. Significantly lower median crisis rate, longer median times to first and second crises, and lower median rate of uncomplicated crises per year were observed with high-dose crizanlizumab compared with placebo following a 52-week treatment period⁹³. Multiple other studies of crizanlizumab are ongoing (Table 2).

Purified poloxamer 188, a nonionic block copolymer surfactant with hemorheologic and antithrombotic properties, was previously shown to significantly decrease the duration of pain episodes, especially in children and patients on hydroxyurea⁹⁴. However, a more recent phase III study in children and adults reported no significant effect of purified poloxamer 188 (vepoloxamer or MST-188) on duration of vaso-occlusive crises compared with placebo⁹⁵. Rivipansel sodium (formerly GMI-1070) is a small-molecule pan-selectin inhibitor that binds to E-, P-, and L-selectin⁹⁶. In a randomized, double-blind, adaptive, phase II trial, treatment with rivipansel during pain crisis produced a significant reduction in the mean cumulative intravenous opioid analgesic use compared to placebo⁹⁷. However, despite the promising phase II trial results, the recently completed multicenter phase III RESET trial failed to meet its primary (time to readiness for discharge) and key secondary (time to discharge, cumulative intravenous opioid utilization, and time to discontinuation of intravenous opioids) efficacy endpoints⁹⁸.

Heparins inhibit adhesive interactions via P-selectin^{99,100}. Tinzaparin, a low-molecular-weight (LMW) heparin, at therapeutic dose reportedly decreased the number of hospital days, the number of days with pain crisis, and the number of days with the most severe pain scores compared with placebo¹⁰¹. Sevuparin, a derivative of LMW heparin, which retains the P-selectin-binding domain of heparin but largely lacks anticoagulant properties, binds to P- and L-selectins,

Table 2. Ongoing clinical trials of anti-adhesion agents and antioxidants in sickle cell disease.

Mechanism	Drug	Sponsor	NCT number (study acronym)	Clinical phase/status	Intervention	Number/age	Objective
P-selectin antagonist	Crizanlizumab	Novartis Pharmaceuticals	NCT03264989 (SOLACE-adults)	Phase II Active, not recruiting	IV infusion, open-label	57/16–70 years	Pharmacokinetics, pharmacodynamics, safety, and efficacy
			NCT03814746 (STAND)	Phase III Recruiting	IV infusion every 2 weeks for 1 st month and then monthly for 1 year	240/≥12 years	Compare efficacy and safety of 5 mg/kg and 7.5 mg/kg doses with placebo
			NCT04053764 (STEADFAST)	Phase II Recruiting	IV infusion every 2 weeks for 1 st month and then every 4 weeks for 51 weeks + SoC vs. SoC alone	170/≥16 years	Evaluating effect on kidney function (albumin-creatinine ratio, protein-creatinine ratio, estimated glomerular filtration rate)
			NCT03938454 (SPARTAN)	Phase II Recruiting	IV infusion every 2 weeks for 1 st month and then every 4 weeks x 51 weeks	56/≥16 years	Evaluate efficacy in priapism, uncomplicated VOC events
			NCT03474965	Phase II Recruiting	IV infusion every 2 weeks for 1 st month and then every 4 weeks	100/6 months–<18 years	Evaluate pharmacokinetics, pharmacodynamics, safety, and effect on VOC events
Blockade of fcyrlII receptors	IVIg	Albert Einstein College of Medicine	NCT01757418	Phase I–II Recruiting	Single dose of IVIg vs. placebo given within 24 hours of hospitalization	94/12–65 years	Length of VOC, total opioid use, time to end of VOC, <i>in vitro</i> adhesion studies
Antioxidant (increased glutathione)	NAC	Bloodworks	NCT01800526	Phase I/II Enrolling by invitation	IV or oral, NAC Part 1: two doses of IV infusion over 8 hours 4 weeks apart or oral NAC for 4 weeks Part 2 (during VOC): IV infusion every 6 hours for 5 days	20/≥18 years	Evaluate effect on vWF activity, measures of hemolysis and oxidation In Part 2, evaluate efficacy during VOC

IV, intravenous; IVIg, intravenous gammaglobulin; NAC, N-acetylcysteine; SoC, standard of care; VOC, vaso-occlusive crisis; vWF activity, von Willebrand factor activity.

thrombospondin, fibronectin, and von Willebrand factor, inhibits the adhesion of sickle RBCs to stimulated cultured endothelial cells *in vitro*, and prevents vaso-occlusion with normalization of blood flow in a mouse model of vaso-occlusion¹⁰². A recently completed randomized, double-blind, placebo-controlled trial found that, although safe, sevuparin did not significantly reduce the time to resolution of vaso-occlusive crisis¹⁰³. A phase II feasibility study of therapeutic-dose unfractionated heparin (NCT02098993) was recently terminated for poor enrollment, but a phase III, randomized, placebo-controlled study of tinzaparin in acute chest syndrome is ongoing (NCT02580773).

Intravenous immunoglobulin (IVIG) binds to FcγRIII receptors, inhibits neutrophil adhesion to the endothelium, reduces RBC capture by leukocytes, and reduces Mac-1 activity due to recruitment of Src homology 2-containing tyrosine phosphatase-1¹⁰⁴. IVIG was tolerated at doses of up to 800 mg/kg during acute vaso-occlusive crises and decreased Mac-1 function from baseline¹⁰⁵. A phase II clinical trial of IVIG during acute pain episodes is ongoing (NCT01757418).

Anti-inflammatory agents. Multiple approaches to down-regulate inflammatory pathways have been evaluated. Initial clinical responses with the use of steroids for pain crisis and acute chest syndrome were followed by “rebound” pain episodes and re-admissions after treatment discontinuation^{106,107}. Inhaled corticosteroids did not significantly reduce the morbidity of acute chest syndrome in a retrospective cohort study¹⁰⁸, but inhaled mometasone significantly reduced daily pain diary scores and levels of soluble vascular adhesion molecule-1 over the 16-week treatment period in another single-center, placebo-controlled trial¹⁰⁹.

The activation of iNKT cells is downregulated by the activation of adenosine A_{2A} receptors (A_{2A}R)¹¹⁰. Regadenoson, a partially selective A_{2A}R agonist, was reportedly safe but did not decrease length of hospital stay, total opioid use, or pain scores compared to placebo during pain crises^{111,112}. In addition, there were no significant differences between regadenoson and placebo regarding the number of patients exhibiting a greater than 30% reduction in activated iNKT cells. The anti-iNKT cell monoclonal antibody NKTT120 resulted in rapid, specific, and continuous iNKT cell depletion in a single-ascending-dose study¹¹³, but further studies are required to determine its long-term safety and efficacy.

Statins have benefits independent of their cholesterol-lowering effect. In a pilot study, simvastatin was well tolerated, increased levels of NO metabolites, and decreased C-reactive protein (CRP) and interleukin-6 levels¹¹⁴. Daily treatment with simvastatin for 3 months significantly reduced pain crises, oral analgesic use, and levels of high-sensitivity CRP, soluble E-selectin, soluble ICAM-1, soluble ICAM-3, and VEGF¹¹⁵, providing supporting data for conduct of a placebo-controlled trial. Atorvastatin was well tolerated in a pilot study but did not improve endothelial function or decrease albuminuria¹¹⁶.

Long-chain omega-3 polyunsaturated fatty acids such as eicosapentaenoic acid (EPA [n-3]) and docosahexaenoic acid (DHA [n-6]) provide benefit to individuals with chronic inflammatory disorders. In a randomized, single-center study, daily omega-3 administered in capsule form significantly lowered the occurrence of vaso-occlusive events, severe anemia, blood transfusion requirements, and school absence compared with placebo¹¹⁷. More recently, a randomized, placebo-controlled study of different doses of SC411, a novel DHA ethyl ester formulation with high DHA bioavailability, found it was well tolerated, with significant reductions in d-dimer and E-selectin and an increase in hemoglobin¹¹⁸. SC411 also significantly reduced electronic diary-recorded pain episodes, analgesic use at home, and days absent from school but did not significantly lower pain crises in the pooled active groups compared to placebo.

Preliminary results suggest that the monoclonal anti-IL1β antibody canakinumab (ACZ885) is well tolerated and not associated with major side effects in SCD¹¹⁹.

Anticoagulant and antiplatelet therapies. With abundant evidence that SCD is a hypercoagulable state²¹, combined with data that coagulation and platelet activation may play roles in disease pathophysiology^{120–124}, multiple studies have evaluated the effects of anticoagulants and antiplatelet agents in SCD. Although treatment with the LMW heparin tinzaparin produced more rapid resolution of pain crisis and shorter duration of hospitalization than placebo¹⁰¹, it is uncertain whether the reported beneficial effects were a result of its anticoagulant or anti-adhesive effects. The results of a pilot study of the direct oral anticoagulant rivaroxaban in SCD are awaited (NCT02072668).

Multiple relatively small studies of aspirin have reported modest benefit at best^{125–127}, although one study reported an increase in oxygen affinity, hemoglobin level, and RBC lifespan¹²⁶. A pilot study of the glycoprotein IIb/IIIa inhibitor eptifibatide showed that it was safe during acute pain episodes but did not reduce the time to resolution of such episodes¹²⁸. Ticlopidine, a P_{2Y}₁₂ ADP-receptor antagonist, reportedly decreased the number of pain episodes, mean duration of pain episodes, and severity of such episodes¹²⁹. However, a more recent phase III trial of prasugrel, a newer generation P_{2Y}₁₂ receptor blocker, showed no significant reduction of the frequency of pain episodes in children with SCD¹³⁰. Ticagrelor, a reversible P_{2Y}₁₂ receptor blocker, was well tolerated in children¹³¹ and young adults¹³² but had no beneficial effects on pain. A phase III study to determine the efficacy of ticagrelor in reducing vaso-occlusive crises in children is ongoing¹³³ (Table 3).

Nitric oxide and related agents. With the role of hemolysis in NO scavenging and subsequent endothelial dysfunction, NO and related agents may be beneficial in SCD¹³⁴. Inhaled NO did not improve time to resolution of pain crisis, length of hospitalization, opioid usage, or rate of acute chest syndrome compared with placebo¹³⁵. In another study, inhaled NO did not reduce the rate of treatment failure in adult patients with mild

Table 3. Ongoing clinical trials of anticoagulants, antiplatelet agents, anti-inflammatory agents, and nitric oxide-related drugs in sickle cell disease.

Mechanism	Drug	Sponsor	NCT number (study acronym)	Clinical phase/status	Intervention	Number/age	Objective
Anticoagulant, anti-adhesive (P-selectin blocker)	Tinzaparin (low-molecular-weight heparin)	Assistance Publique –Hôpitaux de Paris, LEO pharma	NCT02580773 (TASC)	Phase III Unknown status	Subcutaneous therapeutic vs. prophylactic dose for 7 days	200/≥18 years	Time to ACS resolution, bleeding events, hospital mortality
	Ticagrelor	AstraZeneca	NCT03615924 (HESTIA3)	Phase III Active, not recruiting	Oral ticagrelor vs. placebo for at least 12 months	193/2–17 years	Evaluate efficacy in decreasing VOC, acute chest syndrome, duration of VOC, QOL measures
Antithrombotic, anti-inflammatory	Defibrotide	New York Medical College	NCT03805581	Phase II Recruiting	Intravenous, every 6 hours during ACS event for maximum of 7 days	20/2–40 years	Safety (allergic reaction, bleeding), improvement in clinical signs of ACS
Anti-inflammatory	Mometasone	Jeffery Glassberg	NCT03758950 (IMPROVE2)	Phase II Recruiting	Inhaled, one puff daily vs. placebo for 48 weeks	80/≥18 years	Change in soluble VCAM-1, hemoglobin, markers of hemolysis, leukocyte count, QOL measures
	Canakinumab (anti-IL-1 β antibody)	Novartis	NCT02961218	Phase II Active, not recruiting	Subcutaneous, monthly for 6 months vs. placebo, followed by 6-month open-label extension	49/8–20 years	Reduction of average daily pain, days missed from school, blood counts, pharmacokinetics, hs-CRP, measures of hemolysis
	Docosahexaenoic acid and eicosapentaenoic acid	Thomas Jefferson University	NCT01202812	Phase II Not yet recruiting	Oral, daily for 6 months vs. placebo	48/10–19 years	Decrease inflammatory biomarkers, QOL
	Docosahexaenoic acid (SC411)	Micelle BioPharma	NCT02973360 (SCOT study)	Phase II Active, not recruiting	Oral for 8 weeks vs. placebo followed by 49-month open-label extension	68/5–17 years	Safety, pharmacokinetics, effect on VOC
	Omega-3-fatty acid	Sultan Qaboos University	NCT02525107	Phase III Recruitment status unknown	Oral, daily for 52 weeks vs. placebo	280/13–70 years	Effect on VOC, duration of hospitalization, RBC membrane fatty acid profile
	Rifaximin (antibiotic, decrease aged neutrophils)	New York Medical College	NCT03719729	Phase II Recruiting	Oral, twice daily for 1 year	20/18–70 years	Safety, frequency of hospitalizations for VOCs, duration of hospitalizations, RBC transfusions, QOL measures
Increased NO production	Arginine	Emory University	NCT02447874	Phase I/II Enrolling by invitation	Intravenous, three times a day for maximum of 7 days	21/7–21 years	Pharmacokinetics, NO metabolites
			NCT02536170	Phase II Recruiting	Intravenous, three times a day through duration of hospitalization	114/3–21 years	Effect on VOC, including total parenteral opioid use, length of hospital stay, resolution of VOC
Vasodilator	Olinicuat (IW-1701) (soluble guanylate cyclase stimulator)	Cyclerion therapeutics	NCT03285178	Phase II Recruiting	Oral, daily 12 weeks vs. placebo	88/16–70 years	Safety and tolerability, pharmacokinetics, pharmacodynamics
	Riociguat (soluble guanylate cyclase stimulator)	Mark Gladwin	NCT02633397	Phase II Recruiting	Oral, three times daily for 12 weeks	100/≥18 years	Safety, changes in pain intensity, SCD-related complications, functional capacity, blood pressure, plasma NT-proBNP, changes in laboratory measures, TRV
	IMR-687 (selective phosphodiesterase-9 inhibitor)	Imara	NCT03401112	Phase IIa Recruiting	Oral, daily for 24 weeks vs. placebo	70/18–55 years	Safety, pharmacokinetics
			NCT04053803	Phase II Enrolling by invitation	Oral, daily for 49 months, open-label extension	70/≥18 years	Safety, change in hemoglobin, HbF, soluble adhesion markers, renal and cardiac function, QOL
	Ambrisentan (endothelin receptor A antagonist)	Augusta University	NCT02712346	Phase I Active, not recruiting	Oral, once daily for 12 weeks	26/18–65 years	Safety and tolerability, microalbuminuria/proteinuria, TRV, inflammatory markers, pain questionnaire

ACS, acute chest syndrome; HbF, fetal hemoglobin; hs-CRP, high-sensitivity C-reactive protein; NO, nitric oxide; NT-proBNP, N-terminal probrain natriuretic peptide; QOL, quality of life; TRV, echocardiography-derived tricuspid regurgitant jet velocity; VCAM-1, vascular cell adhesion molecule-1; VOC, vaso-occlusive crisis; vWF, von Willebrand factor.

to moderate acute chest syndrome¹³⁶. However, L-arginine, an NO precursor, significantly decreased total parenteral opioid use and pain scores at discharge compared to placebo in children, although there was no difference in length of hospital stay¹³⁷. Furthermore, supplementation of adult patients on hydroxyurea with oral arginine significantly increased the level of NO metabolites and reduced the frequency of pain crises versus placebo following 4 months of treatment¹³⁸. Sildenafil, a phosphodiesterase-5 inhibitor which increases NO-mediated effects by inhibiting the degradation of cyclic guanosine monophosphate (cGMP), was associated with more frequent serious adverse events, predominantly hospitalization for pain, compared with placebo¹³⁹. No evidence of a treatment effect of sildenafil was seen on evaluated study outcomes. Studies of soluble guanylate cycle stimulators, riociguat (NCT02633397) and olinciguat/IW-1701 (NCT03285178), and the phosphodiesterase-9

inhibitor IMR-687 (NCT03401112; NCT04053803) are ongoing (Table 3).

Summary

Recent advances in drug development for SCD raise important issues related to the use and availability of these agents. With limited availability of allogeneic bone marrow transplantation and gene therapy, especially in low- and middle-income countries, the number of recently approved drugs offers hope for further improved outcomes in SCD. Choice of initial drug therapy may be guided by a patient's clinical sub-phenotype as well as the cost of the drug (Table 4). Patients with frequent vaso-occlusive complications (acute pain episodes, acute chest syndrome) may benefit from hydroxyurea, L-glutamine, and crizanlizumab, while those with increased hemolytic anemia (or quite possibly complications related to hemolytic anemia) may benefit from

Table 4. Summary characteristics of FDA-approved drugs for sickle cell disease.

	Hydroxyurea	L-Glutamine	Crizanlizumab	Voxelotor
Age (years)	≥2	≥5	≥16	≥12
Genotypes	HbSS, HbSβ ⁰ thalassemia	All genotypes (only studied in HbSS, HbSβ ⁰ thalassemia)	All genotypes	All genotypes
Mechanism of action	Multiple, but primarily by increasing HbF production	Uncertain, but thought to reduce NAD redox potential, possible decrease in cell adhesion	Anti P-selectin inhibitor (decreases adhesion of WBCs and RBCs to endothelium)	Decreases HbS polymerization by increasing Hb-oxygen affinity
Route of administration	Oral (capsules/tablets)	Oral (powder)	Intravenous	Oral (tablets)
Clinical effects of therapy	Decreased frequency of VOC, decreased frequency of ACS, decreased hospitalization, decreased RBC transfusion requirement, decreased stroke risk	Decreased frequency of VOC, decreased frequency of ACS, decreased hospitalization	Decreased frequency of VOC	Increased hemoglobin
Effect size for primary endpoint (% change and IRR)	44% decrease in VOC per year (median from 4.5 to 2.5), IRR = 0.56	25% decrease in VOC in 48 weeks (median from 4 to 3), IRR = 0.75	45% decrease in crisis rate per year (median from 3 to 1.6), IRR = 0.55	5.5-fold increase in the hemoglobin responders (9% to 59%) at 24 weeks, incidence proportion ratio = 6.6*
Common toxicities	Myelosuppression, skin hyperpigmentation, nail discoloration, teratogenicity, decreased sperm counts, nausea and vomiting	Constipation, nausea, headaches, abdominal pain	Nausea, arthralgia	Headache, diarrhea, nausea
Pharmacokinetics	Excreted via kidneys Adjust dose for eGFR <60 mL/min/1.73 m ²	Use with caution with hepatic and renal impairment, but no recommended dose adjustment	No dosage adjustments in manufacturer labeling for renal and hepatic impairment (not tested in ESRD)	No dosage adjustment for renal impairment, but not yet studied in ESRD requiring dialysis Dose reduction for severe liver disease (Child Pugh class C)
Cost	\$	\$\$\$	\$\$\$\$\$	\$\$\$\$\$

ACS, acute chest syndrome; eGFR, estimated glomerular filtration rate; ESRD, end-stage renal disease; FDA, US Food and Drug Administration; Hb, hemoglobin; HbF, fetal hemoglobin; HbS, sickle hemoglobin; IRR, incidence rate ratio; NAD, nicotinamide adenine dinucleotide; RBC, red blood cell; VOC, vaso-occlusive crisis; WBC, white blood cell.

* Patients treated with 1,500 mg of voxelotor had 6.6 times increased proportion of hemoglobin responders (>1 g/dL increase from baseline at 24 weeks)

hydroxyurea and voxelotor. However, more studies are required before definite recommendations can be made regarding the effect of voxelotor on SCD-related complications. As with other chronic disease conditions, it is important to consider the cost of the drug when deciding on initial therapy. Hydroxyurea is much cheaper than L-glutamine, crizanlizumab, and voxelotor and is likely more cost effective as initial therapy for vaso-occlusive complications and hemolytic anemia. However, with the complex pathophysiology of SCD and the limited clinical efficacy of available pharmacological therapies, it is unlikely that a single drug will ameliorate all SCD-related complications. As such, the availability of multiple drugs offers an opportunity for combination therapy based on different mechanisms of action and non-overlapping side effect profiles. Although more studies are required to evaluate the effect of combinations of these drugs, all of the recently approved agents were tested in combination with hydroxyurea. These studies showed benefit whether or not participants received concomitant hydroxyurea, with no increased toxicity. Despite potential benefits, combination therapy involving the use of multiple drugs increases the likelihood of decreased medication adherence owing to polypharmacy. Like in other chronic diseases, however, this problem may be somewhat ameliorated by the development of pills that contain more than one active drug.

It is particularly important that ongoing trials of novel drugs for SCD involve individuals in resource-limited countries where

the burden of disease is high. This offers patients access to potentially disease-modifying therapies but, perhaps equally as important, contributes substantially to capacity building by improving local infrastructure, strengthening health systems, involving local leaders as collaborators, and training local providers. Furthermore, trials of novel agents in sub-Saharan Africa provide data on the effects of these therapies in a setting with fewer resources and unique challenges, including a high prevalence of malnutrition and infections such as malaria. With the high cost of recently approved agents, mechanisms to ensure affordable access to these drugs should be established. A situation where novel agents are tested in resource-limited countries but patients in these countries are unable to afford such drugs following approval is unacceptable. Finally, studies evaluating the effects of recently approved drugs on organ damage are needed. Demonstration that these agents prevent or decrease end-organ damage would provide additional evidence for the role of drug therapies in improving outcomes in SCD.

Acknowledgements

The authors thank Dr. Joacy Gerard Matthias and Dr. Matthew P. Smeltzer (Both from the Division of Epidemiology, Biostatistics, and Environmental Health, School of Public Health, University of Memphis, Memphis, Tennessee, USA) for helping to ascertain the drug effect sizes.

References

- Piel FB, Hay SI, Gupta S, *et al.*: **Global burden of sickle cell anaemia in children under five, 2010-2050: Modelling based on demographics, excess mortality, and interventions.** *PLoS Med.* 2013; 10(7): e1001484.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Grosse SD, Odam L, Atrash HK, *et al.*: **Sickle cell disease in Africa: A neglected cause of early childhood mortality.** *Am J Prev Med.* 2011; 41(6 Suppl 4): S398–405.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Telfer P, Coen P, Chakravorty S, *et al.*: **Clinical outcomes in children with sickle cell disease living in England: A neonatal cohort in East London.** *Haematologica.* 2007; 92(7): 905–12.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Quinn CT, Rogers ZR, McCavit TL, *et al.*: **Improved survival of children and adolescents with sickle cell disease.** *Blood.* 2010; 115(17): 3447–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Couque N, Girard D, Ducrocq R, *et al.*: **Improvement of medical care in a cohort of newborns with sickle-cell disease in North Paris: Impact of national guidelines.** *Br J Haematol.* 2016; 173(6): 927–37.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Elmariyah H, Garrett ME, de Castro LM, *et al.*: **Factors associated with survival in a contemporary adult sickle cell disease cohort.** *Am J Hematol.* 2014; 89(5): 530–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gardner K, Douiri A, Drasar E, *et al.*: **Survival in adults with sickle cell disease in a high-income setting.** *Blood.* 2016; 128(10): 1436–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Maitra P, Caughey M, Robinson L, *et al.*: **Risk factors for mortality in adult patients with sickle cell disease: A meta-analysis of studies in North America and Europe.** *Haematologica.* 2017; 102(4): 626–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- DeBaun MR, Ghafari DL, Rodeghier M, *et al.*: **Decreased median survival of adults with sickle cell disease after adjusting for left truncation bias: A pooled analysis.** *Blood.* 2019; 133(6): 615–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fitzhugh CD, Abraham AA, Tisdale JF, *et al.*: **Hematopoietic stem cell transplantation for patients with sickle cell disease: Progress and future directions.** *Hematol Oncol Clin North Am.* 2014; 28(6): 1171–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ribeil JA, Hachein-Bey-Abina S, Payen E, *et al.*: **Gene Therapy in a Patient with Sickle Cell Disease.** *N Engl J Med.* 2017; 376(9): 848–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Epstein FH, Bunn HF: **Pathogenesis and Treatment of Sickle Cell Disease.** *N Engl J Med.* 1997; 337(11): 762–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rees DC, Williams TN, Gladwin MT: **Sickle-cell disease.** *Lancet.* 2010; 376(9757): 2018–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
- Lawson V: **Turned on by danger: Activation of CD1d-restricted invariant natural killer T cells.** *Immunology.* 2012; 137(1): 20–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Reiter CD, Wang X, Tanus-Santos JE, *et al.*: **Cell-free hemoglobin limits nitric oxide bioavailability in sickle-cell disease.** *Nat Med.* 2002; 8(12): 1383–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Repka T, Heibel RP: **Hydroxyl radical formation by sickle erythrocyte membranes: Role of pathologic iron deposits and cytoplasmic reducing agents.** *Blood.* 1991; 78(10): 2753–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- de Caterina R, Libby P, Peng HB, *et al.*: **Nitric oxide decreases cytokine-induced endothelial activation. Nitric oxide selectively reduces endothelial expression of adhesion molecules and proinflammatory cytokines.** *J Clin Invest.* 1995; 96(1): 60–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Palmer RMJ, Ashton DS, Moncada S: **Vascular endothelial cells synthesize nitric oxide from L-arginine.** *Nature.* 1988; 333(6174): 664–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kato GJ, Gladwin MT, Steinberg MH: **Deconstructing sickle cell disease: Reappraisal of the role of hemolysis in the development of clinical subphenotypes.** *Blood Rev.* 2007; 21(1): 37–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Manwani D, Frenette PS: **Vaso-occlusion in sickle cell disease: Pathophysiology and novel targeted therapies.** *Hematology Am Soc Hematol*

- Educ Program. 2013; **2013**: 362–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Noubouossie D, Key NS, Ataga KI: **Coagulation Abnormalities of Sickle Cell Disease: Relationship With Clinical Outcomes and the Effect of Disease Modifying Therapies.** *Blood Rev.* 2016; **30**(4): 245–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 22. **F** Kato GJ, Piel FB, Reid CD, *et al.*: **Sickle cell disease.** *Nat Rev Dis Primers.* 2018; **4**: 18010.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 23. Charache S, Terrin ML, Moore RD, *et al.*: **Effect of hydroxyurea on the frequency of painful crises in sickle cell anemia. Investigators of the Multicenter Study of Hydroxyurea in Sickle Cell Anemia.** *N Engl J Med.* 1995; **332**(20): 1317–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
 24. <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-disco-l-glutamine-sickle-cell-disease>
 25. <https://www.novartis.com/news/media-releases/new-novartis-medicine-adakveo-crizanlizumab-approved-fda-reduce-frequency-pain-crisis-individuals-living-sickle-cell-disease>
 26. <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-approves-voxelotor-sickle-cell-disease>
 27. Eaton WA, Bunn HF: **Treating sickle cell disease by targeting HbS polymerization.** *Blood.* 2017; **129**(20): 2719–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 28. Platt OS, Brambilla DJ, Rosse WF, *et al.*: **Mortality in Sickle Cell Disease. Life Expectancy and Risk Factors for Early Death.** *N Engl J Med.* 1994; **330**(23): 1639–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
 29. Castro O, Brambilla DJ, Thorington B, *et al.*: **The acute chest syndrome in sickle cell disease: Incidence and risk factors. The Cooperative Study of Sickle Cell Disease.** *Blood.* 1994; **84**(2): 643–9.
[PubMed Abstract](#)
 30. Mabaera R, West RJ, Conine SJ, *et al.*: **A Cell Stress Signaling Model of Fetal Hemoglobin Induction: What Doesn't Kill Red Blood Cells May Make Them Stronger.** *Exp Hematol.* 2008; **36**(9): 1057–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
 31. Alcaín FJ, Löw H, Crane FL, *et al.*: **Iron chelators hydroxyurea and bathophenanthroline disulfonate inhibit DNA synthesis by different pathways.** *Biochem Mol Biol Int.* 1994; **34**(2): 273–9.
[PubMed Abstract](#)
 32. Cokic VP, Smith RD, Beleslin-Cokic BB, *et al.*: **Hydroxyurea induces fetal hemoglobin by the nitric oxide-dependent activation of soluble guanylyl cyclase.** *J Clin Invest.* 2003; **111**(2): 231–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 33. Huang J, Yakubu M, Kim-Shapiro DB, *et al.*: **Rat liver-mediated metabolism of hydroxyurea to nitric oxide.** *Free Radic Biol Med.* 2006; **40**(9): 1675–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
 34. McGann PT, Ware RE: **Hydroxyurea for sickle cell anemia: What have we learned and what questions still remain?** *Curr Opin Hematol.* 2011; **18**(3): 158–65.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 35. Bridges KR, Barabino GD, Brugnara C, *et al.*: **A multiparameter analysis of sickle erythrocytes in patients undergoing hydroxyurea therapy.** *Blood.* 1996; **88**(12): 4701–10.
[PubMed Abstract](#)
 36. Hillery CA, Du MC, Wang WC, *et al.*: **Hydroxyurea therapy decreases the *in vitro* adhesion of sickle erythrocytes to thrombospondin and laminin.** *Br J Haematol.* 2000; **109**(2): 322–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 37. Odievre MH, Bony V, Benkerrou M, *et al.*: **Modulation of erythroid adhesion receptor expression by hydroxyurea in children with sickle cell disease.** *Haematologica.* 2008; **93**(4): 502–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
 38. Laurance S, Lansiaux P, Pellay FX, *et al.*: **Differential modulation of adhesion molecule expression by hydroxycarbamide in human endothelial cells from the micro- and macrocirculation: Potential implications in sickle cell disease vasoocclusive events.** *Haematologica.* 2011; **96**(4): 534–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 39. Gladwin MT, Shelhamer JH, Ognibene FP, *et al.*: **Nitric oxide donor properties of hydroxyurea in patients with sickle cell disease.** *Br J Haematol.* 2002; **116**(2): 436–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
 40. Glover RE, Ivy ED, Orringer EP, *et al.*: **Detection of nitrosyl hemoglobin in venous blood in the treatment of sickle cell anemia with hydroxyurea.** *Mol Pharmacol.* 1999; **55**(6): 1006–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. **F** Wang WC, Ware RE, Miller ST, *et al.*: **Hydroxycarbamide in Very Young Children With Sickle-Cell Anaemia: A Multicentre, Randomised, Controlled Trial (BABY HUG).** *Lancet.* 2011; **377**(9778): 1663–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 42. Dehury S, Purohit P, Patel S, *et al.*: **Low and Fixed Dose of Hydroxyurea Is Effective and Safe in Patients With HbS(+) Thalassemia With IVS1-5(G→C) Mutation.** *Pediatr Blood Cancer.* 2015; **62**(6): 1017–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
 43. Jain DL, Apte M, Colah R, *et al.*: **Efficacy of Fixed Low Dose Hydroxyurea in Indian Children With Sickle Cell Anemia: A Single Centre Experience.** *Indian Pediatr.* 2013; **50**(10): 929–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
 44. **F** Ware RE, Davis BR, Schultz WH, *et al.*: **Hydroxycarbamide Versus Chronic Transfusion for Maintenance of Transcranial Doppler Flow Velocities in Children With Sickle Cell anaemia-TCD With Transfusions Changing to Hydroxyurea (TWITCH): A Multicentre, Open-Label, Phase 3, Non-Inferiority Trial.** *Lancet.* 2016; **387**(10019): 661–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 45. Bartolucci P, Habibi A, Stehlé T, *et al.*: **Six Months of Hydroxyurea Reduces Albuminuria in Patients With Sickle Cell Disease.** *J Am Soc Nephrol.* 2016; **27**(6): 1847–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 46. Lobo CLdC, Pinto JFC, Nascimento EM, *et al.*: **The Effect of Hydroxycarbamide Therapy on Survival of Children With Sickle Cell Disease.** *Br J Haematol.* 2013; **161**(6): 852–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
 47. Steinberg MH, Barton F, Castro O, *et al.*: **Effect of Hydroxyurea on Mortality and Morbidity in Adult Sickle Cell Anemia: Risks and Benefits Up to 9 Years of Treatment.** *JAMA.* 2003; **289**(13): 1645–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
 48. **F** Steinberg MH, McCarthy WF, Castro O, *et al.*: **The Risks and Benefits of Long-Term Use of Hydroxyurea in Sickle Cell Anemia: A 17.5 Year Follow-Up.** *Am J Hematol.* 2010; **85**(6): 403–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 49. **F** Voskaridou E, Christoulas D, Bilalis A, *et al.*: **The Effect of Prolonged Administration of Hydroxyurea on Morbidity and Mortality in Adult Patients With Sickle Cell Syndromes: Results of a 17-year, Single-Center Trial (LaSHS).** *Blood.* 2010; **115**(12): 2354–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 50. **F** Ware RE: **Optimizing Hydroxyurea Therapy for Sickle Cell Anemia.** *Hematology Am Soc Hematol Educ Program.* 2015; **2015**: 436–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 51. **F** McGann PT, Niss O, Dong M, *et al.*: **Robust clinical and laboratory response to hydroxyurea using pharmacokinetically guided dosing for young children with sickle cell anemia.** *Am J Hematol.* 2019; **94**(8): 871–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 52. Estep JH, Smeltzer MP, Kang G, *et al.*: **A clinically meaningful fetal hemoglobin threshold for children with sickle cell anemia during hydroxyurea therapy.** *Am J Hematol.* 2017; **92**(12): 1333–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 53. **Hydroxyurea Optimization Through Precision Study (HOPS).** *Clinical Trials.* 2020.
[Reference Source](#)
 54. **F** Nazon C, Sabo AN, Becker G, *et al.*: **Optimizing Hydroxyurea Treatment for Sickle Cell Disease Patients: The Pharmacokinetic Approach.** *J Clin Med.* 2019; **8**(10): 1701.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 55. Sauntharajah Y, Molokie R, Saraf S, *et al.*: **Clinical effectiveness of decitabine in severe sickle cell disease.** *Br J Haematol.* 2008; **141**(1): 126–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 56. Molokie R, Lavelle D, Gowhari M, *et al.*: **Oral tetrahydrouridine and decitabine for non-cytotoxic epigenetic gene regulation in sickle cell disease: A randomized phase 1 study.** *PLoS Med.* 2017; **14**(9): e1002382.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 57. Atweh GF, Sutton M, Nassif I, *et al.*: **Sustained induction of fetal hemoglobin by pulse butyrate therapy in sickle cell disease.** *Blood.* 1999; **93**(6): 1790–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 58. Hebbel RP, Vercellotti GM, Pace BS, *et al.*: **The HDAC inhibitors trichostatin A and suberoylanilide hydroxamic acid exhibit multiple modalities of benefit for the vascular pathobiology of sickle transgenic mice.** *Blood.* 2010; **115**(12): 2483–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 59. Ikuta T, Atweh G, Boosalis V, *et al.*: **Cellular and Molecular Effects of a Pulse Butyrate Regimen and New Inducers of Globin Gene Expression and Hematopoiesis.** *Ann N Y Acad Sci.* 1998; **850**: 87–99.
[PubMed Abstract](#) | [Publisher Full Text](#)
 60. Kutlar A, Ataga K, Reid M, *et al.*: **A phase 1/2 trial of HQK-1001, an oral fetal globin inducer, in sickle cell disease.** *Am J Hematol.* 2012; **87**(11): 1017–21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 61. Kutlar A, Reid ME, Inati A, *et al.*: **A dose-escalation phase IIa study of 2,2-dimethylbutyrate (HQK-1001), an oral fetal globin inducer, in sickle cell disease.** *Am J Hematol.* 2013; **88**(11): E255–E260.
[PubMed Abstract](#) | [Publisher Full Text](#)
 62. Okam MM, Esrick EB, Mandell E, *et al.*: **Phase 1/2 trial of vorinostat in patients**

- with sickle cell disease who have not benefitted from hydroxyurea. *Blood*. 2015; **125**(33): 3668–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
63. **F** Reid ME, El Beshlawy A, Inati A, *et al.*: A double-blind, placebo-controlled phase II study of the efficacy and safety of 2,2-dimethylbutyrate (HQK-1001), an oral fetal globin inducer, in sickle cell disease. *Am J Hematol*. 2014; **89**(7): 709–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 64. Shearstone JR, Golonzhka O, Chonkar A, *et al.*: Chemical Inhibition of Histone Deacetylases 1 and 2 Induces Fetal Hemoglobin through Activation of GATA2. *PLoS One*. 2016; **11**: e0153767.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 65. Rivers A, Vaitkus K, Ruiz MA, *et al.*: RN-1, a potent and selective lysine-specific demethylase 1 inhibitor, increases γ -globin expression, F reticulocytes, and F cells in a sickle cell disease mouse model. *Exp Hematol*. 2015; **43**(7): 546–553. e3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 66. Cui S, Lim KC, Shi L, *et al.*: The LSD1 inhibitor RN-1 induces fetal hemoglobin synthesis and reduces disease pathology in sickle cell mice. *Blood*. 2015; **126**(3): 386–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 67. **F** Rivers A, Vaitkus K, Jagadeeswaran R, *et al.*: Oral administration of the LSD1 inhibitor ORY-3001 increases fetal hemoglobin in sickle cell mice and baboons. *Exp Hematol*. 2018; **67**: 60–64.e2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 68. **F** Meiler SE, Wade M, Kutlar F, *et al.*: Pomalidomide augments fetal hemoglobin production without the myelosuppressive effects of hydroxyurea in transgenic sickle cell mice. *Blood*. 2011; **118**(4): 1109–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 69. Kutlar A, Swerdlow PS, Meiler SE, *et al.*: Pomalidomide In Sickle Cell Disease: Phase I Study Of a Novel Anti-Switching Agent. *Blood*. 2013; **122**(21): 777.
[Publisher Full Text](#)
 70. **F** Zhang Y, Paikari A, Sumazin P, *et al.*: Metformin induces FOXO3-dependent fetal hemoglobin production in human primary erythroid cells. *Blood*. 2018; **132**(3): 321–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 71. Krishnamoorthy S, Pace B, Gupta D, *et al.*: Dimethyl fumarate increases fetal hemoglobin, provides heme detoxification, and corrects anemia in sickle cell disease. *JCI Insight*. 2017; **2**(20): e96409.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 72. Dai Y, Chen T, Ijaz H, *et al.*: SIRT1 activates the expression of fetal hemoglobin genes. *Am J Hematol*. 2017; **92**(11): 1177–1186.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 73. Oksenberg D, Dufu K, Patel MP, *et al.*: GBT440 increases haemoglobin oxygen affinity, reduces sickling and prolongs RBC half-life in a murine model of sickle cell disease. *Br J Haematol*. 2016; **175**(1): 141–53.
[PubMed Abstract](#) | [Publisher Full Text](#)
 74. **F** Howard J, Hemmaway CJ, Telfer P, *et al.*: A phase 1/2 ascending dose study and open-label extension study of voxelotor in patients with sickle cell disease. *Blood*. 2019; **133**(17): 1865–1875.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 75. **F** Dufu K, Patel M, Oksenberg D, *et al.*: GBT440 improves red blood cell deformability and reduces viscosity of sickle cell blood under deoxygenated conditions. *Clin Hemorheol Microcirc*. 2018; **70**(1): 95–105.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 76. Vichinsky E, Hoppe CC, Ataga KI, *et al.*: A Phase 3 Randomized Trial of Voxelotor in Sickle Cell Disease. *N Engl J Med*. 2019; **381**(6): 509–519.
[PubMed Abstract](#) | [Publisher Full Text](#)
 77. Kato GJ, Lawrence MP, Mendelsohn LG, *et al.*: Phase 1 Clinical Trial Of The Candidate Anti-Sickling Agent Aes-103 In Adults With Sickle Cell Anemia. *Blood*. 2013; **122**: 1009.
[Publisher Full Text](#)
 78. Kalfa TA, Kuypers FA, Telen MJ, *et al.*: Phase 1 Single (SAD) and Multiple Ascending Dose (MAD) Studies of the Safety, Tolerability, Pharmacokinetics (PK) and Pharmacodynamics (PD) of FT-4202, an Allosteric Activator of Pyruvate Kinase-R, in Healthy and Sickle Cell Disease Subjects. *Blood*. 2019; **134**(Supplement_1): 616.
[Publisher Full Text](#)
 79. Belcher JD, Young M, Chen C, *et al.*: MP4CO, a pegylated hemoglobin saturated with carbon monoxide, is a modulator of HO-1, inflammation, and vaso-occlusion in transgenic sickle mice. *Blood*. 2013; **122**(15): 2757–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 80. Abuchowski A: SANGUINATE (PEGylated Carboxyhemoglobin Bovine): Mechanism of Action and Clinical Update. *Artif Organs*. 2017; **41**(4): 346–350.
[PubMed Abstract](#) | [Publisher Full Text](#)
 81. Keipert PE: Clinical Evaluation of MP4CO: A Phase 1b Escalating-Dose, Safety and Tolerability Study in Stable Adult Patients with Sickle Cell Disease. *Adv Exp Med Biol*. 2016; **923**: 23–29.
[PubMed Abstract](#) | [Publisher Full Text](#)
 82. Misra H, Bainbridge J, Berryman J, *et al.*: A Phase Ib open label, randomized, safety study of SANGUINATE™ in patients with sickle cell anemia. *Rev Bras Hematol Hemoter*. 2017; **39**(1): 20–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 83. Sunshine HR, Hofrichter J, Eaton WA: Requirement for therapeutic inhibition of sickle haemoglobin gelation. *Nature*. 1978; **275**(5677): 238–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
 84. Ataga KI, Reid M, Ballas SK, *et al.*: Improvements in haemolysis and indicators of erythrocyte survival do not correlate with acute vaso-occlusive crises in patients with sickle cell disease: A phase III randomized, placebo-controlled, double-blind study of the Gardos channel blocker senicapoc (ICA-17043). *Br J Haematol*. 2011; **153**(1): 92–104.
[PubMed Abstract](#) | [Publisher Full Text](#)
 85. Ataga KI, Smith WR, de Castro LM, *et al.*: Efficacy and safety of the Gardos channel blocker, senicapoc (ICA-17043), in patients with sickle cell anemia. *Blood*. 2008; **111**(8): 3991–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 86. Brousseau DC, Scott JP, Badaki-Makun O, *et al.*: A multicenter randomized controlled trial of intravenous magnesium for sickle cell pain crisis in children. *Blood*. 2015; **126**(14): 1651–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 87. Goldman RD, Mounstephen W, Kirby-Allen M, *et al.*: Intravenous magnesium sulfate for vaso-occlusive episodes in sickle cell disease. *Pediatrics*. 2013; **132**(6): e1634–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
 88. de Franceschi L, Bachir D, Galacteros F, *et al.*: Oral magnesium pidolate: Effects of long-term administration in patients with sickle cell disease. *Br J Haematol*. 2000; **108**(2): 284–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 89. Swift R, Abdulmalik O, Chen Q, *et al.*: SCD-101: A New Anti-Sickling Drug Reduces Pain and Fatigue and Improves Red Blood Cell Shape in Peripheral Blood of Patients with Sickle Cell Disease. *Blood*. 2016; **128**(22): 121.
[Publisher Full Text](#)
 90. **F** Nihara Y, Smith WR, Stark CW: A Phase 3 Trial of L-Glutamine in Sickle Cell Disease. *N Engl J Med*. 2018; **379**(19): 1880.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 91. Nur E, Brandjes DP, Teerlink T, *et al.*: N-acetylcysteine reduces oxidative stress in sickle cell patients. *Ann Hematol*. 2012; **91**(7): 1097–105.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 92. **F** Sins JWR, Fijnvandraat K, Rijnveld AW, *et al.*: Effect of N-acetylcysteine on pain in daily life in patients with sickle cell disease: A randomised clinical trial. *Br J Haematol*. 2018; **182**(3): 444–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 93. Ataga KI, Kutlar A, Kanter J, *et al.*: Crizanlizumab for the Prevention of Pain Crises in Sickle Cell Disease. *N Engl J Med*. 2017; **376**(5): 429–39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 94. Orringer EP, Casella JF, Ataga KI, *et al.*: Purified poloxamer 188 for treatment of acute vaso-occlusive crisis of sickle cell disease: A randomized controlled trial. *JAMA*. 2001; **286**(17): 2099–106.
[PubMed Abstract](#) | [Publisher Full Text](#)
 95. <https://www.prnewswire.com/news-releases/mast-therapeutics-reports-top-line-results-from-phase-3-study-in-sickle-cell-disease-300331289.html>
 96. **F** Chang J, Patton JT, Sarkar A, *et al.*: GMI-1070, a novel pan-selectin antagonist, reverses acute vascular occlusions in sickle cell mice. *Blood*. 2010; **116**(10): 1779–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 97. **F** Telen MJ, Wun T, McCavit TL, *et al.*: Randomized phase 2 study of GMI-1070 in SCD: Reduction in time to resolution of vaso-occlusive events and decreased opioid use. *Blood*. 2015; **125**(17): 2656–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 98. https://www.pfizer.com/news/press-release/press-release-detail/pfizer_announces_phase_3_top_line_results_for_rivipansel_in_patients_with_sickle_cell_disease_experiencing_a_vaso_occlusive_crisis
 99. Matsui NM, Varki A, Embury SH: Heparin inhibits the flow adhesion of sickle red blood cells to P-selectin. *Blood*. 2002; **100**(10): 3790–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 100. Embury SH, Matsui NM, Ramanujam S, *et al.*: The contribution of endothelial cell P-selectin to the microvascular flow of mouse sickle erythrocytes *in vivo*. *Blood*. 2004; **104**(10): 3378–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
 101. Qari MH, Aljaouni SK, Alardawi MS, *et al.*: Reduction of Painful Vaso-occlusive Crisis of Sickle Cell Anaemia by Tinzaparin in a Double-blind Randomized Trial. *Thromb Haemost*. 2017; **98**(02): 392–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 102. Telen MJ, Batchvarova M, Shan S, *et al.*: Sevuparin Binds to Multiple Adhesive Ligands and Reduces Sickle Red Blood Cell-induced Vaso-occlusion. *Br J*

- Haematol.* 2016; **175**: 935–48.
[PubMed Abstract](#) | [Publisher Full Text](#)
103. Biemond BJ, Tombak A, Kilinc Y, *et al.*: Efficacy and Safety of Sevuparin, a Novel Non-Anti-Coagulant Heparinoid, in Patients with Acute Painful Vaso-Occlusive Crisis: A Global, Multicenter Double-Blind, Randomized, Placebo-Controlled Phase 2 Trial (TVOC01). *Blood*. 2019; **134**(suppl 1): 614.
[Publisher Full Text](#)
 104. Jang JE, Hidalgo A, Frenette PS: Intravenous immunoglobulins modulate neutrophil activation and vascular injury through FcγRIII and SHP-1. *Circ Res*. 2012; **110**(8): 1057–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 105. Manwani D, Chen G, Carullo V, *et al.*: Single-dose Intravenous Gammaglobulin Can Stabilize Neutrophil Mac-1 Activation in Sickle Cell Pain Crisis. *Am J Hematol*. 2015; **90**(5): 381–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 106. Griffin TC, McIntire D, Buchanan GR: High-dose intravenous methylprednisolone therapy for pain in children and adolescents with sickle cell disease. *N Engl J Med*. 1994; **330**(11): 733–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 107. Bernini JC, Rogers ZR, Sandler ES, *et al.*: Beneficial Effect of Intravenous Dexamethasone in Children With Mild to Moderately Severe Acute Chest Syndrome Complicating Sickle Cell Disease. *Blood*. 1998; **92**(9): 3082–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 108. **F** Leonard A, Godiwala N, Herrera N, *et al.*: Early Initiation of Inhaled Corticosteroids does not Decrease Acute Chest Syndrome Morbidity in Pediatric Patients with Sickle Cell Disease. *Blood Cells Mol Dis*. 2018; **71**: 55–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 109. **F** Glassberg J, Minniti C, Cromwell C, *et al.*: Inhaled Steroids Reduce Pain and sVCAM Levels in Individuals with Sickle Cell Disease: A Triple-blind, Randomized trial. *Am J Hematol*. 2017; **92**: 622–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 110. Wallace KL, Linden J: Adenosine A2A Receptors Induced on iNKT and NK Cells Reduce Pulmonary Inflammation and Injury in Mice with Sickle Cell Disease. *Blood*. 2010; **116**: 5010–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 111. **F** Field JJ, Lin G, Okam MM, *et al.*: Sickle cell vaso-occlusion causes activation of iNKT cells that is decreased by the adenosine A2A receptor agonist regadenoson. *Blood*. 2013; **121**(17): 3329–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 112. Field JJ, Majerus E, Gordeuk VR, *et al.*: Randomized phase 2 trial of regadenoson for treatment of acute vaso-occlusive crises in sickle cell disease. *Blood Adv*. 2017; **1**(20): 1645–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 113. Field JJ, Majerus E, Ataga KI, *et al.*: NNKT120, an anti-iNKT cell monoclonal antibody, produces rapid and sustained iNKT cell depletion in adults with sickle cell disease. *PLoS One*. 2017; **12**(2): e0171067.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 114. Hoppe C, Kuypers F, Larkin S, *et al.*: A pilot study of the short-term use of simvastatin in sickle cell disease: Effects on markers of vascular dysfunction. *Br J Haematol*. 2011; **153**(5): 655–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 115. Hoppe C, Jacob E, Styles L, *et al.*: Simvastatin reduces vaso-occlusive pain in sickle cell anaemia: A pilot efficacy trial. *Br J Haematol*. 2017; **177**(4): 620–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 116. Ataga KI, Wichlan D, Elsherif L, *et al.*: A pilot study of the effect of atorvastatin on endothelial function and albuminuria in sickle cell disease. *Am J Hematol*. 2019; **94**(11): E299–E301.
[PubMed Abstract](#) | [Publisher Full Text](#)
 117. Daak AA, Ghebremeskel K, Hassan Z, *et al.*: Effect of omega-3 (n-3) fatty acid supplementation in patients with sickle cell anemia: Randomized, double-blind, placebo-controlled trial. *Am J Clin Nutr*. 2013; **97**(1): 37–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
 118. **F** Daak AA, Dampier CD, Fuh B, *et al.*: Double-blind, randomized, multicenter phase 2 study of SC411 in children with sickle cell disease (SCOT trial). *Blood Adv*. 2018; **2**(15): 1969–79.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 119. Rees DC, Kilinc Y, Unal S, *et al.*: Double-Blind, Randomized Study of Canakinumab Treatment in Pediatric and Young Adult Patients with Sickle Cell Anemia. *Blood*. 2019; **134**: 615.
[Publisher Full Text](#)
 120. Gavins FNE, Russell J, Senchenkova EL, *et al.*: Mechanisms of enhanced thrombus formation in cerebral microvessels of mice expressing hemoglobin-S. *Blood*. 2011; **117**(15): 4125–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 121. Chanrathammachart P, Mackman N, Sparkenbaugh E, *et al.*: Tissue Factor Promotes Activation of Coagulation and Inflammation in a Mouse Model of Sickle Cell Disease. *Blood*. 2012; **120**(3): 636–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 122. Arumugam PI, Mullins ES, Shanmukhappa SK, *et al.*: Genetic diminution of circulating prothrombin ameliorates multiorgan pathologies in sickle cell disease mice. *Blood*. 2015; **126**(15): 1844–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 123. Ataga KI, Key NS: Hypercoagulability in Sickle Cell Disease: New Approaches to an Old Problem. *Hematology Am Soc Hematol Educ Program*. 2007; **2007**: 91–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 124. Pakbaz Z, Wun T: Role of the Hemostatic System on Sickle Cell Disease Pathophysiology and Potential Therapeutics. *Hematol Oncol Clin North Am*. 2014; **28**(2): 355–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 125. Greenberg J, Ohene-Frempong K, Halus J, *et al.*: Trial of Low Doses of Aspirin as Prophylaxis in Sickle Cell Disease. *J Pediatr*. 1983; **102**(5): 781–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
 126. Osamo NO, Photiades DP, Famodu AA: Therapeutic effect of aspirin in sickle cell anaemia. *Acta Haematol*. 1981; **66**(2): 102–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 127. Zago MA, Costa FF, Ismael SJ, *et al.*: Treatment of Sickle Cell Diseases With Aspirin. *Acta Haematol*. 1984; **72**(1): 61–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
 128. **F** Desai PC, Brittain JE, Jones SK, *et al.*: A Pilot Study of Eptifibatide for Treatment of Acute Pain Episodes in Sickle Cell Disease. *Thromb Res*. 2013; **132**(3): 341–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 129. Cabannes R, Lonsdorfer J, Castaigne JP, *et al.*: Clinical and Biological Double-Blind-Study of Ticlopidine in Preventive Treatment of Sickle-Cell Disease Crises. *Agents Actions Suppl*. 1984; **15**: 199–212.
[PubMed Abstract](#)
 130. Heeney MM, Hoppe CC, Abboud MR, *et al.*: A Multinational Trial of Prasugrel for Sickle Cell Vaso-Occlusive Events. *N Engl J Med*. 2016; **374**(7): 625–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
 131. **F** Hsu LL, Sarnaik S, Williams S, *et al.*: A Dose-Ranging Study of Ticagrelor in Children Aged 3-17 Years With Sickle Cell Disease: A 2-part Phase 2 Study. *Am J Hematol*. 2018; **93**(12): 1493–500.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 132. **F** Kanter J, Abboud MR, Kaya B, *et al.*: Ticagrelor Does Not Impact Patient-Reported Pain in Young Adults With Sickle Cell Disease: A Multicentre, Randomised Phase IIb Study. *Br J Haematol*. 2019; **184**(2): 269–78.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 133. Heeney MM, Abboud MR, Amilon C, *et al.*: Ticagrelor Versus Placebo for the Reduction of Vaso-Occlusive Crises in Pediatric Sickle Cell Disease: Rationale and Design of a Randomized, Double-Blind, Parallel-Group, Multicenter Phase 3 Study (HESTIA3). *Contemp Clin Trials*. 2019; **85**: 105835.
[PubMed Abstract](#) | [Publisher Full Text](#)
 134. Kato GJ, Steinberg MH, Gladwin MT: Intravascular Hemolysis and the Pathophysiology of Sickle Cell Disease. *J Clin Invest*. 2017; **127**(3): 750–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 135. Gladwin MT, Kato GJ, Weiner D, *et al.*: Nitric Oxide for Inhalation in the Acute Treatment of Sickle Cell Pain Crisis: A Randomized Controlled Trial. *JAMA*. 2011; **305**(9): 893–902.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 136. Maitre B, Djibre M, Katsahian S, *et al.*: Inhaled Nitric Oxide for Acute Chest Syndrome in Adult Sickle Cell Patients: A Randomized Controlled Study. *Intensive Care Med*. 2015; **41**(12): 2121–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 137. **F** Morris CR, Kuypers FA, Lavrisa L, *et al.*: A Randomized, Placebo-Controlled Trial of Arginine Therapy for the Treatment of Children With Sickle Cell Disease Hospitalized With Vaso-Occlusive Pain Episodes. *Haematologica*. 2013; **98**(9): 1375–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 138. **F** Eleutério RMN, Nascimento FO, Araújo TG, *et al.*: Double-Blind Clinical Trial of Arginine Supplementation in the Treatment of Adult Patients With Sickle Cell Anaemia. *Adv Hematol*. 2019; **2019**: 4397150.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 139. Machado RF, Barst RJ, Yovetich NA, *et al.*: Hospitalization for Pain in Patients With Sickle Cell Disease Treated With Sildenafil for Elevated TRV and Low Exercise Capacity. *Blood*. 2011; **118**(4): 855–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Yogen Sauntharajah

Department of Hematology and Oncology, Taussig Cancer Institute, Cleveland Clinic, Cleveland, OH, USA

Competing Interests: YS has patents, owns equity in, and serves on the board of a company EpiDestiny that is developing drug therapy for sickle cell disease.

2. Daniel E. Bauer

¹ Division of Hematology/Oncology, Boston Children's Hospital, Boston, MA, USA

² Department of Pediatric Oncology, Dana-Farber Cancer Institute, Boston, MA, USA

³ Department of Pediatrics, Harvard Medical School, Boston, MA, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research