RESEARCH ARTICLE

High vaccination coverage, inadequate knowledge and high vector density: Findings from a community-based cross-sectional study on Japanese Encephalitis in Yangon, Myanmar

[version 1; peer review: 2 approved with reservations]

Pyae Phyo Kyaw¹ , Hemant Deepak Shewade²⁻⁴ , Nang Thu Thu Kyaw^{2,5}, Khaing Hnin Phyo^{2,5}, Htar Htar Lin⁶, Aye Mon Mon Kyaw⁷, Mg Mg Mya¹, Sein Thau¹, Yan Naung Maung Maung¹

¹Department of Medical Research, Ministry of Health and Sports, Yangon, 11191, Myanmar

²International Union against Tuberculosis and Lung Disease (The Union), Paris, 75006, France

³The Union South East Asia Office, New Delhi, 110016, India

⁴Karuna Trust, Bengaluru, 560041, India

⁵The Union Myanmar Country Office, Mandalay, 05021, Myanmar

⁶Expanded Programme on Immunization, Ministry of Health and Sports, Nay Pyi Taw, Myanmar

⁷Vector Borne Disease Control Program, Ministry of Health and Sports, Yangon, 11211, Myanmar

V1 First published: 06 Jan 2020, 9:6
<https://doi.org/10.12688/f1000research.21702.1>
 Second version: 30 Jun 2020, 9:6
<https://doi.org/10.12688/f1000research.21702.2>
 Latest published: 08 Sep 2020, 9:6
<https://doi.org/10.12688/f1000research.21702.3>

Abstract

Background: Japanese encephalitis (JE) is a mosquito-borne disease with high case fatality and no specific treatment. Little is known about the community's (especially parents/guardians of children) awareness regarding JE and its vaccine in Yangon region, which bears the highest JE burden in Myanmar.

Methods: We conducted a community-based cross-sectional study in Yangon region (2019) to explore the knowledge and perception of parents/guardians of 1-15 year-old children about JE disease, its vaccination and to describe JE vaccine coverage among 1-15 year-old children. We followed multi-stage random sampling (three stages) to select the 600 households with 1-15 year-old children from 30 clusters in nine townships. Analyses were weighted (inverse probability sampling) for the multi-stage sampling design.

Results: Of 600 parents/guardians, 38% exhibited good knowledge of JE, 55% perceived JE as serious in children younger than 15 years and 59% perceived the vaccine to be effective. Among all the children in the 600 households, the vaccination coverage was 97% (831/855).

Conclusion: In order to reduce JE incidence in the community, focus on an intensified education program is necessary to sustain the high

Open Peer Review

Approval Status

	1	2
version 3 (revision) 08 Sep 2020		 view
version 2 (revision) 30 Jun 2020	 view 	 view
version 1 06 Jan 2020	 view	 view

1. **Jing An** , Capital Medical University, Beijing, China

2. **Babasaheb V. Tandale** , ICMR-National Institute of Virology, Pune, India

Pravin Deshmukh, ICMR - National Institute of Virology, Pune, India

vaccine coverage in the community.

Keywords

JE vaccine, knowledge and perception, community-based survey, caregiver's knowledge, SORT IT

Any reports and responses or comments on the article can be found at the end of the article.

This article is included in the **Médecins Sans Frontières** gateway.

This article is included in the **TDR** gateway.

Corresponding author: Pyae Phyo Kyaw (drppk.dmr@gmail.com)

Author roles: **Kyaw PP:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Shewade HD:** Conceptualization, Formal Analysis, Methodology, Software, Supervision, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Kyaw NTT:** Conceptualization, Formal Analysis, Software, Writing – Original Draft Preparation, Writing – Review & Editing; **Hnin Phyo K:** Conceptualization, Methodology; **Lin HH:** Conceptualization, Writing – Review & Editing; **Kyaw AMM:** Conceptualization, Writing – Review & Editing; **Mya MM:** Conceptualization, Data Curation, Formal Analysis, Methodology, Software, Supervision, Validation, Writing – Review & Editing; **Thaung S:** Data Curation, Visualization; **Maung Maung YN:** Conceptualization, Data Curation, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This research was conducted through the Structured Operational Research and Training Initiative (SORT IT), a global partnership led by the Special Programme for Research and Training in Tropical Diseases at the World Health Organization (WHO/TDR). The model is based on a course developed jointly by the International Union Against Tuberculosis and Lung Disease (The Union) and Médecins Sans Frontières (MSF/Doctors Without Borders). The specific SORT IT programme which resulted in this publication was jointly organized and implemented by The Centre for Operational Research, The Union, Paris, France; Department of Medical Research, Ministry of Health and Sports, Yangon; Department of Public Health, Ministry of Health and Sports, Nay Pyi Taw; The Union Country Office, Mandalay, Myanmar; The Union South-East Asia Office, New Delhi, India and London School of Hygiene and Tropical Medicine, London, UK. The training programme, within which this paper was developed, was funded by the Department for International Development (DFID), London, UK. We received funding support (USD \$1200) for data collection from the Department of Medical Research under Ministry of Health and Sports, Myanmar.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2020 World Health Organisation. This is an open access article distributed under the terms of the **Creative Commons Attribution IGO License**, which permits copying, adaptation and distribution in any medium or format for any purpose, provided the original work is properly cited, a link is provided to the license, and any changes made are indicated. Any such copying, adaptation and distribution must not in any way suggest that World Health Organisation endorses you or your use.

How to cite this article: Kyaw PP, Shewade HD, Kyaw NTT *et al.* **High vaccination coverage, inadequate knowledge and high vector density: Findings from a community-based cross-sectional study on Japanese Encephalitis in Yangon, Myanmar [version 1; peer review: 2 approved with reservations]** F1000Research 2020, 9:6 <https://doi.org/10.12688/f1000research.21702.1>

First published: 06 Jan 2020, 9:6 <https://doi.org/10.12688/f1000research.21702.1>

Introduction

Japanese encephalitis (JE) is a zoonotic disease caused by Japanese encephalitis virus (JEV). JEV exists in a transmission cycle between mosquitoes, pigs and/or water birds and is transmitted to humans through bites from infected mosquitoes of the *Culex* species (mainly *Culex tritaeniorhynchus*)¹. JE usually presents as acute encephalitis syndrome (AES) and is confirmed by serology.

JE is a disease of public health importance as billions of people are at risk of getting infected by JEV and children below 15 years are more susceptible¹. A systematic review (2011) reported that 67,900 clinical cases of JE occur annually in 24 Asian and western Pacific countries despite the widespread availability of the vaccine, with approximately 13,600 to 20,400 deaths. While the overall incidence of JE is 1.8 per 100,000 per year in endemic countries, it is 5.4 among 1–15 year-old children². The infection can lead to severe complications with high case fatality. There is no specific treatment to date. JE is not easily prevented by protection from mosquito control and mosquito bites. Hence, vaccination is the most effective form of prevention¹. Globally, live attenuated SA 14-14-2 JE vaccine is the most commonly used JE vaccine. Vaccine efficacy is reported to be between 80% and 99% following single-dose vaccination and 98% or greater with two doses³.

In Myanmar, *Culex tritaeniorhynchus* is the main JE vector⁴. In 2012, there were only 14 confirmed JE cases, which increased to 151 in 2015 and more than 380 in 2016 and 2017, indicative of improved surveillance in the country⁵.

In 2017, the Expanded Programme on Immunization (EPI) under the Department of Public Health carried out a nationwide JE vaccination catch-up campaign supported by the GAVI, WHO, the United Nations Children's Fund and PATH. In November and December, nearly 14 million children (aged nine months to 15 years) were targeted and vaccinated with the WHO live attenuated (SA 14-14-2) JE vaccine. Alongside the campaign there were extensive advocacy and sensitization sessions provided to schools and communities⁶. Since January 2018, immediately after the JE campaign, routine immunization in Myanmar has included JE live attenuated vaccine, given at the age of nine months together with the measles-rubella vaccine.

Adequate knowledge of JE and a positive perception of the JE vaccine are important for the adoption of preventive measures⁷. In addition, high coverage of the JE vaccine in populations at risk of disease is required to reduce the JE cases because JE vaccination would not provide herd immunity⁸. A study conducted in one township in northern Shan state in Myanmar (2018) showed the level of awareness of JE and its vaccine was low but the perception towards knowledge of JE was generally positive⁹. The vaccination coverage was 93% among 391 study participants.

Little is known about the community's (especially parents/guardians of children) awareness regarding JE and its vaccine in Yangon region, which bears the highest JE burden in the

country⁶. This data, alongside vaccination coverage data for children, may help the regional vector borne disease control (VBDC) programme and EPI to develop a new coordinated strategic plan to successfully reduce JE transmission in Yangon region.

Therefore, in Yangon region, we aimed to describe the i) knowledge and perception of the parents/guardians of children (1–15 years old) towards JE disease and vaccine, and ii) JE vaccine coverage among 1–15 year-old children.

Methods

Ethical statement

The Ethics Review Committee, Department of Medical Research, Myanmar (Ethics/DMR/2018/102EA/2019/028) and the Union Ethics Advisory Group of the International Union against Tuberculosis and Lung Disease (The Union), Paris, France (EAG 04/19) approved the study. Written informed consent for participation in the survey was taken from the parents/guardians of children aged 1–15 years old and the consent process was approved by the ethics committees.

Study design

This was a community-based cross-sectional survey involving primary data collection.

Study setting

Myanmar is located in the Southeast Asia region, neighboring Laos to the east, Bangladesh to the west, Thailand to the southeast, the Republic of China to the north and northeast and India to the northwest. Myanmar has a population of 51 million with an urban:rural population ratio of 30:70. The country has 14 states and regions including Nay Pyi Taw council territory. It consists of 74 districts, 330 townships, 398 towns, 3065 wards, 13,619 village tracts and 64,134 villages¹⁰.

Yangon is the economic capital of Myanmar with four districts, 46 townships, 743 wards and 628 village tracts. The population of Yangon region is the highest in size when compared with other states and regions in the country. The urban:rural population ratio is 70:30¹⁰.

Study population

The study population included parents/guardians of children (1–15 years old) living in Yangon region for the first objective and children (1–15 years old) for the second objective (February to June 2019).

Sample size and sampling procedure

For the first two objectives, assuming that the prevalence of community awareness of JE and its vaccination among parents/guardians of children (1–15 years old) and vaccination coverage of children (1–15 years old) in Yangon being 50% with 5% precision at 95% CI, the calculated sample size was 384. Assuming a non-response rate of 5% and a design effect of 1.5, the final sample size was 600 households with children (1–15 years old) We used the conservative assumption of 50% prevalence as there is no previous data on community

awareness and vaccination coverage in Yangon region and 5% non-response rate based on field experience.

We used three stage random sampling to sample the 600 households with children (1–15 years) from 30 clusters (ward or village tract) in nine townships in Yangon region (see [Figure 1](#) and [Figure 2](#)). In the first two stages, we used stratified random sampling and in the third stage, we used systematic

random sampling. First, we randomly selected the nine townships after stratifying them into urban, rural and mixed based on the classification used in population census in Myanmar¹⁰. Then, in the second stage, we randomly selected 30 clusters after stratifying them into wards and village tracts in each selected township. Within each township, trained field assistants went directly to the general administration department for a list of the households and map of the selected wards/village tracts. The

Figure 1. Multi-stage random sampling (three stage) to select households with 1–15 year-old children in the Japanese encephalitis survey, Yangon region, Myanmar (2019). *Predominantly urban (>70% urban population); **Predominantly rural (>70% rural population).

Figure 2. Randomly selected nine townships for the Japanese encephalitis survey, Yangon region, Myanmar (2019).

trained field assistants conducted systematic random sampling to select 20 households with children aged 1–15 years old within each cluster. The trained field assistants chose a random starting point using the map and then selected the first household. Then, they went to next household using the sampling interval until the required sample size was reached. Sampling interval was calculated by dividing the total number of household in the selected wards or village tracts by the required sample size for that selected wards or village tracts. If the selected household did not have any children aged 1–15 years old, field assistants went to the next adjacent household with a child in this age range. If the selected house was locked or there was no parent or guardian, field assistants followed the same procedure as above. In case the selected house was an apartment building, we selected one household randomly. We interviewed the parent or guardian of the child available at the time of survey, preferably the mother.

Data collection and tools

During February-June 2019, at the selected households, field assistants conducted face to face interviews with parents/guardians using a pretested structured questionnaire (see Annex 1 and Annex 3, *Extended data*)^{11,12}. The field assistant asked the questions verbally to the participants and completed the questionnaires. The structured questionnaires were tested during a pilot survey and were revised according to feedback received during the pilot survey.

The questionnaire consisted of four sections. The first section highlighted the socio-demographic information. The second section was comprised of 12 questions that assessed the knowledge of participants about JE disease and prevention (including vaccination). The third section assessed perception. The fourth section included information about vaccination status of the children. If the selected household had more than one child aged 1–15 years old, we asked the parent/guardian about the vaccination status of all the children in this age range. Vaccination status was based on parents' or guardians' recall. Field assistants also asked about the presence or absence of a vaccination card.

Data management and analysis

Data from the survey forms were double-entered and validated using EpiData entry software (version 3.1, EpiData Association, Odense, Denmark). Data were analyzed using STATA (version 12.1, STATA Corp., College Station, TX, USA). There were no missing data in the study.

We provided weighted estimates as the analyses were weighted (inverse probability sampling) for the multi-stage sampling design. We used frequency and proportion to summarize the characteristics of the study participants. We assigned a knowledge score to each participant based on the number of correct or appropriate responses. Each appropriate answer was assigned one point and incorrect responses or "do not know" were assigned zero points. The scores were further dichotomized into poor or good (0–6 as poor and 7–12 as good). No overall score was calculated for perception. Vaccine coverage was calculated by the number of children that received JE vaccination

(either during campaign or routine immunization) divided by the total number of children and presented as proportion and 95% confidence interval (CI).

Results

The socio-demographic characteristics of 600 parents/guardians are presented in Table 1¹³. Among them, 1% were aged ≤ 20 years, 4% were aged >60 years and 74% were female.

Table 1. Socio-demographic characteristics of participants* in the Japanese encephalitis survey, Yangon region, Myanmar (2019).**

Variable	N	(%)***
Total	600	(100.0)
Age in years		
≤20	6	(1.0)
21–40	293	(48.8)
41–60	280	(46.6)
>60	21	(3.6)
Sex		
Male	155	(25.9)
Female	445	(74.1)
Education level		
Graduate	138	(22.9)
High School	165	(27.4)
Middle School	163	(27.2)
Primary School	105	(17.6)
Can read and write	24	(3.9)
Illiterate	5	(1.0)
Occupation		
Government staff	29	(4.9)
Private employee	186	(30.9)
Own business	150	(24.9)
Dependent	215	(35.9)
Others	20	(3.4)
Type of family		
Nuclear	342	(56.2)
Extended	258	(43.8)
Family members suffered JE		
Yes	0	0.0
No	599	(99.2)
Do not know	1	(0.8)
Family income per month (USD)*		
<95	8	(1.3)
95 – 189	101	(16.8)
190 – 285	316	(52.7)
>285	175	(29.2)

JE, Japanese encephalitis; USD, United States dollar; one USD = 1525 Myanmar Kyats.

*Parent or guardian of the 1–15 year-old children available at time of survey, preferably mother. **Weighted estimates given taking into account the sampling design.

*** Column percentage.

A total of 50% had a high school or graduate level education and 29% had a monthly family income of more than \$285 USD.

Knowledge and perception

Overall, 37.6% exhibited good knowledge of JE. We have depicted the knowledge of respondents regarding cause, transmission, symptoms, prevention and treatment in [Table 2](#). Among 600 parents/guardians, 49.3% had correct knowledge that JE is a fatal disease. Although 65.3% knew that the JE vaccine was available locally, only 26.8% correctly answered that vaccination is the most effective means to protect against JE.

It was found that 23% did not know the symptoms of JE, 16% responded incorrectly that JE has specific treatment and 58.8% responded that they did not know whether there is a treatment for JE or not. Participants responded that they used mosquito nets (31.3%), mosquito coils (14.1%) and spray or fumigation methods (10.7%) to avoid mosquito bites.

Over half (55%) of participants perceived JE as serious in children younger than 15 years, 59% perceived the vaccine to be effective and 25% perceived JE to be harmful for pig farmers ([Figure 3](#)). Health care staff (25%) and television (17%) were the main sources of information about JE disease ([Figure 4](#)). The source of information on JE routine vaccination or campaigns was from school (35.1%), health worker visits (33.1%), announcements made using microphones in the neighborhood (13.9%) and volunteer visits (5.3%).

Vaccine coverage

Among all the children in the 600 households (n=855), 831 were vaccinated. The vaccination coverage was 97.2% (95% CI: 95.9-98.1). Of 831 vaccinated children, 423 (50.9%) were cross-checked through a vaccination card. Of 831 children, 516 (62.1%) received the vaccination during the JE campaign, 234 (28.2%) during routine immunization and 67 (7.8%) received the vaccine twice during both campaign and routine vaccination. Among the 24 children that did not receive vaccination, the main reasons were: parents or guardians did not realize the importance of JE vaccination (n=4), the child was sick at the time of immunization (n=3), the parents/guardians did not know about the JE vaccination (n=2) and travel (n=6).

Discussion

In this region-wide survey on JE and its vaccine coverage in Yangon, Myanmar, the majority of parents or guardians did not have good knowledge of JE. Perception of the seriousness of JE disease was poor in half of participants. Vaccination coverage was excellent.

Strengths and limitations

Data were robust as double entry and validation minimized data entry errors. Only one attempt was made to visit each household, which may impact the generalizability of the results as the households with parents and guardians who were working during the survey visit might be missed. Half of parents/guardians did not produce a vaccination card (JE

Table 2. Knowledge of participants* regarding cause, transmission, symptoms, prevention and treatment in the Japanese encephalitis survey, Yangon region, Myanmar (2019).**

Question	Response	Total	
		n	(%)***
Total		600	(100.0)
Have you ever heard about JE?	Yes	409	(68.2)
What is the cause of JE?	Virus	99	(16.5)
How does the disease get transmitted?	Mosquito	237	(39.5)
What are the common symptoms of JE?	High fever and convulsion	189	(31.5)
Is JE prevalent in children under 15 years?	Yes	264	(44.0)
Where does JE mostly prevail?	Rural areas and paddy fields	114	(19.0)
Are pigs the amplifier host?	Yes	129	(21.5)
Is JE a fatal disease?	Yes	296	(49.3)
Is there any specific treatment for JE?	No	151	(25.2)
Is JE a preventable disease?	Yes	331	(55.2)
Is the JE vaccine available locally?	Yes	392	(65.3)
How can one protect against JE?	JE vaccine	161	(26.8)

JE, Japanese encephalitis.

*Parent or guardian of the 1–15 year-old children available at time of survey, preferably mother.

**Weighted estimates given taking into account the sampling design.

*** Column percentage.

Figure 3. Perception of participants* in the Japanese encephalitis survey, Yangon region, Myanmar (2019) **. The variable "JE is serious if it occur in you" was included because sometimes adults think that some kinds of mosquito borne diseases such as Japanese encephalitis and Dengue are more serious in children and not serious when those diseases occur in them. Researchers in this study wanted to know whether participants had these perceptions. JE, Japanese encephalitis. Y axis as percentage (responded 'yes'). *Parent or guardian of the 1–15 year-old children available at time of survey, preferably mother. **Weighted estimates given taking into account the sampling design.

Figure 4. Source of information of participants* about Japanese encephalitis in the Japanese encephalitis survey, Yangon region, Myanmar (2019) **. Y axis as percentage (responded 'yes'). *Parent or guardian of the 1–15 year-old children available at time of survey, preferably mother. **Weighted estimates given taking into account the sampling design.

routine immunization or campaign) and recall bias cannot be ruled out. However, as JE vaccination is through an injection (subcutaneous) at a specific site (in the right upper arm) and the vaccination campaign was implemented only once in 2017, we do not think this is a serious limitation.

Key findings and comparison with other studies

We found that parents/guardians had poor knowledge of JE. This finding is similar to the study conducted in a slum of Kolkata by Dasgupta *et al.* (2016), which showed that 56.7% of mothers had poor knowledge¹⁴. Our study showed that 39.5% of participants knew that JE is caused by mosquito bites, which is higher than the 25.6% reported by Dasgupta *et al.*¹⁴. In our study, the majority of participants showed a lack of

knowledge about treatment of the disease, which is similar to a study from India (2015) by Ahmad *et al.*⁷.

More than half of the participants perceived JE as serious in children younger than 15 years old, but they did not perceive pig farming as contributing to the threat of JE. This is similar to the study in India, where respondents did not perceive pig farming as contributing to the threat of JE¹⁵. Possible reasons for poor knowledge and perception of participants was that communities did not think of JE as a possible threat like other vector-borne diseases such as dengue and malaria.

In our study, although only 55.2% of participants knew that JE is a preventable disease and 65.3% knew the most effective

means to protect JE (vaccination), almost all parents/guardians vaccinated their children. It reflects their reliance on existing health care delivery systems and EPI. JE vaccine coverage was high in this study and an effective JE vaccination catch-up campaign by EPI might have contributed to the high vaccine coverage. Coverage is similar to the study from the district of Ambala, Haryana in north India, which demonstrated a high level (93.9%) of JE vaccine coverage in the study area¹⁶. Similarly, JE vaccine coverage in Hsipaw Township, Northern Shan State, Myanmar was 93.0% in 2018⁹.

The most important vector is *Culex tritaeniorhynchus*, which feeds on vertebrate hosts, primarily pigs in preference to humans¹⁷. Both larva and adult mosquito of *Culex tritaeniorhynchus* can spread JEV and were collected for an entomological and vector bionomics study of JE transmission done in four villages of Letpadan Township, Bago region (Myanmar), showing a high proportion (69.8%) of *Culex tritaeniorhynchus* in areas with high pig farming¹⁸.

Implications

JE cases in Yangon region have declined in 2018 and 2019 compared to earlier years. In January – August 2019, there were only 244 AES cases and 12 confirmed JE cases¹⁹. To sustain this decline, high vaccination coverage, health education on JE and effective vector control activity should be maintained. There is a possibility that the high coverage may not be maintained in the long term. This is because within 16–20 months of JE campaign, many were not aware of JE or its vaccine. Hence, steps should be taken in this direction.

Authorities should encourage retention of cards or records of vaccination so unvaccinated children can be identified and vaccinated in future.

Conclusions

JE vaccination coverage was excellent in Yangon region, Myanmar, despite the majority of parents/guardians having poor knowledge and perception of JE disease, its prevention and vaccination. In order to reduce JE incidence in community, a focus on an intensified education program is necessary to sustain the high vaccine coverage in the community.

Data availability

Underlying data

Figshare: Annex 2. <https://doi.org/10.6084/m9.figshare.10548623.v1>¹³.

Extended data

Figshare: Annex 1. <https://doi.org/10.6084/m9.figshare.10552088.v1>¹¹.

Figshare: Annex 3. <https://doi.org/10.6084/m9.figshare.11458437.v1>¹².

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC0 1.0 Public domain dedication).

Acknowledgements

We acknowledge the regional health department, Vector Borne Disease Control Program, Expanded Programme on Immunization, staff in medical entomology research division, the township administrative offices and basic health staff in Yangon for their support while collecting data.

Disclaimer: The views expressed in this paper are of the authors and do not represent the views of the authors' organizations.

References

- World Health Organization WHO: **Japanese encephalitis, Immunization, Vaccines and Biologicals, estimate of the global disease burden**. 2015. (accessed on Sep 20, 2019).
[Reference Source](#)
- Campbell GL, Hills SL, Fischer M, *et al.*: **Estimated global incidence of Japanese encephalitis: a systematic review**. *Bull World Health Organ*. 2011; **89**(10): 766–74, 774A–774E.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- World Health Organization: **Japanese encephalitis/live attenuated vaccine**. 2006. (accessed on May 20, 2006).
[Reference Source](#)
- World Health Organization: **The sixth Bi-regional meeting on Prevention and control of Japanese encephalitis**. Bangkok, Thailand, 2014.
[Reference Source](#)
- Central Epidemiology Unit: **Monthly Epidemiology Bulletin**. 2018.
[Reference Source](#)
- Vector Borne Disease Control Unit: **Annual VBDC report**. NayPyi Taw, Myanmar, 2014.
- Ahmad A, Khan MU, Gogoi LJ, *et al.*: **Japanese Encephalitis in Assam, India: Need to Increase Healthcare Workers' Understanding to Improve Health Care**. *PLoS One*. 2015; **10**(8): e0135767.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hegde NR, Gore MM: **Japanese encephalitis vaccines: Immunogenicity, protective efficacy, effectiveness, and impact on the burden of disease**. *Hum Vaccin Immunother*. 2017; **13**(6): 1–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Thinn T, Lin TP, Thauung Y, *et al.*: **Awareness, Perception and Vaccination of Japanese Encephalitis Vaccine among Care Givers in Hsipaw Township, Northern Shan State**. 2018. (accessed on Sep 20, 2019).
[Reference Source](#)
- Ministry of Immigration and Population: **The 2014 Myanmar Population and Housing Census**. NayPyi Taw, 2015.
[Reference Source](#)
- Shewade HD: **Annex 1. figshare**.
<http://www.doi.org/10.6084/m9.figshare.10552088.v1>
- Shewade H: **Annex 3. figshare**. 2019.
<http://www.doi.org/10.6084/m9.figshare.11458437.v1>
- Shewade HD: **Annex 2. figshare**.
<http://www.doi.org/10.6084/m9.figshare.10548623.v1>
- Aparajita D, Atanu D, Bobby P, *et al.*: **Perception of Prevention of Japanese Encephalitis with Emphasis on Its Vaccination Programme: A Community Based Study In a Slum of Kolkata**. *Int J Public Heal Res*. 2017; **7**(2): 845–851.
[Reference Source](#)
- Chaturvedi S, Sharma N, Kakkar M: **Perceptions, practices and health seeking behaviour constrain JE/AES interventions in high endemic district of North India**. *BMC Public Health*. 2017; **17**(1): 645.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kumar D, Singh AJ: **Response of Various Stakeholders towards Newly**

- Introduced Japanese Encephalitis Vaccine in a North Indian State. *J Vaccines*. 2014; 1–4.
[Publisher Full Text](#)
17. Gupta N, Rao PV: **Transcriptomic profile of host response in Japanese encephalitis virus infection**. *Virology*. 2011; 8: 92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Mya MM, Lin Z, Min Hein ZN, *et al.*: **Vector bionomics of Culex species responsible for Japanese Encephalitis transmission in 4 villages of Letpadan Township Bago region, Myanmar**. *Myanmar Heal Res J*. 2017; 48: 37–39.
19. Central Epidemiology Unit: **AFP surveillance Indicators by State and Region, Fever with Rash Surveillance**. 2019.
[Reference Source](#)

Open Peer Review

Current Peer Review Status: ? ?

Version 1

Reviewer Report 29 May 2020

<https://doi.org/10.5256/f1000research.23924.r63433>

© 2020 Tandale B et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Babasaheb V. Tandale

ICMR-National Institute of Virology, Pune, Maharashtra, India

Pravin Deshmukh

ICMR - National Institute of Virology, Pune, India

- The manuscript is composed well and has been presented in a very simple way making it very easy to understand by the readers. As indicated by the other reviewer, more in-depth analysis and comparison to explain the similarities and differences in knowledge and perception of JE disease and JE vaccine may be considered along with the explanatory variables for coverage of JE vaccination. The comparisons would thus help identify the important aspects that could be targeted for improvements.
- The sampling design could be better explained with sampling ratios planned at different phases of sampling.
- Vector distribution / density has been indicated in title, however it is not dealt in abstract, objectives, methods and results. Therefore, this aspect may be dropped from title also. It could be mentioned in discussion section as the secondary data and indicated for its importance for understanding of study findings.
- The higher coverage of vaccination in spite of low level of knowledge and perception of JE disease and JE vaccine is difficult to understand. It may be highlighted and discussed in details.
- The objectives may be clearly mentioned at the end of background section.
- The potential sources of bias in selection, verification of information and interpretation of information given by respondents needs to be critically appraised and discussed. The confounders also need to be addressed properly.
- The verification of vaccination card for ascertaining vaccination recall needs to be addressed

with efforts made for data quality.

- The sample size and statistical analysis plan could be included with specific details on the basis of assumptions and proposed analysis. The sample size estimation could have been based on reported coverage of vaccination to be 93% in small survey findings earlier. The weighting approach needs to be provided clearly with its application in crude/unadjusted and adjusted/corrected coverage.
- The survey currently presented is mostly descriptive in nature and needs to be made analytical with comparisons that could be made and presented.
- The questionnaire validity may be presented with variable responses for similar aspects.
- Non-responses and its handling in analysis may be included.
- The baseline characteristics of respondents may be compared with good or bad knowledge, acceptable perception or otherwise and vaccinated and unvaccinated.
- Limitations of study may be discussed with interpretations based on the same.
- Generalisability of study findings may be considered and discussed.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Epidemiology of virus diseases causing outbreaks.

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however we have

significant reservations, as outlined above.

Author Response 25 Jun 2020

Pyae Phyo Kyaw, Ministry of Health and Sport, Myanmar, Yangon, Myanmar

1. The manuscript is composed well and has been presented in a very simple way making it very easy to understand by the readers. As indicated by the other reviewer, more in-depth analysis and comparison to explain the similarities and differences in knowledge and perception of JE disease and JE vaccine may be considered along with the explanatory variables for coverage of JE vaccination. The comparisons would thus help identify the important aspects that could be targeted for improvements.

Response: Thank you for your suggestion and we performed more in-depth analysis to determine the socio-demographic characteristics associated with good knowledge score and presented in the table-3. (line 218-234) We also performed an in-depth analysis to determine the good knowledge score with the child being vaccinated and presented in the table-4. (line 239-244)

2. The sampling design could be better explained with sampling ratios planned at different phases of sampling.

Response: We added a detailed explanation of the sampling design and procedure in the method section. (line 81-105)

3. Vector distribution / density has been indicated in title, however it is not dealt in abstract, objectives, methods and results. Therefore, this aspect may be dropped from title also. It could be mentioned in discussion section as the secondary data and indicated for its importance for understanding of study findings.

Response: We have revised the title according to your suggestion.

4. The higher coverage of vaccination in spite of low level of knowledge and perception of JE disease and JE vaccine is difficult to understand. It may be highlighted and discussed in details.

Response: We have added a discussion in the revised manuscript. (line 276-282)

5. The objectives may be clearly mentioned at the end of background section.

Response: We revised the paragraph at the end of introduction section and stated the objectives clearly. (line 42-44)

6. The potential sources of bias in selection, verification of information and interpretation of information given by respondents needs to be critically appraised and discussed. The confounders also need to be addressed properly.

Response: We have discussed the potential for recall bias as half the respondents could not

produce the immunization or JE campaign card (for details see line 251-258). We performed more in-depth analysis to assess factors associated. (line 145-148) (line 218-226)

7. The verification of vaccination card for ascertaining vaccination recall needs to be addressed with efforts made for data quality.

Response: We were able to verify 50% of the vaccinated children with their vaccination card. We acknowledged that the respondents who did not have vaccination card would subject to recall bias. (line 254-255)

8. The sample size and statistical analysis plan could be included with specific details on the basis of assumptions and proposed analysis. The sample size estimation could have been based on reported coverage of vaccination to be 93% in small survey findings earlier. The weighting approach needs to be provided clearly with its application in crude/unadjusted and adjusted/corrected coverage.

Response: We agree that we should have calculated the sample size based on a previous estimate. But we decided to assume a coverage of 50% that would give a maximum sample size. We have added a detailed description on sample size calculation and statistical analysis in the revised manuscript. We provided weighted estimates as the analyses were weighted (inverse probability sampling) for the multi-stage sampling design, this was done both for the descriptive and analytical calculations (line 82-89, 136-137)

9. The survey currently presented is mostly descriptive in nature and needs to be made analytical with comparisons that could be made and presented.

Response: We performed more in-depth analysis to determine the socio-demographic characteristics associated with good knowledge score and presented in the table-3. (line 218-240) We also performed an in-depth analysis to determine the good knowledge score with the child being vaccinated and presented in the table-4. (line 239-244)

10. The questionnaire validity may be presented with variable responses for similar aspects.

Response: Thank you for the comment. Unfortunately, we do not have information on questionnaire validity.

11. Non-responses and its handling in analysis may be included.

Response: There was no non-responses in our study because we went to the household using the sampling interval until the required sample size was reached. We added this information in the revised manuscript. (line 104-105)

12. The baseline characteristics of respondents may be compared with good or bad knowledge, acceptable perception or otherwise and vaccinated and unvaccinated.

Response: We have added results on the association between the baseline characteristics of respondents with knowledge level and vaccination status. (line 218-240)

13. Limitations of study may be discussed with interpretations based on the same.

Response: We have added this discussion in our revised manuscript. (line 250-258)

14. Generalisability of study findings may be considered and discussed.

Response: We have added this discussion in our revised manuscript. (line 250-258)

Competing Interests: No competing interests were disclosed.

Reviewer Report 27 April 2020

<https://doi.org/10.5256/f1000research.23924.r62066>

© 2020 An J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Jing An

Department of Microbiology, School of Basic Medical Sciences, Capital Medical University, Beijing, China

The authors here conducted a set of survey data to reflect the public awareness on JEV in community. A great deal of data collection was presented. These contribute to show the significant progresses of the JE campaign in Yangon. However, there are several questions to be answered.

Some Comments:

1. The whole article was simple as a scientific research. More in-depth analysis of data is recommended. More comparison should be conducted to reflect the details in multi-aspects. For example, "parents/guardians had poor knowledge of JE" is sentenced. Since the details of the guardians were investigated, the correlation should be conducted, like with age? or occupations? or else? or none. Besides, are there any differences in awareness between urban and rural places?...
2. To fully reflect the effect of public awareness in Yangon, authors should not choose nine townships randomly. Especially the locations are concentrated. The choices need to be more meaningful, like the widest vector distribution ones, the most severely suffered ones. OR, at least, the details of these nine townships should be presented, like incidence cases.
3. The complex relationship between JEV and congeneric members, like DENV and ZIKV, is widely proposed. Since dengue is prevalent locally, the vaccination against Flavivirus members or infections should be also considered in the questionnaire.

4. In table 1, (1) is single parent family excluded? (2) what does "Do not know family members suffered JE" exactly mean? do not know the symptoms or do not know the children's conditions?

5. "23% did not know the symptoms of JE" is sentenced. So, how will these parents know their kids are not suffered from JE? Is the "0" morbidity presented by parents credible?

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Interaction between viruses and hosts; mosquito-borne flavivirus; vaccine

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 25 Jun 2020

Pyae Phyo Kyaw, Ministry of Health and Sport, Myanmar, Yangon, Myanmar

1. The whole article was simple as a scientific research. More in-depth analysis of data is recommended. More comparison should be conducted to reflect the details in multi-aspects. For example, "parents/guardians had poor knowledge of JE" is sentenced. Since the details of the guardians were investigated, the correlation should be conducted, like with age? or occupations? or else? or none. Besides, are there any differences in awareness between urban and rural places?...

Response : Thank you for your suggestion and we performed more in-depth analysis to determine the socio-demographic characteristics associated with good knowledge score and presented in the table-3. (line 218-234) We also performed an in-depth analysis to

determine the good knowledge score with the child being vaccinated and presented in the table-4. (line 239-244)

2. To fully reflect the effect of public awareness in Yangon, authors should not choose nine townships randomly. Especially the locations are concentrated. The choices need to be more meaningful, like the widest vector distribution ones, the most severely suffered ones. OR, at least, the details of these nine townships should be presented, like incidence cases.

Response : We believe that random selection of township would provide generalizable information for the Yangon region. If we selected based on the incidence rate of JE cases or vector distribution, the knowledge, perception and vaccination coverage would not be representative to the population in Yangon. We hope this is fine. (line 78-90)

3. The complex relationship between JEV and congeneric members, like DENV and ZIKV, is widely proposed. Since dengue is prevalent locally, the vaccination against Flavivirus members or infections should be also considered in the questionnaire.

Response: In Myanmar, we have no vaccination programme against Dengue and Flavivirus infections and we did not have the opportunity to explore about this.

4. In table 1, (1) is single parent family excluded? (2) what does "Do not know family members suffered JE" exactly mean? do not know the symptoms or do not know the children's conditions?

Response:

(1) Single parent family was not excluded, considered under nuclear family. In type of family, Nuclear family means family which has father or mother with their children. Extended family means family which has either father, mother or grandfather, grandmother or uncle, aunt with children. We added this in the footnote of the table. (line 159-161)

(2) "Do not know family members suffered JE" means parents/guardians did not know their children have previous history of infection with JE virus. (We removed that variable from table-1 as the author group feels it does not add valuable information)

5. "23% did not know the symptoms of JE" is sentenced. So, how will these parents know their kids are not suffered from JE? Is the "0" morbidity presented by parents credible?

Response: The answer by the participant to the question on JE morbidity by asking whether the family member suffered JE before was based on the clinic or hospital record. So it is possible that the participant can report whether or not their children suffered JE even they did not know the symptoms of JE. Anyway, I removed that variable from table 1 as the author group feels it does not add valuable information.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research