REVIEW

An overview of mammalian p38 mitogen-activated protein kinases, central regulators of cell stress and receptor signaling [version 1; peer review: 2 approved]

Jiahuai Han¹, Jianfeng Wu¹, John Silke^{2,3}

¹State Key Laboratory of Cellular Stress Biology, Innovation Center for Cell Biology, School of Life Sciences, Xiamen University, Xiamen, Fujian, 361102, China

²The Walter and Eliza Hall Institute, 1G Royal Parade, Parkville, Victoria, 3052, Australia

³Department of Medical Biology, University of Melbourne, Parkville, Victoria, 3050, Australia

V1 First published: 29 Jun 2020, 9(F1000 Faculty Rev):653
<https://doi.org/10.12688/f1000research.22092.1>

Latest published: 29 Jun 2020, 9(F1000 Faculty Rev):653
<https://doi.org/10.12688/f1000research.22092.1>

Abstract

The p38 family is a highly evolutionarily conserved group of mitogen-activated protein kinases (MAPKs) that is involved in and helps co-ordinate cellular responses to nearly all stressful stimuli. This review provides a succinct summary of multiple aspects of the biology, role, and substrates of the mammalian family of p38 kinases. Since p38 activity is implicated in inflammatory and other diseases, we also discuss the clinical implications and pharmaceutical approaches to inhibit p38.

Keywords

p38, MAPK, inflammation, signalling

Open Peer Review

Approval Status

	1	2
version 1		
29 Jun 2020		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Jonathan D. Ashwell**, National Cancer Institute, National Institutes of Health, Bethesda, USA

2. **Guadalupe Sabio**, Centro Nacional de Investigaciones Cardiovasculares (CNIC), Madrid, Spain

Any comments on the article can be found at the end of the article.

Corresponding authors: Jiahuai Han (jiahuaihan@xmu.edu.cn), John Silke (silke@wehi.edu.au)

Author roles: **Han J:** Writing – Original Draft Preparation, Writing – Review & Editing; **Wu J:** Writing – Original Draft Preparation; **Silke J:** Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the National Natural Science Foundation of China (81788101, 31420103910 and 81630042 to J.H.), the 111 Project (B12001 to J.H.) and by an NHMRC fellowship (1107149) to JS.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2020 Han J *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Han J, Wu J and Silke J. **An overview of mammalian p38 mitogen-activated protein kinases, central regulators of cell stress and receptor signaling [version 1; peer review: 2 approved]** F1000Research 2020, 9(F1000 Faculty Rev):653 <https://doi.org/10.12688/f1000research.22092.1>

First published: 29 Jun 2020, 9(F1000 Faculty Rev):653 <https://doi.org/10.12688/f1000research.22092.1>

p38 mitogen-activated protein kinases

p38 α (originally named p38) was identified and cloned as a 38 kDa protein that was tyrosine-phosphorylated in response to LPS stimulation in mammalian cells^{1,2}. Sequence comparison, on the day p38 α was cloned, revealed that it belonged to the mitogen-activated protein kinase (MAPK) family and that a *Saccharomyces cerevisiae* osmotic response protein kinase HOG1 was a p38 α homologue^{3–5}. p38 α was also named cytokine suppressive drug binding protein (CSBP) because it was identified as the target of a series of anti-inflammatory pyridinyl-imidazole compounds and as reactivating kinase (RK) because it phosphorylated and activated MK2^{3–5}. There are four members of the p38 group of MAPKs encoded by four different genes in mammals: p38 α (*MAPK14*, chromosome 6p21.31 in humans), p38 β (*MAPK11*, *SAPK2b*, Chr22q13.33)⁶, p38 γ (*MAPK12*, *ERK6*, *SAPK3*, Chr22q13.33)^{7,8}, and p38 δ (*MAPK13*, *SAPK4*, *Serk4*, Chr6p21.31)^{9,10}. As can be surmised from their chromosomal locations, *MAPK14*/p38 α and *MAPK13*/p38 δ are physically close and separated by just over 15 kb, as are *MAPK12*/p38 β and *MAPK11*/p38 γ , which are separated by less than 2 kb. All the p38s contain a conserved Thr–Gly–Tyr (TGY) dual phosphorylation motif within the kinase activation loop, and both Thr and Tyr phosphorylation are necessary to fully activate the kinase¹¹. However, monophosphorylated p38 α Thr¹⁸⁰ has some kinase activity *in vitro*, but a different substrate specificity, when compared with dual-site phosphorylated p38 α ¹². p38 group members are expressed ubiquitously, but p38 γ and p38 δ are enriched in certain cell types and tissues, such as p38 γ in skeletal muscle and p38 δ in the salivary, pituitary, and adrenal glands¹³. p38 β shares more amino acid sequence identity with p38 α (~70%), while p38 γ and p38 δ share ~60% identity with p38 α . p38 γ and p38 δ also share high sequence homology with cyclin-dependent kinases (CDKs) and are sensitive to some CDK inhibitors¹⁴.

Activation and inactivation of p38

p38 α is involved in the response to almost all stressful stimuli, including LPS, UV light, heat shock, osmotic shock, inflammatory cytokines, T cell receptor ligation, glucose starvation, and oncogene activation^{2,4,5,15–20}. Under certain circumstances, it is also activated upon growth factor stimulation. It should be noted that the activation of p38 in some cases is cell type specific, since an activating stimulus in one cell type may inhibit p38 in other cell types²¹. The study of p38 group members other than p38 α has been less intensive; however, where it has been examined, the other p38s are frequently co-activated with p38 α ²².

Like other MAPK signaling pathways, the activation of all p38s is mediated by a kinase cascade: MAPKKK (MAP3K), which activates MAPKK (MAP2K), which in turn activates MAPK. The MAP2K kinases MKK3 and MKK6 are the major upstream kinases for p38 activation^{23–25}. Although MKK3 and MKK6 phosphorylate most p38 isoforms *in vitro*, selective activation and substrate specificity have been observed *in vivo*²⁶. MKK4 has also been reported to phosphorylate p38 α and p38 δ in specific

cell types⁹. A number of MAP3Ks have been reported to participate in p38 activation including TAK1²⁷, ASK1²⁸, DLK²⁹, and MEKK4^{29,30}. Low-molecular-weight GTP-binding proteins in the Rho family, such as Rac1 and Cdc42, can activate p38 through binding to MEK1 or MLK1, which function as upstream activators of MAP3K^{31,32}.

p38 α can also be activated by MAP2K-independent mechanisms. TAB1 (TAK1-binding protein 1) directly interacts with p38 α and can promote trans autophosphorylation on Thr¹⁸⁰ and Tyr¹⁸² and thus full activation of p38 α ³³. A subsequent study revealed that autophosphorylation of Thr¹⁸⁰ and Tyr¹⁸² requires a conserved Thr¹⁸⁵ residue³⁴. TAB1-dependent p38 α activation has been implicated in ischemic myocardial injury and T cell anergy^{35,36}. TAB1 is also claimed to play a role in Sestrin-mediated p38 α activation¹². Another MAP2K-independent activation is mediated by ZAP70 after T cell receptor ligation. ZAP70 can directly phosphorylate p38 α /p38 β on Tyr^{323/18}, leading to autophosphorylation on Thr¹⁸⁰, one of the dual phosphorylation sites. As discussed, mono-Thr¹⁸⁰ phosphorylated p38 still has some kinase activity³⁷, and loss of ZAP70-mediated p38 activation in p38 α β ^{Y323F} double knock-in mice reduces autoimmunity and inflammation in several autoimmune disease models^{38–40}. Interestingly, p38 α also phosphorylates ZAP70, resulting in a decrease in the size and persistence of the T cell receptor signaling complex, and therefore acts as a feedback regulator of ZAP70⁴¹.

Conversely, de-phosphorylation of both threonine and tyrosine residues in the activation loop inactivates MAPKs, and this is mainly carried out by dual-specificity phosphatases of the MAPK phosphatase (MKP)/dual specificity phosphatase (DUSP) family⁴². Although several MKPs have been reported to dephosphorylate p38 α , MKP1/DUSP1, MKP5/DUSP10, MKP8/DUSP26, and DUSP8 are more potent inhibitors of p38 α and JNK than ERK⁴³. A recent report showed that DUSP12 is also a p38 α phosphatase⁴⁴. While there are a number of p38 α DUSPs, no DUSP for p38 γ or p38 δ has been reported, and these two p38s are resistant to several known p38 α MKPs such as MKP1, 3, 5, and 7⁴⁵. p38 α -dependent upregulation of MKP1 was reported and is believed to be part of a negative feedback loop of p38 α activation⁴⁶. Other types of phosphatases have also been reported to target p38 MAPKs, such as CacyBP/SIP⁴⁷, Wip1⁴⁸, and PP2C^{49,50}. The substrate specificity between p38 and phosphatases and the related physiological functions *in vivo* still need further investigation. p38 γ has also been reported to be degraded by a p38/JNK/ubiquitin-proteasome-dependent pathway, which represents an additional mechanism by which p38 kinases may cross regulate each other⁵¹. Yet other ways of regulating p38 are suggested from studies in *Caenorhabditis elegans*, where a genetic screen for resistance against bacterial infection identified RIOK-1, an atypical serine kinase and human RIO kinase homolog, as a suppressor of the p38 pathway⁵². As RIOK-1 is a transcriptional target of the p38 pathway in *C. elegans*, this suggests that RIOK-1 is part of a negative feedback loop. A brief summary of the p38 pathway is shown in [Figure 1](#).

Figure 1. A diagram of the p38 pathway. MKP, mitogen-activated protein kinase phosphatase; TAB1, TAK1-binding protein 1; Tyr, tyrosine.

Downstream substrates of p38

Protein kinases

The p38 MAPK cascade does not end at p38. Members of the MAPK-activated protein kinase (MAPKAPK) family such as MK2, MK3, and MK5 (PRAK) are all p38 substrates^{3,4,53–55}. The MKs have a broad range of substrates that extend the range of functions regulated by p38 kinases. Mitogen- and stress-activated protein kinase-1/2 (MSK1/2), which are important for CREB activation and chromosome remodeling, have also been identified as substrates of p38 α ⁵⁶. MNK1/2, kinases that phosphorylate the eukaryotic initiation factor-4e (eIF-4E), are phosphorylated by p38 α ^{57,58}. p38 α has also been reported to inactivate murine GSK3 β by phosphorylating Ser³⁸⁹, and since GSK3 β is required for the continuous degradation of β -catenin in the Wnt signaling pathway, this can lead to an accumulation of β -catenin^{59,60}. It was also reported that p38 δ negatively regulates insulin secretion by catalyzing an inhibitory phosphorylation of PKD1⁶¹. A number of p38 protein kinase substrates are summarized in Table 1.

Transcription factors

p38 targets a large number of transcription factors, including myocyte-specific enhancer factor 2 (MEF2) family members, cyclic AMP-dependent transcription factor 1, 2, and 6 (ATF-1/2/6), CHOP (growth arrest and DNA damage inducible gene 153, or GADD153), p53, C/EBP β , MITF1, DDIT3, ELK1/4, NFAT, and STAT1/4. p38 phosphorylation of transcription factors predominantly leads to enhanced transcriptional activity. However, in some cases, it represses transcription, and this is summarized in Table 2. Transcription factor phosphorylation by p38 is often stimulus and cell type dependent and plays a role in the cellular response to inflammation, DNA damage, metabolic stress, and many other stresses^{62–76}. The effects of p38 on transcription seem to constitute the major part of p38's responses to stress stimuli.

Transcriptional regulators

A large number of transcriptional regulators, including epigenetic enzymes, are substrates of p38, and these are summarized

Table 1. Substrates of p38 group members – kinases.

Substrate	Kinase	Function	References
MAPKAPK2 (MK2)	p38 α , p38 β , p38 γ , p38 δ	Activates the kinase substrate	Freshney NW <i>et al.</i> , <i>Cell</i> , 1994 ⁴ Rouse J <i>et al.</i> , <i>Cell</i> , 1994 ³
MAPKAPK3 (MK3)	p38 α , p38 β , p38 γ , p38 δ	Activates the kinase substrate	McLaughlin MM <i>et al.</i> , <i>J Biol Chem</i> , 1996 ⁵⁴
MNK1/2	p38 α	Activates the kinase substrate	Fukunaga R <i>et al.</i> , <i>EMBO J</i> , 1997 ⁵⁸ Waskiewicz AJ <i>et al.</i> , <i>EMBO J</i> , 1997 ⁵⁷
MSK1/2	p38 α	Activates the kinase substrate	Deak M <i>et al.</i> , <i>EMBO J</i> , 1998 ⁵⁶ Pierrat B <i>et al.</i> , <i>J Biol Chem</i> , 1998 ⁷⁷
PAK6	p38 α	Activates the kinase substrate	Kaur R <i>et al.</i> , <i>J Biol Chem</i> , 2005 ⁷⁸
PIP4Kb	p38 α	Inactivates the kinase substrate	Jones DR <i>et al.</i> , <i>Mol Cell</i> , 2006 ⁷⁹
RPAK (MK5)	p38 α , p38 β	Activates the kinase substrate	New L <i>et al.</i> , <i>EMBO J</i> , 1998 ⁵⁵
PKC ϵ	p38 α , p38 β	Completes cytokinesis	Saurin AT <i>et al.</i> , <i>Nat Cell Biol</i> , 2008 ⁸⁰
GSK3 β	p38 α	Inactivates the kinase substrate, activates Wnt pathway.	Bikkavilli RK <i>et al.</i> , <i>J Cell Sci</i> , 2008 ⁶⁰ Thornton TM <i>et al.</i> , <i>Science</i> , 2008 ⁵⁹

GSK3 β , glycogen synthase kinase 3 beta; MAPKAPK, mitogen-activated protein kinase activated protein kinase; MSK1/2, mitogen- and stress-activated protein kinase; PAK6, p21-activated kinase 6; PIP4Kb, phosphatidylinositol 5 phosphate 4-kinase; PKC ϵ , protein kinase C epsilon type.

Table 2. Substrates of p38 group members – transcription factors.

Substrate	Kinase	Function	References
ATF2	p38 α , p38 β , p38 γ , p38 δ	Enhances transcriptional activity	Cuenda A <i>et al.</i> , <i>EMBO J</i> , 1997 ⁸¹ Jiang Y <i>et al.</i> , <i>J Biol Chem</i> , 1997 ⁹
C/EBP α	p38 α	Enhances transcriptional activity	Qiao L <i>et al.</i> , <i>J Biol Chem</i> , 2006 ⁸²
C/EBP β	p38 α	Enhances transcriptional activity	Engelman JA <i>et al.</i> , <i>J Biol Chem</i> , 1998 ⁸³
C/EBP ϵ	p38 α	Enhances transcriptional activity	Williamson EA <i>et al.</i> , <i>Blood</i> , 2005 ⁸⁴
CHOP	p38 α , p38 β	Enhances transcriptional activity	Wang XZ <i>et al.</i> , <i>Science</i> , 1996 ⁶⁸
E2F4	p38 α	Enhances transcriptional activity	Morillo SM <i>et al.</i> , <i>Mol Cell Biol</i> , 2012 ⁸⁵
Elk-1	p38 α	Enhances transcriptional activity in specific cell types	Janknecht R <i>et al.</i> , <i>EMBO J</i> , 1997 ⁶⁷ Whitmarsh AJ <i>et al.</i> , <i>Mol Cell Biol</i> , 1997 ⁶⁶
ER α	p38 α	Enhances nuclear localization and transcriptional activity	Lee H <i>et al.</i> , <i>Mol Cell Biol</i> , 2002 ⁸⁶
Fos	p38 α , p38 β , p38 γ , p38 δ	Enhances transcriptional activity	Tanos T <i>et al.</i> , <i>J Biol Chem</i> , 2005 ⁸⁷
FOXO3a	p38 α	Enhances nuclear relocalization	Ho KK <i>et al.</i> , <i>J Biol Chem</i> , 2012 ⁸⁸
GR	p38 α	Enhances transcriptional activity	Miller AL <i>et al.</i> , <i>Mol Endocrinol</i> , 2005 ⁸⁹
IUF1	p38 α , p38 β	Enhances transcriptional activity	Macfarlane WM <i>et al.</i> , <i>J Biol Chem</i> , 1997 ⁹⁰
JDP2	p38 α	N/D	Katz S <i>et al.</i> , <i>Biochem J</i> , 2002 ⁹¹
c-JUN	p38 α , p38 β , p38 γ	Enhances transcriptional activity	Humar M <i>et al.</i> , <i>Int J Biochem Cell Biol</i> , 2007 ⁹²

Substrate	Kinase	Function	References
MafA	p38 α , p38 β , p38 γ , p38 δ	Enhances transcriptional activity	Sii-Felice K <i>et al.</i> , <i>FEBS Lett</i> , 2005 ⁹³
MEF2A	p38 α , p38 β , p38 δ	Enhances transcriptional activity	Zhao M <i>et al.</i> , <i>Mol Cell Biol</i> , 1999 ⁹⁴
MEF2C	p38 α , p38 β , p38 γ , p38 δ	Enhances transcriptional activity	Han J <i>et al.</i> , <i>Nature</i> , 1997 ⁹²
MEF2D	p38 α	Enhances recruitment of Ash2L to muscle-specific promoters	Zhao M <i>et al.</i> , <i>Mol Cell Biol</i> , 1999 ⁹⁴ Rampalli S <i>et al.</i> , <i>Nat Struct Mol Biol</i> , 2007 ⁷³
MITF	p38 α	Enhances transcriptional activity	Mansky KC <i>et al.</i> , <i>J Biol Chem</i> , 2002 ⁹⁵
MRF4	p38 α	Represses transcriptional activity	Suelves M <i>et al.</i> , <i>EMBO J</i> , 2004 ⁹⁶
NFATc1	p38 α	Enhances transcriptional activity and interaction with PU.1	Matsumoto M <i>et al.</i> , <i>J Biol Chem</i> , 2004 ⁹⁷
NFATc4	p38 α , p38 β , p38 γ	Represses nuclear localization and transcriptional activity	Yang TT <i>et al.</i> , <i>Mol Cell Biol</i> , 2002 ⁹⁸
NR4A	p38 α	Enhances transcriptional activity	Sekine Y <i>et al.</i> , <i>J Cell Sci</i> , 2011 ⁹⁹
Nur77	p38 α	Disrupts interaction with p65 and represses transcriptional activity	Li L <i>et al.</i> , <i>Nat Chem Biol</i> , 2015 ¹⁰⁰
Osterix	p38 α	Enhances recruitment of coactivators	Ortuño MJ <i>et al.</i> , <i>J Biol Chem</i> , 2010 ¹⁰¹
p53	p38 α	Increases protein stability and apoptosis	Bulavin DV <i>et al.</i> , <i>EMBO J</i> , 1999 ⁹⁹
Pax6	p38 α	Enhances transcriptional activity	Mikkola I <i>et al.</i> , <i>J Biol Chem</i> , 1999 ¹⁰²
PPAR α	p38 α	Enhances transcriptional activity	Barger PM <i>et al.</i> , <i>J Biol Chem</i> , 2001 ¹⁰³
SAP1	p38 α , p38 β , p38 γ , p38 δ	Enhances transcriptional activity	Janknecht R <i>et al.</i> , <i>EMBO J</i> , 1997 ⁶⁷
Smad3	p38 α	Enhances nuclear translocation	Hayes SA <i>et al.</i> , <i>Oncogene</i> , 2003 ¹⁰⁴
Snail	p38 α	Increases protein stability and transcriptional activity	Ryu KJ <i>et al.</i> , <i>Cancer Res</i> , 2019 ¹⁰⁵
STAT1	p38 α , p38 β	Enhances transcriptional activity	Kovarik P <i>et al.</i> , <i>Proc Natl Acad Sci U S A</i> , 1999 ¹⁰⁶
STAT4	p38 α	Enhances transcriptional activity	Visconti R <i>et al.</i> , <i>Blood</i> , 2000 ¹⁰⁷
TEAD4	p38 α	Enhances cytoplasmic translocation and suppresses transcriptional activity	Lin KC <i>et al.</i> , <i>Nat Cell Biol</i> , 2017 ⁷⁶
Twist1	p38 α	Increases protein stability and transcriptional activity	Hong J <i>et al.</i> , <i>Cancer Res</i> , 2011 ¹⁰⁸
USF1	p38 α	Enhances transcriptional activity	Galibert MD <i>et al.</i> , <i>EMBO J</i> , 2001 ⁷¹
Xbp1s	p38 α	Enhances nuclear translocation and transcriptional activity	Lee J <i>et al.</i> , <i>Nat Med</i> , 2011 ⁷⁵

ATF2, activating transcription factor 2; C/EBP, CCAAT/enhancer binding protein; CHOP, CCAAT/enhancer-binding protein homologous protein; ER, estrogen receptor; GR, glucocorticoid receptor; IUF1, insulin upstream factor 1; JDP2, Jun dimerization protein 2; MEF, myocyte-specific enhancer factor; MITF, microphthalmia transcription factor; MRF, muscle regulatory factor; NFAT, nuclear factor of activated T cells; Pax6, paired box 6; PPAR α , peroxisome proliferator-activated receptor alpha; TEAD4, TEA domain family transcription factor 4; USF1, upstream transcription factor 1; Xbp1s, spliced form of X-box binding protein 1.

in **Table 3**. The SWI–SNF complex subunit BAF60 is phosphorylated and inactivated by p38 during skeletal myogenesis^{109,110}, and EZH2, the catalytic component of the Polycomb Repressive Complex 2 (PRC2), was also found to be phosphorylated by p38, particularly in ER-negative breast cancer samples¹¹¹. Besides its transcriptional function, dATF-2 is also involved in heterochromatin formation, and stress-induced phosphorylation of dATF-2 by p38 disrupts heterochromatin in *Drosophila*¹¹².

Other substrates

Given the wide range of responses that p38 is involved in, it is not surprising that many p38 substrates cannot be so easily categorized into groups, and these miscellaneous substrates are summarized in **Table 4**. Some of them are involved in metabolism such as Raptor phosphorylation by p38 β , which enhances mTORC1 activity in response to arsenite-stress¹¹³, and DEPTOR (mTOR-inhibitory protein) phosphorylation by p38 γ and p38 δ , leading to its degradation and mTOR hyperactivation¹¹⁴. p38 α phosphorylation of Tip60 at Thr¹⁵⁸ promotes senescence and DNA-damage-induced apoptosis^{115,116}. Some p38 substrates are

cell death regulators. In the ER stress response, p38 α locates to the lysosome and phosphorylates the chaperone-mediated autophagy (CMA) receptor LAMP2A, leading to activation of CMA and thus protecting cells from ER stress-induced death¹¹⁷.

Biological functions of the p38 pathway

Embryo development

p38 α is required for embryo development, since the mouse *Mapk14*^{-/-} embryo dies between embryonic days (E) 10.5 and 12.5^{118–121}. Mutant mice with a single Thr¹⁸⁰ to Ala mutation or with the double T180A Y182F mutation are also embryonic lethal^{122,123}. Surprisingly, given the importance of the dual phosphorylation for complete p38 activation, substitution of Tyr¹⁸² with Phe results in mice that have reduced p38 signaling but are nevertheless viable¹²³, although this is consistent with previous studies showing that the p38 phosphorylated on Thr¹⁸⁰ alone retains some activity *in vitro*³⁷. Histological analysis demonstrates that p38 α is required for placental angiogenesis, but not embryonic cardiovascular development, and tetraploid rescue of the placental defect in *Mapk14*^{-/-} embryos confirmed that p38 α is

Table 3. Substrates of p38 group members – transcriptional regulators.

	Substrate	Kinase	Function	References
Chromatin remodeling regulators	BAF60c	p38 α , p38 β	Activates transcription of MyoD-target genes	Simone C <i>et al.</i> , <i>Nat Genet</i> , 2004 ¹⁰⁹ Forcales SV <i>et al.</i> , <i>EMBO J</i> , 2012 ¹¹⁰
	RNF2	p38 α	Modulates gene expression and histone 2B acetylation	Rao PS <i>et al.</i> , <i>Proteomics</i> , 2009 ¹²⁴
	EZH2	p38 α	Promotes cytoplasmic localization	Anwar T <i>et al.</i> , <i>Nat Commun</i> , 2018 ¹¹¹
	dAFF2	p38 α , p38 β	Disrupts heterochromatin formation	Seong K-H <i>et al.</i> , <i>Cell</i> , 2011 ¹¹²
Other regulators	CRTC2	p38 α	Enhances nucleocytoplasmic transport and represses transcription activity	Ma H <i>et al.</i> , <i>Mol Cell Biol</i> , 2019 ¹²⁵
	E47	p38 α , p38 β	Enhances the formation of MyoD/E47 heterodimers	Page JL <i>et al.</i> , <i>J Biol Chem</i> , 2004 ¹²⁶ Lluís F <i>et al.</i> , <i>EMBO J</i> , 2005 ¹²⁷
	HBP1	p38 α	Increases protein stability and represses transcription	Xiu M <i>et al.</i> , <i>Biol</i> , 2003 ¹²⁸
	p18(Hamlet)	p38 α , p38 β	Increases protein stability and enhances transcription	Cuadrado A <i>et al.</i> , <i>EMBO J</i> , 2007 ¹²⁹
	PGC-1 α	p38 α , p38 β	Increases protein stability and enhances transcription	Puigserver P <i>et al.</i> , <i>Mol Cell</i> , 2001 ¹³⁰
	Rb1	p38 α , p38 γ	Induces Rb degradation and cell death; suppresses Rb activity and promotes the G0-to-G1 transition	Delston RB <i>et al.</i> , <i>Oncogene</i> , 2011 ¹³¹ Tomás-Loba A <i>et al.</i> , <i>Nature</i> , 2019 ¹⁴
	SRC-3	p38 α	Induces SRC-3 degradation and suppresses RAR α -dependent transcription	Gianni M <i>et al.</i> , <i>EMBO J</i> , 2006 ¹³²

CRTC2, CREB-regulated transcription coactivator 2; HBP1, HMG-box transcription factor 1; PGC-1 α , peroxisome proliferator-activated receptor gamma co-activator 1 alpha; RAR, retinoic acid receptor; RNF2, ring finger protein 2.

Table 4. Substrates of p38 group members – others.

	Substrate	Kinase	Function	References
Cell-cycle regulators	Cdc25A	p38 α	Increases protein stability	Goloudina A <i>et al.</i> , <i>Cell Cycle</i> , 2003 ¹³³
	Cdc25B	p38 α	Increases protein stability	Lemaire M <i>et al.</i> , <i>Cell Cycle</i> , 2006 ¹³⁴
	Cyclin D1	p38 α	Causes ubiquitination and degradation of cyclin D1	Casanovas O <i>et al.</i> , <i>J Biol Chem</i> , 2000 ¹³⁵
	Cyclin D3	p38 α , p38 β p38 γ , p38 δ	Causes ubiquitination and degradation of cyclin D3	Casanovas O <i>et al.</i> , <i>Oncogene</i> , 2004 ¹³⁶
	p57kip2	p38 α	Enhances interaction with CDKs and inhibits CDKs	Joaquin M <i>et al.</i> , <i>EMBO J</i> , 2012 ¹³⁷
Cell-death regulators	Bax	p38 α	Prevents Bcl-2–Bax heterodimer formation, enhances apoptosis	Min H <i>et al.</i> , <i>Mol Carcinog</i> , 2012 ¹³⁸
	BimEL	p38 α	Enhances apoptosis	Cai B <i>et al.</i> , <i>J Biol Chem</i> , 2006 ¹³⁹
	Caspase-3	p38 α	Inhibits caspase-3 activity and apoptosis	Alvarado-Kristensson M <i>et al.</i> , <i>J Exp Med</i> , 2004 ¹⁴⁰
	Caspase-8	p38 α	Inhibits caspase-8 activity and apoptosis	Alvarado-Kristensson M <i>et al.</i> , <i>J Exp Med</i> , 2004 ¹⁴⁰
	Caspase-9	p38 α	Inhibits caspase-9 activity and apoptosis	Seifert A <i>et al.</i> , <i>Cell Signal</i> , 2009 ¹⁴¹
DNA/RNA binding proteins	Cdt1	p38 α , p38 β	Increases protein stability	Chandrasekaran S <i>et al.</i> , <i>Mol Cell Biol</i> , 2011 ¹⁴²
	Drosha	p38 α	Enhances nuclear export and degradation	Yang Q <i>et al.</i> , <i>Mol Cell</i> , 2015 ¹⁴³
	FBP2	p38 α	Promotes prothrombin mRNA 3' end processing	Danckwardt S <i>et al.</i> , <i>Mol Cell</i> , 2011 ¹⁴⁴
	FBP3	p38 α	Promotes prothrombin mRNA 3' end processing	Danckwardt S <i>et al.</i> , <i>Mol Cell</i> , 2011 ¹⁴⁴
	H2AX	p38 α , p38 β	Promotes serum starvation-induced apoptosis	Lu C <i>et al.</i> , <i>FEBS Lett</i> , 2008 ¹⁴⁵
	H3	p38 α	N/D	Zhong SP <i>et al.</i> , <i>J Biol Chem</i> , 2000 ¹⁴⁶
	HuR	p38 α , p38 β	Enhances cytoplasmic accumulation and increases mRNA stability	Lafarga V <i>et al.</i> , <i>Mol Cell Biol</i> , 2009 ¹⁴⁷
	KSRP	p38 α , p38 β	Prevents KSRP-mediated ARE-directed mRNA decay	Briata P <i>et al.</i> , <i>Mol Cell</i> , 2005 ¹⁴⁸
	Rps27	p38 α	N/D	Knight JD <i>et al.</i> , <i>Skelet Muscle</i> , 2012 ¹⁴⁹
	SPF45	p38 α	Inhibits Fas alternative splicing (exon 6 exclusion)	Al-Ayoubi AM <i>et al.</i> , <i>Mol Cell Biol</i> , 2012 ¹⁵⁰
Endocytosis regulators	EEA1	p38 α	Promotes recruitment to endocytic membranes and enhances MOR endocytosis	Macé G <i>et al.</i> , <i>EMBO J</i> , 2005 ¹⁵¹
	Rabenosyn-5	p38 α	Promotes recruitment to endocytic membranes and enhances MOR endocytosis	Macé G <i>et al.</i> , <i>EMBO J</i> , 2005 ¹⁵¹
	GDI-2	p38 α	Enhances GDI:Rab5 complex formation and modulates endocytosis	Cavalli V <i>et al.</i> , <i>Mol Cell</i> , 2001 ¹⁵²
MAPK pathway regulator	JIP4	p38 α	Enhances p38 activity	Kelkar N <i>et al.</i> , <i>Mol Cell Biol</i> , 2005 ¹⁵³
	Tip60	p38 α	Enhances the pro-senescent function of Tip60	Zheng H <i>et al.</i> , <i>Mol Cell</i> , 2013 ¹¹⁵
	TAB1	p38 α	Inhibits TAK1 activity	Cheung PC <i>et al.</i> , <i>EMBO J</i> , 2003 ¹⁵⁴
	TAB3	p38 α	Inhibits TAK1 activity	Mendoza H <i>et al.</i> , <i>Biochem J</i> , 2008 ¹⁵⁵
	FRS2	p38 α	Downregulates FGF1-induced signaling	Zakrzewska M <i>et al.</i> , <i>Int J Mol Sci</i> , 2019 ¹⁵⁶

	Substrate	Kinase	Function	References
Membrane proteins	EGFR	p38 α	Induces EGFR internalization	Winograd-Katz SE <i>et al.</i> , <i>Oncogene</i> , 2006 ¹⁵⁷
	FGFR1	p38 α	Regulates translocation of exogenous FGF1 into the cytosol/nucleus	Sørensen V <i>et al.</i> , <i>Mol Cell Biol</i> , 2008 ¹⁵⁸
	Nav1.6	p38 α	Promotes interaction with NEDD-4 and protein degradation	Gasser A <i>et al.</i> , <i>J Biol Chem</i> , 2010 ¹⁵⁹
	NHE1	p38 α	Induces intracellular alkalization	Khaled AR <i>et al.</i> , <i>Mol Cell Biol</i> , 2001 ¹⁶⁰
	PLA2	p38 α	N/D	Börsch-Haubold AG <i>et al.</i> , <i>J Biol Chem</i> , 1998 ¹⁶¹
	TACE	p38 α , p38 β	Increases TACE-mediated ectodomain shedding and TGF- α family ligand release	Xu P <i>et al.</i> , <i>Mol Cell</i> , 2010 ¹⁶²
	ZAP70	p38 α	Phosphorylation of ZAP70 increases stability of T cell receptor	Giardino Torchia ML <i>et al.</i> , <i>Proc Natl Acad Sci U S A</i> , 2018 ⁴¹
Structure proteins	Caldesmon	p38 α	N/D	Hedges JC <i>et al.</i> , <i>Am J Physiol</i> , 1998 ¹⁶³
	Hsp27	p38 α	N/D	Knight JD <i>et al.</i> , <i>Skelet Muscle</i> , 2012 ¹⁴⁹
	Keratin 8	p38 α	Regulates cellular keratin filament reorganization	Ku NO <i>et al.</i> , <i>J Biol Chem</i> , 2002 ¹⁶⁴
	Lamin B1	p38 α	Enhances lamin B1 accumulation	Barascu A <i>et al.</i> , <i>EMBO J</i> , 2012 ¹⁶⁵
	Paxillin	p38 α	Required for NGF-induced neurite extension of PC-12 cells	Huang C <i>et al.</i> , <i>J Cell Biol</i> , 2004 ¹⁶⁶
	Stathmin	p38 δ	N/D	Parker CG <i>et al.</i> , <i>Biochem Biophys Res Commun</i> , 1998 ¹⁶⁷
	SAP97	p38 γ	Modulating the association of this protein with other cytoskeleton proteins	Sabio G <i>et al.</i> , <i>EMBO J</i> , 2005 ¹⁶⁸
	Tau	p38 α , p38 γ , p38 δ	Enhances formation of paired helical filaments Inhibits amyloid- β toxicity in Alzheimer's mice	Reynolds CH <i>et al.</i> , <i>J Neurochem</i> , 1997 ¹⁶⁹ Ittner A <i>et al.</i> , <i>Science</i> , 2016 ¹⁷⁰
	Tensin1	p38 α	Regulates the binding specificity of tensin1 to different proteins	Hall EH <i>et al.</i> , <i>Mol Cell Proteomics</i> , 2010 ¹⁷¹
Others	DEPTOR	p38 γ , p38 δ	Enhances degradation and mTOR hyperactivation	González-Terán B <i>et al.</i> , <i>Nat Commun</i> , 2016 ¹¹⁴
	GS	p38 β	Required for subsequent phosphorylation to inhibit enzyme activity	Kuma Y <i>et al.</i> , <i>Biochem J</i> , 2004 ¹⁷²
	LAMP2A	p38 α	Activates chaperone-mediated autophagy	Li W <i>et al.</i> , <i>Nat Commun</i> , 2017 ¹¹⁷
	Parkin	p38 α	Decreases its interaction with PINK1 and suppresses mitophagy	Chen J <i>et al.</i> , <i>Cell Death Dis</i> , 2018 ¹⁷³
	p47 ^{phox}	p38 α	Promotes NADPH oxidase activation and superoxide production	Makni-Maalej K <i>et al.</i> , <i>J Immunol</i> , 2012 ¹⁷⁴
	p62	p38 γ , p38 δ	Enhances mTORC1 activity	Linares JF <i>et al.</i> , <i>Cell Rep</i> , 2015 ¹⁷⁵ Koh A <i>et al.</i> , <i>Cell</i> , 2018 ¹⁷⁶
	Raptor	p38 β	Enhances mTORC1 activity in response to arsenite stress	Wu X-N <i>et al.</i> , <i>J Biol Chem</i> , 2011 ¹¹³
	Rpn2	p38 α	Inhibits proteasome activity	Lee SH <i>et al.</i> , <i>J Biol Chem</i> , 2010 ¹⁷⁷
	Siah2	p38 α	Increases Siah2-mediated degradation of PHD3	Khurana A <i>et al.</i> , <i>J Biol Chem</i> , 2006 ¹⁷⁸

CDK, cyclin-dependent kinase; EGFR, epidermal growth factor receptor; FBP1, far upstream binding protein; FGF1, fibroblast growth factor 1; FGFR1, fibroblast growth factor receptor 1; FRS2, fibroblast growth factor receptor substrate 2; GDI, GDP dissociation inhibitor; KSRP, hnRNPK-homology type splicing regulatory protein; MAPK, mitogen-activated protein kinase; mTORC1, mammalian target of rapamycin complex 1; NADPH, nicotinamide adenine dinucleotide phosphate; NGF, nerve growth factor; NHE1, Na⁺/H⁺ exchanger isoform 1; PHD3, prolyl hydroxylase 3; PLA2, phospholipase A2; SAP97, synapse-associated protein 97; TAB, transforming growth factor- β -activated protein kinase-1-binding protein; TACE, tumor necrosis factor- α -converting enzyme; TAK1, transforming growth factor β -activated kinase 1; TGF, transforming growth factor.

essential for extraembryonic development^{120,121}. Given the important role that p38 and MK2 plays in regulating TNF-induced cell death^{179–182}, it is intriguing that the *Mapk14*^{−/−} embryonic lethal phenotype is very similar to that observed in other mice with defects in the TNF death pathway. Caspase-8, FADD, and cFLIP knock-out mice also die at E10.5, and this is due to TNF-dependent endothelial cell death and disruption of the vasculature in the yolk sac^{183,184}. Other p38 isoforms are not necessary for embryo development, but p38 α and p38 β have overlapping functions, as *Mapk14*^{loxP/loxP}*Mapk11*^{−/−}*Sox2-Cre* embryos die before E16.5 with spina bifida that correlates with neural hyperproliferation and increased apoptosis in the liver, which was not observed in *Mapk14*^{Δ/Δ}*Sox2-Cre* embryos¹⁸⁵. Remarkably, p38 α appears to have a very specific function during embryogenesis because when p38 α was replaced by p38 β in the *Mapk14* chromosomal locus, which thereby placed p38 β under the control of the endogenous p38 α promoter, it was unable to rescue the embryonic lethality induced by loss of p38 α ¹⁸⁵.

Immune responses

p38 is activated by many inflammatory stimuli, and its activity is important for inflammatory responses. Macrophage-specific deletion of *Mapk14* inhibits inflammatory cytokine production and protects mice from CLP-induced sepsis¹⁸⁶. p38 α controls the production of inflammatory cytokines, such as TNF and IL-6, at many levels. It directly phosphorylates transcription factors, such as MEF2C^{62,186}, and regulators of mRNA stability, such as hnRNP-K-homology (KH) type splicing regulatory protein (KSRP)¹⁸⁷. MEF2C appears to play an anti-inflammatory role in endothelial cells *in vivo*¹⁸⁸. Via MK2/MK3, p38 also upregulates cytokine mRNA transcription by the serum response transcription factor (SRF)¹⁸⁹, and similarly, via MK2/MK3, p38 regulates mRNA stability by phosphorylating and inactivating TTP/Zfp36, a protein that promotes rapid turnover of AU-rich mRNAs, many of which are cytokine mRNAs^{187,190}. p38 activation also induces the expression of inflammatory mediators such as COX-2, MMP9, iNOS, and VCAM-1, which are involved in tissue remodeling and oxidation regulation^{191–194}. The p38 pathway also regulates adaptive immunity. p38 α participates in antigen processing in CD8⁺ cDCs¹⁹⁵, and ZAP70-mediated p38 α / β activation is important for T cell homeostasis and function¹⁸. In B cells, p38 α is important for CD40-induced gene expression and proliferation of B cells¹⁹⁶, and the p38 α –MEF2c axis is believed to be necessary for germinal center B (GCB) cell proliferation and survival^{197,198}. Excessive activation of p38 α has been observed in many inflammatory diseases, such as inflammatory bowel disease (IBD), asthma, rheumatoid arthritis, and steatohepatitis^{199–201}. The other members of the p38 family also play roles in immune responses. For example, p38 γ and p38 δ are required for neutrophil migration to damaged liver in non-alcoholic fatty liver disease²⁰² and inhibition of eukaryotic elongation factor 2 in LPS-induced liver damage²⁰³. p38 δ is required for neutrophil accumulation in acute lung injury²⁰⁴. These observations, and the role that p38s play in TNF production, led to enormous pharmaceutical efforts to develop p38 inhibitors to treat chronic inflammatory diseases. However, unfortunately, these drugs were not efficacious in these diseases²⁰⁵.

Cell cycle

p38 has been implicated in G1 and G2/M phases of the cell cycle in several studies. The addition of activated recombinant p38 α caused mitotic arrest *in vitro*, and an inhibitor of p38 α / β suppressed activation of the checkpoint by nocodazole in NIH3T3 cells²⁰⁶. G1 arrest caused by Cdc42 overexpression is also dependent on p38 α in NIH3T3 cells²⁰⁷. Besides, p38 γ is specially required for gamma-irradiation-induced G2 arrest²⁰⁸. The link between p38 and cell cycle control has been proposed through the regulation of several p38 substrates. Both p38 α and p38 γ regulate cell cycle progression via Rb but in opposite directions^{14,209}. HBP1 represses the expression of cell cycle regulatory genes during cell cycle arrest in a p38-dependent manner²¹⁰; p53 and p21 activation by p38 α prevented G1 progression through blockade of CDK activity^{211,212}. The p38 pathway is also involved in cell cycle progress, as it is essential for self-renewal of mouse male germline stem cells²¹³ and its regulation of G1-length plays a role in cell size uniformity²¹⁴.

Cell differentiation

Participation of p38 in cell differentiation has been reported in certain cell types. p38 α activity is essential for neuronal differentiation in PC-12 cells and EPO-induced differentiation in SKT6 cells^{20,215}. Treatment of 3T3-L1 fibroblasts with specific p38 α / β inhibitors prevents their differentiation into adipocytes by reducing C/EBP β phosphorylation⁸³, and p38 α -dependent phosphorylation of MEF2C and BAF60 is critical for myogenic differentiation^{110,216}. Intestinal epithelial cell-specific deletion of p38 α also influences goblet cell differentiation in a Notch-dependent manner²⁰⁰.

Cell metabolism

p38 group members participate in many cellular events related to metabolism. The p38 β –PRAK axis specifically phosphorylates Rheb and suppresses mTORC1 activity under energy depletion conditions²². DEPTOR, an inhibitor of mTORC, can be phosphorylated by p38 γ and p38 δ , leading to its degradation¹²³. Meanwhile, p38 δ directly phosphorylates p62 to enhance mTORC1 activity in response to amino acids¹⁷⁵. In brown adipocytes, p38 α functions as a central mediator in β -adrenergic-induced UCP1 expression^{217,218}, while in white adipocytes, p38 α inactivation leads to elevated white-to-beige adipocyte reprogramming and resistance to diet-induced obesity^{219,220}. In hepatocytes, p38 α controls lipolysis and protects against nutritional steatohepatitis. Thus, mice with hepatocyte-specific loss of p38 α developed more severe steatohepatitis than wild type mice when fed high-fat or -cholesterol diets. Intriguingly, macrophage specific deletion of p38 had the opposite effect in the same high-fat diets and resulted in less steatohepatitis than in wild type mice, which probably reflects the inflammatory role of p38 in macrophages¹⁹⁹. p38 α also directly phosphorylates Xbp1s to enhance its nuclear migration for maintaining glucose homeostasis in obesity⁷⁵. However, p38 α also functions as a negative regulator of AMPK signaling in maintaining gluconeogenesis, and hepatic p38 α could be a drug target for hyperglycemia²²¹. It was also reported that p38 γ directly phosphorylates p62 under imidazole propionate stimulation to promote mTORC1 activity in hepatocytes¹⁷⁶.

Interestingly, AMPK also triggers the recruitment of p38 α to scaffold protein TAB1 for p38 α autoactivation in human T cells²²².

Cell senescence

p38 α appears to play a pivotal role in senescence. Constitutive activation of the p38 pathway by active MKK3 or MKK6 induces senescence in several cell types^{223,224}, and p38 α activity is responsible for senescence induced by multiple stimuli, such as telomere shortening^{225,226}, H₂O₂ exposure^{227,228}, and chronic oncogene activation^{19,223,229}. p38 α / β -specific inhibitors have been successfully used to prevent cellular senescence in cultivated human corneal endothelial cells²³⁰. Since cellular senescence is considered a defense strategy against oncogene activation, the p38 pathway plays important roles in tumorigenesis²³¹. Meanwhile, p38 α activity is important for senescence-associated secretory phenotype (SASP), and its inhibition markedly reduces the secretion of most SASP factors, suggesting multiple roles for the p38 pathway in senescence^{232–235}.

Cell survival and death

The role of the p38 pathway in cell fate is cell type and stimulus dependent. For example, p38 α becomes activated upon NGF withdrawal in PC-12 cells, and p38 α activated by overexpression of MKK3 induced apoptosis in NGF differentiated PC-12 cells²¹¹. Similarly, inhibition of p38 with PD169316 blocked NGF withdrawal-induced apoptosis in PC-12 cells^{236,237}. The interplay between the p38 pathway and caspases, the central regulators/executors of apoptosis, is complicated because p38 α activity can be elevated in a caspase-dependent manner in death stimulus treated cells^{238,239}, and caspase activity can also be elevated in MKK6E (dominant active form) overexpressed cells^{239,240}. In contrast, inhibition of caspase-8 and caspase-3 by p38 α -mediated phosphorylation in neutrophils was also reported¹⁴⁰. Recent studies show that p38-activated MK2 directly phosphorylates RIPK1 in TNF-treated cells or pathogen-infected cells, limiting TNF-induced cell death^{180–182}. This represents an interesting link between cytokine production induced by TNF and

cell death because TNF-induced MK2/MK3 phosphorylation of tristetraprolin/Zfp36 inactivates it and leads to increased stability of cytokine mRNAs¹⁹⁰. Aberrant p38 α activity is observed in many tumor cells, and inhibition of p38 α / β enhances cell death in these cells^{241,242}.

Perspectives

p38 is one of the most researched of all proteins, let alone kinases, and a search in PubMed for p38 MAPK or p38 kinase returns more than 36,000 publications, which is a higher number than some proteins listed in a review of the “top 10” most studied genes²⁴³. By contrast, searches for the kinases Raf and Src return about 17,000 and 25,000 hits, respectively. In 2018, there were more than 2,000 publications that mention p38, and it is clearly impractical to summarize such a vast amount of literature. As might be surmised from the preceding commentary, the studies are on a wide range of topics; however, the publications are more concentrated in some areas than others. The role of the p38 pathway in cancers (>10,000)^{244–246}, inflammation (>8,000)^{247–249}, and infections (>3,600)^{250,251} was intensively studied. About 1,600 publications include the specific term “p38 inhibitor”. This reflects the previously mentioned enormous interest of the pharmaceutical industry in developing p38 inhibitors to treat chronic inflammatory diseases, such as rheumatoid arthritis. Yet other publications report natural products that can activate or inhibit p38, with the ultimate aim of using them clinically^{252–258}. In 2011, the European Commission approved Esbriet (pirfenidone), which was described as a p38 γ inhibitor, for the treatment of idiopathic pulmonary fibrosis²⁵⁹. However, when this drug was approved by the FDA in 2014 for treating the same disease, it was described as a compound that acts on multiple pathways. In 2008, there were 27 clinical trials listed testing the use of p38 inhibitors in inflammatory disease settings²⁰⁵, while a search today for p38 inhibitors in clinicaltrials.gov returns 44 studies for conditions as diverse as pain, asthma, cognitive impairment, rheumatoid arthritis, cancer, myelodysplastic syndrome, and depression (Table 5). This indicates that there remains clinical interest in targeting the pathway

Table 5. Clinical trials of p38 inhibitors.

Drug	Target	Condition or disease	Status	NCT number
ARRY-371797	p38	Ankylosing spondylitis	Phase 2	NCT00811499
ARRY-371797	p38	Dental pain	Phase 2	NCT00542035 NCT00663767
ARRY-371797	p38	Healthy	Phase 1	NCT00790049
ARRY-371797	p38	LMNA-related dilated cardiomyopathy	Phase 2	NCT02351856 NCT02057341
ARRY-371797	p38	Osteoarthritis of the knee	Phase 2	NCT01366014
ARRY-371798	p38	Rheumatoid arthritis	Phase 1	NCT00729209
ARRY-614	p38 and Tie2	Myelodysplastic syndromes	Phase 1	NCT01496495 NCT00916227
AZD7624	p38	Corticosteroid-resistant asthma	Phase 2	NCT02753764
BIRB 796 BS	p38	Healthy	Phase 1	NCT02211170
BMS-582949	p38 α	Rheumatoid arthritis	Phase 2	NCT00605735

Drug	Target	Condition or disease	Status	NCT number
BMS-582949	p38 α	Vascular diseases (atherosclerosis)	Phase 2	NCT00570752
CHF6297	p38 α	Chronic obstructive pulmonary disease	Phase 1/2	NCT02815488
Losmapimod (GS856553)	p38 α/β	Acute coronary syndrome	Phase 1/2/3	NCT01756495 NCT02145468 NCT00910962
Losmapimod (GS856553)	p38 α/β	Chronic obstructive pulmonary disease	Phase 2	NCT00642148 NCT01541852
Losmapimod (GS856553)	p38 α/β	Depressive disorder, major	Phase 2	NCT00976560 NCT00569062
Losmapimod (GS856553)	p38 α/β	Glomerulosclerosis, focal segmental	Phase 2	NCT02000440
Losmapimod (GS856553)	p38 α/β	Pain, neuropathic	Phase 2	NCT01110057 NCT00969059
LY3007113	p38	Metastatic cancer	Phase 1	NCT01463631
Neflamapimod (VX-745)	p38 α	Alzheimer's disease	Phase 2	NCT03402659 NCT02423200 NCT02423122
Neflamapimod (VX-745)	p38 α	Dementia with Lewy bodies	Recruiting	NCT04001517
P38 inhibitor (4)	p38	Rheumatoid arthritis	Phase 2	NCT00303563 NCT00316771
PF-03715455	p38 α	Asthma	Phase 2	NCT02219048
PF-03715455	p38 α	Chronic obstructive pulmonary disease	Phase 2	NCT02366637
PF-03715455	p38 α	Healthy	Phase 1	NCT01226693
PH-797804	p38 α/β	Rheumatoid arthritis	Phase 2	NCT00383188 NCT00620685
Ralimetinib (LY2228820)	p38 α/β	Adult glioblastoma	Phase 1/2	NCT02364206
Ralimetinib (LY2228820)	p38 α/β	Advanced cancer	Phase 1	NCT01393990
Ralimetinib (LY2228820)	p38 α/β	Epithelial ovarian cancer Fallopian tube cancer Primary peritoneal cancer	Phase 1/2	NCT01663857
Ralimetinib (LY2228820)	p38 α/β	Postmenopausal metastatic breast cancer	Phase 2	NCT02322853
SB-681323	p38	Acute lung injury	Phase 2	NCT00996840
SB-681323	p38	Coronary heart disease	Phase 2	NCT00291902
SB-681323	p38	Chronic obstructive pulmonary disease	Phase 1/2	NCT00564746 NCT00144859
SB-681323	p38	Pain, neuropathic	Phase 2	NCT00390845
SB-681323	p38	Rheumatoid arthritis Inflammation	Phase 1/2	NCT00419809 NCT00439881 NCT00134693
Talmapimod (SCIO-469)	p38 α	Bone marrow diseases Myelodysplastic syndromes Hematologic diseases Bone marrow neoplasms	Phase 2	NCT00113893
Talmapimod (SCIO-469)	p38 α	Multiple myeloma	Phase 2	NCT00095680 NCT00087867
Talmapimod (SCIO-469)	p38 α	Rheumatoid arthritis	Phase 2	NCT00043732 NCT00089921
VX-702	p38 α	Rheumatoid arthritis	Phase 2	NCT00395577 NCT00205478

and that there is therefore a need for more specific inhibitors of each of the p38 group members and more basic research to fully understand how the pathway, especially how each member of the p38 family, is utilized and regulated.

One consequence of the massive pharmaceutical effort over the last 20 years is a large number of very specific, well-tolerated, and readily bioavailable drugs that can enable such basic research. For example, one study using a boutique panel of kinase inhibitors was able to demonstrate that 11 potent and specific p38 inhibitors synergized with Smac-mimetic drugs to kill a subset of AML leukemias, providing the strongest evidence implicating p38 in Smac-mimetic-induced killing¹⁷⁹. Since several of these p38 inhibitors had already been clinically trialed, this presents an opportunity to fast-track such combinations into

the clinic. In our opinion, it is likely that this is where the future of p38 research and p38 inhibitors lies, in revealing the intricate web of inter-connections and inter-dependencies of this core and central regulator of cell stress. We also believe that greater efforts to genetically assess the role of p38 and p38 isoforms in the pathophysiology of inflammatory and other diseases need to be made in order to push forward the clinical application of our burgeoning knowledge.

Acknowledgments

We thank Peipei Zhang and Ye-hsuan Sun (School of Life Sciences, Xiamen University, China) for their help in preparing this manuscript.

References

- Han J, Lee JD, Tobias PS, *et al.*: Endotoxin induces rapid protein tyrosine phosphorylation in 702/3 cells expressing CD14. *J Biol Chem.* 1993; **268**(33): 25009–25014.
[PubMed Abstract](#)
- Han J, Lee JD, Bibbs L, *et al.*: A MAP kinase targeted by endotoxin and hyperosmolarity in mammalian cells. *Science.* 1994; **265**(5173): 808–811.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rouse J, Cohen P, Trigon S, *et al.*: A novel kinase cascade triggered by stress and heat shock that stimulates MAPKAP kinase-2 and phosphorylation of the small heat shock proteins. *Cell.* 1994; **78**(6): 1027–1037.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Freshney NW, Rawlinson L, Guesdon F, *et al.*: Interleukin-1 activates a novel protein kinase cascade that results in the phosphorylation of Hsp27. *Cell.* 1994; **78**(6): 1039–1049.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lee JC, Laydon JT, McDonnell PC, *et al.*: A protein kinase involved in the regulation of inflammatory cytokine biosynthesis. *Nature.* 1994; **372**(6508): 739–746.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jiang Y, Chen C, Li Z, *et al.*: Characterization of the structure and function of a new mitogen-activated protein kinase (p38beta). *J Biol Chem.* 1996; **271**(30): 17920–17926.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lechner C, Zahalka MA, Giot JF, *et al.*: ERK6, a mitogen-activated protein kinase involved in C2C12 myoblast differentiation. *Proc Natl Acad Sci U S A.* 1996; **93**(9): 4355–4359.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Li Z, Jiang Y, Ulevitch RJ, *et al.*: The primary structure of p38 gamma: a new member of p38 group of MAP kinases. *Biochem Biophys Res Commun.* 1996; **228**(2): 334–340.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jiang Y, Gram H, Zhao M, *et al.*: Characterization of the structure and function of the fourth member of p38 group mitogen-activated protein kinases, p38delta. *J Biol Chem.* 1997; **272**(48): 30122–30128.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kumar S, McDonnell PC, Gum RJ, *et al.*: Novel homologues of CSBP/p38 MAP kinase: activation, substrate specificity and sensitivity to inhibition by pyridinyl imidazoles. *Biochem Biophys Res Commun.* 1997; **235**(3): 533–538.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Raingeaud J, Gupta S, Rogers JS, *et al.*: Pro-inflammatory cytokines and environmental stress cause p38 mitogen-activated protein kinase activation by dual phosphorylation on tyrosine and threonine. *J Biol Chem.* 1995; **270**(13): 7420–7426.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lanna A, Gomes DC, Muller-Durovic B, *et al.*: A sestrin-dependent Erk-Jnk-p38 MAPK activation complex inhibits immunity during aging. *Nat Immunol.* 2017; **18**(3): 354–363.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Obata T, Brown GE, Yaffe MB: MAP kinase pathways activated by stress: the p38 MAPK pathway. *Crit Care Med.* 2000; **28**(4 Suppl): N67–77.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tomas-Loba A, Manieri E, Gonzalez-Teran B, *et al.*: p38γ Is Essential for Cell Cycle Progression and Liver Tumorigenesis. *Nature.* 2019; **568**(7753): 557–560.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
- Gong X, Liu A, Ming X, *et al.*: UV-induced interaction between p38 MAPK and p53 serves as a molecular switch in determining cell fate. *FEBS Lett.* 2010; **584**(23): 4711–4716.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zauberman A, Zipori D, Krupsky M, *et al.*: Stress activated protein kinase p38 is involved in IL-6 induced transcriptional activation of STAT3. *Oncogene.* 1999; **18**(26): 3886–3893.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Foltz IN, Lee JC, Young PR, *et al.*: Hemopoietic growth factors with the exception of interleukin-4 activate the p38 mitogen-activated protein kinase pathway. *J Biol Chem.* 1997; **272**(6): 3296–3301.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Salvador JM, Mittelstadt PR, Guszczynski T, *et al.*: Alternative p38 activation pathway mediated by T cell receptor-proximal tyrosine kinases. *Nat Immunol.* 2005; **6**(4): 390–395.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
- Sun P, Yoshizuka N, New L, *et al.*: PRAK is essential for ras-induced senescence and tumor suppression. *Cell.* 2007; **128**(2): 295–308.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Nagata Y, Takahashi N, Davis RJ, *et al.*: Activation of p38 MAP kinase and JNK but not ERK is required for erythropoietin-induced erythroid differentiation. *Blood.* 1998; **92**(6): 1859–1869.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zarubin T, Han J: Activation and signaling of the p38 MAP kinase pathway. *Cell Res.* 2005; **15**(1): 11–18.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zheng M, Wang YH, Wu XN, *et al.*: Inactivation of Rheb by PRAK-mediated phosphorylation is essential for energy-depletion-induced suppression of mTORC1. *Nat Cell Biol.* 2011; **13**(3): 263–272.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Raingeaud J, Whitmarsh AJ, Barrett T, *et al.*: MKK3- and MKK6-regulated gene expression is mediated by the p38 mitogen-activated protein kinase signal transduction pathway. *Mol Cell Biol.* 1996; **16**(3): 1247–1255.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Derijard B, Raingeaud J, Barrett T, *et al.*: Independent human MAP-kinase signal transduction pathways defined by MEK and MKK isoforms. *Science.* 1995; **267**(5198): 682–685.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Han J, Lee JD, Jiang Y, *et al.*: Characterization of the structure and function of a novel MAP kinase kinase (MKK6). *J Biol Chem.* 1996; **271**(6): 2886–2891.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Enslin H, Raingeaud J, Davis RJ: Selective activation of p38 mitogen-activated protein (MAP) kinase isoforms by the MAP kinase kinases MKK3 and MKK6. *J Biol Chem.* 1998; **273**(3): 1741–1748.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Moriguchi T, Kuroyanagi N, Yamaguchi K, *et al.*: A novel kinase cascade mediated by mitogen-activated protein kinase kinase 6 and MKK3. *J Biol Chem.* 1996; **271**(23): 13675–13679.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ichijo H, Nishida E, Irie K, *et al.*: Induction of apoptosis by ASK1, a mammalian MAPKKK that activates SAPK/JNK and p38 signaling pathways. *Science.* 1997; **275**(5296): 90–94.
[PubMed Abstract](#) | [Publisher Full Text](#)

29. Hirai S, Katoh M, Terada M, *et al.*: **MST/MLK2, a member of the mixed lineage kinase family, directly phosphorylates and activates SEK1, an activator of c-Jun N-terminal kinase/stress-activated protein kinase.** *J Biol Chem.* 1997; 272(24): 15167–15173.
[PubMed Abstract](#) | [Publisher Full Text](#)
30. Takekawa M, Posas F, Saito H: **A human homolog of the yeast Ssk2/Ssk22 MAP kinase kinase kinases, MTK1, mediates stress-induced activation of the p38 and JNK pathways.** *EMBO J.* 1997; 16(16): 4973–4982.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Bagrodia S, Derijard B, Davis RJ, *et al.*: **Cdc42 and PAK-mediated signaling leads to Jun kinase and p38 mitogen-activated protein kinase activation.** *J Biol Chem.* 1995; 270(47): 27995–27998.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Martin GA, Bollag G, McCormick F, *et al.*: **A novel serine kinase activated by rac1/CDC42Hs-dependent autophosphorylation is related to PAK65 and STE20.** *EMBO J.* 1995; 14(9): 1970–1978.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. **F** Ge B, Gram H, Di Padova F, *et al.*: **MAPKK-independent activation of p38alpha mediated by TAB1-dependent autophosphorylation of p38alpha.** *Science.* 2002; 295(5558): 1291–1294.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
34. **F** Thapa D, Nichols C, Bassi R, *et al.*: **TAB1-Induced Autoactivation of p38alpha Mitogen-Activated Protein Kinase Is Crucially Dependent on Threonine 185.** *Mol Cell Biol.* 2018; 38(5): e00409-17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
35. Ohkusu-Tsukada K, Tominaga N, Udono H, *et al.*: **Regulation of the maintenance of peripheral T-cell anergy by TAB1-mediated p38 alpha activation.** *Mol Cell Biol.* 2004; 24(16): 6957–6966.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Tanno M, Bassi R, Gorog DA, *et al.*: **Diverse mechanisms of myocardial p38 mitogen-activated protein kinase activation: evidence for MKK-independent activation by a TAB1-associated mechanism contributing to injury during myocardial ischemia.** *Circ Res.* 2003; 93(3): 254–261.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Mittelstadt PR, Yamaguchi H, Appella E, *et al.*: **T cell receptor-mediated activation of p38(alpha) by mono-phosphorylation of the activation loop results in altered substrate specificity.** *J Biol Chem.* 2009; 284(23): 15469–15474.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Jirmanova L, Sarma DN, Jankovic D, *et al.*: **Genetic disruption of p38alpha Tyr323 phosphorylation prevents T-cell receptor-mediated p38alpha activation and impairs interferon-gamma production.** *Blood.* 2009; 113(10): 2229–2237.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. **F** Jirmanova L, Giardinio Torchia ML, *et al.*: **Lack of the T cell-specific alternative p38 activation pathway reduces autoimmunity and inflammation.** *Blood.* 2011; 118(12): 3280–3289.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
40. Alam MS, Gaida MM, Debnath S, *et al.*: **Unique properties of TCR-activated p38 are necessary for NFAT-dependent T-cell activation.** *PLoS Biol.* 2018; 16(1): e2004111.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. **F** Giardinio Torchia ML, Dutta D, *et al.*: **Intensity and duration of TCR signaling is limited by p38 phosphorylation of ZAP-70^{T293} and destabilization of the signalosome.** *Proc Natl Acad Sci U S A.* 2018; 115(9): 2174–2179.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
42. Sun H, Charles CH, Lau LF, *et al.*: **MKP-1 (3CH134), an immediate early gene product, is a dual specificity phosphatase that dephosphorylates MAP kinase in vivo.** *Cell.* 1993; 75(3): 487–493.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Nunes-Xavier C, Roma-Mateo C, Rios P, *et al.*: **Dual-specificity MAP kinase phosphatases as targets of cancer treatment.** *Anticancer Agents Med Chem.* 2011; 11(1): 109–132.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Cho SSL, Han J, James SJ, *et al.*: **Dual-Specificity Phosphatase 12 Targets p38 MAP Kinase to Regulate Macrophage Response to Intracellular Bacterial Infection.** *Front Immunol.* 2017; 8: 1259.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
45. Tanoue T, Yamamoto T, Maeda R, *et al.*: **A Novel MAPK phosphatase MKP-7 acts preferentially on JNK/SAPK and p38 alpha and beta MAPKs.** *J Biol Chem.* 2001; 276(28): 26629–26639.
[PubMed Abstract](#) | [Publisher Full Text](#)
46. Manetsch M, Che W, Seidel P, *et al.*: **MKP-1: a negative feedback effector that represses MAPK-mediated pro-inflammatory signaling pathways and cytokine secretion in human airway smooth muscle cells.** *Cell Signal.* 2012; 24(4): 907–913.
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Topolska-Wos AM, Rosinska S, Filippek A: **MAP kinase p38 is a novel target of CacyBP/SIP phosphatase.** *Amino Acids.* 2017; 49(6): 1069–1076.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. Liu G, Hu X, Sun B, *et al.*: **Phosphatase Wip1 negatively regulates neutrophil development through p38 MAPK-STAT1.** *Blood.* 2013; 121(3): 519–529.
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Takekawa M, Maeda T, Saito H: **Protein phosphatase 2Alpha inhibits the human stress-responsive p38 and JNK MAPK pathways.** *EMBO J.* 1998; 17(16): 4744–4752.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Maeda T, Wurgler-Murphy SM, Saito H: **A two-component system that regulates an osmosensing MAP kinase cascade in yeast.** *Nature.* 1994; 369(6477): 242–245.
[PubMed Abstract](#) | [Publisher Full Text](#)
51. Qi X, Pohl NM, Loesch M, *et al.*: **p38alpha antagonizes p38gamma activity through c-Jun-dependent ubiquitin-proteasome pathways in regulating Ras transformation and stress response.** *J Biol Chem.* 2007; 282(43): 31398–31408.
[PubMed Abstract](#) | [Publisher Full Text](#)
52. **F** Chen YW, Ko WC, Chen CS, *et al.*: **RIOK-1 Is a Suppressor of the p38 MAPK Innate Immune Pathway in *Caenorhabditis elegans*.** *Front Immunol.* 2018; 9: 774.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
53. Bird TA, Schule HD, Delaney P, *et al.*: **The interleukin-1-stimulated protein kinase that phosphorylates heat shock protein hsp27 is activated by MAP kinase.** *FEBS Lett.* 1994; 338(1): 31–36.
[PubMed Abstract](#) | [Publisher Full Text](#)
54. McLaughlin MM, Kumar S, McDonnell PC, *et al.*: **Identification of mitogen-activated protein (MAP) kinase-activated protein kinase-3, a novel substrate of CSBP p38 MAP kinase.** *J Biol Chem.* 1996; 271(14): 8488–8492.
[PubMed Abstract](#) | [Publisher Full Text](#)
55. New L, Jiang Y, Zhao M, *et al.*: **PRAK, a novel protein kinase regulated by the p38 MAP kinase.** *EMBO J.* 1998; 17(12): 3372–3384.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. Deak M, Clifton AD, Lucocq LM, *et al.*: **Mitogen- and stress-activated protein kinase-1 (MSK1) is directly activated by MAPK and SAPK2/p38, and may mediate activation of CREB.** *EMBO J.* 1998; 17(15): 4426–4441.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
57. Waskiewicz AJ, Flynn A, Proud CG, *et al.*: **Mitogen-activated protein kinases activate the serine/threonine kinases Mnk1 and Mnk2.** *EMBO J.* 1997; 16(8): 1909–1920.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
58. Fukunaga R, Hunter T: **MNK1, a new MAP kinase-activated protein kinase, isolated by a novel expression screening method for identifying protein kinase substrates.** *EMBO J.* 1997; 16(8): 1921–1933.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
59. **F** Thornton TM, Pedraza-Alva G, Deng B, *et al.*: **Phosphorylation by p38 MAPK as an alternative pathway for GSK3beta inactivation.** *Science.* 2008; 320(5876): 667–670.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
60. Bikkavilli RK, Feigin ME, Malbon CC: **p38 mitogen-activated protein kinase regulates canonical Wnt-beta-catenin signaling by inactivation of GSK3beta.** *J Cell Sci.* 2008; 121(Pt 21): 3598–3607.
[PubMed Abstract](#) | [Publisher Full Text](#)
61. **F** Sumara G, Formentini I, Collins S, *et al.*: **Regulation of PKD by the MAPK p38delta in insulin secretion and glucose homeostasis.** *Cell.* 2009; 136(2): 235–248.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
62. Han J, Jiang Y, Li Z, *et al.*: **Activation of the transcription factor MEF2C by the MAP kinase p38 in inflammation.** *Nature.* 1997; 386(6622): 296–299.
[PubMed Abstract](#) | [Publisher Full Text](#)
63. Tan Y, Rouse J, Zhang A, *et al.*: **FGF and stress regulate CREB and ATF-1 via a pathway involving p38 MAP kinase and MAPKAP kinase-2.** *EMBO J.* 1996; 15(17): 4629–4642.
[PubMed Abstract](#) | [Free Full Text](#)
64. Barsyte-Lovejoy D, Galanis A, Sharrocks AD: **Specificity determinants in MAPK signaling to transcription factors.** *J Biol Chem.* 2002; 277(12): 9896–9903.
[PubMed Abstract](#) | [Publisher Full Text](#)
65. Hazzalin CA, Cano E, Cuenda A, *et al.*: **p38/RK is essential for stress-induced nuclear responses: JNK/SAPKs and c-Jun/ATF-2 phosphorylation are insufficient.** *Curr Biol.* 1996; 6(8): 1028–1031.
[PubMed Abstract](#) | [Publisher Full Text](#)
66. Whitmarsh AJ, Yang SH, Su MS, *et al.*: **Role of p38 and JNK mitogen-activated protein kinases in the activation of ternary complex factors.** *Mol Cell Biol.* 1997; 17(5): 2360–2371.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
67. Janknecht R, Hunter T: **Convergence of MAP kinase pathways on the ternary complex factor Sap-1a.** *EMBO J.* 1997; 16(7): 1620–1627.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
68. Wang XZ, Ron D: **Stress-induced phosphorylation and activation of the transcription factor CHOP (GADD153) by p38 MAP Kinase.** *Science.* 1996; 272(5266): 1347–1349.
[PubMed Abstract](#) | [Publisher Full Text](#)
69. Bulavin DV, Saito S, Hollander MC, *et al.*: **Phosphorylation of human p53 by p38 kinase coordinates N-terminal phosphorylation and apoptosis in response to UV radiation.** *EMBO J.* 1999; 18(23): 6845–6854.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
70. Huang C, Ma WY, Maxiner A, *et al.*: **p38 kinase mediates UV-induced**

- phosphorylation of p53 protein at serine 389. *J Biol Chem.* 1999; **274**(18): 12229–12235.
[PubMed Abstract](#) | [Publisher Full Text](#)
71. Galibert MD, Carreira S, Goding CR: The Usf-1 transcription factor is a novel target for the stress-responsive p38 kinase and mediates UV-induced Tyrosinase expression. *EMBO J.* 2001; **20**(17): 5022–5031.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 72. Gomez del Arco P, Martinez-Martinez S, Maldonado JL, et al.: A role for the p38 MAP kinase pathway in the nuclear shuttling of NFATp. *J Biol Chem.* 2000; **275**(18): 13872–13878.
[PubMed Abstract](#) | [Publisher Full Text](#)
 73. Rampalli S, Li L, Mak E, et al.: p38 MAPK signaling regulates recruitment of Ash2L-containing methyltransferase complexes to specific genes during differentiation. *Nat Struct Mol Biol.* 2007; **14**(12): 1150–1156.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 74. Ramsauer K, Sadzak I, Porras A, et al.: p38 MAPK enhances STAT1-dependent transcription independently of Ser-727 phosphorylation. *Proc Natl Acad Sci U S A.* 2002; **99**(20): 12859–12864.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 75. Lee J, Sun C, Zhou Y, et al.: p38 MAPK-mediated regulation of Xbp1s is crucial for glucose homeostasis. *Nat Med.* 2011; **17**(10): 1251–1260.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 76. Lin KC, Morioishi T, Meng Z, et al.: Regulation of Hippo pathway transcription factor TEAD by p38 MAPK-induced cytoplasmic translocation. *Nat Cell Biol.* 2017; **19**(8): 996–1002.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 77. Pierrat B, Correia JS, Mary JL, et al.: RSK-B, a novel ribosomal S6 kinase family member, is a CREB kinase under dominant control of p38alpha mitogen-activated protein kinase (p38alphaMAPK). *J Biol Chem.* 1998; **273**(45): 29661–29671.
[PubMed Abstract](#) | [Publisher Full Text](#)
 78. Kaur R, Liu X, Gjerup O, et al.: Activation of p21-activated kinase 6 by MAP kinase kinase 6 and p38 MAP kinase. *J Biol Chem.* 2005; **280**(5): 3323–3330.
[PubMed Abstract](#) | [Publisher Full Text](#)
 79. Jones DR, Bultsma Y, Keune WJ, et al.: Nuclear PtdIns5P as a transducer of stress signaling: an *in vivo* role for PIP4Kbeta. *Mol Cell.* 2006; **23**(5): 685–695.
[PubMed Abstract](#) | [Publisher Full Text](#)
 80. Saurin AT, Durgan J, Cameron AJ, et al.: The regulated assembly of a PKCepsilon complex controls the completion of cytokinesis. *Nat Cell Biol.* 2008; **10**(8): 891–901.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 81. Cuenda A, Cohen P, Buee-Scherrer V, et al.: Activation of stress-activated protein kinase-3 (SAPK3) by cytokines and cellular stresses is mediated via SAPK3 (MKK6); comparison of the specificities of SAPK3 and SAPK2 (RK/p38). *EMBO J.* 1997; **16**(2): 295–305.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 82. Qiao L, MacDougald OA, Shao J: CCAAT/enhancer-binding protein alpha mediates induction of hepatic phosphoenolpyruvate carboxykinase by p38 mitogen-activated protein kinase. *J Biol Chem.* 2006; **281**(34): 24390–24397.
[PubMed Abstract](#) | [Publisher Full Text](#)
 83. Engelman JA, Lisanti MP, Scherer PE: Specific inhibitors of p38 mitogen-activated protein kinase block 3T3-L1 adipogenesis. *J Biol Chem.* 1998; **273**(48): 32111–32120.
[PubMed Abstract](#) | [Publisher Full Text](#)
 84. Williamson EA, Williamson IK, Chumakov AM, et al.: CCAAT/enhancer binding protein epsilon: changes in function upon phosphorylation by p38 MAP kinase. *Blood.* 2005; **105**(10): 3841–3847.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 85. Morillo SM, Abanto EP, Roman MJ, et al.: Nerve growth factor-induced cell cycle reentry in newborn neurons is triggered by p38^{MAPK}-dependent E2F4 phosphorylation. *Mol Cell Biol.* 2012; **32**(14): 2722–2737.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 86. Lee H, Bai W: Regulation of estrogen receptor nuclear export by ligand-induced and p38-mediated receptor phosphorylation. *Mol Cell Biol.* 2002; **22**(16): 5835–5845.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 87. Tanos T, Marinissen MJ, Leskow FC, et al.: Phosphorylation of c-Fos by members of the p38 MAPK family. Role in the AP-1 response to UV light. *J Biol Chem.* 2005; **280**(19): 18842–18852.
[PubMed Abstract](#) | [Publisher Full Text](#)
 88. Ho KK, McGuire VA, Koo CY, et al.: Phosphorylation of FOXO3a on Ser-7 by p38 promotes its nuclear localization in response to doxorubicin. *J Biol Chem.* 2012; **287**(2): 1545–1555.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 89. Miller AL, Webb MS, Copik AJ, et al.: p38 Mitogen-activated protein kinase (MAPK) is a key mediator in glucocorticoid-induced apoptosis of lymphoid cells: correlation between p38 MAPK activation and site-specific phosphorylation of the human glucocorticoid receptor at serine 211. *Mol Endocrinol.* 2005; **19**(6): 1569–1583.
[PubMed Abstract](#) | [Publisher Full Text](#)
 90. Macfarlane WM, Smith SB, James RF, et al.: The p38/reactivating kinase mitogen-activated protein kinase cascade mediates the activation of the transcription factor insulin upstream factor 1 and insulin gene transcription by high glucose in pancreatic beta-cells. *J Biol Chem.* 1997; **272**(33): 20936–20944.
[PubMed Abstract](#) | [Publisher Full Text](#)
 91. Katz S, Aronheim A: Differential targeting of the stress mitogen-activated protein kinases to the c-Jun dimerization protein 2. *Biochem J.* 2002; **368**(Pt 3): 939–945.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 92. Humar M, Loop T, Schmidt R, et al.: The mitogen-activated protein kinase p38 regulates activator protein 1 by direct phosphorylation of c-Jun. *Int J Biochem Cell Biol.* 2007; **39**(12): 2278–2288.
[PubMed Abstract](#) | [Publisher Full Text](#)
 93. Sii-Felice K, Pouponnot C, Gillet S, et al.: MafA transcription factor is phosphorylated by p38 MAP kinase. *FEBS Lett.* 2005; **579**(17): 3547–3554.
[PubMed Abstract](#) | [Publisher Full Text](#)
 94. Zhao M, New L, Kravchenko VV, et al.: Regulation of the ME2 family of transcription factors by p38. *Mol Cell Biol.* 1999; **19**(1): 21–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 95. Mansky KC, Sankar U, Han J, et al.: Microphthalmia transcription factor is a target of the p38 MAPK pathway in response to receptor activator of NF-kappa B ligand signaling. *J Biol Chem.* 2002; **277**(13): 11077–11083.
[PubMed Abstract](#) | [Publisher Full Text](#)
 96. Suelves M, Lluís F, Ruiz V, et al.: Phosphorylation of MRF4 transactivation domain by p38 mediates repression of specific myogenic genes. *EMBO J.* 2004; **23**(2): 365–375.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 97. Matsumoto M, Kogawa M, Wada S, et al.: Essential role of p38 mitogen-activated protein kinase in cathepsin K gene expression during osteoclastogenesis through association of NFATc1 and PU.1. *J Biol Chem.* 2004; **279**(44): 45969–45979.
[PubMed Abstract](#) | [Publisher Full Text](#)
 98. Yang TT, Xiong Q, Enslin H, et al.: Phosphorylation of NFATc4 by p38 mitogen-activated protein kinases. *Mol Cell Biol.* 2002; **22**(11): 3892–3904.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 99. Sekine Y, Takagahara S, Hatanaka R, et al.: p38 MAPKs regulate the expression of genes in the dopamine synthesis pathway through phosphorylation of NR4A nuclear receptors. *J Cell Sci.* 2011; **124**(Pt 17): 3006–3016.
[PubMed Abstract](#) | [Publisher Full Text](#)
 100. Li L, Liu Y, Chen HZ, et al.: Impeding the interaction between Nur77 and p38 reduces LPS-induced inflammation. *Nat Chem Biol.* 2015; **11**(5): 339–346.
[PubMed Abstract](#) | [Publisher Full Text](#)
 101. Ortuno MJ, Ruiz-Gaspa S, Rodriguez-Carballo E, et al.: p38 regulates expression of osteoblast-specific genes by phosphorylation of osterix. *J Biol Chem.* 2010; **285**(42): 31985–31994.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 102. Mikkola I, Bruun JA, Bjorkoy G, et al.: Phosphorylation of the transactivation domain of Pax6 by extracellular signal-regulated kinase and p38 mitogen-activated protein kinase. *J Biol Chem.* 1999; **274**(21): 15115–15126.
[PubMed Abstract](#) | [Publisher Full Text](#)
 103. Barger PM, Browning AC, Garner AN, et al.: p38 mitogen-activated protein kinase activates peroxisome proliferator-activated receptor alpha: a potential role in the cardiac metabolic stress response. *J Biol Chem.* 2001; **276**(48): 44495–44501.
[PubMed Abstract](#) | [Publisher Full Text](#)
 104. Hayes SA, Huang X, Kambhampati S, et al.: p38 MAP kinase modulates Smad-dependent changes in human prostate cell adhesion. *Oncogene.* 2003; **22**(31): 4841–4850.
[PubMed Abstract](#) | [Publisher Full Text](#)
 105. Ryu KJ, Park SM, Park SH, et al.: p38 Stabilizes Snail by Suppressing DYRK2-Mediated Phosphorylation That Is Required for GSK3beta-betaTrCP-Induced Snail Degradation. *Cancer Res.* 2019; **79**(16): 4135–4148.
[PubMed Abstract](#) | [Publisher Full Text](#)
 106. Kovarik P, Stoiber D, Eysers PA, et al.: Stress-induced phosphorylation of STAT1 at Ser727 requires p38 mitogen-activated protein kinase whereas IFN-gamma uses a different signaling pathway. *Proc Natl Acad Sci U S A.* 1999; **96**(24): 13956–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 107. Visconti R, Gadinia M, Chiariello M, et al.: Importance of the MKK6/p38 pathway for interleukin-12-induced STAT4 serine phosphorylation and transcriptional activity. *Blood.* 2000; **96**(5): 1844–52.
[PubMed Abstract](#)
 108. Hong J, Zhou J, Fu J, et al.: Phosphorylation of serine 68 of Twist1 by MAPKs stabilizes Twist1 protein and promotes breast cancer cell invasiveness. *Cancer Res.* 2011; **71**(11): 3980–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 109. Simone C, Forcales SV, Hill DA, et al.: p38 pathway targets SWI-SNF chromatin-remodeling complex to muscle-specific loci. *Nat Genet.* 2004; **36**(7): 738–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 110. Forcales SV, Albini S, Giordani L, et al.: Signal-dependent incorporation of MyoD-BAF60c into Brg1-based SWI/SNF chromatin-remodelling complex. *EMBO J.* 2012; **31**(2): 301–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 111. Anwar T, Arellano-Garcia C, Ropa J, et al.: p38-mediated phosphorylation

- at T367 induces EZH2 cytoplasmic localization to promote breast cancer metastasis. *Nat Commun.* 2018; 9(1): 2801.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
112. **F** Seong KH, Li D, Shimizu H, *et al.*: Inheritance of stress-induced, ATF-2-dependent epigenetic change. *Cell.* 2011; 145(7): 1049–1061.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 113. Wu XN, Wang XK, Wu SQ, *et al.*: Phosphorylation of Raptor by p38beta participates in arsenite-induced mammalian target of rapamycin complex 1 (mTORC1) activation. *J Biol Chem.* 2011; 286(36): 31501–31511.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 114. **F** Gonzalez-Teran B, Lopez JA, Rodriguez E, *et al.*: p38gamma and delta promote heart hypertrophy by targeting the mTOR-inhibitory protein DEPTOR for degradation. *Nat Commun.* 2016; 7: 10477.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 115. Zheng H, Seit-Nebi A, Han X, *et al.*: A posttranslational modification cascade involving p38, Tip60, and PRAK mediates oncogene-induced senescence. *Mol Cell.* 2013; 50(5): 699–710.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 116. Xu Y, Liao R, Li N, *et al.*: Phosphorylation of Tip60 by p38alpha regulates p53-mediated PUMA induction and apoptosis in response to DNA damage. *Oncotarget.* 2014; 5(24): 12555–12572.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 117. Li W, Zhu J, Dou J, *et al.*: Phosphorylation of LAMP2A by p38 MAPK couples ER stress to chaperone-mediated autophagy. *Nat Commun.* 2017; 8(1): 1763.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 118. Allen M, Svensson L, Roach M, *et al.*: Deficiency of the stress kinase p38alpha results in embryonic lethality: characterization of the kinase dependence of stress responses of enzyme-deficient embryonic stem cells. *J Exp Med.* 2000; 191(5): 859–870.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 119. Tamura K, Sudo T, Sentfleben U, *et al.*: Requirement for p38alpha in erythropoietin expression: a role for stress kinases in erythropoiesis. *Cell.* 2000; 102(2): 221–231.
[PubMed Abstract](#) | [Publisher Full Text](#)
 120. Adams RH, Porras A, Alonso G, *et al.*: Essential role of p38alpha MAP kinase in placental but not embryonic cardiovascular development. *Mol Cell.* 2000; 6(1): 109–116.
[PubMed Abstract](#) | [Publisher Full Text](#)
 121. Mudgett JS, Ding J, Guh-Siesel L, *et al.*: Essential role for p38alpha mitogen-activated protein kinase in placental angiogenesis. *Proc Natl Acad Sci U S A.* 2000; 97(19): 10454–10459.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 122. Wong ES, Le Guezennec X, Demidov ON, *et al.*: p38MAPK controls expression of multiple cell cycle inhibitors and islet proliferation with advancing age. *Dev Cell.* 2009; 17(1): 142–149.
[PubMed Abstract](#) | [Publisher Full Text](#)
 123. Brichkina A, Bertero T, Loh HM, *et al.*: p38MAPK builds a hyaluronan cancer niche to drive lung tumorigenesis. *Genes Dev.* 2016; 30(23): 2623–2636.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 124. Rao PS, Satelli A, Zhang S, *et al.*: RNF2 is the target for phosphorylation by the p38 MAPK and ERK signaling pathways. *Proteomics.* 2009; 9(10): 2776–2787.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 125. Ma H, Liu Z, Zhong CQ, *et al.*: Inactivation of Cyclic AMP Response Element Transcription Caused by Constitutive p38 Activation Is Mediated by Hyperphosphorylation-Dependent CRT2 Nucleocytoplasmic Transport. *Mol Cell Biol.* 2019; 39(9): e00554–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 126. Page JL, Wang X, Sordillo LM, *et al.*: MEK1 signaling through p38 leads to transcriptional inactivation of E47 and repression of skeletal myogenesis. *J Biol Chem.* 2004; 279(30): 30966–30972.
[PubMed Abstract](#) | [Publisher Full Text](#)
 127. Lluís F, Ballestar E, Suelves M, *et al.*: E47 phosphorylation by p38 MAPK promotes MyoD/E47 association and muscle-specific gene transcription. *EMBO J.* 2005; 24(5): 974–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 128. Xiu M, Kim J, Sampson E, *et al.*: The transcriptional repressor HBP1 is a target of the p38 mitogen-activated protein kinase pathway in cell cycle regulation. *Mol Cell Biol.* 2003; 23(23): 8890–901.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 129. Cuadrado A, Lafarga V, Cheung PCF, *et al.*: A new p38 MAP kinase-regulated transcriptional coactivator that stimulates p53-dependent apoptosis. *EMBO J.* 2007; 26(8): 2115–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 130. Puigserver P, Rhee J, Lin J, *et al.*: Cytokine stimulation of energy expenditure through p38 MAP kinase activation of PPARgamma coactivator-1. *Mol Cell.* 2001; 8(5): 971–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
 131. Delston RB, Matatall KA, Sun Y, *et al.*: p38 phosphorylates Rb on Ser567 by a novel, cell cycle-independent mechanism that triggers Rb-Hdm2 interaction and apoptosis. *Oncogene.* 2011; 30(5): 588–99.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 132. Gianni M, Parrella E, Raska I Jr, *et al.*: P38MAPK-dependent phosphorylation and degradation of SRC-3/AIB1 and RARalpha-mediated transcription. *EMBO J.* 2006; 25(4): 739–751.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 133. Goloudina A, Yamaguchi H, Chervyakova DB, *et al.*: Regulation of human Cdc25A stability by Serine 75 phosphorylation is not sufficient to activate a S phase checkpoint. *Cell Cycle.* 2003; 2(5): 473–478.
[PubMed Abstract](#) | [Publisher Full Text](#)
 134. Lemaire M, Froment C, Boutros R, *et al.*: CDC25B phosphorylation by p38 and MK-2. *Cell Cycle.* 2006; 5(15): 1649–1653.
[PubMed Abstract](#) | [Publisher Full Text](#)
 135. Casanovas O, Miro F, Estanyol JM, *et al.*: Osmotic stress regulates the stability of cyclin D1 in a p38SAPK2-dependent manner. *J Biol Chem.* 2000; 275(45): 35091–35097.
[PubMed Abstract](#) | [Publisher Full Text](#)
 136. Casanovas O, Jaumot M, Paules AB, *et al.*: P38SAPK2 phosphorylates cyclin D3 at Thr-283 and targets it for proteasomal degradation. *Oncogene.* 2004; 23(45): 7537–7544.
[PubMed Abstract](#) | [Publisher Full Text](#)
 137. Joaquin M, Gubern A, Gonzalez-Nunez D, *et al.*: The p57 CDKi integrates stress signals into cell-cycle progression to promote cell survival upon stress. *EMBO J.* 2012; 31(13): 2952–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 138. Min H, Ghatnekar GS, Ghatnekar AV, *et al.*: 2-Methoxyestradiol induced Bax phosphorylation and apoptosis in human retinoblastoma cells via p38 MAPK activation. *Mol Carcinog.* 2012; 51(7): 576–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
 139. Cai B, Chang SH, Becker EBE, *et al.*: p38 MAP kinase mediates apoptosis through phosphorylation of BimEL at Ser-65. *J Biol Chem.* 2006; 281(35): 25215–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
 140. Alvarado-Kristensson M, Melander F, Leandersson K, *et al.*: p38-MAPK signals survival by phosphorylation of caspase-8 and caspase-3 in human neutrophils. *J Exp Med.* 2004; 199(4): 449–58.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 141. Seifert A, Clarke PR: p38alpha- and DYRK1A-dependent phosphorylation of caspase-9 at an inhibitory site in response to hyperosmotic stress. *Cell Signal.* 2009; 21(11): 1626–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
 142. Chandrasekaran S, Tan TX, Hall JR, *et al.*: Stress-stimulated mitogen-activated protein kinases control the stability and activity of the Cdt1 DNA replication licensing factor. *Mol Cell Biol.* 2011; 31(22): 4405–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 143. Yang Q, Li W, She H, *et al.*: Stress induces p38 MAPK-mediated phosphorylation and inhibition of Drosha-dependent cell survival. *Mol Cell.* 2015; 57(4): 721–734.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 144. Dankwardt S, Gantzer AS, Macher-Goeppinger S, *et al.*: p38 MAPK controls prothrombin expression by regulated RNA 3' end processing. *Mol Cell.* 2011; 41(3): 298–310.
[PubMed Abstract](#) | [Publisher Full Text](#)
 145. Lu C, Shi Y, Wang Z, *et al.*: Serum starvation induces H2AX phosphorylation to regulate apoptosis via p38 MAPK pathway. *FEBS Lett.* 2008; 582(18): 2703–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 146. Zhong SP, Ma WY, Dong Z: ERKs and p38 kinases mediate ultraviolet B-induced phosphorylation of histone H3 at serine 10. *J Biol Chem.* 2000; 275(28): 20980–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
 147. Lafarga V, Cuadrado A, Lopez de Silanes I, *et al.*: p38 Mitogen-activated protein kinase- and HuR-dependent stabilization of p21^{Cip1} mRNA mediates the G₁/S checkpoint. *Mol Cell Biol.* 2009; 29(16): 4341–4351.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 148. Briata P, Forcales SV, Ponassi M, *et al.*: p38-dependent phosphorylation of the mRNA decay-promoting factor KSRP controls the stability of select myogenic transcripts. *Mol Cell.* 2005; 20(6): 891–903.
[PubMed Abstract](#) | [Publisher Full Text](#)
 149. Knight JD, Tian R, Lee RE, *et al.*: A novel whole-cell lysate kinase assay identifies substrates of the p38 MAPK in differentiating myoblasts. *Skelet Muscle.* 2012; 2: 5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 150. Al-Ayoubi AM, Zheng H, Liu Y, *et al.*: Mitogen-activated protein kinase phosphorylation of splicing factor 45 (SPF45) regulates SPF45 alternative splicing site utilization, proliferation, and cell adhesion. *Mol Cell Biol.* 2012; 32(14): 2880–2893.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 151. Mace G, Miaczynska M, Zerial M, *et al.*: Phosphorylation of EEA1 by p38 MAP kinase regulates mu opioid receptor endocytosis. *EMBO J.* 2005; 24(18): 3235–3246.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 152. Cavalli V, Vilbois F, Corti M, *et al.*: The stress-induced MAP kinase p38 regulates endocytic trafficking via the GDI:Rab5 complex. *Mol Cell.* 2001; 7(2): 421–432.
[PubMed Abstract](#) | [Publisher Full Text](#)
 153. Kelkar N, Standen CL, Davis RJ: Role of the JIP4 scaffold protein in the regulation of mitogen-activated protein kinase signaling pathways. *Mol Cell Biol.* 2005; 25(7): 2733–2743.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

154. Cheung PC, Campbell DG, Nebreda AR, *et al.*: **Feedback control of the protein kinase TAK1 by SAPK2a/p38alpha**. *EMBO J*. 2003; **22**(21): 5793–5805.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
155. Mendoza H, Campbell DG, Burness K, *et al.*: **Roles for TAB1 in regulating the IL-1-dependent phosphorylation of the TAB3 regulatory subunit and activity of the TAK1 complex**. *Biochem J*. 2008; **409**(3): 711–722.
[PubMed Abstract](#) | [Publisher Full Text](#)
156. Zakrzewska M, Opalinski L, Haugsten EM, *et al.*: **Crosstalk between p38 and Erk 1/2 in Downregulation of FGF1-Induced Signaling**. *Int J Mol Sci*. 2019; **20**(8): 1826.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
157. Winograd-Katz SE, Levitzki A: **Cisplatin induces PKB/Akt activation and p38^{MAPK} phosphorylation of the EGF receptor**. *Oncogene*. 2006; **25**(56): 7381–7390.
[PubMed Abstract](#) | [Publisher Full Text](#)
158. Sorensen V, Zhen Y, Zakrzewska M, *et al.*: **Phosphorylation of fibroblast growth factor (FGF) receptor 1 at Ser777 by p38 mitogen-activated protein kinase regulates translocation of exogenous FGF1 to the cytosol and nucleus**. *Mol Cell Biol*. 2008; **28**(12): 4129–4141.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
159. Gasser A, Cheng X, Gilmore ES, *et al.*: **Two Nedd4-binding motifs underlie modulation of sodium channel Na_v1.6 by p38 MAPK**. *J Biol Chem*. 2010; **285**(34): 26149–26161.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
160. Khaled AR, Moor AN, Li A, *et al.*: **Trophic factor withdrawal: p38 mitogen-activated protein kinase activates NHE1, which induces intracellular alkalization**. *Mol Cell Biol*. 2001; **21**(22): 7545–7557.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
161. Börsch-Haubold AG, Bartoli F, Asselin J, *et al.*: **Identification of the phosphorylation sites of cytosolic phospholipase A2 in agonist-stimulated human platelets and HeLa cells**. *J Biol Chem*. 1998; **273**(8): 4449–4458.
[PubMed Abstract](#) | [Publisher Full Text](#)
162. **F** Xu P, Derynck R: **Direct activation of TACE-mediated ectodomain shedding by p38 MAP kinase regulates EGF receptor-dependent cell proliferation**. *Mol Cell*. 2010; **37**(4): 551–566.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
163. Hedges JC, Yamboliev IA, Ngo M, *et al.*: **p38 mitogen-activated protein kinase expression and activation in smooth muscle**. *Am J Physiol*. 1998; **275**(2): C527–534.
[PubMed Abstract](#) | [Publisher Full Text](#)
164. Ku NO, Azhar S, Omary MB: **Keratin 8 phosphorylation by p38 kinase regulates cellular keratin filament reorganization: modulation by a keratin 1-like disease causing mutation**. *J Biol Chem*. 2002; **277**(13): 10775–10782.
[PubMed Abstract](#) | [Publisher Full Text](#)
165. Barascu A, Le Chalony C, Pennarun G, *et al.*: **Oxidative stress induces an ATM-independent senescence pathway through p38 MAPK-mediated lamin B1 accumulation**. *EMBO J*. 2012; **31**(5): 1080–1094.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
166. Huang C, Borchers CH, Schaller MD, *et al.*: **Phosphorylation of paxillin by p38MAPK is involved in the neurite extension of PC-12 cells**. *J Cell Biol*. 2004; **164**(4): 593–602.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
167. Parker CG, Hunt J, Diener K, *et al.*: **Identification of stathmin as a novel substrate for p38 delta**. *Biochem Biophys Res Commun*. 1998; **249**(3): 791–796.
[PubMed Abstract](#) | [Publisher Full Text](#)
168. **F** Sabio G, Arthur JS, Kuma Y, *et al.*: **p38gamma regulates the localisation of SAP97 in the cytoskeleton by modulating its interaction with GKAP**. *EMBO J*. 2005; **24**(6): 1134–1145.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
169. Reynolds CH, Nebreda AR, Gibb GM, *et al.*: **Reactivating kinase/p38 phosphorylates tau protein in vitro**. *J Neurochem*. 1997; **69**(1): 191–198.
[PubMed Abstract](#) | [Publisher Full Text](#)
170. Ittner A, Chua SW, Bertz J, *et al.*: **Site-specific phosphorylation of tau inhibits amyloid- β toxicity in Alzheimer's mice**. *Science*. 2016; **354**(6314): 904–908.
[PubMed Abstract](#) | [Publisher Full Text](#)
171. Hall EH, Balsbaugh JL, Rose KL, *et al.*: **Comprehensive analysis of phosphorylation sites in Tensin1 reveals regulation by p38MAPK**. *Mol Cell Proteomics*. 2010; **9**(12): 2853–2863.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
172. Kuma Y, Campbell DG, Cuenda A: **Identification of glycogen synthase as a new substrate for stress-activated protein kinase 2b/p38beta**. *Biochem J*. 2004; **379**(Pt 1): 133–139.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
173. Chen J, Ren Y, Gui C, *et al.*: **Phosphorylation of Parkin at serine 131 by p38 MAPK promotes mitochondrial dysfunction and neuronal death in mutant A53T α -synuclein model of Parkinson's disease**. *Cell Death Dis*. 2018; **9**(6): 700.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
174. Makni-Maalej K, Boussetta T, Hurtado-Nedelec M, *et al.*: **The TLR7/8 agonist CL097 primes N-formyl-methionyl-leucyl-phenylalanine-stimulated NADPH oxidase activation in human neutrophils: critical role of p47phox phosphorylation and the proline isomerase Pin1**. *J Immunol*. 2012; **189**(9): 4657–4665.
[PubMed Abstract](#) | [Publisher Full Text](#)
175. Linares JF, Duran A, Reina-Campos M, *et al.*: **Amino Acid Activation of mTORC1 by a PB1-Domain-Driven Kinase Complex Cascade**. *Cell Rep*. 2015; **12**(8): 1339–1352.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
176. **F** Koh A, Molinaro A, Stahlman M, *et al.*: **Microbially Produced Imidazole Propionate Impairs Insulin Signaling through mTORC1**. *Cell*. 2018; **175**(4): 947–961.e917.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
177. Lee SH, Park Y, Yoon SK, *et al.*: **Osmotic stress inhibits proteasome by p38 MAPK-dependent phosphorylation**. *J Biol Chem*. 2010; **285**(53): 41280–41289.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
178. **F** Khurana A, Nakayama K, Williams S, *et al.*: **Regulation of the ring finger E3 ligase Slah2 by p38 MAPK**. *J Biol Chem*. 2006; **281**(46): 35316–35326.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
179. Lalaoui N, Hanggi K, Brumatti G, *et al.*: **Targeting p38 or MK2 Enhances the Anti-Leukemic Activity of Smac-Mimetics**. *Cancer Cell*. 2016; **30**(3): 499–500.
[PubMed Abstract](#) | [Publisher Full Text](#)
180. **F** Jaco I, Annibaldi A, Lalaoui N, *et al.*: **MK2 Phosphorylates RIPK1 to Prevent TNF-Induced Cell Death**. *Mol Cell*. 2017; **66**(5): 698–710.e5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
181. Dondelinger Y, Delanghe T, Rojas-Rivera D, *et al.*: **MK2 phosphorylation of RIPK1 regulates TNF-mediated cell death**. *Nat Cell Biol*. 2017; **19**(10): 1237–1247.
[PubMed Abstract](#) | [Publisher Full Text](#)
182. Menon MB, Gropengießer J, Fischer J, *et al.*: **p38^{MAPK}/MK2-dependent phosphorylation controls cytotoxic RIPK1 signalling in inflammation and infection**. *Nat Cell Biol*. 2017; **19**(10): 1248–1259.
[PubMed Abstract](#) | [Publisher Full Text](#)
183. Silke J, Rickard JA, Gerlic M: **The diverse role of RIP kinases in necroptosis and inflammation**. *Nat Immunol*. 2015; **16**(7): 689–697.
[PubMed Abstract](#) | [Publisher Full Text](#)
184. Dillon CP, Tummers B, Baran K, *et al.*: **Developmental checkpoints guarded by regulated necrosis**. *Cell Mol Life Sci*. 2016; **73**(11-12): 2125–2136.
[PubMed Abstract](#) | [Publisher Full Text](#)
185. del Barco Barrantes I, Coya JM, Maina F, *et al.*: **Genetic analysis of specific and redundant roles for p38alpha and p38beta MAPKs during mouse development**. *Proc Natl Acad Sci U S A*. 2011; **108**(31): 12764–12769.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
186. Kang YJ, Chen J, Otsuka M, *et al.*: **Macrophage deletion of p38alpha partially impairs lipopolysaccharide-induced cellular activation**. *J Immunol*. 2008; **180**(7): 5075–5082.
[PubMed Abstract](#) | [Publisher Full Text](#)
187. Tiedje C, Holtmann H, Gaestel M: **The role of mammalian MAPK signaling in regulation of cytokine mRNA stability and translation**. *J Interferon Cytokine Res*. 2014; **34**(4): 220–232.
[PubMed Abstract](#) | [Publisher Full Text](#)
188. Xu Z, Yoshida T, Wu L, *et al.*: **Transcription factor MEF2C suppresses endothelial cell inflammation via regulation of NF-kappaB and KLF2**. *J Cell Physiol*. 2015; **230**(6): 1310–1320.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
189. Ronkina N, Menon MB, Schwermann J, *et al.*: **Stress induced gene expression: a direct role for MAPKAP kinases in transcriptional activation of immediate early genes**. *Nucleic Acids Res*. 2011; **39**(7): 2503–2518.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
190. **F** Ronkina N, Shushakova N, Tiedje C, *et al.*: **The Role of TTP Phosphorylation in the Regulation of Inflammatory Cytokine Production by MK2/3**. *J Immunol*. 2019; **203**(8): 2291–2300.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
191. Guan Z, Buckman SY, Pentland AP, *et al.*: **Induction of cyclooxygenase-2 by the activated MEK1 $\rightarrow$ SEK1/MKK4 $\rightarrow$ p38 mitogen-activated protein kinase pathway**. *J Biol Chem*. 1998; **273**(21): 12901–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
192. Badger AM, Cook MN, Lark MW, *et al.*: **SB 203580 inhibits p38 mitogen-activated protein kinase, nitric oxide production, and inducible nitric oxide synthase in bovine cartilage-derived chondrocytes**. *J Immunol*. 1998; **161**(1): 467–73.
[PubMed Abstract](#)
193. Da Silva J, Pierrat B, Mary JL, *et al.*: **Blockade of p38 mitogen-activated protein kinase pathway inhibits inducible nitric-oxide synthase expression in mouse astrocytes**. *J Biol Chem*. 1997; **272**(45): 28373–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
194. Wiehler S, Cuvelier SL, Chakrabarti S, *et al.*: **p38 MAP kinase regulates rapid matrix metalloproteinase-9 release from eosinophils**. *Biochem Biophys Res Commun*. 2004; **315**(2): 463–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
195. Zhou Y, Wu J, Liu C, *et al.*: **p38alpha has an important role in antigen cross-presentation by dendritic cells**. *Cell Mol Immunol*. 2018; **15**(3): 246–259.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
196. Craxton A, Shu G, Graves JD, *et al.*: **p38 MAPK is required for CD40-induced gene expression and proliferation in B lymphocytes**. *J Immunol*. 1998; **161**(7): 3225–36.
[PubMed Abstract](#)
197. Khiem D, Cyster JG, Schwarz JJ, *et al.*: **A p38 MAPK-MEF2C pathway regulates**

- B-cell proliferation.** *Proc Natl Acad Sci U S A.* 2008; **105**(44): 17067–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
198. Wilker PR, Kohyama M, Sandau MM, *et al.*: Transcription factor Mef2c is required for B cell proliferation and survival after antigen receptor stimulation. *Nat Immunol.* 2008; **9**(6): 603–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 199. Zhang X, Fan L, Wu J, *et al.*: Macrophage p38 α promotes nutritional steatohepatitis through M1 polarization. *J Hepatol.* 2019; **71**(1): 163–174.
[PubMed Abstract](#) | [Publisher Full Text](#)
 200. **F** Otsuka M, Kang YJ, Ren J, *et al.*: Distinct effects of p38 α deletion in myeloid lineage and gut epithelia in mouse models of inflammatory bowel disease. *Gastroenterology.* 2010; **138**(4): 1255–65.1265.e1-9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 201. Liang L, Li F, Bao A, *et al.*: Activation of p38 mitogen-activated protein kinase in ovalbumin and ozone-induced mouse model of asthma. *Respirology.* 2013; **18** Suppl 3: 20–29.
[PubMed Abstract](#) | [Publisher Full Text](#)
 202. Gonzalez-Teran B, Matesanz N, Nikolic I, *et al.*: p38 γ and p38 δ Reprogram Liver Metabolism by Modulating Neutrophil Infiltration. *EMBO J.* 2016; **35**(5): 536–552.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 203. Gonzalez-Teran B, Cortes JR, Manieri E, *et al.*: Eukaryotic Elongation Factor 2 Controls TNF- α Translation in LPS-induced Hepatitis. *J Clin Invest.* 2013; **123**(1): 164–178.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 204. Ittner A, Block H, Reichel CA, *et al.*: Regulation of PTEN Activity by p38 δ -PKD1 Signaling in Neutrophils Confers Inflammatory Responses in the Lung. *J Exp Med.* 2012; **209**(12): 2229–2246.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 205. Genovese MC: Inhibition of p38: has the fat lady sung? *Arthritis Rheum.* 2009; **60**(12): 317–320.
[PubMed Abstract](#) | [Publisher Full Text](#)
 206. Takenaka K, Moriguchi T, Nishida E: Activation of the protein kinase p38 in the spindle assembly checkpoint and mitotic arrest. *Science.* 1998; **280**(5363): 599–602.
[PubMed Abstract](#) | [Publisher Full Text](#)
 207. Molnar A, Theodoras AM, Zon LI, *et al.*: Cdc42Hs, but not Rac1, inhibits serum-stimulated cell cycle progression at G₁/S through a mechanism requiring p38/RK. *J Biol Chem.* 1997; **272**(20): 13229–13235.
[PubMed Abstract](#) | [Publisher Full Text](#)
 208. Wang X, McGowan CH, Zhao M, *et al.*: Involvement of the MKK6-p38 γ cascade in gamma-radiation-induced cell cycle arrest. *Mol Cell Biol.* 2000; **20**(13): 4543–4552.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 209. **F** Gubern A, Joaquin M, Marquès M, *et al.*: The N-Terminal Phosphorylation of RB by p38 Bypasses Its Inactivation by CDKs and Prevents Proliferation in Cancer Cells. *Mol Cell.* 2016; **64**(1): 25–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 210. Yee AS, Paulson EK, McDevitt MA, *et al.*: The HBP1 transcriptional repressor and the p38 MAP kinase: unlikely partners in G1 regulation and tumor suppression. *Gene.* 2004; **336**(1): 1–13.
[PubMed Abstract](#) | [Publisher Full Text](#)
 211. Kumari G, Ulrich T, Gaubatz S: A role for p38 in transcriptional elongation of p21^{Cip1} in response to Aurora B inhibition. *Cell Cycle.* 2013; **12**(13): 2051–2060.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 212. Saha K, Adhikary G, Kanade SR, *et al.*: p38 δ regulates p53 to control p21^{Cip1} expression in human epidermal keratinocytes. *J Biol Chem.* 2014; **289**(16): 11443–11453.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 213. Niu Z, Mu H, Zhu H, *et al.*: p38 MAPK pathway is essential for self-renewal of mouse male germline stem cells (mGSCs). *Cell Prolif.* 2017; **50**(1): e12314.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 214. **F** Liu S, Ginzberg MB, Patel N, *et al.*: Size uniformity of animal cells is actively maintained by a p38 MAPK-dependent regulation of G1-length. *eLife.* 2018; **7**: e26947.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 215. Morooka T, Nishida E: Requirement of p38 mitogen-activated protein kinase for neuronal differentiation in PC12 cells. *J Biol Chem.* 1998; **273**(38): 24285–24288.
[PubMed Abstract](#) | [Publisher Full Text](#)
 216. Li Y, Jiang B, Ensign WY, *et al.*: Myogenic differentiation requires signalling through both phosphatidylinositol 3-kinase and p38 MAP kinase. *Cell Signal.* 2000; **12**(110–12): 751–757.
[PubMed Abstract](#) | [Publisher Full Text](#)
 217. Cao W, Medvedev AV, Daniel KW, *et al.*: β -Adrenergic activation of p38 MAP kinase in adipocytes: cAMP induction of the uncoupling protein 1 (UCP1) gene requires p38 MAP kinase. *J Biol Chem.* 2001; **276**(29): 27077–27082.
[PubMed Abstract](#) | [Publisher Full Text](#)
 218. Cao W, Daniel KW, Robidoux J, *et al.*: p38 mitogen-activated protein kinase is the central regulator of cyclic AMP-dependent transcription of the brown fat uncoupling protein 1 gene. *Mol Cell Biol.* 2004; **24**(7): 3057–3067.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 219. **F** Matesanz N, Nikolic I, Leiva M, *et al.*: p38 α blocks brown adipose tissue thermogenesis through p38 δ inhibition. *PLoS Biol.* 2018; **16**(7): e2004455.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 220. **F** Zhang S, Cao H, Li Y, *et al.*: Metabolic benefits of inhibition of p38 α in white adipose tissue in obesity. *PLoS Biol.* 2018; **16**(5): e2004225.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 221. Jing Y, Liu W, Cao H, *et al.*: Hepatic p38 α regulates gluconeogenesis by suppressing AMPK. *J Hepatol.* 2015; **62**(6): 1319–1327.
[PubMed Abstract](#) | [Publisher Full Text](#)
 222. **F** Lanna A, Henson SM, Escors D, *et al.*: The kinase p38 activated by the metabolic regulator AMPK and scaffold TAB1 drives the senescence of human T cells. *Nat Immunol.* 2014; **15**(10): 965–972.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 223. **F** Wang W, Chen JX, Liao R, *et al.*: Sequential activation of the MEK-extracellular signal-regulated kinase and MKK3/6-p38 mitogen-activated protein kinase pathways mediates oncogenic ras-induced premature senescence. *Mol Cell Biol.* 2002; **22**(10): 3389–3403.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 224. **F** Haq R, Brenton JD, Takahashi M, *et al.*: Constitutive p38HOG mitogen-activated protein kinase activation induces permanent cell cycle arrest and senescence. *Cancer Res.* 2002; **62**(17): 5076–5082.
[PubMed Abstract](#) | [Faculty Opinions Recommendation](#)
 225. Tivey HS, Brook AJ, Rokicki MJ, *et al.*: p38^{MAPK} stress signalling in replicative senescence in fibroblasts from progeroid and genomic instability syndromes. *Biogerontology.* 2013; **14**(1): 47–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 226. Naka K, Tachibana A, Ikeda K, *et al.*: Stress-induced premature senescence in hTERT-expressing ataxia telangiectasia fibroblasts. *J Biol Chem.* 2004; **279**(3): 2030–2037.
[PubMed Abstract](#) | [Publisher Full Text](#)
 227. Dimozi A, Mavrogenatou E, Sklirova A, *et al.*: Oxidative stress inhibits the proliferation, induces premature senescence and promotes a catabolic phenotype in human nucleus pulposus intervertebral disc cells. *Eur Cell Mater.* 2015; **30**: 89–102.
[PubMed Abstract](#) | [Publisher Full Text](#)
 228. Borodkina AV, Shatrova AN, Nikolsky NN, *et al.*: Role of P38 Map-Kinase in the Stress-Induced Senescence Progression of Human Endometrium-Derived Mesenchymal Stem Cells. *Tsitologiya.* 2016; **58**(6): 429–435.
[PubMed Abstract](#)
 229. Harada G, Neng Q, Fujiki T, *et al.*: Molecular mechanisms for the p38-induced cellular senescence in normal human fibroblast. *J Biochem.* 2014; **156**(5): 283–290.
[PubMed Abstract](#) | [Publisher Full Text](#)
 230. Hongo A, Okumura N, Nakahara M, *et al.*: The Effect of a p38 Mitogen-Activated Protein Kinase Inhibitor on Cellular Senescence of Cultivated Human Corneal Endothelial Cells. *Invest Ophthalmol Vis Sci.* 2017; **58**(9): 3325–3334.
[PubMed Abstract](#) | [Publisher Full Text](#)
 231. Han J, Sun P: The pathways to tumor suppression via route p38. *Trends Biochem Sci.* 2007; **32**(8): 364–371.
[PubMed Abstract](#) | [Publisher Full Text](#)
 232. **F** Freund A, Patil CK, Campisi J: p38MAPK is a novel DNA damage response-independent regulator of the senescence-associated secretory phenotype. *EMBO J.* 2011; **30**(8): 1536–1548.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 233. Zhang B, Fu D, Xu Q, *et al.*: The senescence-associated secretory phenotype is potentiated by feedforward regulatory mechanisms involving Zscan4 and TAK1. *Nat Commun.* 2018; **9**(1): 1723.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 234. Alimbetov D, Davis T, Brook AJ, *et al.*: Suppression of the senescence-associated secretory phenotype (SASP) in human fibroblasts using small molecule inhibitors of p38 MAP kinase and MK2. *Biogerontology.* 2016; **17**(2): 305–315.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 235. Coppe JP, Desprez PY, Krtolica A, *et al.*: The senescence-associated secretory phenotype: the dark side of tumor suppression. *Annu Rev Pathol.* 2010; **5**: 99–118.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 236. Xia Z, Dickens M, Raingeaud J, *et al.*: Opposing effects of ERK and JNK-p38 MAP kinases on apoptosis. *Science.* 1995; **270**(5240): 1326–1331.
[PubMed Abstract](#) | [Publisher Full Text](#)
 237. Kummer JL, Rao PK, Heidenreich KA: Apoptosis induced by withdrawal of trophic factors is mediated by p38 mitogen-activated protein kinase. *J Biol Chem.* 1997; **272**(33): 20490–20494.
[PubMed Abstract](#) | [Publisher Full Text](#)
 238. Cahill MA, Peter ME, Kischkel FC, *et al.*: CD95 (APO-1/Fas) induces activation of SAP kinases downstream of ICE-like proteases. *Oncogene.* 1996; **13**(10): 2087–2096.
[PubMed Abstract](#)
 239. Huang S, Jiang Y, Li Z, *et al.*: Apoptosis signaling pathway in T cells is composed

- of ICE/Ced-3 family proteases and MAP kinase kinase 6b. *Immunity*. 1997; 6(6): 739–749.
[PubMed Abstract](#) | [Publisher Full Text](#)
240. Cardone MH, Salvesen GS, Widmann C, *et al.*: The regulation of anoikis: MEKK-1 activation requires cleavage by caspases. *Cell*. 1997; 90(2): 315–323.
[PubMed Abstract](#) | [Publisher Full Text](#)
241. Tan W, Yu HG, Luo HS: Inhibition of the p38 MAPK pathway sensitizes human gastric cells to doxorubicin treatment *in vitro* and *in vivo*. *Mol Med Rep*. 2014; 10(6): 3275–3281.
[PubMed Abstract](#) | [Publisher Full Text](#)
242. Slawinska-Brych A, Zdzisinska B, Mizerska-Dudka M, *et al.*: Induction of apoptosis in multiple myeloma cells by a statin-thalidomide combination can be enhanced by p38 MAPK inhibition. *Leuk Res*. 2013; 37(5): 586–594.
[PubMed Abstract](#) | [Publisher Full Text](#)
243. Dolgin E: The most popular genes in the human genome. *Nature*. 2017; 551(7681): 427–431.
[PubMed Abstract](#) | [Publisher Full Text](#)
244. Alam MS, Gaida MM, Bergmann F, *et al.*: Selective inhibition of the p38 alternative activation pathway in infiltrating T cells inhibits pancreatic cancer progression. *Nat Med*. 2015; 21(11): 1337–1343.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
245. Luk ST, Ng KY, Zhou L, *et al.*: Deficiency in Embryonic Stem Cell Marker Reduced Expression 1 Activates Mitogen-Activated Protein Kinase Kinase 6-Dependent p38 Mitogen-Activated Protein Kinase Signaling to Drive Hepatocarcinogenesis. *Hepatology*. 2019.
[PubMed Abstract](#) | [Publisher Full Text](#)
246. Salome M, Magee A, Yalla K, *et al.*: A Trib2-p38 axis controls myeloid leukaemia cell cycle and stress response signalling. *Cell Death Dis*. 2018; 9(5): 443.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
247. Gibbs KL, Kalmar B, Rhymes ER, *et al.*: Inhibiting p38 MAPK alpha rescues axonal retrograde transport defects in a mouse model of ALS. *Cell Death Dis*. 2018; 9(6): 596.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
248. Fang C, Wu B, Le NTT, *et al.*: Prions activate a p38 MAPK synaptotoxic signaling pathway. *PLoS Pathog*. 2018; 14(9): e1007283.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
249. Obergasteiger J, Frapporti G, Pramstaller PP, *et al.*: A new hypothesis for Parkinson's disease pathogenesis: GTPase-p38 MAPK signaling and autophagy as convergence points of etiology and genomics. *Mol Neurodegener*. 2018; 13(1): 40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
250. Wang PY, Hsu PI, Wu DC, *et al.*: SUMOs Mediate the Nuclear Transfer of p38 and p-p38 during *Helicobacter Pylori* Infection. *Int J Mol Sci*. 2018; 19(9): 2482.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
251. Zhang F, Zhao Q, Tian J, *et al.*: Effective Pro-Inflammatory Induced Activity of GALT, a Conserved Antigen in *A. Pleuropneumoniae*, Improves the Cytokines Secretion of Macrophage via p38, ERK1/2 and JNK MAPKs Signal Pathway. *Front Cell Infect Microbiol*. 2018; 8: 337.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
252. Zhu J, Yu W, Liu B, *et al.*: Escin induces caspase-dependent apoptosis and autophagy through the ROS/p38 MAPK signalling pathway in human osteosarcoma cells *in vitro* and *in vivo*. *Cell Death Dis*. 2017; 8(10): e3113.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
253. Fan H, Gao Z, Ji K, *et al.*: The *in vitro* and *in vivo* anti-inflammatory effect of osthole, the major natural coumarin from *Cnidium monnieri* (L.) Cuss, via the blocking of the activation of the NF-kappaB and MAPK/p38 pathways. *Phytomedicine*. 2019; 58: 152864.
[PubMed Abstract](#) | [Publisher Full Text](#)
254. Li Y, Xu B, Xu M, *et al.*: 6-Gingerol protects intestinal barrier from ischemia/reperfusion-induced damage via inhibition of p38 MAPK to NF-kappaB signalling. *Pharmacol Res*. 2017; 119: 137–148.
[PubMed Abstract](#) | [Publisher Full Text](#)
255. Zhang X, Wang X, Wu T, *et al.*: Isolensinine induces apoptosis in triple-negative human breast cancer cells through ROS generation and p38 MAPK/JNK activation. *Sci Rep*. 2015; 5: 12579.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
256. Bachegowda L, Morrone K, Winski SL, *et al.*: Pexmetinib: A Novel Dual Inhibitor of Tie2 and p38 MAPK with Efficacy in Preclinical Models of Myelodysplastic Syndromes and Acute Myeloid Leukemia. *Cancer Res*. 2016; 76(16): 4841–4849.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
257. O'Donoghue ML, Glaser R, Aylward PE, *et al.*: Rationale and design of the LosmApimod To Inhibit p38 MAP kinase as a Therapeutic target and modify outcomes after an acute coronary syndrome trial. *Am Heart J*. 2015; 169(5): 622–630.e6.
[PubMed Abstract](#) | [Publisher Full Text](#)
258. Emami H, Vucic E, Subramanian S, *et al.*: The effect of BMS-582949, a P38 mitogen-activated protein kinase (P38 MAPK) inhibitor on arterial inflammation: a multicenter FDG-PET trial. *Atherosclerosis*. 2015; 240(2): 490–496.
[PubMed Abstract](#) | [Publisher Full Text](#)
259. Moran N: p38 kinase inhibitor approved for idiopathic pulmonary fibrosis. *Nat Biotechnol*. 2011; 29(4): 301.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Guadalupe Sabio

Centro Nacional de Investigaciones Cardiovasculares (CNIC), Madrid, Spain

Competing Interests: No competing interests were disclosed.

2. Jonathan D. Ashwell

Laboratory of Immune Cell Biology, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD, 20892, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research