Check for updates

RESEARCH ARTICLE

Bibliometric assessment and implications for practice of sporotrichosis research (1945-2018) [version 1; peer review: 3 approved with reservations]

Priscila Costa Albuquerque¹, Bruna de Paula Fonseca e Fonseca¹, Fabio Zicker¹, Rosely Maria Zancopé-Oliveira², Rodrigo Almeida-Paes ²

¹Center for Technological Development in Health (CDTS), Oswaldo Cruz Foundation, Rio de Janeiro, RJ, 21040361, Brazil

²Mycology Laboratory, Evandro Chagas National Institute of Infectious Diseases (INI), Oswaldo Cruz Foundation, Rio de Janeiro, RJ, 21040900, Brazil

v1 First published: 29 Jun 2020, 9:654
<https://doi.org/10.12688/f1000research.24250.1>

Latest published: 09 Apr 2021, 9:654
<https://doi.org/10.12688/f1000research.24250.2>

Abstract

Background: Sporotrichosis has recently emerged as an important mycosis worldwide, with diverse transmission and epidemiologic profiles. For instance, in Brazil most cases are related to zoonotic transmission from naturally infected cats, and the majority of cases in China are due to external injury with environmental materials. Publications on sporotrichosis and on its etiologic agent may guide the direction of the research in this field. It can also define priorities for future studies.

Methods: In this study, we evaluated the trends of global research in *Sporothrix* and sporotrichosis, based on publications records retrieved from Scopus and Web of Science databases for the period of 1945 to 2018. The overall productivity in the field, its geographical and temporal distribution, research themes, co-authorship networks, funding sources, and the implications of research findings for health practice were assessed using bibliometric approaches.

Results: A total of 4,007 unique publications involving 99 countries were retrieved, most of them published after 2000. Authors based on institutions from the United States of America and Brazil accounted for 57.4% of the publications. Brazil was the leading country in terms of research collaboration and networking, with co-authorship with 45 countries. The thematic mapping revealed a temporal shift from clinical to applied research. Despite the large number of countries publishing in this field, most of funded studies came from Brazil, Mexico, China, South Africa, or the United States of America. The analysis of content identified few specific public health recommendations for prevention, case-management, or research. Moreover, most papers do not have a clearly defined intended audience.

Conclusion: As the research in this field is emerging in several

Open Peer Review

Approval Status

	1	2	3
version 2			
(revision)
09 Apr 2021
version 1			
29 Jun 2020

1. **Max Carlos Ramírez-Soto** , Universidad Peruana Cayetano Heredia, Lima, Peru
2. **Maria Simone Menezes Alencar** , Federal University of the State of Rio de Janeiro, Rio de Janeiro, Brazil
3. **Daniel Zamith-Miranda** , Albert Einstein College of Medicine, New York, USA

Any reports and responses or comments on the article can be found at the end of the article.

countries, with the generation of a large amount of data, it is necessary that scientists strengthen efforts to translate the research results into practice to curb this neglected infection.

Keywords

Bibliometrics, Network, Scientometry, Sporothrix, Sporotrichosis

This article is included in the **Research on Research, Policy & Culture** gateway.

This article is included in the **Neglected Tropical Diseases** collection.

Corresponding author: Rodrigo Almeida-Paes (rodrigo.paes@ini.fiocruz.br)

Author roles: **Albuquerque PC:** Conceptualization, Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Fonseca e Fonseca BdP:** Funding Acquisition, Methodology, Writing – Review & Editing; **Zicker F:** Funding Acquisition, Methodology, Resources, Writing – Review & Editing; **Zancopé-Oliveira RM:** Conceptualization, Funding Acquisition, Supervision, Writing – Review & Editing; **Almeida-Paes R:** Conceptualization, Formal Analysis, Investigation, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: Software licenses were funded by the Instituto Nacional de Ciência e Tecnologia de Inovação em Doenças de Populações Negligenciadas (INCT-IDPN). RMZ-O was supported in part by Conselho Nacional de Desenvolvimento Científico e Tecnológico [CNPq 302796/2017-7] and Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro [FAPERJ E-26/202.527/2019]. This work was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2020 Albuquerque PC *et al.* This is an open access article distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Albuquerque PC, Fonseca e Fonseca BdP, Zicker F *et al.* **Bibliometric assessment and implications for practice of sporotrichosis research (1945-2018) [version 1; peer review: 3 approved with reservations]** F1000Research 2020, 9:654 <https://doi.org/10.12688/f1000research.24250.1>

First published: 29 Jun 2020, 9:654 <https://doi.org/10.12688/f1000research.24250.1>

Introduction

Sporotrichosis is a subcutaneous mycotic infection caused by dimorphic species of fungi belonging to the genus *Sporothrix*¹. It has a worldwide distribution, a broad range of clinical presentations, and can be fatal as an opportunistic infection in immunosuppressed patients². For more than one century, the etiological agent of sporotrichosis was identified as the sole species *Sporothrix schenckii*³. However, in last years, using molecular biology techniques, it was possible to identify other sibling species that also cause sporotrichosis: *Sporothrix brasiliensis*, *Sporothrix globosa*, and, to a lesser extent, *Sporothrix luriei*, *Sporothrix pallida*, *Sporothrix mexicana*, and *Sporothrix chilensis*^{4–7}. These agents can be found in the environment¹ and they present different clinical manifestations⁸, virulence⁹, drug susceptibility¹⁰, and phenotypic characteristics¹¹.

The research about fungal diseases has been relatively neglected worldwide by public health authorities. The initiative “Global Action Fund for Fungal Infections” (GAFFI), an international non-governmental organization dedicated to combating fungal disease, was created to highlight gaps in diagnostics and treatments for fungal diseases as well as to fund raise and lobby global health agencies¹². GAFFI has identified some fungal infections as its highest priorities and sporotrichosis, along with other deep mycoses, is included¹³. This organization has claimed that sporotrichosis, paracoccidioidomycosis, and fungal keratitis should be included in the WHO’s Neglected Tropical Diseases portfolio. This would be a big step towards increasing research funding and better care for patients with these serious diseases.

Bibliometric studies are frequently used to describe the global dynamics of knowledge generation and provide useful information on research discoveries, pointing at the strengths and weakness of new findings¹⁴. Through bibliometric analyses, based on quantitative and qualitative indicators, it is possible to assess progress and collaboration in science, technology and innovation¹⁵. For instance, the co-authorship of scientific publications reveals collaborative patterns between individuals, organizations, or countries¹⁶ and represents a formal statement of interaction between two or more researchers, being widely used to understand and assess collaboration profiles^{17,18}. Bibliometric methods, however, have rarely been applied to the mycology field and not much is known about the extension and trends of *Sporothrix* research. Therefore, an assessment of the characteristics of the research focusing on sporotrichosis and its etiological agents is necessary to evaluate the progress of the findings in this field and the implications and impacts of sporotrichosis research for health practice.

Driven by the continued expansion of sporotrichosis in some countries such as Brazil, China, Mexico, and India¹, we analyzed the global scientific publications and scientific collaboration on the *Sporothrix* and sporotrichosis research field, with emphasis on the impact of the zoonotic epidemic of sporotrichosis in Brazil² and on the discovery of the new species genetically related to *S. schenckii*⁵. We combined bibliometrics and social network analysis to generate evidence of the dynamics of the research community. Also, implications for practice of the

publications were assessed, to check whether the research results were being translated into actions to curb sporotrichosis.

Methods

Source of data

Publications were retrieved from the [Web of Science](#) (WoS) and [Scopus](#) databases searching for the terms “*Sporothrix*” or “sporotrichosis” on the title, abstract, and keyword fields. The review included all scientific publications records from 1945 to 2018 (WoS Core Collection). For comparison purposes, the number of publications about other mycoses caused by dimorphic fungi (paracoccidioidomycosis, histoplasmosis, coccidioidomycosis, and blastomycosis) was also obtained, using the genus of the fungus and the name of the mycosis as the search query terms.

Cleaning and standardization of data

The data retrieved was imported into the text mining software [VantagePoint](#) 10.0 (Search Technology Inc. Norcross, GA, USA) and duplicate records were excluded. Names of institutions, countries, funding organizations, and journals were standardized using the VantagePoint list cleanup tool, and further manual processing. An open-access alternative for the use of VantagePoint 10.0 is the software [OpenRefine](#) 3.3.

Co-authorship network analyses

After cleaning and standardization, the data was formatted into adjacency matrixes to map co-authorship between countries and institutions based on authors’ professional affiliations. The matrixes were imported into the open-source software [Gephi](#) 0.9.1 for network visualization and calculation of metrics¹⁹. Degree centrality was used to identify the most central institutions in the network, reflecting the significance of a network member (node) relative to all other nodes in the network. This metric takes into account the diverse means in which a node interacts and communicates with the rest of the network. The most important, or central ones, have a strategic impact in the network. The degree centrality can be explained as the number of direct links that a node has with other nodes. The more relational ties a node has, more power or prestige it may present in a network²⁰. Betweenness centrality was used to recognize organizations that mediated the connection between other institutions and their capacity to control the flow of information in the network²⁰. This metric reveals the extent to which a node works as a bridge among the other nodes in the network, which would otherwise be disconnected. For the spatial visualization of international collaboration, the authors’ professional affiliation country was manually geocoded and processed using the “[GeoLayout](#)” 0.9.1.2 and “[Map of Countries](#)” 1.5.1 plugins available within Gephi. In these networks, nodes represent countries or an institution, and two or more countries/institutions were connected if their researchers shared the authorship of one or more papers. As co-authorship involves reciprocal collaboration, all connections were considered as non-directional.

Mapping research themes

Term maps were created using the [VOSviewer](#) 1.6.6 software (Leiden University, the Netherlands) using terms obtained from

titles and abstracts of all publications in the database. Each term was graphically denoted by a circle whose diameter and label size were directly proportional to their frequency. The software positions the circles closer to each other according to the power of the relationship and co-occurrence between terms. The mapping allowed a cluster analysis by research themes using a weighted and parameterized variant of modularity-based clustering to recognize groups of correlated terms²¹.

Funding data

Funding acknowledgments on publications were only available as a searchable field in WoS and Scopus from late 2008. In order to achieve reliable coverage, only papers published from 2012 onwards were selected for this purpose. Funding agencies were identified, their names standardized (whenever possible), and the number of publications per funding agency summarized.

Implications for practice

Abstracts of the publications were reviewed to identify statements or recommendations for sporotrichosis research, case-management, and prevention. Titles and abstracts were tabulated on a Microsoft Excel 2010 spreadsheet, and searched for terms related to statements and recommendations for research, practice or public health. A set of words related to tentative language (“may”, “might”, “speculate”, “suggest”, or “potentially”), prescriptive language (“must”, “propose”, “should”, “stress”, or “recommend”) or related to minimal advice (“consider”, “advise”, “notify”, or “inform”) was used in this process²². The audiences

to whom the recommendations were directed to (medical doctors, nurses, laboratory staff, or veterinarians) were also identified.

Statistical analysis

The software **GraphPad Prism 5** was used to build linear regressions, to check the frequency of publications over time and to compare slopes of different best-fit lines. The Chi-square test was used to test for differences in proportions of “tentative”, “prescriptive”, and “minimal advice” from research results according to the country of origin in an attempt to identify how explicit was the research message to the scientific community, practitioners and public health professionals. A $p < 0.05$ was considered to be significant.

Results

The scientific literature on *Sporothrix* and sporotrichosis

Sporotrichosis was found to be the mycosis caused by dimorphic fungi with the lowest number of publications in both databases used in this work (**Table 1**). The literature search on the *Sporothrix*/sporotrichosis field retrieved 1,868 publications from the WoS database and 3,660 publications from the Scopus database for the period of 1945 to 2018. After removing duplicate references present on both databases, new totals were 1,866 and 3,599 papers, respectively, with 1,458 publications present in both databases. The final data set resulted in 4,007 unique publications (**Figure 1**; Underlying data²³). As depicted in **Figure 2a**, the overall number of publications on *Sporothrix*/sporotrichosis has increased steadily over the years. The

Table 1. Number of publications on dimorphic fungi and their respective mycoses retrieved from Web of Science and Scopus databases (1945 – 2018).

Fungus / Disease	Number of publications per database	
	Web of Science	Scopus
<i>Sporothrix</i> / Sporotrichosis	1,868	3,660
<i>Paracoccidioides</i> / Paracoccidioidomycosis	2,747	3,692
<i>Histoplasma</i> / Histoplasmosis	7,111	11,520
<i>Blastomyces</i> / Blastomycosis	2,513	6,491
<i>Coccidioides</i> / Coccidiomycosis	3,312	5,786

Figure 1. Process of publication acquisition in the *Sporothrix*/sporotrichosis field from the studied databases.

Figure 2. Global scientific production in the *Sporothrix* and *Sporotrichosis* field. (a) Overall publication numbers per year (1945–2018). Trends in the number of publications on the *Sporothrix*/sporotrichosis field are represented in a linear regression form for the following time periods: (b) 1945–1962; (c) 1963–1991; (d) 1992–2002; (e) 2003–2018. The continuous line represents the best-fit line for the linear regression. Dashed lines represent the 95% confidence interval of the best-fit line.

frequency of publications reflecting the interest in the studied subject varied over time. Four periods could be identified: 1945 to 1962 (Figure 2b), with an average of 10.8 ± 3.7 publications per year; 1963 to 1991 (Figure 2c), 47.2 ± 11.3 papers per year; 1992 to 2002 (Figure 2d), with 63.1 ± 10.3 publications per year; and in the final period, 2003 to 2018 (Figure 2e), 108.9 ± 20.3 papers per year. In two of these periods, 1945–1962 and 1992–2002, the amount of publications per year was approximately constant (slopes of 0.2157 ± 0.1646 and -0.1364 ± 0.9980 , p values of 0.2085 and 0.8943, respectively), whereas the two remaining periods, 1963–1991 and 2003–2018, presented increases in the publication numbers per year (slopes of 0.7113 ± 0.2095 and 3.494 ± 0.6573 , p values of 0.0021 and 0.0001, respectively). The differences between the slopes were found to be extremely significant ($p < 0.0001$).

Authorship by country

Authors from 99 countries accounted for the 4,007 publications on the *Sporothrix*/sporotrichosis field. Table 2 lists the 20 most productive countries during the entire period studied and Figure 3 depicts the annual trends of the top 15 countries. The United States of America (USA) and Brazil were the leading countries, publishing together 2,300 publications (57.4% of all publications). The frequency of publications by these two countries differed considerably. After 1973, the USA showed a regular pattern of publications with a range of 15–44 and a median of 25 papers per year, while Brazil had up to 10 publications per year until 1999, with an exponential increase in the number of publications noticed after 2000. Overall, the frequency of publications from the Japan, France, and Canada remained stable throughout the studied period, whereas a significant increase was also observed for Mexico from 2007 onwards, and China from 2009 onwards.

Scientific journals

Table 3 lists the top 15 journals (out of 1,182) with the highest number of articles on *Sporothrix*/sporotrichosis, along with their respective impact factors. Overall, the journals with most publications were Mycopathologia, Medical Mycology, Mycoses, the International Journal of Dermatology, and JAMA Dermatology. Since Brazil is the current leader on annual publications in this field (Figure 3), the journals harboring these publications were also evaluated separately. When only publications with at least one Brazilian author were considered, in addition to Mycopathologia, Medical Mycology, Mycoses, and the International Journal of Dermatology, the journals PLoS Neglected Tropical Diseases, Frontiers in Microbiology, and four Brazilian scientific journals also appear listed among the top ten journals.

Research trends

The analysis of term maps revealed five broad knowledge areas of the *Sporothrix*/sporotrichosis research (Figure 4a): description of clinical aspects (green), treatment (yellow), epidemiology and taxonomy (blue), cellular biology (red), and susceptibility assays (pink). The most frequent topics throughout the period studied are represented in a heat map format (Figure 4b), which shows that terms about clinical aspects were more frequent, followed by terms on taxonomy and cellular

biology. The analysis in two time periods - 1945–1999 (Figure 4c) and 2000–2018 (Figure 4d) revealed, in this last period of highly increasing publication numbers, a shift from research focused on treatment to other knowledge areas such as epidemiology and taxonomy.

Network of countries

The international research network on *Sporothrix* and sporotrichosis was mapped according to the country affiliation of all authors (Figure 5a). The network is formed by 99 countries, with a Brazilian leadership on collaborations. In fact, Brazilian authors have co-authorship with authors from 45 of the countries that compose the network (Figure 5b). The most frequent Brazilian collaborations were with institutions from the USA (84 publications), followed by the Netherlands (44 publications), Mexico (31 publications), Spain (23 publications), and France (16 publications).

Network evolution among Brazilian institutions

To further understand the dynamics of the *Sporothrix*/sporotrichosis research in the current leader country of publications, the evolution of *Sporothrix* research networks in Brazil was analyzed in the two-time intervals, 1945–1999 and 2000–2018, that is, before and after the beginning of the Brazilian zoonotic

Table 2. Top 20 countries in the *Sporothrix*/sporotrichosis field (1945–2018) according to authors professional affiliation.

Rank	Country	Number of publications
1	United States of America	1,315
2	Brazil	985
3	Japan	417
4	India	309
5	Mexico	275
6	China	190
7	South Africa	155
8	France	137
9	Spain	135
10	Canada	130
11	The Netherlands	106
12	Germany	105
13	United Kingdom	96
14	Italy	87
15	Australia	74
16	Peru	64
17	Venezuela	55
18	Poland	49
18	South Korea	49
19	Colombia	46
20	Greece	44

Figure 3. Annual publication from the top 15 countries on numbers of articles in the *Sporothrix/sporotrichosis* field. The diameter of each circle is directly proportional to the annual number of publications. USA: United States of America; BRA: Brazil; JPN: Japan; IND: India; MEX: Mexico; CHN: China; SAF: South Africa; FRA: France; SPN: Spain; CAN: Canada; GER: Germany; NTL: The Netherlands; GBR: Great Britain; ITA: Italy; AUS: Australia.

Table 3. Top scientific journals publishing articles on the *Sporothrix*/sporotrichosis field (1945–2018).

Journal	Country	JIF Trend 2017/18	All Publications		Brazilian Publications	
			Rank	Papers (n)	Rank	Papers (n)
Mycopathologia	Netherlands	1.476	1	283	2	68
Medical Mycology	United Kingdom	2.799	2	260	1	93
Mycoses	Germany	2.793	3	141	4	33
International Journal of Dermatology	USA	1.541	4	120	10	14
JAMA Dermatology	USA	8.107	5	96	N/A	0
Clinical Infectious Diseases	USA	9.117	6	80	13	11
Journal of Clinical Microbiology	USA	4.054	7	69	13	11
Journal of the American Academy of Dermatology	USA	6.898	8	57	21	3
Revista Iberoamericana de Micologia	Spain	0.989	9	51	14	10
Anais Brasileiros de Dermatologia	Brazil	0.884	10	50	3	37
Revista da Sociedade Brasileira de Medicina Tropical	Brazil	1.358	22	26	5	26
PLoS Neglected Tropical Diseases	USA	4.367	24	19	6	19
Frontiers in Microbiology	Switzerland	4.019	23	18	7	18
Memórias do Instituto Oswaldo Cruz	Brazil	2.833	21	16	8	16
Revista do Instituto de Medicina Tropical de São Paulo	Brazil	1.489	26	15	9	15

JIF – Journal Impact Factor

N/A – Not available

endemic of sporotrichosis (Figure 6). In total, 29 institutions, mostly from the Southeast region, were present in the first period. The network core (giant component) was formed by 16 institutions: nine universities, one public foundation, and one private laboratory, all from the Southeast region, two universities and one hospital from the South region, one university from the Midwest region and one national research center. The network degree average was 1.5, indicating low connectivity in the network (Figure 6a). The first three institutions with higher degree centrality in that period were federal universities located in the Brazilian Southeast region (Table 4). For the second period, the number of institutions increased almost six times. The network was composed by 169 institutions, 135 of them within the giant component. The degree average increased four times (degree average: 4.426), showing a gain in the network connectivity (Figure 6b). A national research center appears as the most central institution in the network, followed by three federal universities, central actors in the sporotrichosis network throughout the studied period. The ten most central organizations in these networks are shown in Table 4. Notably, most of them are from the Southeast region of the country (70% and 60% for 1945–1999 and 2000–2018, respectively).

Research funding

The analysis of funding acknowledgments was used as proxy information for research funding. A total of 457 (34%) articles

out of 1,310 publications acknowledged funding during the period 2012 to 2018. Table 5 shows the top 10 funding agencies and the number of publications out of a total 236 funding organizations and initiatives. Five of them are from Brazil, two from Mexico, one from China, one from the USA, and one from South Africa.

Statements about implications for practice

After excluding 1,262 articles with no published abstract, the type of language and the intended audience of the publications were assessed for 2,745 articles (Table 6). Since Brazil was the current leading country publishing in the *Sporothrix*/sporotrichosis field, we compared the frequency of statements related to research and practice on papers with at least one Brazilian author with those of authors from other countries exclusively. The proportion of papers presenting language compatible with tentative ($p = 0.0003$) or minimal advice ($p = 0.0139$) statements for sporotrichosis research was higher in the papers co-authored by Brazilians. The specific audience for the statements was mentioned in 167 abstracts. As depicted in Figure 7, most of the implications for practice were directed to medical doctors, followed by veterinarians, nurses, and laboratory technicians. Only 16 publications presented more than one audience, as indicated at the abstract. Some examples of actionable messages and their audiences are listed in Table 7.

Figure 4. Thematic maps of *Sporothrix/sporotrichosis* research generated with the VOSviewer software. (a) Terms extracted from titles and abstracts clustered into five major research areas: cellular biology, epidemiology and taxonomy, clinical aspects, treatment, and susceptibility assays. Colors indicate clusters of terms that have co-occurred more frequently in the dataset. **(b)** The heat map shows the most frequent terms in the period analyzed. The frequency was graded from blue to red; where red indicates a higher frequency. **(c)** Research trends from 1945 to 1999. **(d)** Research trends from 2000 to 2018. The diameters of the circles on **a**, **c**, and **d** panels are directly proportional to the occurrence of each term. Lines between different circles represent relationships between terms. The colors in **c** and **d** thematic maps indicate the occurrence of a term in each period. Blue represents a low occurrence, green an average occurrence, and red a high occurrence.

Figure 5. Co-authorship map between *Sporothrix*/sporotrichosis researchers. (a) Global research network among 99 countries publishing in the *Sporothrix*/sporotrichosis field from 1945–2018. (b) Brazilian network of scientific collaborations on the period studied. Country links were mapped based on the authors' affiliations. Each node represents one country and two countries were considered connected if their researchers shared the authorship of a paper. Only relationships between the first author and their co-authors are shown. Links are color-coded according to the continent of the first author: North America – blue; Africa – dark green; Europe – yellow; South America – light green; Asia – red; Oceania – pink.

Figure 6. Evolution of *Sporothrix*/sporotrichosis research networks involving Brazilian institutions. (a) Research network from 1945 to 1999. (b) Research network from 2000 to 2018. Each node represents one institution and two institutions were considered connected if their members shared the authorship of a paper. Nodes are color coded according to the geographic location of the institutions: North region - green; Northeast region - orange; Center-West region- yellow; Southeast region- blue; and South region - lilac. Multicampus National institutions are colored in gray. The size of the nodes is proportional to their centrality degree. For visualization purposes, only the giant component is shown. The top ten Brazilian organizations with highest degree centrality are labeled in each panel.

Table 4. Centrality Index of institutions in the Brazilian sporotrichosis collaboration network.

Rank	Period I (1945–1999)				Period II (2000–2017)			
	Institutions	Degree	Betweenness centrality	Coauthored Publications	Institutions	Degree	Betweenness centrality	Coauthored Publications
1	UN1 (UFRJ)	6	58.0	38	RC 1 (Fiocruz)	44	3,041.99	313
2	UN2 (USP)	6	57.0	26	UN 3 (UNIFESP)	43	2,092.34	133
3	UN3 (UNIFESP)	4	15.0	16	UN 1 (UFRJ)	37	1,810.33	115
4	UN4 (UNESP)	4	7.0	9	UN 2 (USP)	32	797.79	93
5	UN5 (UFSM)	3	26.0	6	UN 8 (UFMG)	22	851.92	34
6	UN6 (UERJ)	3	14.0	10	UN 10 (UFRGS)	21	1,555.58	85
7	UN7 (UFF)	3	0.0	4	UN 9 (UNB)	18	694.66	27
8	UN8 (UFMG)	2	14.0	10	UN 19 (UFPR)	18	650.58	21
9	UN9 (UNB)	2	26.0	2	UN 6 (UERJ)	18	394.06	105
10	UN10 (UFRGS)	2	0.0	2	UN 4 (UNESP)	16	766.62	75

UFRJ: Universidade Federal do Rio de Janeiro, Southeast Brazil; USP: Universidade de São Paulo, Southeast Brazil; UNIFESP: Universidade Federal de São Paulo, Southeast Brazil; UNESP: Universidade Estadual Paulista, Southeast Brazil; UFSM: Universidade Federal de Santa Maria, South Brazil; UERJ: Universidade do Estado do Rio de Janeiro, Southeast Brazil; UFF: Universidade Federal Fluminense, Southeast Brazil; UFMG: Universidade Federal de Minas Gerais, Southeast Brazil; UNB: Universidade de Brasília, Center Western Brazil; UFRGS: Universidade Federal do Rio Grande do Sul, South Brazil; UFPR: Universidade Federal do Paraná, South Brazil; Fiocruz: Fundação Oswaldo Cruz, Brazil.

Table 5. Top 10 funding organizations supporting publications on *Sporothrix*/sporotrichosis (2012 – 2018).

Rank	Major Funding Organization	Number of funded publications (%)	Country
1	CNPq - Conselho Nacional de Desenvolvimento Científico e Tecnológico	145 (11)	Brazil
2	CAPES - Coordenação de Aperfeiçoamento de Pessoal de Nível Superior	84 (6.4)	Brazil
3	FAPERJ - Fundação Carlos Chagas de Amparo à Pesquisa do Estado do Rio de Janeiro	64 (4.8)	Brazil
4	FAPESP - Fundação de Amparo à Pesquisa do Estado de São Paulo	61 (4.6)	Brazil
5	National Natural Science Foundation of China	36 (2.7)	China
6	UG - Universidad de Guanajuato	34 (2.6)	Mexico
7	CONACyT - Consejo Nacional de Ciencia y Tecnología	33 (2.5)	Mexico
8	FIOCRUZ - Fundação Oswaldo Cruz	32 (2.4)	Brazil
9	NIH - National Institutes of Health	22 (1.6)	United States of America
10	DST/NRF Centre of Excellence in Tree Health Biotechnology (CTHB)	15 (1.1)	South Africa

Table 6. Type of language statements and identified audience from *Sporothrix* and sporotrichosis articles (1945–2018).

Statements with	Brazil	Other countries	p-value
Prescriptive language	52 (9.5%)	226 (10.3%)	0.6576
Tentative language	195 (35.7%)	613 (27.9%)	0.0003
Minimal advice	78 (14.3%)	230 (10.4%)	0.0139
Audience specified	37 (6.8%)	122 (5.5%)	0.2714

Figure 7. Identification of specified audiences in publications about *Sporothrix* or sporotrichosis. The Venn diagram was constructed based on four different audiences observed in the abstracts of the publications studied. Numbers on diagrams indicate how many abstracts reporting advice for medical doctors, veterinary doctors, nurses, and/or lab technicians were identified.

Discussion

The results herein presented demonstrate that research on *Sporothrix* and sporotrichosis is increasing worldwide. This increase could be even greater, because some regional or specialized journals may possibly not be indexed in the databases reviewed. However, considering the proposed focus, the authors believe that the study material was a comprehensive representation of the scientific production in the field.

The apparent fast engagement of authors from several countries in the late 90s can be explained by the global increase in the incidence of sporotrichosis¹, especially in Brazil² and China²⁴, two of the major countries currently publishing on this field. Historically, Brazil and China are also the leading countries reporting sporotrichosis cases, followed by South Africa, all of them with more than 3,000 human and animal cases reported²⁵. In this study, South Africa ranked as the seventh country on number of publications with no increasing trend. Most of the cases in this country occurred during an outbreak in the 1940's²⁶. Although a new mine-related outbreak occurred more recently²⁷, there is no evidence that there is a re-emergence of sporotrichosis occurring in South Africa, which may explain its apparently constant number of publications in this field.

The number of publications on the *Sporothrix*/sporotrichosis field usually followed crucial events that have driven to a shift on typical clinical cases of sporotrichosis. In the 1980s, with the development of ketoconazole, the first oral antifungal azole drug²⁸, and the emergence of AIDS, there was an increase in papers reporting results of sporotrichosis treatment^{29,30} and unusual severe forms of sporotrichosis related to immunosuppression^{31–33}, coherent with the predominant thematic at the time. After 2007, the advances in polyphasic taxonomy of

Table 7. Examples of actionable messages found in article abstracts from papers evaluated in this study.

Title of publication	PMID	Type of language	Audience Specified	actionable message
Histopathology of canine sporotrichosis: A morphological study of 86 cases from Rio de Janeiro	19360480	Prescriptive	No	Specific staining of serial sections is recommended in the case of dogs with skin lesions whose histopathological presentation is consistent with sporotrichosis.
Sporotrichosis in cats: ABCD guidelines on prevention and management	23813827	Prescriptive	Yes	Professionals must wear gloves when handling cats with skin nodules and ulcers and dealing with diagnostic samples
Molecular cloning, characterization and differential expression of DRK1 in <i>Sporothrix schenckii</i>	23175272	Tentative	No	SsDRK1 may be involved in the dimorphic switch in <i>S. schenckii</i> .
Disseminated sporotrichosis with extensive cutaneous involvement in a patient with AIDS	10025867	Tentative	Yes	It is important that clinicians be aware of the presentation of this unusual opportunistic infection
Sporotrichosis in Himachal Pradesh (north India)	10492787	Minimal advice	No	This study identifies Kangra district and adjoining areas in Himachal Pradesh as an endemic region for sporotrichosis and highlights the need for evaluation of geo-climatically similar areas.
Unsuspected sporotrichosis in childhood	11332673	Minimal advice	Yes	We urge clinicians to consider sporotrichosis in the differential diagnosis of a solitary skin nodule.

Sporothrix spp., driven by the worldwide increasing numbers of clinical cases⁵, resulted in the description of several new pathogenic *Sporothrix* species⁵⁻⁷, which may explain the shift for epidemiology and taxonomy occurred in this century.

Authors from Brazil and the USA authored around 57% of the papers on the subject studied. Also, a high-level of collaboration between these two countries was seen, as occurred in other knowledge areas^{34,35}. While the number of publications authored by researchers from the USA showed a consistent trend in the studied period, Brazilian authors have emerged as very productive in this field. Some factors that may have influenced the strong commitment of Brazilian researchers in the *Sporothrix*/sporotrichosis field include: (i) the zoonotic sporotrichosis epidemic, that begun in Rio de Janeiro state in 2000³⁶ and now is spreading to almost the totality of the Brazilian territory³⁷⁻³⁹; (ii) the increasing numbers of national and international Brazilian collaboration networks⁴⁰; and (iii) the beginning and still discrete recognition of the relevance of this research field by Brazilian funding agencies, which was detected in the funding analysis of this study.

The evaluation of co-authorship networks in sporotrichosis identified structural patterns of research involving Brazilian scientists. USA, Netherlands, Mexico, Spain, and France were Brazil's most frequent collaborators. Government collaboration and research programs supported by the USA are in place since 1950's, explaining the robust scientific collaboration between the countries³⁴.

The Brazilian sporotrichosis research network has grown, almost six times in size, from the first (initial 54 years of low productivity) to the second period (last 18 years with a high level of publications), over the seven decades evaluated. This fact, together with the increase in the average degree and size of the giant component, indicated a strengthening of network cohesion for collaboration over the years. It is worth noting that the measure of centrality is not related to the volume of publications, but to the capacity to aggregate collaboration. The centrality analysis of the interinstitutional network highlighted the current role of one national research center in promoting collaboration and knowledge spreading in the sporotrichosis field. This same research center also has a centrality on the research of other infectious diseases^{15,41}. High degree centrality from a research center indicated a strong collaborative pattern in research¹⁵. Together with other institutions, the Federal Universities in Southeast Brazil had a vital role in maintaining the connection between the overall research network and in ensuring that the less well-connected organizations gained access to new knowledge and information on sporotrichosis during the period studied. It is interesting to note that just recently, sporotrichosis has spread to the Northeast³⁹ and Center-Western regions of the country³⁸, which means that, in the future, institutions from these regions may have a more important role in the collaboration network.

The network evolution was accompanied by a shift in the research trends. When comparing the first period to the 2000 onwards, the basic biomedical research profile gained more importance and became most frequent. The diversity of research

trends may be related to the continued increase of institutions engagement in the *Sporothrix*/sporotrichosis network, providing new insights through new collaborations, showing the effectiveness of the research network in knowledge generation, sharing and diffusion. A similar scenario was observed in dengue research networks⁴².

The current funding scenario for fungal infections has a negative perspective. Only cryptococcal meningitis was classified within the most poorly funded neglected diseases by the G-Finder survey (a reference publication for research funding flows on neglected infectious), receiving less than 0.5% of global funding⁴³. Other clinically relevant fungal infections (paracoccidioidomycosis, mycetoma, sporotrichosis, and chromoblastomycosis) were not even quoted in the G-Finder report. In fact, a recent study noticed that some fungal diseases, including sporotrichosis, have received negligible funding⁴⁴. Our study corroborates these findings, showing that, with the exception of Brazil, possibly because of the expanding sporotrichosis endemic areas³⁷⁻³⁹, other countries are not strongly committed to sporotrichosis funding research.

In our analysis, publications co-authored by Brazilians researchers have more tentative and minimal advice or recommendations for practice than papers written by authors from other countries. This could be related to a concern of Brazilian researchers to address the major public health problem that sporotrichosis has become in the country^{2,37}, which has certainly impacted the overall number of publications. The review based solely on abstracts is a limitation of this study. The fact that some journals have a specific audience (for instance, four of the top ten journals are directed to dermatologists) may bias the nature of recommendations. However, as abstracts are the first contact for the readers with a publication, we believe mentioning statements or recommendations for practice in this section can improve the visibility of the research. Also, the analysis of contents revealed that a minor proportion of publications is directed to more than one specific audience, which is one indicator of a poor translational research in the studied field.

Despite cases being reported all around the globe¹, sporotrichosis remains a neglected disease in terms of research interest, funding and medical attention. The growth of research on sporotrichosis needs to be translated into practical applications on diagnosis, treatment, and prevention, given the limited tools available for rapid tests², the cost-effective treatment⁴⁵, and the lack of effective vaccines⁴⁶. The challenge is to share and advance knowledge to curb this disease. The funding agencies have a critical role to play in this context.

Data availability

Underlying data generated in this study are available at Open Science Framework: Bibliometric assessment of sporotrichosis research. <https://doi.org/10.17605/OSF.IO/MXU6V>²³

This project contains the following underlying data:

- (F1_T1) Countries publication by years.xlsx (All countries with sporotrichosis publications)
- (F2) MapVoS.txt (Map file for VOSViewer software)

- (F2) NetVoS.txt (Network file for VOSViewer software)
- (F3) CountryNet.gephi (Country map file for Gephi software)
- (F4) Period I.gephi (Network file for Gephi Software)
- (F4) Period II.gephi (Network file for Gephi Software)
- (T2) Source.xlsx (All Journals with retrieved publications about sporotrichosis)

- (T3) Funding BR 12_18.xlsx (Funding agencies supporting sporotrichosis research)
- (T4) Actionable messages.xlsx (implications for practice on retrieved articles)

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC0 1.0 Public domain dedication).

References

1. Chakrabarti A, Bonifaz A, Gutierrez-Galhardo MC, *et al.*: **Global epidemiology of sporotrichosis.** *Med Mycol.* 2015; **53**(1): 3–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Orofino-Costa R, Macedo PM de, Rodrigues AM, *et al.*: **Sporotrichosis: an update on epidemiology, etiopathogenesis, laboratory and clinical therapeutics.** *An Bras Dermatol.* 2017; **92**(5): 606–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. Lopes-Bezerra LM, Schubach A, Costa RO: ***Sporothrix schenckii* and sporotrichosis.** *An Acad Bras Cienc.* 2006; **78**(2): 293–308.
[Reference Source](#)
4. Morrison AS, Lockhart SR, Bromley JG, *et al.*: **An environmental *Sporothrix* as a cause of corneal ulcer.** *Med Mycol Case Rep.* 2013; **2**: 88–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Marimon R, Cano J, Gené J, *et al.*: ***Sporothrix brasiliensis*, *S. globosa*, and *S. mexicana*, three new *Sporothrix* species of clinical interest.** *J Clin Microbiol.* 2007; **45**(10): 3198–206.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
6. Marimon R, Gené J, Cano J, *et al.*: ***Sporothrix luriei*: a rare fungus from clinical origin.** *Med Mycol.* 2008; **46**(6): 621–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Rodrigues AM, Cruz Choappa R, Fernandes GF, *et al.*: ***Sporothrix chilensis* sp. nov. (Ascomycota: Ophiostomatales), a soil-borne agent of human sporotrichosis with mild-pathogenic potential to mammals.** *Fungal Biol.* 2016; **120**(2): 246–64.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Almeida-Paes R, de Oliveira MME, Freitas DFS, *et al.*: **Sporotrichosis in Rio de Janeiro, Brazil: *Sporothrix brasiliensis* is associated with atypical clinical presentations.** *PLoS Negl Trop Dis.* 2014; **8**(9): e3094.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. Arrillaga-Moncrieff I, Capilla J, Mayayo E, *et al.*: **Different virulence levels of the species of *Sporothrix* in a murine model.** *Clin Microbiol Infect.* 2009; **15**(7): 651–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Marimon R, Serena C, Gené J, *et al.*: ***In vitro* antifungal susceptibilities of five species of *Sporothrix*.** *Antimicrob Agents Chemother.* 2008; **52**(2): 732–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Almeida-Paes R, de Oliveira LC, Oliveira MME, *et al.*: **Phenotypic characteristics associated with virulence of clinical isolates from the *Sporothrix* complex.** *BioMed Res Int.* 2015; **2015**: 212308.
[Publisher Full Text](#)
12. Global Action Fund for Fungal Infections: **95–95 by 2025. Improving outcomes for patients with fungal infections across the world: a roadmap for the next decade.** 2015.
[Reference Source](#)
13. Global Action Fund for Fungal Infections: **Gaffi - fact sheet: sporotrichosis.** [cited 2019 Aug 14]. 2018.
[Reference Source](#)
14. Joshi MA: **Bibliometric indicators for evaluating the quality of scientific publications.** *J Contemp Dent Pract.* 2014; **15**(2): 258–62.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Bruna de Paula F, da Silva MV, de Araújo KM, *et al.*: **Network analysis for science and technology management: Evidence from tuberculosis research in Fiocruz, Brazil.** *PLoS One.* 2017; **12**(8): e0181870.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Melin G, Persson O: **Studying research collaboration using co-authorships.** *Scientometrics.* 1996; **36**(3): 363–77.
[Publisher Full Text](#)
17. Fonseca BPF, Sampaio RB, Fonseca MVA, *et al.*: **Co-authorship network analysis in health research: method and potential use.** *Health Res Policy Syst.* 2016; **14**(1): 34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Sampaio RB, Fonseca BPF, Bahulkar A, *et al.*: **Network analysis to support public health: evolution of collaboration among leishmaniasis researchers.** *Scientometrics.* 2017; **111**(3): 2001–21.
[Publisher Full Text](#)
19. Bastian M, Heymann S, Jacomy M: **Gephi: An Open Source Software for Exploring and Manipulating Networks.** 2.
[Reference Source](#)
20. Freeman LC: **Centrality in social networks conceptual clarification.** *Soc Netw.* 1978; **1**(3): 215–39.
[Publisher Full Text](#)
21. Waltman L, van Eck NJ, Noyons ECM: **A unified approach to mapping and clustering of bibliometric networks.** *J Informetr.* 2010; **4**(4): 629–35.
[Publisher Full Text](#)
22. Lynn J, Owens AP, Bartunek JM: **Clarity and strength of implications for practice in medical journal articles: an exploratory analysis.** *BMJ Qual Saf.* 2011; **20**(Suppl 1): i52–57.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Albuquerque PC, Almeida-Paes R: **Bibliometric assessment of sporotrichosis research.** 2020.
<http://www.doi.org/10.17605/OSF.IO/MXU6V>
24. Chen M, Xu Y, Hong N, *et al.*: **Epidemiology of fungal infections in China.** *Front Med.* 2018; **12**(1): 58–75.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Zhang Y, Hagen F, Stielow B, *et al.*: **Phylogeography and evolutionary patterns in *Sporothrix* spanning more than 14 000 human and animal case reports.** *Persoonia.* 2015; **35**: 1–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Quintal D: **Sporotrichosis infection on mines of the Witwatersrand.** *J Cutan Med Surg.* 2000; **4**(1): 51–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Govender NP, Maphanga TG, Zulu TG, *et al.*: **An outbreak of lymphocutaneous sporotrichosis among mine-workers in South Africa.** *PLoS Negl Trop Dis.* 2015; **9**(9): e0004096.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Gupta AK, Lyons DCA: **The rise and fall of oral ketoconazole.** *J Cutan Med Surg.* 2015; **19**(4): 352–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Martins JE, Mendonça PR, Cucé LC: **Tratamento da esporotricose com cetoconazol.** *Rev Hosp Clin.* 1982; **37**(2): 92–4.
30. Samorodin CS, Sina B: **Ketoconazole-treated sporotrichosis in a veterinarian.** *Cutis.* 1984; **33**(5): 487–8.
[PubMed Abstract](#)
31. Fitzpatrick JE, Eubanks S: **Acquired immunodeficiency syndrome presenting as disseminated cutaneous sporotrichosis.** *Int J Dermatol.* 1988; **27**(6): 406–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Lipstein-Kresch E, Isenberg HD, Singer C, *et al.*: **Disseminated *Sporothrix schenckii* infection with arthritis in a patient with acquired immunodeficiency syndrome.** *J Rheumatol.* 1985; **12**(4): 805–8.
[PubMed Abstract](#)
33. Penn CC, Goldstein E, Bartholomew WR: ***Sporothrix schenckii* meningitis in a patient with AIDS.** *Clin Infect Dis.* 1992; **15**(4): 741–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Albuquerque PC, Fonseca BPF, Girard-Dias W, *et al.*: **Mapping the Brazilian microscopy landscape: A bibliometric and network analysis.** *Micron.* 2019; **116**: 84–92.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. González-Alcalde G, Salinas A, Ramos JM: **Scientometrics analysis of research activity and collaboration patterns in Chagas cardiomyopathy.** *PLoS Negl Trop Dis.* 2018; **12**(6): e0006602.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. de Lima Barros MB, Schubach TM, Galhardo MC, *et al.*: **Sporotrichosis: an**

- emergent zoonosis in Rio de Janeiro. *Mem Inst Oswaldo Cruz*. 2001; **96**(6): 777–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Gutierrez-Galhardo MC, Freitas DFS, do Valle ACF, *et al.*: **Epidemiological aspects of sporotrichosis epidemic in Brazil**. *Curr Fungal Infect Rep*. 2015; **9**(4): 238–45.
[Publisher Full Text](#)
 38. Araújo LTR, Silva WA, Juliano RS: **Região fronteriza e epidemiologia: Estudo da esporotricose e sua relação na dinâmica da fronteira Brasil-Bolívia**. *Rev GeoPantanal*. 2017; **12**: 97–105.
[Reference Source](#)
 39. Silva GM, Howes JCF, Leal CAS, *et al.*: **Surto de esporotricose felina na região metropolitana do Recife**. *Pesq Vet Bras* 2018; **38**(9): 1767–71.
[Publisher Full Text](#)
 40. Lang PB, Gouveia FC, Leta J: **Health research networks on the web: an analysis of the Brazilian presence**. *Cad Saude Publica*. 2014; **30**(2): 369–78.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. Mota FB, Fonseca BPFE, Galina AC, *et al.*: **Mapping the dengue scientific landscape worldwide: a bibliometric and network analysis**. *Mem Inst Oswaldo Cruz*. 2017; **112**(5): 354–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 42. Fonseca BPFE, Zicker F: **Dengue research networks: building evidence for policy and planning in Brazil**. *Health Res Policy Syst*. 2016; **14**(1): 80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 43. Chapman DN, Doubell A, Oversteegen L, *et al.*: **Policy cures research**. 121.
 44. Rodrigues ML, Albuquerque PC: **Searching for a change: The need for increased support for public health and research on fungal diseases**. *PLoS Negl Trop Dis*. 2018; **12**(6): e0006479.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 45. Mahajan VK: **Sporotrichosis: An Overview and Therapeutic Options**. *Dermatol Res Pract*. 2014; **2014**: 272376.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 46. García-Carnero LC, Pérez-García LA, Martínez-Álvarez JA, *et al.*: **Current trends to control fungal pathogens: exploiting our knowledge in the host-pathogen interaction**. *Infect Drug Resist*. 2018; **11**: 903–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status: ? ? ?

Version 1

Reviewer Report 24 February 2021

<https://doi.org/10.5256/f1000research.26752.r77886>

© 2021 Zamith-Miranda D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Daniel Zamith-Miranda

Division of Infectious Diseases, Department of Medicine, Albert Einstein College of Medicine, New York, NY, USA

The manuscript "Bibliometric assessment and implications for practice of sporotrichosis research (1945-2018)" from Albuquerque and colleagues shows how research on sporotrichosis has been conducted in the mentioned time range. The authors raised relevant data to conclude where and how this research has been done. Besides usual typos, parts of the text, as well as some figures should be remade (comments below).

Introduction

When citing reference #15, the authors should highlight that the cited bibliometric analyses were about tuberculosis. Otherwise, the reader may think that those analyses were already made for sporotrichosis and that would decrease the impact of the current work.
End of third paragraph: "impact" is more appropriate than "impacts".

Methods

Is citation #19 correctly referenced? Isn't the publishing year supposed to be included at least?

Results

What is reference 23? Is it a preprint? Is it the same work as this manuscript published somewhere else?

Table 3: Why does the Rank of all publications, after going from 1 – 10, go to 22, 24, 23, 21, and 26? Isn't that supposed to be the top 15 journals?

Figure 4: From A to D, most of what is written is way too small. Even when zooming in, some words are difficult to read. I recommend the authors to at least increase the resolution, but also try to rearrange the panels in order to make it easier for the reader. Even 4c and 4D have distinct resolution, even though the text is supposed to be identical among them.

Fig. 4 Legend: "(b) The heat map shows the most frequent terms in the period analyzed". Please specify the period.

Fig 5A: I am not sure if this figure is adding information or even helping the understanding of the

text. The authors should consider to change the format, or even take it out of the manuscript.
Fig 6: It seems that, besides the research networks, this figure is also about the distribution of scientific institutions in Brazil. Merging the current figure with a map would make it easier to visualize.

Table 7: Consider to keep it as supplementary information.

Discussion

In "...This fact, together with the increase in the average degree and size of the giant component...". Please clarify what is the giant component.

In "...It is interesting to note that just recently, sporotrichosis has spread to the Northeast and Center-Western regions of the country...". For the sake of being consistent with the nomenclature used in Results, please change Center-Western to Midwest.

In "...In our analysis, publications co-authored by Brazilians researchers have more tentative and minimal...", change to Brazilian researchers.

I agree with the other reviewer that conclusions regarding implications in practice cannot be taken based on the method used.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Microbiology (mycology) and immunology.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 29 Mar 2021

Rodrigo Almeida-Paes

The authors thank the careful revision of our manuscript. Our point-by-point response to the reviewers comments are presented below:

Reviewer comment: When citing reference #15, the authors should highlight that the cited bibliometric analyses were about tuberculosis. Otherwise, the reader may think that those analyses were already made for sporotrichosis and that would decrease the impact of the current work. End of third paragraph: "impact" is more appropriate than "impacts".

Our response and action taken: Thank you for the corrections. We have added reference #15 to emphasize the potential of network analysis. We have clarified the article findings as suggested.

Reviewer comment: Methods: Is citation #19 correctly referenced? Isn't the publishing year supposed to be included at least?

Our response and action taken: Thank you for the corrections. All references were carefully reviewed and corrected as suggested. The correct year is 2009.

Reviewer comment: Results: What is reference 23? Is it a preprint? Is it the same work as this manuscript published somewhere else?

Our response and action taken: This reference refers to the raw data generated in this study that are available at Open Science Framework: Bibliometric assessment of sporotrichosis research. <https://doi.org/10.17605/OSF.IO/MXU6V> This was a requirement of the journal.

Reviewer comment: Table 3: Why does the Rank of all publications, after going from 1 – 10, go to 22, 24, 23, 21, and 26? Isn't that supposed to be the top 15 journals?

Our response: Our intention with this table is to show the top 10 global journals and the top 10 journals publishing Brazilian research. Sometimes a journal within the Brazilian top 10 is not within the top 10 global journals, PLoS Neglected Tropical Diseases, for instance (top 6 Brazilian and 24th global).

Action taken: Table 3 caption was amended to include the description of rank comparison.

Reviewer comment: Figure 4: From A to D, most of what is written is way too small. Even when zooming in, some words are difficult to read. I recommend the authors to at least increase the resolution, but also try to rearrange the panels in order to make it easier for the reader. Even 4c and 4D have distinct resolution, even though the text is supposed to be identical among them.

Fig. 4 Legend: "(b) The heat map shows the most frequent terms in the period analyzed". Please specify the period.

Our response and action taken: Regarding Figure 4 resolution, we gave our best to increase font letters as requested but, to do it, figure would be larger than the page size. However, all terms that generated the figure are presented in the (F2) NetVoS.txt (Network file for VOSViewer software) underlying data, available at <https://doi.org/10.17605/OSF.IO/MXU6V>. The legend of the figure was rewritten, to comply with the reviewer's request.

Reviewer comment: Fig 5A: I am not sure if this figure is adding information or even helping the understanding of the text. The authors should consider to change the format,

or even take it out of the manuscript.

Our response and action taken: According to the reviewer suggestion, the figure was removed from the manuscript.

Reviewer comment: Fig 6: It seems that, besides the research networks, this figure is also about the distribution of scientific institutions in Brazil. Merging the current figure with a map would make it easier to visualize.

Our response: Unfortunately, we were not able to merge this figure with a map. We apologize for this situation.

Reviewer comment: Table 7: Consider to keep it as supplementary information.

Our response: F1000Research does not accept supplementary material, so we kept the table within the main article.

Reviewer comment: In "...This fact, together with the increase in the average degree and size of the giant component...". Please clarify what is the giant component.

Our response and action taken: The giant component is the network core (Mentioned in the results section - Network evolution among Brazilian institutions). This information was added to the text.

Reviewer comment: In "...It is interesting to note that just recently, sporotrichosis has spread to the Northeast and Center-Western regions of the country...". For the sake of being consistent with the nomenclature used in Results, please change Center-Western to Midwest.

Our response and action taken: The text was changed accordingly.

Reviewer comment: In "...In our analysis, publications co-authored by Brazilians researchers have more tentative and minimal...", change to Brazilian researchers.

Our response and action taken: The text was changed accordingly.

Reviewer comment: I agree with the other reviewer that conclusions regarding implications in practice cannot be taken based on the method used.

Our response and action taken: As requested by all reviewers, the text has been rewritten to demonstrate the proposal for key message recovery methodology and not to conclude implications for practice.

Competing Interests: No competing interests were disclosed.

Reviewer Report 15 February 2021

<https://doi.org/10.5256/f1000research.26752.r77889>

© 2021 Menezes Alencar M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Maria Simone Menezes Alencar

Federal University of the State of Rio de Janeiro, Rio de Janeiro, Brazil

The topic of the work is relevant, and the bibliometric approach is adequate. The content analysis proposal is innovative and deserves to be highlighted.

It is strongly recommended that searches in bibliometric studies be conducted in specialized databases. Although the used methodology which includes data from several sources is strong, the study should include other data sources in the field. I also suggest the inclusion of the Medline database in this study.

I suggest that only articles published in journals be used, as they are more relevant documents that report completed research. Peer-reviewed journals provide scientific credibility and are not as biased as other indexed documents, such as letters, notes, editorials, etc.

Although the methodological stage "Implications for practice" shows an interesting innovation in its approach, it should be interpreted with care. To validate the results, I suggest that this stage be improved with two independent experts analyzing a statistical percentage of the sample.

Although the analysis of studies on the subject in the four periods shows different behaviors, these ranges are not used in other analyses. In this way, this presentation is somewhat disconnected with the general context of the article.

The other results are well presented and consistently discussed.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Scientific and technological information acting mainly on the following subjects: bibliometrics, technological foresight, patent analysis and open science.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 29 Mar 2021

Rodrigo Almeida-Paes

We would like to thank this reviewer for her helpful comments that helped to improve the quality of our manuscript.

Our point-by-point responses are presented below:

Reviewer comment: It is strongly recommended that searches in bibliometric studies be conducted in specialized databases. Although the used methodology which includes data from several sources is strong, the study should include other data sources in the field. I also suggest the inclusion of the Medline database in this study. I suggest that only articles published in journals be used, as they are more relevant documents that report completed research. Peer-reviewed journals provide scientific credibility and are not as biased as other indexed documents, such as letters, notes, editorials, etc.

Our response: The authors chose as methodological approach not to use regional bases as SciELO and LILACS, due to the difficulty to obtain total equivalence for a global analysis, since other regional bases should be included and the difficulty of data harmonization. Scopus includes a more expanded spectrum of journals (covers 100% of Medline and Embase) and Web of Science covers the oldest publications. Also, metadata for multiple affiliations for each author or contributor are included in Pubmed only at the end of 2014.

Action taken: The sentence in methods was altered in order to clarify the point raised by the reviewer identifying the different types of articles analyzed (original articles, reviews, letters and editorials). Please see Source of data section.

Reviewer comment: Although the methodological stage “Implications for practice” shows an interesting innovation in its approach, it should be interpreted with care. To validate the results, I suggest that this stage be improved with two independent experts analyzing a statistical percentage of the sample.

Our response: We appreciate the suggestion. The proposal of this research was to evaluate a methodology to identify messages and technical recommendations for action in the abstracts of scientific publications on sporotrichosis. In this first moment, the authors do not intend to analyze the messages, but the validation of the recovery methodology. And, in fact, two of the authors reviewed the data. In the further analysis, in order to validate the recovery of recommendations, two independently experts will always review the recommendations.

Action taken: The text has been modified to clarify that this was a methodological evaluation proposal, rather than a comprehensive analysis of implications for practice. We also clarified that two evaluators collected the data.

Reviewer comment: Although the analysis of studies on the subject in the four periods shows different behaviors, these ranges are not used in other analyses. In this way, this presentation is somewhat disconnected with the general context of the article. The other results are well presented and consistently discussed.

Our response: The reviewer raised an important point. Our initial analysis of the four periods revealed that two of them, 1945-1962 and 1992-2002, publications remained without significant variation. The strongest variation occurred after 2002, which coincides with the start of publications about the Brazilian zoonotic epidemics of sporotrichosis.

Therefore, for subsequent analysis, we chose to separate the articles only in two periods: before and after the beginning of the Brazilian sporotrichosis epidemics.

Action taken: The issue raised by the reviewer was clarified in the research trends section of results and further addressed in the discussion.

Competing Interests: No competing interests were disclosed.

Reviewer Report 24 September 2020

<https://doi.org/10.5256/f1000research.26752.r71160>

© 2020 Ramírez-Soto M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Max Carlos Ramírez-Soto

School of Public Health and Administration, Universidad Peruana Cayetano Heredia, Lima, Peru

Congratulations to the authors. Despite sporotrichosis remains a neglected disease in terms of research interest, funding and medical attention, to which the authors have made significant contributions. However, this study has several major observations:

Methods.

The literature search is limited to two databases. Fig. 1. Present a flow, and follow the guidelines of a scoping review.

MedLine / Pubmed and Abstracts of Scientific Conference should be reviewed.

There are several published studies on sporotrichosis in journals indexed in SCIELO, LiLACS, etc. These papers have not been included in the study.

I don't think that this article contains enough robust data to evidence the implications for practice. The strategy and organization of the information about Implications for practice is not appropriate. A set of words related to tentative language is biased. Furthermore, only the abstracts have been reviewed. To evaluate the implications in practice, it is necessary to organize the different types of published studies: case series, observational study, clinical trials and systematic review, diagnostic test, etc. Furthermore, studies not related to the clinical practice of sporotrichosis should be excluded.

Results:

Paragraph 1. I suggest include in Results section include the statistical test.

Scientific production by hyperendemic areas of sporotrichosis should be presented.

Table 3. I suggest to make a comparison between Brazil, USA, Mexico, China, or hyperendemic areas of sporotrichosis.

The authors should include the different types of published articles: review, original, short communication, case reports, clinical trial, systematic, review, correspondence, Research Letters, etc.

Research trends. It would be interesting to know how these trends vary between Latin America, Europe and Asia.

Implications for practice

I suggest organizing the published studies: case series, observational study, clinical trials and systematic review, diagnostic test, etc. Although there are very few clinical trials and systematic reviews on sporotrichosis, this evidence should be presented in tables. Furthermore, studies not related to the clinical practice of sporotrichosis should be excluded. Finally, sporotrichosis is not limited to dermatology.

Discussion

In the "Discussion" section I would have wished to see more information on implication practice. Limitations should be described and discussed such as: the number of databases included in the study, the studies were not classified, to differentiate the type of scientific evidence for clinical practice, etc.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

No

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Medical Mycology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have

significant reservations, as outlined above.

Author Response 29 Mar 2021

Rodrigo Almeida-Paes

We would like to thank this reviewer for his thoughtful comments. Our point-by-point response is presented below:

Reviewer comment: Methods. The literature search is limited to two databases. Fig. 1. Present a flow, and follow the guidelines of a scoping review. MedLine / Pubmed and Abstracts of Scientific Conference should be reviewed. There are several published studies on sporotrichosis in journals indexed in SCIELO, LILACS, etc. These papers have not been included in the study.

Our response: The authors used a bibliometric analysis approach to observe the evolution of the world scientific literature and to identify research trends in the sporotrichosis field, focusing on the Brazilian scenario. The authors did not intend to assess the state of the art in the area, but rather to perform a bibliometric analysis with the research effort, which differs from a scoping review.

The authors chose as methodological approach not to use regional bases such as SciELO and LILACS, due to the difficulty to obtain total equivalence for a global analysis, since other regional bases should be included, making data harmonization difficult.

Scopus includes a more expanded spectrum of journals (covers 100% of Medline and Embase) and Web of Science covers the oldest publications. Also, metadata for multiple affiliations for each author or contributor are included in Pubmed only at the end of 2014. The authors understand the relevance and challenges of integrating different databases in bibliometric analysis to provide a reliable research panorama. The inclusion of limitations will clarify any methodological bias.

Action taken: We included a paragraph in the discussion addressing this limitation of study.

Reviewer comment: I don't think that this article contains enough robust data to evidence the implications for practice. The strategy and organization of the information about Implications for practice is not appropriate. A set of words related to tentative language is biased. Furthermore, only the abstracts have been reviewed. To evaluate the implications in practice, it is necessary to organize the different types of published studies: case series, observational study, clinical trials and systematic review, diagnostic test, etc. Furthermore, studies not related to the clinical practice of sporotrichosis should be excluded.

Our response and action taken: The text was amended to describe in the methodology the recovery of recommendation messages for practice, research and public health in abstracts based on a wording recovery. Several authors have already demonstrated the importance of abstracts, since they are a condensed version of a scientific research and the most often read section of a research article.

Searching abstracts are convenient because they contain most of the article's relevant keywords and appear to be a useful tool to inform to practitioners as a resource for guiding clinical decisions.

The tentative set of words was already validated by Lynn J et al 2011, where research articles from three leading clinical journals (*N Engl J Med*, *JAMA*, and *Ann Intern Med*) were examined to identify terms that articulated implications (eg, words such as 'may', 'speculate', 'suggest',

'reinforce' and 'potentially').

Reviewer comment: Paragraph 1. I suggest include in Results section include the statistical test.

Our response and action taken: This information was included.

Reviewer comment: Table 3. I suggest to make a comparison between Brazil, USA, Mexico, China, or hyperendemic areas of sporotrichosis. Scientific production by hyperendemic areas of sporotrichosis should be presented.

Our response and action taken: Table 2 was replaced for an improved version as suggested.

To address the reviewer's suggestion, the authors included the proportion of publications compared to the total number of articles for each country in the table 1, which shows Top 20 most active countries in the *Sporothrix*/sporotrichosis field according to author's country of professional affiliation what allows comparison between hyperendemic countries.

Reviewer comment: The authors should include the different types of published articles: review, original, short communication, case reports, clinical trial, systematic, review, correspondece, Research Letters, etc.

Our response: The sentence in methods was altered in order to clarify the point raised by the reviewer identifying the different types of articles analyzed (original articles, reviews, letters and editorials).

Action taken: See Souce of data section.

Reviewer comment: Research trends. It would be interesting to know how these trends vary between Latin America. Europe and Asia.

Our response: Thank you for this suggestion. The authors agree that this analysis would show a bigger picture. However, for a robust and reproducible analysis, the quantity of publications must be consistent to construct a map based on a co-occurrence matrix. To assess how these trends vary between Latin America, Europe and Asia, the small size of text corpus of each region may not be able, unfortunately, to represent nuances in the research scenario.

Reviewer comment: Implications for practice: I suggest organizing the published studies: case series, observational study, clinical trials and systematic review, diagnostic test, etc. Although there are very few clinical trials and systematic reviews on sporotrichosis, this evidence should be presented in tables. Furthermore, studies not related to the clinical practice of sporotrichosis should be excluded.

Our response: The text was amended to describe the methodology to recovery recommendations messages for practice, research and public health in abstracts based on wording recovery, rather than a comprehensive search for recommendations for practice. Our original intention with this research was to evaluate a methodology to identify messages and technical recommendations for action in the scientific publications on sporotrichosis. In this first moment, the authors do not intend to analyze the messages, but the validation of the recovery methodology. Further analysis with a broader sample of medical journals to examine how researchers report evidence for action, aiming to assess how they framing their research findings to address implications for practice and public

health is needed.

Action taken: Changes in the title, abstract (methods section), introduction (last paragraph), results (statements about practice section), and discussion

Reviewer comment: Finally, sporotrichosis is not limited to dermatology.

Our response and action taken: Thank you for the reminder. The authors were referring to journals with the scopus in dermatology. The text was amended to: "for instance, four of the top ten journals are directed to medical doctors with interest in dermatology".

Reviewer comment: Discussion: In the "Discussion" section I would have wished to see more information on impicance practic. Limitations should be described and discussed such as: the number of databases included in the study, the studies were not classified, to differentiate the type of scientific evidence for clinical practice, etc.

Our response and action taken: Thank you for this observation. To clarify the limitations the text was amended and a limitations section was included in the end of the discussion. To address the reviewer's suggestion, a paragraph describing the applicability of the methodology proposed to the retrieval of the key messages was also included in the discussion section.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research