REVIEW

Recent advances in managing axial spondyloarthritis [version 1; peer review: 2 approved]

Priyanka Agrawal¹, Pedro M. Machado¹⁻³

¹Department of Rheumatology, Northwick Park Hospital, London North West University NHS Trust, London, UK

²Department of Rheumatology, University College London Hospitals NHS Foundation Trust, London, UK

³Centre for Rheumatology, Division of Medicine & Department of Neuromuscular Diseases, Institute of Neurology, University College London, London, UK

V1 First published: 13 Jul 2020, 9(Faculty Rev):697
<https://doi.org/10.12688/f1000research.22577.1>

Latest published: 13 Jul 2020, 9(Faculty Rev):697
<https://doi.org/10.12688/f1000research.22577.1>

Abstract

Axial spondyloarthritis (axSpA) is a chronic inflammatory rheumatic disease that predominantly affects the axial skeleton. The advent of biologic drugs has transformed the management of patients with axSpA. However, non-steroidal anti-inflammatory drugs remain the first-line drug treatment for axSpA. The optimal management of patients with axSpA requires a combination of pharmacological and non-pharmacological treatment modalities, namely exercise and physical therapy. This review looks at novel therapeutic options in patients with axSpA. It also summarises current evidence regarding radiographic progression and treat-to-target in axSpA.

Keywords

Axial spondyloarthritis, ankylosing spondylitis, treatment, management, biologics, outcomes.

Open Peer Review

Approval Status

	1	2
version 1
13 Jul 2020		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Daniel Wendling**, CHRU Besançon, Besançon, France

2. **Peter Cheung**, National University Hospital, Singapore, Singapore
 National University Singapore, Singapore, Singapore

Any comments on the article can be found at the end of the article.

Corresponding author: Pedro M. Machado (p.machado@ucl.ac.uk)

Author roles: Agrawal P: Writing – Original Draft Preparation; Machado PM: Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: PMM has received consulting/speaker's fees from Abbvie, BMS, Celgene, Eli Lilly, Janssen, MSD, Novartis, Pfizer, Roche and UCB. PA declares that she has no competing interests.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Agrawal P and Machado PM. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Agrawal P and Machado PM. **Recent advances in managing axial spondyloarthritis [version 1; peer review: 2 approved]** F1000Research 2020, 9(Faculty Rev):697 <https://doi.org/10.12688/f1000research.22577.1>

First published: 13 Jul 2020, 9(Faculty Rev):697 <https://doi.org/10.12688/f1000research.22577.1>

Introduction

Axial spondyloarthritis (axSpA) belongs to a clinically heterogeneous group of inflammatory rheumatic diseases that share common genetic, histological, and clinical features. It encompasses ankylosing spondylitis (AS) (or radiographic axSpA [r-axSpA]) and non-radiographic axSpA (nr-axSpA)¹. Over the last couple of decades, we have witnessed remarkable advances in the pathogenesis, management and treatment of axSpA. The discovery of anti-tumour necrosis factor (anti-TNF) has revolutionised the treatment of this chronic condition. More recently, interleukin-17 (IL-17) has been discovered as an alternative therapeutic target and there is promise from small molecules such as Janus kinase (JAK) inhibitors².

In this article, we shall discuss the management of axSpA with a primary focus on the recent advances. We shall review potential new biologics on the horizon. We shall also consider the use of biosimilars, radiographic progression in axSpA and the controversy surrounding the treat-to-target (T2T) approach.

Management of axial spondyloarthritis

AxSpA can interfere with patients' daily activities, including schooling, work, and social life^{3,4}. The goals of treatment are to reduce disease activity (signs and symptoms), to prevent disability and structural damage, and to maintain work productivity, health-related quality of life, and social participation⁵. Non-steroidal anti-inflammatory drugs (NSAIDs) and physical therapy remain the mainstays of treatment of axSpA. The Assessment of Spondylarthritis International Society (ASAS) and European League Against Rheumatism (EULAR) (2016 update) recommend that patients with pain and stiffness should use an NSAID as first-line drug treatment up to the maximum dose while taking risks and benefits into account. For patients who respond well to NSAIDs, continuous use of this medication is preferred in case on-demand use results in worsening of symptoms⁶. The 2019 American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network (ACR/SAA/SPARTAN) recommendations similarly advise that adults with active axSpA receive continuous NSAIDs over on-demand NSAIDs. However, in adults with stable axSpA, on-demand treatment with NSAIDs is recommended over continuous treatment⁵.

There is controversy regarding the role of NSAIDs in preventing radiographic progression in axSpA. Slower radiographic progression was observed in AS patients taking celecoxib continuously (that is, daily) for a 2-year period, compared with patients taking it on demand, in a study by Wanders *et al.*⁷. A post-hoc analysis of this study showed that this effect was more pronounced in patients with elevated acute-phase reactants or in patients with a high or very high Ankylosing Spondylitis Disease Activity Score (ASDAS), which includes C-reactive protein (CRP) as one of the variables⁸. However, this inhibitory effect on new bone formation in the spine of patients with AS was not observed in a more recent randomised multicentre trial (ENRADAS) comparing continuous diclofenac intake over the course of 2 years versus on-demand

treatment⁹. Conversely, a 2-year observational study in patients with AS showed a reduction in the progression of structural damage of the spine of patients with a high NSAID intake compared with those with a low NSAID intake. This protective effect was seen nearly exclusively in patients with syndesmophytes and elevated CRP at baseline¹⁰. More recently, it was suggested that continuous use of NSAIDs reduces radiographic progression in sacroiliac joints in patients with early axSpA¹¹.

Non-pharmacological treatment modalities are important in the management of patients with axSpA. ASAS-EULAR recommend that patients should be educated about axSpA and encouraged to exercise on a regular basis and stop smoking; physical therapy should be considered⁶. The inclusion of aerobic components, cardiorespiratory exercises, and educational programs in traditional programs of exercises may lead to improved clinical outcomes, although the most effective exercise protocol remains unclear¹². Promising effects of cardiorespiratory and strength exercises on emotional distress, fatigue, and ability to do a full day's activities were shown in a small pilot Scandinavian study in patients with axSpA¹³.

Biologics in the treatment of axial spondyloarthritis

Until very recently there were five licensed anti-TNF drugs (adalimumab, certolizumab, etanercept, golimumab and infliximab) for the indication of AS and four (adalimumab, etanercept, certolizumab and golimumab) for the indication of nr-axSpA (in the US, only certolizumab was approved for the indication of nr-axSpA). The IL-17 blocker secukinumab has been approved by both the European Medicines Agency (EMA) and the US Food and Drug Administration (FDA) for the indication of AS. Ixekizumab was initially (2019) approved by the FDA for AS, and in July 2020, the FDA expanded the approval of ixekizumab to include nr-axSpA; almost simultaneously, the EMA also approved ixekizumab for the treatment of both AS and nr-axSpA.

ASAS-EULAR-recommended disease activity cutoffs to start anti-TNF treatment are either a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score of at least 4 or an ASDAS of at least 2.1 after treatment with two different NSAIDs for at least 4 weeks in total⁶. British Society of Rheumatology (BSR) and British Health Professionals in Rheumatology recommendations¹⁴ define high disease activity as BASDAI and spinal pain visual analogue scale (VAS) scores of at least 4. According to the BSR, BASDAI and VAS scores should be measured on two occasions at least 4 weeks apart and patients need to have failed two NSAIDs for at least 2 weeks each unless contraindicated.

The 2019 ACR-SAA-SPARTAN recommendations also advise anti-TNF drugs in patients with axSpA when activity persists despite NSAID treatment. Anti-TNF monoclonal antibodies should be preferred in patients with concomitant inflammatory bowel disease (IBD) or recurrent iritis. According to these recommendations, IL-17 blockers are advised for patients with active disease who have heart failure or demyelinating disease as a contraindication to anti-TNF and in primary non-responders to anti-TNF. Secukinumab and ixekizumab

are not recommended in patients with IBD or recurrent uveitis. In the 2019 ACR-SAA-SPARTAN recommendations, tofacitinib, a JAK inhibitor currently not approved for axSpA, is highlighted as a potential second-line option for patients with contraindications to a TNF inhibitor (TNFi) other than infections. Recommendations regarding tofacitinib may change pending the results of larger clinical trials⁵. Owing to the likelihood for symptom recurrence, discontinuation of biologics is not recommended by ACR-SAA-SPARTAN. If tapering

is considered, patients should be counselled regarding the potential for increased disease activity⁵. In [Table 1](#), an overview of biological disease-modifying anti-rheumatic drugs (DMARDs) and targeted synthetic DMARDs in axSpA and related chronic inflammatory conditions is presented.

Biologic drugs that have recently shown efficacy

The efficacy of ixekizumab, a high-affinity monoclonal antibody (mAb) against IL-17A, has been demonstrated in both

Table 1. Overview of biological disease-modifying anti-rheumatic drugs (DMARDs) and targeted synthetic DMARDs in axial spondyloarthritis and related chronic inflammatory conditions.

Target	Drug	Axial spondyloarthritis	Psoriatic arthritis	Psoriasis	Crohn's disease	Rheumatoid arthritis	Uveitis
TNF	Adalimumab (mAb to TNF)						
	Certolizumab (mAb to TNF)						
	Etanercept (fusion protein against TNF)						
	Golimumab (mAb to TNF)						
	Infliximab (mAb to TNF)						
IL-1	Anakinra (IL-1R antagonist)						
B cells	Rituximab (mAb to CD20)						
T cells	Abatacept (inhibitor of T-cell co-stimulation)						
IL-6	Tocilizumab (mAb to IL-6R)						
	Sarilumab (mAb to IL-6R)						
IL-17	Secukinumab (mAb to IL-17A)						
	Ixekizumab (mAb to IL-17A)						
	Brodalumab (mAb to IL-17R)						
	Bimekizumab (mAb to IL-17A and IL-17F)						
	Netakimab/BCD-085 (mAb to IL-17R)						
IL-12 and IL-23	Ustekinumab (mAb to IL-12/23p40)						
	Guselkumab (mAb to IL-23p19)						
	Tildrakizumab (mAb to IL23p19)						
	Risankizumab (mAb to IL23p19)						
PDE4	Apremilast (PDE4 inhibitor)						
JAK	Tofacitinib (JAK1/3 inhibitor)						
	Filgotinib (JAK1 inhibitor)						
	Upadacitinib (JAK1 inhibitor)						
GM-CSF	Namylumab (mAb to GM-CSF)						

Key:

	Efficacy shown in at least one randomised controlled trial
	Some data from pilot or proof-of-concept trials suggest a positive effect
	Some data from pilot or proof-of-concept trials suggest a lack of effect
	Lack of efficacy shown in at least one randomised controlled trial
	No conclusive data available about efficacy

GM-CSF, granulocyte macrophage-colony-stimulating factor; IL, interleukin; JAK, Janus kinase; mAb, monoclonal antibody; PDE4, phosphodiesterase 4; TNF, tumour necrosis factor

biologic-naïve and TNFi-experienced patients with r-axSpA (NCT02696785 and NCT02696798)^{15–17}. More recently, the first study demonstrating the efficacy of ixekizumab in nr-axSpA was also published¹⁸. Positive secukinumab (NCT02696031) data in nr-axSpA were also recently presented in abstract form, but the manuscript has yet to be published¹⁹.

Bimekizumab, a mAb that potently and selectively neutralises both IL-17A and IL-17F, has demonstrated substantial improvements in both musculoskeletal and skin outcomes²⁰. Brodalumab, a human anti-IL-17 receptor A mAb has also demonstrated efficacy in phase 3 studies in both AS and nr-axSpA²¹. The bimekizumab and brodalumab studies have been published in abstract form only.

Netakimab/BCD-085, a humanised mAb against IL-17 with genetically modified Fc- and CDR-regions, has shown efficacy in active AS in a dose-finding phase 2 clinical trial (abstract publication²²).

JAK inhibitors are emerging as an effective therapeutic approach in patients with axSpA. Both tofacitinib (JAK1–3 inhibitor) and filgotinib (selective JAK1 inhibitor) have shown efficacy and safety in phase 2 placebo-controlled trials of patients with AS. Tofacitinib was superior to placebo in reducing signs, symptoms and objective endpoints of patients with active AS and had a safety profile similar to what has been reported in the literature²³. More recently, van der Heijde *et al.* showed that filgotinib was efficacious and safe for patients with active AS and inadequate response or intolerance to NSAIDs²⁴. Therefore, further investigation of filgotinib in AS is warranted²⁴ (NCT03117270). Finally, results of the first randomised trial of upadacitinib (selective JAK1 inhibitor) versus placebo in patients with AS were recently published; upadacitinib was efficacious and well tolerated in patients with active AS who had not responded to or had a contraindication to treatment with NSAIDs. These data support further investigation of upadacitinib for the treatment of axSpA²⁵.

Biologic drugs that have not shown efficacy

Uncontrolled or controlled trials with anakinra (IL-1 blocker)^{26,27}, abatacept (inhibitor of T-cell co-stimulation)²⁸, rituximab (B-cell depletion agent)^{29,30} and tocilizumab (IL-6 blocker)^{31,32} have shown no consistent efficacy in patients with AS. Of note, conventional synthetic DMARDs have also failed to show efficacy in axial disease.

Apremilast is a small-molecule inhibitor of phosphodiesterase 4 (PDE4) that modulates the inflammatory response. In a small (n = 38), 12-week, double-blind, placebo-controlled, phase 2 study of symptomatic AS patients with positive magnetic resonance imaging (MRI)³³, apremilast did not meet its primary endpoint (change in BASDAI score at week 12) but was associated with numerically greater improvement in all clinical assessments compared with placebo. A subsequent double-blind, placebo-controlled, phase 3 study assessing the efficacy and safety of apremilast in active AS was completed (NCT01583374). Preliminary online reports suggest that this was

a negative study (primary endpoint not met: ASAS20 at week 16) (source: NCT01583374). This study has not been published yet.

In a randomised, double-blind, placebo-controlled, dose-finding phase 2 study, treatment with risankizumab (a humanised mAb targeting IL-23A) did not show evidence of clinically meaningful improvements compared with placebo in patients with active AS; the primary endpoint, ASAS40 response at week 12, was not met. This study suggested that despite a genetic association with the IL-23 pathway, IL-23 may not be a relevant driver of disease pathogenesis and clinical manifestations in AS (NCT02047110)³⁴.

In three placebo-controlled trials, the efficacy of ustekinumab (an anti-IL-12/23 antibody) in the treatment of axial SpA was not demonstrated. The safety profile was consistent with that of studies in other indications³⁵ (NCT02437162, NCT02438787 and NCT02407223).

Siebert *et al.* recently published an editorial on the reasons for failure of IL-23p19 inhibition in AS³⁶. The authors hypothesise that established AS may have “transitioned pathogenetically” to a mature type 17 phenotype, making it unresponsive to IL-23p19 blockade and other upstream treatment strategies, such as IL-6 inhibition, that would normally modulate the IL-17 response³⁶. Therefore, neutralising upstream molecules would be less effective than specifically targeting IL-17A. Furthermore, other effector pathologic pathways driving the IL-17 response are probably still unknown and need to be identified³⁶. In Table 2, ongoing axSpA drug trials are presented.

Biosimilars

The World Health Organization has defined a “similar biotherapeutic product” (also called “biosimilar”) as a biotherapeutic product that is similar in terms of quality, safety and efficacy to an already licensed original or “reference biotherapeutic product” (also called “originator” drug)³⁷. In adults with stable AS receiving an originator anti-TNF, ACR-SAA-SPARTAN strongly recommends continuing treatment with the originator anti-TNF over mandated switching to its biosimilar⁵. The BSR published a position statement³⁸ on the use of biosimilars in clinical practice. The key principles of this document are that (1) all biologics and biosimilars should be prescribed by brand name rather than by international non-proprietary name, (2) clinical effectiveness and patient safety should be the overriding principles for prescribing any biologic agent, (3) substitution should be done only with the consent of the prescribing clinician, (4) decisions should be made in partnership with the patients, and (5) registration with the BSR Biologic Registers or other appropriate UK register is recommended. Therefore, switching should be based on a shared decision-making process between patients and rheumatologists, should be a clinically informed decision not made solely for economic reasons, and should take contextual factors of the health-care system into account³⁹.

Biosimilars are now often prescribed as the first biologic drug treatment, and many patients have been successfully switched

Table 2. Ongoing clinical trials in axial spondyloarthritis.

Drug	Mechanism of action	Study design	Patient group	Trial number
Secukinumab (intravenous)	Human monoclonal IgG1 kappa antibody against IL-17	Phase 3, multicentre, DB RCT	axSpA	NCT04156620
Ixekizumab	Humanised mAb against IL-17A	Long Term Extension Study of a Phase 3, multicentre, DB RCT	axSpA	NCT03129100
Brodalumab	Human mAb that binds to the IL-17 receptor	Phase 3, multicentre, DB RCT	axSpA	NCT02985983
Bimekizumab	Humanised monoclonal IgG1 antibody against IL-7A and IL-17F	Phase 3, multicentre, DB RCT	r-axSpA nr-axSpA	NCT03928743 NCT03928704
Tildrakizumab	Human mAb against IL-23	Phase 3, multicentre, open label, non-randomised	axSpA	NCT03552276
Netakimab/BCD-085	Humanised mAb against IL-17 with genetically modified Fc- and CDR-regions	Phase 3, multicentre, DB RCT	r-axSpA	NCT03447704
Upadacitinib	JAK1 selective inhibitor	Phase 3, DB RCT	axSpA	NCT04169373
Namilumab	Human monoclonal IgG1 kappa antibody against GM-CSF	Phase 2a, proof-of-concept, DB RCT	axSpA	NCT03622658

axSpA, axial spondyloarthritis; DB RCT, Double-Blind Placebo-Controlled Randomised Clinical Trial; GM-CSF, granulocyte-macrophage colony-stimulating factor; IL, interleukin; mAb, monoclonal antibody; nr-axSpA, non-radiographic axial spondyloarthritis; r-axSpA, radiographic axial spondyloarthritis

from originator drugs to the biosimilar. They have resulted in substantial savings on the use of biologic therapy, providing greater accessibility to patients. There are now biosimilars for adalimumab (Amgevita, Cyltezo, Hyrimoz and Imraldi), etanercept (Benepali, Erelzi and Eticovo) and infliximab (Inflectra, Ixifi, Remicade, Remsima and Renflexis) and many more being produced/tested.

Role of biologics in the inhibition of radiographic progression

The clinical efficacy of biologics in axSpA has been robustly demonstrated. However, it is still unclear whether they can prevent the progression of structural damage, particularly new bone formation. There are also methodological challenges that have contributed to this uncertainty, namely the fact that radiographic progression is a slow process that requires at least 2 years in order to be reliably detected and the ethical impossibility of performing a comparative study with a 2-year placebo arm (given the known clinical efficacy of biologics in patients who do not respond to conventional treatments)².

Importantly, radiographic progression is a heterogeneous process that varies significantly not only between patients but also within the same patient (bursts of radiographic progression can be followed by periods of a relatively low rate of progression or even no progression). Some risk factors for radiographic progression have been identified: the presence of syndesmophytes (the strongest risk factor; previous damage predicts further damage), male gender, HLA-B27 positivity, long disease/symptom duration, high CRP, MRI activity, high disease activity and smoking. It is hypothesised that more intensive T2T approaches might contribute to decreasing the rate of radiographic progression, particularly in high-risk groups⁴⁰.

Karmacharya *et al.*⁴⁰ recently undertook a systematic review and meta-analysis on the effect of different therapies on radiographic progression in axSpA and suggested a protective effect of TNFi treatment on spinal radiographic progression of AS after at least 4 years of treatment in studies with low risk of bias. Observational data therefore suggest that anti-TNF treatment might prevent new bone formation in axSpA. Regarding NSAIDs (also discussed above) and secukinumab, only data up to 2 years were available, and no effect was found.

SURPASS is an ongoing trial comparing the effect of secukinumab on radiographic progression in AS as compared with an adalimumab biosimilar⁴¹. The primary endpoint of this first head-to-head, phase 3b, randomised, biologic-controlled study is “the absence of progression as measured by the modified Stoke Ankylosing Spondylitis Spine Score”⁴¹. Hopefully, this study will give us more information regarding the comparative structural efficacy of TNF blockers versus IL-17 blockers, thus contributing to evidence-based decision-making.

Treat-to-target approach

The concept of T2T strategies is well founded in the management of chronic diseases that have a well-defined and usually very objective measure that is highly associated with future health outcomes and evidence that maintaining the levels of this measure below a certain threshold will lead to better long-term health⁴². The T2T approach in axSpA is indirectly supported by associations between levels of axSpA disease activity (particularly ASDAS) and future radiographic progression but lacks robust direct evidence. The 2019 ACR-SAA-SPARTAN recommendations mention that “focus on a specific target could lead to rapid cycling through all currently available

treatments in some patients⁵. The 2016 update of the ASAS/EULAR recommendations for the management of axSpA recommends that “treatment should be guided according to a predefined treatment target” but controversy remains as to what this target should be⁶. Importantly, this document also states that “the target should be a shared decision between patient and rheumatologist, taking all relevant situational factors into consideration”⁶. However, data from a randomised controlled trial comparing the efficacy of a T2T approach versus standard of care in axSpA are still lacking. Furthermore, the cost effectiveness of a T2T strategy in clinical practice will require testing⁴³.

The 2017 international task force update of recommendations on T2T in axSpA and psoriatic arthritis⁴⁴ advises that the treatment target should be clinical remission/inactive disease of musculoskeletal (arthritis, dactylitis, enthesitis and axial disease) and extra-articular manifestations. In axSpA, the ASDAS is a preferred measure to define the target.

The Tight Control in Spondyloarthritis (TICOSPA) study is a prospective, randomised (cluster) study to evaluate the potential benefit of a T2T approach in comparison with standard of care in patients with axSpA (TICOSPA, NCT03043846). Results from this study were presented in June at the EULAR 2020 conference (THU0370); the authors reported that usual care resulted in a good outcome in a substantial number of patients but the tight control and treat-to-target strategy was not superior for the primary outcome (ASAS Health Index) despite a greater number of bDMARDs prescription; nevertheless, a general trend in favour of the tight control arm was observed, with a comparable safety profile, and was found to be favourable from a societal

health economic perspective. A German 1-year randomised controlled study taking an intense treatment approach versus routine treatment (STRIKE, NCT02897115) was prematurely terminated because of slow recruitment. Another phase 3 clinical trial, the Treat-to-target in Axial Spondyloarthritis (AScalate) study, is currently recruiting and is a randomised, open-label multicentre trial to investigate the efficacy of a T2T treatment strategy with secukinumab as a first-line biologic compared with a standard-of-care treatment over 36 weeks in patients with active axSpA (NCT03906136). A phase 4 trial, the Treat-to-target With Secukinumab in Axial Spondyloarthritis (TRACE) study, is also looking at T2T with secukinumab in axSpA (NCT03639740).

Conclusions

With so many new biologics on the horizon, the future looks promising for managing patients with axSpA. However, there is an unmet need for results from head-to-head studies directly comparing the efficacy and safety of different biologics in axSpA. This would help determine the optimal sequencing of treatments. There are limitations in using disease activity scores and radiographic progression as markers of treatment response and hence research into suitable biomarkers needs to be undertaken.

Acknowledgements

PMM is supported by the National Institute for Health Research (NIHR) University College London Hospitals Biomedical Research Centre. The views expressed are those of the authors and not necessarily those of the National Health Service (UK), the NIHR or the Department of Health (UK).

References

1. Dubash S, McGonagle D, Marzo-Ortega H: **New advances in the understanding and treatment of axial spondyloarthritis: From chance to choice.** *Ther Adv Chronic Dis.* 2018; 9(3): 77–87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
2. Jones A, Ciurtin C, Ismajli M, *et al.*: **Biologics for treating axial spondyloarthritis.** *Expert Opin Biol Ther.* 2018; 18(6): 641–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Boonen A, Sieper J, van der Heijde D, *et al.*: **The burden of non-radiographic axial spondyloarthritis.** *Semin Arthritis Rheum.* 2015; 44(5): 556–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
4. Ward MM: **Quality of life in patients with ankylosing spondylitis.** *Rheum Dis Clin North Am.* 1998; 24(4): 815–27.
[PubMed Abstract](#)
5. Ward MM, Deodhar A, Gensler LS, *et al.*: **2019 Update of the American College of Rheumatology/Spondylitis Association of America/ Spondyloarthritis Research and Treatment Network Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis.** *Arthritis Care Res (Hoboken).* 2019; 71(10): 1285–99.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
6. van der Heijde D, Ramiro S, Landewé R, *et al.*: **2016 update of the ASAS-EULAR management recommendations for axial spondyloarthritis.** *Ann Rheum Dis.* 2017; 76(6): 978–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
7. Wanders A, van der Heijde D, Landewé R, *et al.*: **Nonsteroidal antiinflammatory drugs reduce radiographic progression in patients with ankylosing spondylitis: A randomized clinical trial.** *Arthritis Rheum.* 2005; 52(6): 1756–65.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
8. Kroon F, Landewé R, Dougados M, *et al.*: **Continuous NSAID use reverts the effects of inflammation on radiographic progression in patients with ankylosing spondylitis.** *Ann Rheum Dis.* 2012; 71(10): 1623–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Sieper J, Listing J, Poddubnyy D, *et al.*: **Effect of continuous versus on-demand treatment of ankylosing spondylitis with diclofenac over 2 years on radiographic progression of the spine: Results from a randomised multicentre trial (ENRADAS).** *Ann Rheum Dis.* 2016; 75(8): 1438–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
10. Poddubnyy D, Rudwaleit M, Haibel H, *et al.*: **Effect of non-steroidal anti-inflammatory drugs on radiographic spinal progression in patients with axial spondyloarthritis: Results from the German Spondyloarthritis Inception Cohort.** *Ann Rheum Dis.* 2012; 71(10): 1616–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
11. Rumiantseva D, Dubinina T, Rumyantseva O, *et al.*: **SAT0278 NSAIDs reduces radiographic progression in sacroiliac joints in early axial spondyloarthritis.** *Ann Rheum Dis.* 2018; 77(Suppl 2): 1003.
[Publisher Full Text](#)
12. Perrotta FM, Musto A, Lubrano E: **New Insights in Physical Therapy and**

- Rehabilitation in Axial Spondyloarthritis: A Review.** *Rheumatol Ther.* 2019; 6(4): 479–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
13. Sveaas SH, Berg IJ, Fongen C, *et al.*: High-intensity cardiorespiratory and strength exercises reduced emotional distress and fatigue in patients with axial spondyloarthritis: A randomized controlled pilot study. *Scand J Rheumatol.* 2018; 47(2): 117–21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 14. Hamilton L, Barkham N, Bhalla A, *et al.*: BSR and BHPR guideline for the treatment of axial spondyloarthritis (including ankylosing spondylitis) with biologics. *Rheumatology (Oxford).* 2017; 56(2): 313–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 15. van der Heijde D, Cheng-Chung Wei J, Dougados M, *et al.*: Ixekizumab, an interleukin-17A antagonist in the treatment of ankylosing spondylitis or radiographic axial spondyloarthritis in patients previously untreated with biological disease-modifying anti-rheumatic drugs (COAST-V): 16 week results of a phase 3 randomised, double-blind, active-controlled and placebo-controlled trial. *Lancet.* 2018; 392(10163): 2441–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 16. Deodhar A, Poddubny D, Pacheco-Tena C, *et al.*: Efficacy and Safety of Ixekizumab in the Treatment of Radiographic Axial Spondyloarthritis: Sixteen-Week Results From a Phase III Randomized, Double-Blind, Placebo-Controlled Trial in Patients With Prior Inadequate Response to or Intolerance of Tumor Necrosis Factor Inhibitors. *Arthritis Rheumatol.* 2019; 71(4): 599–611.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 17. Dougados M, Wei JCC, Landewé R, *et al.*: Efficacy and safety of ixekizumab through 52 weeks in two phase 3, randomised, controlled clinical trials in patients with active radiographic axial spondyloarthritis (COAST-V and COAST-W). *Ann Rheum Dis.* 2020; 79(2): 176–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 18. Deodhar A, van der Heijde D, Gensler LS, *et al.*: Ixekizumab for patients with non-radiographic axial spondyloarthritis (COAST-X): A randomised, placebo-controlled trial. *Lancet.* 2020; 395(10217): 53–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 19. Deodhar A, Blanco R, Dokoupilova E, *et al.*: Secukinumab 150 mg Significantly Improved Signs and Symptoms of Non-radiographic Axial Spondyloarthritis: Results from a Phase 3 Double-blind, Randomized, Placebo-controlled Study [abstract]. *Arthritis Rheumatol.* 2019; 71(suppl 10). Accessed November 26.
[Reference Source](#)
 20. Ritchlin CT, Kavanaugh A, Merola JF, *et al.*: Dual Neutralization of IL-17A and IL-17F with Bimekizumab in Patients with Active, PsA: Results from a 48-Week Phase 2b, Randomized, Double Blind, Placebo-Controlled, Dose- Ranging Study. *Abstr Arthritis Rheumatol.* 2018; 70(suppl 1).
[Reference Source](#)
 21. Wei JCC, *et al.*: OP0234 Efficacy and safety of Brodalumab, an anti-interleukin-17 receptor A monoclonal antibody, in patients with axial spondyloarthritis: a 16 week results of a phase 3, multicenter, randomized, double-blind, placebo-controlled study. In *Oral Presentations. Annals of the Rheumatic Diseases.* 2019; 78(Suppl 2): 195.1–195 (BMJ Publishing Group Ltd and European League Against Rheumatism).
[Publisher Full Text](#)
 22. Mazurov V, Erdes S, Kunder E, *et al.*: OP0028 Efficacy and safety of bcd-085, a novel il-17 inhibitor, in ankylosing spondylitis. results of phase 2 clinical study. *Annals of the Rheumatic Diseases.* 2018; 77(Suppl 2): 64.
[Publisher Full Text](#)
 23. van der Heijde D, Deodhar A, Wei JC, *et al.*: Tofacitinib in patients with ankylosing spondylitis: A phase II, 16-week, randomised, placebo-controlled, dose-ranging study. *Ann Rheum Dis.* 2017; 76(8): 1340–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 24. van der Heijde D, Baraliakos X, Gensler LS, *et al.*: Efficacy and safety of filgotinib, a selective Janus kinase 1 inhibitor, in patients with active ankylosing spondylitis (TORTUGA): Results from a randomised, placebo-controlled, phase 2 trial. *Lancet.* 2018; 392(10162): 2378–87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 25. van der Heijde D, Song IH, Pangan AL, *et al.*: Efficacy and safety of upadacitinib in patients with active ankylosing spondylitis (SELECT-AXIS 1): A multicentre, randomised, double-blind, placebo-controlled, phase 2/3 trial. *Lancet.* 2019; 394(10214): 2108–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 26. Tan AL, Marzo-Ortega H, O'Connor P, *et al.*: Efficacy of anakinra in active ankylosing spondylitis: A clinical and magnetic resonance imaging study. *Ann Rheum Dis.* 2004; 63(9): 1041–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 27. Haibel H, Rudwaleit M, Listing J, *et al.*: Open label trial of anakinra in active ankylosing spondylitis over 24 weeks. *Ann Rheum Dis.* 2005; 64(2): 296–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 28. Song IH, Heldmann F, Rudwaleit M, *et al.*: Treatment of active ankylosing spondylitis with abatacept: An open-label, 24-week pilot study. *Ann Rheum Dis.* 2011; 70(6): 1108–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
 29. Song IH, Heldmann F, Rudwaleit M, *et al.*: Different response to rituximab in tumor necrosis factor blocker-naïve patients with active ankylosing spondylitis and in patients in whom tumor necrosis factor blockers have failed: A twenty-four-week clinical trial. *Arthritis Rheum.* 2010; 62(5): 1290–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 30. Song IH, Heldmann F, Rudwaleit M, *et al.*: One-year follow-up of ankylosing spondylitis patients responding to rituximab treatment and re-treated in case of a flare. *Ann Rheum Dis.* 2013; 72(2): 305–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 31. Sieper J, Porter-Brown B, Thompson L, *et al.*: Assessment of short-term symptomatic efficacy of tocilizumab in ankylosing spondylitis: Results of randomised, placebo-controlled trials. *Ann Rheum Dis.* 2014; 73(1): 95–100.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 32. Sieper J, Braun J, Kay J, *et al.*: Sarilumab for the treatment of ankylosing spondylitis: Results of a Phase II, randomised, double-blind, placebo-controlled study (ALIGN). *Ann Rheum Dis.* 2015; 74(6): 1051–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 33. Pathan E, Abraham S, van Rossen E, *et al.*: Efficacy and safety of apremilast, an oral phosphodiesterase 4 inhibitor, in ankylosing spondylitis. *Ann Rheum Dis.* 2013; 72(9): 1475–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 34. Baeten D, Østergaard M, Wei JCC, *et al.*: Risankizumab, an IL-23 Inhibitor, for Ankylosing Spondylitis: Results of a Randomised, Double-Blind, Placebo-Controlled, Proof-Of-Concept, Dose-Finding Phase 2 Study. *Ann Rheum Dis.* 2018; 77(9): 1295–302.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 35. Deodhar A, Gensler LS, Sieper J, *et al.*: Three Multicenter, Randomized, Double-Blind, Placebo-Controlled Studies Evaluating the Efficacy and Safety of Ustekinumab in Axial Spondyloarthritis. *Arthritis Rheumatol.* 2019; 71(2): 258–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 36. Siebert S, Millar NL, McInnes IB: Why Did IL-23p19 Inhibition Fail in AS: A Tale of Tissues, Trials or Translation? *Ann Rheum Dis.* 2019; 78(8): 1015–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 37. Guidelines on Evaluation of Similar Biotechnological Products (SBPs): WHO Expert Committee on Biological Standardization, Sixtieth report (19-23 October 2009). WHO Technical Report Series No. 977, 2013 - Annex 2.
[Reference Source](#)
 38. January 2017 Position statement: biosimilar medicines.
[Reference Source](#)
 39. Kay J, Schoels MM, Dörner T, *et al.*: Consensus-based Recommendations for the Use of Biosimilars to Treat Rheumatological Diseases. *Ann Rheum Dis.* 2018; 77(2): 165–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 40. Karmacharya P, Duarte-Garcia A, Dubreuil M, *et al.*: Effect of Therapy on Radiographic Progression in Axial Spondyloarthritis: A Systematic Review and Meta-Analysis. *Arthritis Rheumatol.* 2020; 72(5): 733–49.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 41. Baraliakos X, Østergaard M, Gensler LS, *et al.*: Comparison of the Effects of Secukinumab and Adalimumab Biosimilar on Radiographic Progression in Patients With Ankylosing Spondylitis: Design of a Randomized, Phase IIb Study (SURPASS). *Clin Drug Investig.* 2020; 40(3): 269–78.
[PubMed Abstract](#) | [Publisher Full Text](#)
 42. Atar D, Birkeland KI, Uhlig T: 'Treat to Target': Moving Targets From Hypertension, Hyperlipidaemia and Diabetes to Rheumatoid Arthritis. *Ann Rheum Dis.* 2010; 69(4): 629–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
 43. Machado PM, Deodhar A: Treat-to-target in Axial Spondyloarthritis: Gold Standard or Fools' Gold? *Curr Opin Rheumatol.* 2019; 31(4): 344–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 44. Smolen JS, Schöls M, Braun J, *et al.*: Treating axial spondyloarthritis and peripheral spondyloarthritis, especially psoriatic arthritis, to target: 2017 update of recommendations by an international task force. *Ann Rheum Dis.* 2018; 77(1): 3–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Peter Cheung

¹ Division of Rheumatology, National University Hospital, Singapore, Singapore

² Department of Medicine, Yong Loo Lin School of Medicine, National University Singapore, Singapore, Singapore

Competing Interests: No competing interests were disclosed.

2. Daniel Wendling

Department of Rheumatology, CHRU Besançon, Besançon, France

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research