RESEARCH ARTICLE

Evaluation of antioxidant and anti-collagenase activity of *Rosa damascena* L. flower petal and receptacle extract [version 1; peer review: 1 approved, 1 approved with reservations]

Elfitriani Elfitriani ¹, Ahmad Raif^{1,2}, Chrismis N. Ginting¹, Refi Ikhtiari ¹

¹Department of Biomedical Sciences, Faculty of Medicine, Universitas Prima Indonesia, Medan, 20118, Indonesia

²Department of Pathology, Faculty of Health Sciences, Universitas Imelda Medan, Medan, 20239, Indonesia

V1 First published: 17 Jul 2020, 9:716
<https://doi.org/10.12688/f1000research.24772.1>
 Latest published: 17 Jul 2020, 9:716
<https://doi.org/10.12688/f1000research.24772.1>

Abstract

Background: *Rosa damascena* L. is the most notable species of the Rosaceae family in the world, and has been used in food, cosmetics, and the pharmaceutical industry. Bioactive compounds in this flower are known to have several activities, such as antioxidant, antimicrobial, and anti-inflammatory. In this study, the antioxidant and collagenase inhibitory activities of *R. damascena* L. petal and receptacle extracts were evaluated.

Methods: Ethanolic extraction of *R. damascena* L. petals (EERP) and *R. damascena* L. receptacles (EERR) were obtained, and bioactive compounds (flavonoids, phenolics, alkaloids, steroids, tannins, terpenoids, and triterpenoids) were classified by phytochemical screening. Antioxidant activities were analyzed by Ferric Reducing Antioxidant Power (FRAP) assay, while anti-collagenase analysis was examined through the inhibition of collagenase.

Results: Phytochemical test revealed the presence of flavonoids, phenolics, alkaloids, steroids, triterpenoids, triterpenes, and tannin. EERP showed higher FRAP activity ($164.23 \pm 1.34 \mu\text{M Fe(II)}$) than EERR ($12.85 \pm 6.19 \mu\text{M Fe(II)}$). EERP also had higher inhibitory activity of collagenase ($\text{IC}_{50} = 115.48 \pm 1.78 \mu\text{M/mL}$) compared to EERR ($\text{IC}_{50} = 141.96 \pm 6.13 \mu\text{M/mL}$).

Conclusions: *R. damascena* L. petal and receptacle ethanol extracts contain several components, such as phenolics, flavonoids, alkaloids, tannins, terpenes, triterpenoids, and steroids. These extracts exhibit antioxidant activity and collagenase inhibition. *R. damascena* L. petal extract showed higher antioxidant activity through FRAP assay and inhibitory activity of collagenase than *R. damascena* L. receptacle extract.

Keywords

Rosa damascena L., Ferric Reducing Antioxidant Power, antioxidant, anti-collagenase

Open Peer Review

Approval Status

	1	2
version 1		
17 Jul 2020	view	view

1. **Ihsan Iswaldi**, Universitas Prasetya Mulya, Tangerang, Indonesia

2. **Oksana Sytar** , Kiev National University of Taras Shevchenko, Kyiv, Ukraine

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Refi Ikhtiari (refiikhhtiari@unprimdn.ac.id)

Author roles: **Elfitriani E:** Conceptualization, Investigation, Methodology, Writing – Original Draft Preparation; **Raif A:** Conceptualization, Investigation, Methodology, Resources, Writing – Original Draft Preparation; **Ginting CN:** Formal Analysis, Supervision, Visualization, Writing – Review & Editing; **Ikhtiari R:** Investigation, Methodology, Project Administration, Supervision, Validation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Elfitriani E *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Elfitriani E, Raif A, Ginting CN and Ikhtiari R. **Evaluation of antioxidant and anti-collagenase activity of *Rosa damascena* L. flower petal and receptacle extract [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2020, 9:716 <https://doi.org/10.12688/f1000research.24772.1>

First published: 17 Jul 2020, 9:716 <https://doi.org/10.12688/f1000research.24772.1>

Introduction

Antioxidants are compounds that can react with Reactive Oxygen Species (ROS) to prevent the oxidation of lipids, DNA, and proteins¹. Several studies report that ROS correlates with several human diseases, such as cancer², inflammation³, and the aging process⁴. In the aging process, ROS play a role in protein unfolding, lipid breakdown, DNA damage, fragmentation, and collagen disintegration⁵. In the body, ROS promote the activation of collagenase and block collagen synthesis, leading to a decrease in skin elasticity and acceleration of skin wrinkles. This mechanism by ROS is the major factor of degeneration collagen fiber⁶.

Rosa damascena L. is the most important species among the Rosaceae family of flowers. This species has many pharmacological properties and has been used in food, herbal medicine, and cosmetics since ancient times⁷. Previous research showed that *R. damascena* L. has anti-viral, anti-microbial, anti-cancer, anti-depressant, anti-inflammatory, anti-convulsant, and antioxidant activities⁸. Several studies report that phenolic compounds and flavonoids contributed to an increase in antioxidant activity^{7,9–11}. In this study, we evaluate the antioxidant and anti-collagenase activity from ethanol extract of *R. damascena* L. petals and receptacles.

Methods

Sample extraction

R. damascena L. flowers were obtained from Bandung, Indonesia (-6.803662, 107.586633). In total, 1400 g and 700 g of petals and receptacles of *R. damascena* L. were dried and mashed, respectively. The dried samples (250 g petals and 90g receptacles) were soaked with 8000 mL and 3000 mL of 70% ethanol, respectively, and macerated. This solvent was replaced every 3 days and the extracted solution from the first to the last (15th day) was mixed and concentrated using a rotary evaporator at 50°C for 4 h. The extracts were labeled as ethanol extract of rose petals (EERP) and ethanol extract of rose receptacles (EERR).

Phytochemical analysis

Phytochemical analysis was done to detect the presence of flavonoids, phenolics, tannins, steroids/triterpenoids, terpenoids and alkaloids. 10 mg of extracts were used for each analysis following the procedures of Widowati *et al.* without modification¹².

Ferric ion reducing antioxidant power (FRAP) activity

The antioxidant activity of EERP and EERR was investigated using FRAP assay following Sricharoen *et al.* with slight modifications¹³. Briefly, ferric chloride hexahydrate (Merck) and 2,4,6-tripyridyl-s-triazine (TPTZ) (Sigma) were mixed to form the FRAP reagent. 142.5 µL FRAP solution was added into 7.5 µL of sample extract and the mixtures were incubated at 37°C for 5 min. The absorbance of the reaction mixture was read at 539 nm. A set of concentrations 0–200 µg/mL of FeSO₄·7H₂O (Sigma) was used for calibration of the standard curve. The results were expressed as µM Fe(II)/µg extract.

Anti-collagenase assay

Anti-collagenase activity of EERP and EERR was assessed by evaluating the inhibition of collagenase, following the procedure of Wittenauer *et al.*¹⁴ 10 µL collagenase from *Clostridium histolyticum* (0.01 U/mL, Sigma), 60 µL of 50 mM Tricine buffer pH 7.5 (containing 10 mM CaCl₂ (Merck) and 400 mM NaCl (TCL Co., Ltd)), and 30 µL of extract (7.81 – 250 µg/mL) were mixed and incubated at 37°C for 20 min. After incubation, 20 µL of 1 mM N-[3-(2-Furyl)acryloyl]-leu-gly-Pro-Ala (FALGPA, Sigma) in Tricine buffer (Sigma) was added to the mixture. The collagenase inhibitory activities were measured at 335 nm by monitoring for 20 min after starting the reaction. The inhibition value was calculated according to Eq. (1). IC₅₀ values were determined from dose-effect curves.

$$\text{Collagenase inhibition activity (\%)} = \frac{A_{\text{negative control}} - A_{\text{extracts}}}{A_{\text{negative control}}} \times 100\% \quad (1)$$

Data analysis

Statistical analysis using SPSS version 23 was carried out by normality test using numeric variable with Levene test and ANOVA and continued with post-hoc honestly significant difference, and Tukey test to the confidence level of 95% ($\alpha = 0.05$). All data are expressed as mean $\pm$ SEM.

Results and Discussion

Phytochemical analysis

Phytochemical investigations of *R. damascena* L. petal and receptacle extracts in ethanol expressed positive results of flavonoids, phenolic compounds, tannins, steroids, terpenoids, and alkaloids. The details are presented in Table 1.

Table 1. Phytochemical test results for ethanol extracts of *Rosa damascena* L. petals and receptacles. EERP, ethanolic extraction of *R. damascena* L. petals; EERR, ethanolic extraction of *R. damascena* L. receptacles.

Test	Result			
	Observation	EERP	Observation	EERR
Flavonoids	Yellow/ Orange	(+)	Yellow/ Orange	(+)
Phenolics	Green/Blue/Black	(+)	Green/Blue/Black	(+)
Tannins	Red/ Orange Layer	(+)	Yellow gold Layer	(+)
Steroids/Triterpenoids	Orange	(+) Triterpenoids	Green	(+) Steroids
Terpenoids	Purple	(+)	Green/ Gray	(-)
Alkaloids	Orange	(+)	Orange	(+)

The results of phytochemical screening showed steroids are detected in EERP by the bluish-green color, while in the same test, EERR gave a red color indicated it contained triterpenes. Although there was a difference between chemical compounds, both samples contained phenol and flavonoids that act as antioxidants and anti-collagenases. In previous research, Patil *et al.*¹⁵ found that the ethanol extract of *R. damascene* Mill L. contains total phenolic and flavonoid compounds at a higher level than acetone, aqueous, and chloroform extracts.

The antioxidant and anti-collagen activity of flowers depends on the compounds it contains and the solvent used. Phenolics are well known to have antioxidant capability^{9-10,16}, as does flavonoids and alkaloids¹⁷. Steroids, terpenoids, and triterpenoids also have been reported to have antioxidant activity¹⁸. Muccilli *et al.*¹⁹ reported that tannins also play a role in blocking oxidative chain reactions. The solvent extract also has a contribution to increasing antioxidant and anti-collagenase activity. Ethanol extract of *R. damascena* L. petals has been shown to have a lower IC_{50} value than acetone when evaluated by DPPH¹⁵. Ethanol extracts also show the highest phenolic content and antioxidant activity than water extracts of *R. spinosissima*⁹.

Antioxidant activity

Total antioxidant activity of EERP and EERR were further evaluated by FRAP assay. The FRAP value was obtained by plotting the standard curve of $FeSO_4$ at a concentration between 0.78 and 50 $\mu\text{g/mL}$ (Figure 1).

The antioxidant activity of both extracts is presented in Figure 2. In this test, the antioxidant activities were obtained based

on the capacity to reduce ferric (III) to ferrous (II). The result demonstrates that EERP has higher antioxidant capacity compared to EERR.

Table 2 showed FRAP values for different concentrations of EERP and EERR. The data showed normal distribution and was significantly different ($p > 0.005$). Both extracts showed higher antioxidant activity with higher concentrations (Figure 2). The highest antioxidant activities were observed at 50 $\mu\text{g/mL}$ (EERP: $164.23 \pm 1.34 \mu\text{M/mL}$; EERR: $12.85 \pm 6.19 \mu\text{M/mL}$).

Anti-collagenase activity

Collagen is the most abundant protein responsible for conferring skin thickness, strength, and elasticity²⁰. This metalloproteinase enzyme has a vital role in collagen degradation, which increases aging signs²¹. The percentage of collagenase inhibition of EERP and EERR found in the present study is presented in Table 3. Anti-collagenase activity showed a concentration-dependent manner, where higher concentrations of the extracts increased the anti-collagenase (and thus antiaging) activity of extracts. At the highest concentration (250 $\mu\text{g/mL}$) EERP showed the highest inhibition activity compared to EERR ($91.52 \pm 3.44 \mu\text{g/mL}$ and $77 \pm 2.03 \mu\text{g/mL}$, respectively).

The IC_{50} percentage can be seen in Table 4. The IC_{50} value of EERR is higher than EERP ($141.96 \pm 6.13 \mu\text{M/mL}$ and $115.48 \pm 1.78 \mu\text{M/mL}$, respectively). This indicates that EERR has less activity to impede collagenase activity than EERP; however, these results show that there is potential for EERR to inhibit collagenase. Park *et al.*²² found that *R. damascena* extract impacted

Figure 1. Standard curve of $FeSO_4$.

Figure 2. Effect of various concentrations of EERP and EERR in FRAP activity. EERP, ethanolic extraction of *Rosa damascena* L. petals; EERR, ethanolic extraction of *R. damascena* L. receptacles.

Table 2. FRAP activities of EERP and EERR.

EERP, ethanolic extraction of *Rosa damascena* L. petals; EERR, ethanolic extraction of *R. damascena* L. receptacles.

Concentration (µg/mL)	FRAP activity (µM Fe(II))	
	EERP	EERR
0.78	23.70±3.88 ^a	33.35±4.83 ^a
1.56	37.83±5.53 ^{ab}	39.25±6.73 ^a
3.13	50.43±6.32 ^{bc}	47.07±7.25 ^a
6.25	56.63±5.66 ^c	51.97±7.33 ^a
12.50	88.50±7.75 ^d	73.08±9.71 ^b
25.00	115.20±8.24 ^e	85.68±9.28 ^b
50.00	164.23±1.34 ^f	112.85±6.19 ^c

Data presented in the form of mean ± SEM. Different lowercase letters indicate a significant difference in $p < 0.05$ (Post Hoc Tukey HSD Test).

Table 3. Anti-collagenase activity of EERP and EERR.

EERP and EERR were diluted in tricine buffer to reach the final concentration of 7.81; 15.63; 31.25; 62.50; 125.00; 250.00 (µg/mL). EERP, ethanolic extraction of *Rosa damascena* L. petals; EERR, ethanolic extraction of *R. damascena* L. receptacles.

Concentration (µg/mL)	Inhibition activity (%)	
	EERP	EERR
7.81	11.81±1.76 ^a	13.32±2.36 ^a
15.63	21.09±1.49 ^a	17.38±2.04 ^{ab}
31.25	21.68±2.50 ^{a,b}	19.08±1.04 ^b
62.50	33.01±2.42 ^b	33.74±2.27 ^c
125.00	56.46±7.67 ^c	46.19±1.92 ^d
250.00	91.52±3.44 ^d	77.04±2.03 ^e

Data presented in the form of mean ± SEM. Different lowercase letters indicate a significant difference in $p < 0.05$ (Post Hoc Tukey HSD Test).

Table 4. IC₅₀ value of the anti-collagenase activity of EERP and EERR.

EERP, ethanolic extraction of *Rosa damascena* L. petals; EERR, ethanolic extraction of *R. damascena* L. receptacles.

Samples	Equation	R ²	IC ₅₀ (µg/mL)
EERP	$y = 0.3208x + 12.944$	0.991	115.48±1.78
EERR	$y = 0.2598x + 13.146$	0.990	141.96±6.13

metalloproteinase transcription by suppressing AP-1 activation. AP-1 is a protein activated by UVB penetration to the skin, and initiates changes of the extracellular matrix, including collagen, elastin, and proteoglycans degradation. Our results show that ethanol extracts of petals and receptacles of *R. damascene* L. possess anti-collagenase activity.

Conclusion

R. damascene L. petal and receptacle ethanol extracts contain several components, such as phenolics, flavonoids, alkaloids, tannins, terpenes, triterpenoids, and steroids. The pharmacological effects of these extracts exhibit antioxidant activity and collagenase inhibition. *R. damascene* L. petal extract showed higher antioxidant activity through FRAP assay and inhibitory

activity of collagenase than *R. damascene* L. receptacle extract.

Data availability

Underlying data

Figshare: Raw data Analysis of anticollagenase an IC50 value.xlsx, <https://doi.org/10.6084/m9.figshare.12585626.v1>²³

Figshare: Raw data FRAP Activity of Extracts of Rosa Damascene Petals and Receptacles, <https://doi.org/10.6084/m9.figshare.12585671.v1>²⁴

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

References

- Bast A, Haenen G: **Nutritional Antioxidants: It Is Time to Categorise**. In *Antioxidants in Sport Nutrition*. Lamprecht, M., Ed. CRC Press/Taylor & Francis © 2015 by Taylor & Francis Group, LLC.: Boca Raton (FL): 2015. [PubMed Abstract](#)
- Aggarwal V, Tuli HS, Varol A, *et al.*: **Role of Reactive Oxygen Species in Cancer Progression: Molecular Mechanisms and Recent Advancements**. *Biomolecules*. 2019; 9(11): 735. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chelombitko MA: **Role of Reactive Oxygen Species in Inflammation: A Minireview**. *Moscow Univ Biol Sci Bull*. 2018; 73(4): 199–202. [Publisher Full Text](#)
- Xu H, Zheng YW, Liu Q, *et al.*: **Reactive Oxygen Species in Skin Repair, Regeneration, Aging, and Inflammation**. In *Reactive Oxygen Species (ROS) in Living Cells*. 2018. [Publisher Full Text](#)
- Hasan RR, Jat D: **Oxidative stress and antioxidants: an overview**. *International Journal of Advanced Research and Review*. 2017; 2(9): 110–119. [Reference Source](#)
- Choi EK, Guo H, Choi JK, *et al.*: **Extraction conditions of white rose petals for the inhibition of enzymes related to skin aging**. *Lab Anim Res*. 2015; 31(3): 148–152. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Al-Garni AA, Rahimulddin SA, Doghaither HAA, *et al.*: **Evaluation of the antioxidant activities of aqueous extracts of fresh madeni rose petals**. *Int J Sci Nat*. 2017; 8(3): 461–468. [Reference Source](#)
- Mahboubi M: **Rosa damascena as holy ancient herb with novel applications**. *J Tradit Complement Med*. 2015; 6(1): 10–6. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Koczka N, Stefanovits-Bányai É, Ombódi A: **Total polyphenol content and antioxidant capacity of rosehips of some rosa species**. *Medicines (Basel)*. 2018; 5(3): 84. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sengul M, Sener D, Ercisli S: **The determination of antioxidant capacities and chemical properties of rosa (*Rosa damascena* Mill.) products**. *Acta Sci Pol Hortorum Cultus*. 2017; 16(4): 63–72. [Publisher Full Text](#)
- Djamaan A, Noviza D, Septianingsih D, *et al.*: **The use of purple sweet potato (*Ipomoea batatas*) starch as binder in mangosteen peel extracts lozenges formulation**. *Der Pharma Chemica*. 2016; 8(2): 410–414. [Reference Source](#)
- Widowati W, Fauziah N, Herdman H, *et al.*: **Antioxidant and anti aging assays of *Oryza sativa* extracts, vanillin and coumaric acid**. *Journal of Natural Remedies*. 2016; 16(3): 88–99. [Publisher Full Text](#)
- Sricharoen P, Techawongstein S, Chanthai S: **A high correlation indicating for an evaluation of antioxidant activity and total phenolics content of various chilli varieties**. *J Food Sci Technol*. 2015; 52(12): 8077–8085. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wittenauer J, Mäckle S, Sußmann D, *et al.*: **Inhibitory effects of polyphenols from grape pomace extract on collagenase and elastase activity**. *Fitoterapia*. 2015; 101: 179–187. [PubMed Abstract](#) | [Publisher Full Text](#)
- Patil PS, Tatke PA, Gabhe SY: **In vitro Antioxidant and Free Radical Scavenging Activity of Extracts of Rosa damascena Flower Petals**. *American Journal of Phytomedicine and Clinical Therapeutics*. 2015; 589–601. [Reference Source](#)
- Nayebi N, Khalilb N, Kamalinejad M, *et al.*: **A systematic review of the efficacy and safety of Rosa damascena**. Mill. with an overview on its phytopharmacological properties. *Complement Ther Med*. 2017; 34: 129–140. [PubMed Abstract](#) | [Publisher Full Text](#)
- Jiang XL, Wang L, Wang EJ, *et al.*: **Flavonoid glycosides and alkaloids from the embryos of Nelumbo nucifera seeds and their antioxidant activity**. *Fitoterapia*. 2018; 125: 184–190. [PubMed Abstract](#) | [Publisher Full Text](#)
- Koduru S, Jimoh FO, Griegson DS, *et al.*: **Antioxidant Activity of Two Steroid Alkaloids Extracted from Solanum aculeastrum**. *J Pharmacol Toxicol*. 2007; 2(2): 160–167. [Publisher Full Text](#)
- Muccilli V, Cardullo N, Spatafora C, *et al.*: **α -Glucosidase inhibition and antioxidant activity of an oenological commercial tannin. Extraction, fractionation and analysis by HPLC/ESI-MS/MS and 1H NMR**. *Food Chem*. 2017; 215: 50–60. [PubMed Abstract](#) | [Publisher Full Text](#)
- Avila Rodríguez MI, Rodríguez Barroso LG, Sánchez ML: **Collagen: A review on its sources and potential cosmetic applications**. *J Cosmet Dermatol*. 2018; 17(1): 20–26. [PubMed Abstract](#) | [Publisher Full Text](#)
- Alipour H, Raz A, Zakeri S, *et al.*: **Therapeutic applications of collagenase (metalloproteases): A review**. *Asian Pac J Trop Biomed*. 2016; 6(11): 975–981. [Publisher Full Text](#)
- Park B, Hwang E, Seo SA, *et al.*: **Dietary Rosa damascena protects against UVB-induced skin aging by improving collagen synthesis via MMPs reduction through alterations of c-Jun and c-Fos and TGF- β 1 stimulation mediated smad2/3 and smad7**. *J Funct Foods*. 2017; 36: 480–489. [Publisher Full Text](#)
- Elfitriani E: **Raw data Analysis of anticollagenase an IC50 value.xlsx**. figshare. Dataset. 2020. <https://www.doi.org/10.6084/m9.figshare.12585626.v1>
- Elfitriani E: **Raw data FRAP Activity of Extracts of Rosa Damascene Petals and Receptacles**. figshare. Dataset. 2020. <https://www.doi.org/10.6084/m9.figshare.12585671.v1>

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 11 January 2022

<https://doi.org/10.5256/f1000research.27327.r118942>

© 2022 Sytar O. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Oksana Sytar

Department of Plant Biology, Institute of Biology, Kiev National University of Taras Shevchenko, Kyiv, Ukraine

Dear Authors,

The article "Evaluation of antioxidant and anti-collagenase activity of *Rosa damascena* L. flower petal and receptacle extract" is original research work and can be indexed after major revision.

Please, see some comments below:

- In Introduction part it would be good to add more information about previous studies with representatives family Rosaceae.
- It would be good to admit that research studies were done with identification of flavonoids and phenolics in the leaves of some representatives family Rosaceae but less current studies about petal biochemical composition (e.g. Sytar *et al.*, 2015¹; Sytar *et al.*, 2018²). Also for *Rosa damascena* was studied well biochemical composition of essential oils but not petal which can be source of essential oil depending from method of extraction (Mahboubi *et al.*, 2011³; Piotrowicz *et al.*, 2021⁴).
- In the part material and methods for line "Phytochemical analysis was done to detect the presence of flavonoids, phenolics, tannins, steroids/triterpenoids, terpenoids and alkaloids." - please add information if it was used spectrophotometrically or other kind of methods.
- Please, admit that evaluation of anti-collagenase activity of *Rosa damascena* petals was done for the first time. Add information why it is important to study anti-collagenase activity.
- In the part where is written about Phytochemical test results for ethanol extracts of *Rosa damascena* L. petals and receptacles, please add a sentence that the method of visual assessment was recommended to use for pre-screening of anthocyanins content and rutin for some plants (Sytar *et al.*, 2014⁵).

References

1. Sytar O, Bruckova K, Hunkova E, Zivcak M, et al.: The application of multiplex fluorimetric sensor for the analysis of flavonoids content in the medicinal herbs family Asteraceae, Lamiaceae, Rosaceae. *Biol Res*. 2015; **48**: 5 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Sytar O, Hemmerich I, Zivcak M, Rauh C, et al.: Comparative analysis of bioactive phenolic compounds composition from 26 medicinal plants. *Saudi J Biol Sci*. 2018; **25** (4): 631-641 [PubMed Abstract](#) | [Publisher Full Text](#)
3. Mahboubi M, Kazempour N, Khamechian T, Fallah M, et al.: Chemical Composition and Antimicrobial Activity of *Rosa damascena* Mill Essential Oil. *Journal of Biologically Active Products from Nature*. 2011; **1** (1): 19-26 [Publisher Full Text](#)
4. Piotrowicz Z, Tabisz Ł, Waligórska M, Pankiewicz R, et al.: Phenol-rich alternatives for *Rosa x damascena* Mill. Efficient phytochemical profiling using different extraction methods and colorimetric assays. *Sci Rep*. 2021; **11** (1): 23883 [PubMed Abstract](#) | [Publisher Full Text](#)
5. Sytar O, Kosyan A, Taran N, Smetanska I: Anthocyanin's as marker for selection of buckwheat plants with high rutin content. *Gesunde Pflanzen*. 2014; **66** (4): 165-169 [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Plant secondary metabolites, especially plant natural compounds with healthy effects

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 12 August 2021

<https://doi.org/10.5256/f1000research.27327.r86477>

© 2021 Iswaldi I. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Ihsan Iswaldi

Food Business Technology Study Programme, Universitas Prasetiya Mulya, Tangerang, Indonesia

This article can be accepted in this form after the affirmation of several issues:

1. Tannins are highly polymerized compounds of polyphenols and flavonoids are also chemically defined families of polyphenols. So, the authors do not need to mention polyphenols in line with flavonoids and tannins because the last two are classified as polyphenols. Unless the authors want to mention the classes of polyphenols, then write them in sequence.
2. The authors should explain why they only used FRAP assay for antioxidant activity. Why did not they carry out DPPH, ABTS, TEAC, ORAC, and TBARS methods as well and compare the results?

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Food Chemistry and Analysis; Bioactive Ingredients; Analytical Separation

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research