SYSTEMATIC REVIEW

Estimating the impact of pneumococcal conjugate vaccines on childhood pneumonia in sub-Saharan Africa: A systematic review [version 1; peer review: 1 approved, 1 approved with reservations]

Chukwuemeka Onwuchekwa ¹, Bassey Edem², Victor Williams³, Emmanuel Oga⁴

¹Institute of Tropical Medicine, Antwerp, Antwerp, 2000, Belgium

²Medical Research Council Unit The Gambia at London School of Hygiene and Tropical Medicine, Serekunda, The Gambia

³School of Public Health, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

⁴Research Triangle Institute (RTI) International, 6110 Executive Boulevard, Rockville, USA

V1 First published: 23 Jul 2020, 9:765
<https://doi.org/10.12688/f1000research.25227.1>
 Latest published: 09 Nov 2020, 9:765
<https://doi.org/10.12688/f1000research.25227.2>

Abstract

Background: This study aimed to summarise the evidence on the impact of routine administration of 10-valent and 13-valent pneumococcal conjugate vaccines on pneumonia in children under five years of age in sub-Saharan Africa.

Methods: A systematic search of the literature was conducted including primary research reporting on the impact of 10- or 13-valent pneumococcal vaccines on childhood pneumonia in a sub-Saharan African country. Case-control, cohort, pre-post and time-series study designs were eligible for inclusion. Thematic narrative synthesis was carried out to summarise the findings.

Results: Eight records were included in the final analysis, 6 records were pre-post or time-series studies, 1 was a case-control study and 1 report combined pre-post and case-control studies. Vaccine impact on clinical pneumonia measured as percentage reduction in risk (%RR) was mostly non-significant. The reduction in risk was more consistent in radiological and pneumococcal pneumonia.

Conclusions: Evidence of the positive impact of routine infant pneumococcal vaccination on clinical pneumonia incidence in sub-Saharan Africa is inconclusive. Ongoing surveillance and further research is required to establish the long term trend in pneumonia epidemiology and aetiology after PCV introduction.

PROSPERO registration: CRD42019142369 30/09/19

Keywords

Pneumonia, Streptococcus pneumoniae, child, sub-Saharan

Open Peer Review

Approval Status

	1	2
version 2 (revision) 09 Nov 2020		 view
version 1 23 Jul 2020	 view	 view

1. **Blandina Theophil Mmbaga** , Kilimanjaro
Clinical Research Institute, Moshi, Tanzania
2. **Dodi Safari**, Eijkman Institute for Molecular
Biology, Jakarta, Indonesia

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding authors: Chukwuemeka Onwuchekwa (emyonwuchekwa@gmail.com), Bassey Edem (bedem@mrc.gm)

Author roles: **Onwuchekwa C:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Edem B:** Data Curation, Formal Analysis, Investigation, Methodology, Validation, Writing – Review & Editing; **Williams V:** Data Curation, Formal Analysis, Methodology, Validation, Writing – Review & Editing; **Oga E:** Data Curation, Formal Analysis, Methodology, Validation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Onwuchekwa C *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Onwuchekwa C, Edem B, Williams V and Oga E. **Estimating the impact of pneumococcal conjugate vaccines on childhood pneumonia in sub-Saharan Africa: A systematic review [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2020, 9:765 <https://doi.org/10.12688/f1000research.25227.1>

First published: 23 Jul 2020, 9:765 <https://doi.org/10.12688/f1000research.25227.1>

Introduction

Pneumonia is one of the leading causes of childhood deaths globally, particularly in sub-Saharan Africa¹. Annually over a 100 million cases of pneumonia are reported in children less than five years of age, mostly in poor countries in Africa and Asia²⁻⁴.

Streptococcus pneumoniae is the major cause of childhood pneumonia deaths and is the leading cause of vaccine-preventable child deaths globally⁵⁻⁷. Pneumococcal pneumonia causes between 1 and 4 million episodes of pneumonia in Africa yearly⁸. There are currently 97 known serotypes of *S. pneumoniae*, characterised by the polysaccharide capsule antigen⁹. The 97 serotypes differ in carriage potential and propensity to cause invasive disease including pneumonia, otitis media and meningitis^{10,11}, with the 13 most common serotypes accounting for 70 – 75% of invasive pneumococcal disease globally^{8,11,12}.

Pneumococcal conjugate vaccines (PCV) have been licensed since 2000; previous polysaccharide-based vaccines were found to have poor immunogenicity in children⁸. An initial 7-valent PCV included serotypes 4, 6B, 9V, 14, 18C, 19F and 23F providing cover against 67% of disease-causing serotypes. The 10-valent and 13-valent PCV include in addition to the 7-valent serotypes 1, 5 and 7F; and 1, 3, 5, 7F, 19A, and 6A. The 10-valent vaccine covers 70% - 84% while the 13-valent vaccine covers about 74% - 88% % of invasive disease-causing serotypes^{8,13,14}. Sub-Saharan African countries like South-Africa and The Gambia introduced the 7-valent PCV into routine infant vaccination schedule in 2009^{15,16}. Many countries in sub-Saharan Africa through the support of Gavi, the vaccine alliance, have incorporated the 10- or 13-valent PCV into their expanded program of immunization (EPI) schedules. PCV vaccination has been associated with a significant decline in invasive pneumococcal disease incidence globally and at individual country level^{9,15,16}. While there is evidence of effectiveness against invasive pneumococcal disease, the impact of vaccination on clinical and radiological pneumonia remains unclear¹⁶.

Objectives

This review aims to summarise the existing evidence on the impact of routine administration of pneumococcal conjugate vaccines on clinical pneumonia, radiological pneumonia and pneumococcal pneumonia in children under five years of age in sub-Saharan Africa.

Methods

Study protocol

The systematic review protocol was developed in accordance with the PRISMA-P guidelines¹⁷, and registered with PROSPERO on 30 September 2019 (CRD42019142369).

Eligibility criteria

We included primary, individual and population-based studies conducted in sub-Saharan Africa evaluating 10-valent or 13-valent PCV impact in children published in English since 1 January 2010. This time was chosen because the earliest countries in this region

to introduce PCV into routine infant vaccine schedule did so in 2009. The eligibility criteria are detailed in [Table 1](#) below.

Studies with invasive pneumococcal disease (IPD) as outcome were considered for inclusion if pneumonia cases were reported separately.

The study designs eligible for inclusion were pre-post quasi-experimental, interrupted time series, Cohort, and Case-control studies. For pre-post studies and interrupted time series, we included only studies where the final outcome assessment was made at least 3 years after vaccine introduction.

Information sources

We conducted a search of peer-reviewed and grey literature relating to the study question. [PubMed](#) search was first conducted on 16 July 2019, with final search on 31 July 2019. [Scopus](#) search was conducted on 20 Jul 2019, [Embase](#) (Ovid) was searched on the 29 Jul 2019. Other peer-review sources include [Africa-Wide Information](#) (29 July 2019) and [African Index Medicus](#) (24 July 2019). Grey literature sources include [OpenGrey](#) (20 July 2019), and [ProQuest Dissertation & theses global](#) (20 July 2019), London School of Hygiene and Tropical Medicine research online (22 July 2019) and University of Edinburgh library (28 July 2019).

Search strategy

The search strategy combined the key concepts of the research question and was based on the PICOS framework. The three components or concepts were: population of interest (children below 5 years of age), intervention being investigated (pneumococcal conjugate vaccine) and the outcome of interest (pneumonia). The Boolean operators AND, and OR were used to combine the search concepts.

Further details of the database search strategy and date of searches can be found as extended data¹⁸.

Study selection process

Two members of the review team conducted the database screening independently. Reading through the titles and abstracts

Table 1. PICO framework for formulating the review question.

Concept	Elaboration
Population of interest	Children between 1 – 59 months of age from any of the 46 countries in sub-Saharan Africa.
Intervention	Routine immunization with 10- or 13-Valent PCV
Comparison (or control)	Control group (either contemporary or historical)
Outcomes	Clinical pneumonia, radiological pneumonia or pneumococcal pneumonia
Study type	Interrupted time-series, pre-post studies, case control and cohort

of the search results we identified records to be included in the full-text screening based on the eligibility criteria. Records for which there was uncertainty or disagreement about eligibility during the title and abstract screening were included for full-text screening. The second stage of the screening involved reading the full-text of the records to determine if they were eligible for inclusion. Finally, we searched through the references of eligible papers for other relevant publications that could be included in the review. The PRISMA flow diagram for the study selection procedure is shown in [Figure 1](#) in the result section.

Data collection process and data items

The following information was collected for each included record: year of publication, study location and country, study setting (hospital-based or population-based), study design, study aim (to assess impact or effectiveness), data sources (clinical or laboratory), study population description, HIV status of participants, type of PCV in current use (PCV 10 or PCV13), year PCV7 introduced, year PCV10 or PCV13 introduced, reported coverage during PCV10 or PCV13 period, baseline and post-vaccination periods (for pre-post and interrupted

Figure 1. PRISMA flow diagram showing review selection process.

time-series), outcome definition, outcome measure and confidence interval if reported. The information was collected directly into an extraction form in Excel by one member of the review team and crosschecked by a second member¹⁹. Disagreement was resolved by consensus after discussion with another member of the review team. No additional information was sought from investigators or authors.

Risk of bias in individual studies

We assessed the quality of case-control studies assessing vaccine effectiveness, we used the National Heart, Lung and Blood Institute Quality assessment tool for case-control studies. We adapted the National Heart, Lung and Blood Institute Quality Assessment Tool for Before-After studies with No Control Group to assess quality in pre-post and interrupted time series analyses²⁰. The study quality assessment tools are available at <https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools> and available as extended data²¹.

Summary measures

The primary outcomes of interest in this review were clinical pneumonia and radiological pneumonia. The secondary outcome was pneumococcal pneumonia. In pre-post and interrupted time-series studies comparing outcome incidence before and after PCV introduction, measures were presented as percentage reduction in incidence and incidence ratios. Where possible incidence ratios were converted to percentage reduction in incidence: $\% \text{ reduction in risk} = (1 - aRR) \times 100\%$; where aRR is the adjusted Risk/Rate ratio for post- and pre-vaccination periods²². In case-control studies we presented adjusted vaccine effectiveness (aVE) as reported by the authors. When adjusted

odd Ratios (aOR) were presented we estimated aVE as: $\text{adjusted vaccine effectiveness} = (1 - aOR) \times 100\%$ ¹⁹; where aOR = adjusted odd ratio. All calculations were made in Microsoft Excel 2016, and graphical presentations were performed in Stata 13²³.

Synthesis of results

Planned quantitative synthesis could not be conducted in this review due to variation in included studies. We therefore, present a narrative synthesis of the impact of PCV on childhood clinical pneumonia, radiological pneumonia and pneumococcal pneumonia.

Results

Study selection

The database search produced 3581 distinct titles, from which we identified 34 relevant records for the full-text review. We excluded a further 26 records after full-text assessment. Eight records were included in the review as shown in Figure 1 below.

Study characteristics

Four of the included studies were conducted in South Africa (23 – 26), two were conducted in Kenya^{24,25}, and one each from Rwanda²⁶, and The Gambia²⁷. There were 7 studies with pre-post or interrupted time-series design and two case-control studies (Mackenzie *et al.* included report on a case-control study). Most of the study locations had 7-valent PCV introduced before transitioning to the 13-valent PCV except in Kenya where the 10-valent PCV was used without a 7-valent PCV period. Table 2 below shows the characteristics of articles included in the review.

Table 2. Characteristics of studies included in review.

Author	Country	Study design	Data sources	Population	PCV	Year PCV7 introduced	Year PCV 10 or 13 introduced	Baseline period	Post-intervention period
Von Mollendorf <i>et al.</i> , 2017 ²⁸	South Africa	Pre-post	Laboratory surveillance	< 60 months	13	2009	2011	2005 – 2008	2012 - 2013
Mackenzie <i>et al.</i> , 2017 ²⁹	URR, The Gambia	Pre-post	Population surveillance	2 to 59 months	13	2009	2011	2008 – 2010	2014 - 2015
Mackenzie <i>et al.</i> , 2017 ^{*29}	URR, The Gambia	Case-control	Clinical	3 to 59 months	13	2009	2011	NA	NA
Hammit <i>et al.</i> , 2019 ²⁴	Kilifi, Kenya	Pre-post	Clinical surveillance	< 60 months	10	NA	2011	1999 – 2010	2012 - 2016
Gatera <i>et al.</i> , 2016 ²⁵	Rwanda (five districts)	Pre-post	Clinical and laboratory surveillance	< 60 months	13	2009	2011	2002 – 2009	2012
Silaba <i>et al.</i> , 2019 ³⁰	Kilifi, Kenya	ITS	Population surveillance	2 to 59 months	10	NA	2011	2002 – 2011	2011 - 2015
Mahdi <i>et al.</i> , 2015 ³¹	Soweto, Cape town, KwaZulu-Natal. South Africa	Case-control	Clinical	2 to 42 months	13	2009	2011	NA	NA
Solomon <i>et al.</i> , 2017 ³²	Soweto, South Africa	Pre-post	Population surveillance	< 60 months	13	2009	2011	2006 – 2008	2014
Tempia <i>et al.</i> , 2015 ³³	Soweto, South	Pre-post	Laboratory surveillance	< 24 months	13	2009	2011	2009 -	2011 – 2012

NA: Not applicable, * Reported along with the pre-post study.

Risk of bias within studies

One of the seven population-based studies were considered to be of poor quality with significant risk of bias. None of the population-based studies were blinded hence there is the possibility of bias in the findings. The two case-control reports were graded as having good quality. Details on the study quality assessment are available as extended data²¹.

Result of individual studies

There was variation in the definition of clinical and radiological pneumonia, and in the method of identification of pneumococcal pneumonia among the included studies. Some studies applied the WHO standards for identification of radiological pneumonia and clinical pneumonia^{25,27,34}. One study based pneumonia diagnosis on ICD-10 coding from clinical notes³⁵. Studies reporting on pneumococcal pneumonia were based on culture with occasional confirmation by polymerase chain reaction (PCR). Table 3 below summarises the findings of the included studies.

Synthesis of results

Impact on clinical pneumonia. Four population-based studies measured the impact of 10- or 13-valent vaccine on clinical pneumonia with inconsistent findings. Silaba *et al.* reported a decline in hospitalization for severe or very severe pneumonia based on WHO clinical definition²⁵. Mackenzie *et al.* showed no significant decline in clinical pneumonia incidence in all under-five age groups three years after 13-valent PCV introduction³⁰. Solomon *et al.* estimated a significant decline in pneumonia hospitalizations in infants and children between 24 – 59 months based on Bayesian methods. This effect was also found among HIV infected and HIV uninfected children³⁵. However, these figures were estimated around a 50% credible interval. Figure 2 graphically displays the reported reduction in clinical pneumonia incidence.

Impact on radiological pneumonia. Two impact studies evaluated radiological pneumonia as outcomes^{25,27}. Mackenzie

Table 3. Summary of effect reported in studies included in the review.

Study	population	Clinical pneumonia	Radiological pneumonia	Pneumococcal pneumonia
		Percent rate reduction (95% CI)	Percent rate reduction (95% CI)	Percent rate reduction (95% CI)
Von Mollendorf <i>et al.</i> , 2017 ²⁸	< 60 months	--	--	65 (64 – 67)
	HIV positive	--	--	51 (49 – 55)
	12 – 59 months	--	--	69 (67 – 71)
Mackenzie <i>et al.</i> , 2017 ²⁹	2 – 11 months	2 (-4 – 8)	23 (7 – 36)	--
	12– 23 months	-6 (-15 – 2)	29 (12 - 42)	--
	24 – 59 months	-7 (-18 – 2)	22 (1 - 39)	--
Hammit <i>et al.</i> , 2019 ²⁴	< 60 months	--	--	85 (66 – 93)
Gatera <i>et al.</i> , 2016 ²⁵	< 60 months	15 (Not reported)	--	--
Silaba <i>et al.</i> , 2019 ³⁰	2 – 59 months	27 (3 – 46)	48 (14 – 68)	--
	2 – 11 months	--	27 (-36 – 61)	--
	12 – 23 months	--	46 (-10 – 73)	--
	24 – 59 months	--	11 (-69 – 53)	--
Solomon <i>et al.</i> , 2017 ^{32**}	HIV infected	33 (6 -52) ¶	--	--
	HIV uninfected or unconfirmed	39 (24 – 50) ¶	--	--
	3 – 11 months	39 (10 – 57) ¶	--	--
	12 – 23 months	15 (-37 – 42) ¶	--	--
	24 – 59 months	45 (17 – 62) ¶	--	--
Tempia <i>et al.</i> , 2015 ³³	HIV uninfected	--	--	66.8 (43.8 – 81.3)
		--	--	64.0 (52.6 – 81.2)
		Adjusted vaccine effectiveness (95% CI)	Adjusted vaccine effectiveness (95% CI)	Adjusted vaccine effectiveness (95% CI)
Mackenzie <i>et al.</i> , 2017 ^{*29}	3 – 59 months	--	28 (-23 – 58)	--
Mahdi <i>et al.</i> , 2015 ³¹	Hospital controls	--	20.0 (-9.3 – 41.6)	--
	Community controls	--	32.1 (4.6 – 51.6)	--

*Based on WHO clinical classification of severe and very severe pneumonia, ** pneumonia hospitalization based on ICD-10 coding, ¶ 50% credible interval based on Bayesian methods (negative values indicate an increase in incidence), € Identification based on polymerase chain reaction, ± pneumonia classified based on WHO criteria or based on abnormal CXR plus C-reactive protein >40mg/L.

Figure 2. Forest plot showing the percentage risk reduction in clinical pneumonia incidence in included studies and population.

et al. reported a decline in WHO defined radiological pneumonia in all age groups with decline in the range of 22 – 29%. This was most pronounced in the 12 to 23-month age group. Silaba *et al.* also reported a 48% decline in radiological pneumonia in the entire under-five population. A similar age-related trend was observed, with children between 12 to 23-month age experienced the largest reduction in radiological pneumonia²⁵.

The case-control study reported by Mackenzie *et al.* using community controls showed a vaccine effectiveness of about 28%, however, this did not reach statistical significance. Mahdi *et al.* reported adjusted vaccine effectiveness measures using both community and hospital controls³⁴. Vaccine effectiveness was significant with community controls (aVE 32.1%, 95% CI 4.6% - 51.6%), but not with hospital controls (aVE 20%, 95% CI -9.3% – 41.6%).

Impact on pneumococcal pneumonia. All three studies that reported on pneumococcal pneumonia were based on microbiological diagnosis with or without PCR confirmation^{24,29,36}. Included studies consistently showed decline in cases of pneumococcal pneumonia after vaccine introduction irrespective of method of pneumococcal identification. Figure 3 shows the reduction in pneumococcal pneumonia incidence reported from included studies.

Discussion

Summary of evidence

This review set out to answer the question: has the introduction of pneumococcal conjugate vaccine resulted in a decline in childhood pneumonia? From our review, we can summarise that the population impact of pneumococcal vaccination on pneumonia depends largely on how pneumonia is classified. We observed

overall that when the outcome is clinical pneumonia, the impact tends to be modest at best. We see this in the reports by Mackenzie *et al.*, Silaba *et al.*, and Solomon *et al.* However, it is important to note that the severity of pneumonia differed in these studies. Mackenzie *et al.* evaluated clinical pneumonia at the population level and showed results that were not statistically significant in all age groups. Silaba *et al.*, however, looked at clinical pneumonia hospitalizations (i.e. severe or very severe pneumonia) and showed significant difference in incidence after vaccine introduction. We also see a subtle decline in hospitalizations for clinical pneumonia based on ICD-10 coding. Clinical pneumonia is a non-specific outcome and therefore may not be ideal for evaluating disease-specific vaccine impact. There was a more positive but still modest impact observed with radiological pneumonia as an outcome. This is likely because radiological pneumonia is more specific for pneumococcal disease. Overall, there was a consistent decline in pneumococcal pneumonia cases in the post-vaccine period in reported impact studies^{24,29,36}. Two studies that reported on vaccine effect in HIV infected children showed that pneumococcal vaccine had similar impact as with HIV uninfected children^{23,25}.

The overall trend in the findings from this review is similar to reports from other parts of the world. Meta-analyses from other regions have found modest decline in clinical and radiological pneumonia hospitalizations. A meta-analysis using global data from 12 pre-post and time series studies produced a pooled reduction in hospitalization rates for clinical pneumonia of 17% and 9% in the under 24 months and 24 – 59 months' age groups. However, they reported a more significant decline in hospitalization rates due to radiological pneumonia of 31% and 24% in the same age groups³⁷. A randomised placebo-controlled trial on 9-valent vaccine showed vaccine effectiveness of 37% based on

Figure 3. Forest plot showing the percentage risk reduction in pneumococcal pneumonia incidence in included studies and population.

per-protocol population²². Overall, it appears that pneumococcal vaccination has its greatest impact in preventing the more severe forms of pneumonia for which children are likely to be hospitalised. This is probably due to the finding that bacterial pathogens are more likely to cause severe pneumonia²⁸. The minimal impact on clinical pneumonia might suggest an increase in other forms of pneumonia or may be due to serotype replacement^{30–32}. We observed an age-related trend in vaccine effect, with the 12 to 23-month age group experiencing the greatest benefit. This might be due in part to a greater proportion of children under 12 months having not completed their vaccination schedules; and potentially waning immunity in the older age groups³³.

Limitations

This review has some limitations that have to be considered when interpreting our findings. First was the inconsistency in the definition of pneumonia outcomes between studies, which made combining the impact measured between studies impractical. While some studies used comparable methods for outcome ascertainment, this was not consistent across studies. The WHO standardised definition of clinical and radiological pneumonia is markedly helpful in this situation as it ensures consistency across studies³⁸. Studies conducted in locations with a functional health and demographic surveillance system like in the Upper River region of The Gambia and in Kilifi, in Kenya, were particularly robust as they combined consistent pneumonia surveillance methods with up-to-date population information^{24,37}. Some studies relied on routine clinical data which is usually of variable quality^{26,35}.

It is also important to note that while we set out to evaluate the impact of 10-valent and 13-valent vaccines, all of the study

locations except in Kenya had a period of 7-valent vaccine use. Therefore, it is impossible to separate the effect of the 7-valent from the 10- and 13-valent vaccines since the 7-valent PCV might have changed the pre-existing disease trend.

Like all time-trend studies - including pre-post studies, phenomena such as regression to the mean, seasonality, trend, and history bias have to be considered in the analysis. By including a control outcome and conducting sensitivity analysis, some of the included studies considered the impact of history bias and trend on their results³⁹.

Publication bias is a potential limitation of this review. However, due to the small number of reports in each outcome category, we did not formally assess for publication bias using a funnel plot⁴⁰. We are therefore unable to comment on publication bias in this review. Finally, due to limited resources we were unable to include studies published in other languages; hence, language bias cannot be ruled out in our review.

Conclusion

To the best of our knowledge this is the first systematic attempt at summarising the impact and effectiveness of routine pneumococcal vaccination on childhood pneumonia from this region. The 10- and 13-valent PCV use as part of infant immunization is effective in preventing the more severe forms of childhood pneumonia. There appears to be a smaller effect on clinical pneumonia especially when all severity spectra are included. There is the need for consistency in pneumonia definition for the purpose of disease surveillance and the WHO clinical definitions provide an appropriate option ensuring ease of implementation and reproducibility. One major issue encountered was that few studies had applied comparable pneumonia

definitions in estimating disease burden prior to pneumococcal vaccine introduction hence making trend analysis difficult. There is the need for the generation of updated information on the causes of pneumonia in this region in this era of extensive pneumococcal vaccine use. Ongoing surveillance is needed to investigate the long-term trend in childhood pneumonia in the PCV era in sub-Saharan Africa.

Data availability

Underlying data

All data underlying the results are available as part of the article and no additional source data are required.

Extended data

This project contains the following extended data:

Figshare: Detailed search <https://doi.org/10.6084/m9.figshare.12656309.v2>¹⁸

Figshare: Data extraction form <https://doi.org/10.6084/m9.figshare.12608264.v1>¹⁹

Figshare: Quality assessment tool <https://doi.org/10.6084/m9.figshare.12608264.v1>²¹

Reporting guidelines

Figshare: PRISMA checklist for 'Estimating the impact of pneumococcal conjugate vaccines on childhood pneumonia in sub-Saharan Africa: A systematic review'. <https://doi.org/10.6084/m9.figshare.12608672.v2>⁴¹

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

Acknowledgment

The authors acknowledge the invaluable input of Dr Kristien Verdonck in reviewing the initial manuscript and guiding interpretation and presentation of the review findings. A previous version of this article is available from bioRxiv⁴².

References

- Black RE, Morris SS, Bryce J: **Where and why are 10 million children dying every year?** *Lancet*. Elsevier; 2003 [cited 2016 Jul 16]; 361(9376): 2226–34. [PubMed Abstract](#) | [Publisher Full Text](#)
- Howie SRC, Murdoch DR: **Global childhood pneumonia: the good news, the bad news, and the way ahead.** *Lancet Glob Health*. 2019; 7(1): e4–e5. [PubMed Abstract](#) | [Publisher Full Text](#)
- Nair H, Simões EAF, Rudan I, *et al.*: **Global and regional burden of hospital admissions for severe acute lower respiratory infections in young children in 2010: A systematic analysis.** *Lancet*. 2013; 381(9875): 1380. [Publisher Full Text](#) | [Free Full Text](#)
- McAllister DA, Liu L, Shi T, *et al.*: **Global, regional, and national estimates of pneumonia morbidity and mortality in children younger than 5 years between 2000 and 2015: a systematic analysis.** *Lancet Glob Heal*. 2019; 7(1): e47–e57. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Benavides JA, Ovalle OO, Salvador GR, *et al.*: **Population-based surveillance for invasive pneumococcal disease and pneumonia in infants and young children in Bogotá, Colombia.** *Vaccine*. 2012; 30(40): 5886–5892. [PubMed Abstract](#) | [Publisher Full Text](#)
- Andrade AL, Oliveira R, Vieira MA, *et al.*: **Population-based surveillance for invasive pneumococcal disease and pneumonia in infants and young children in Goiânia, Brazil.** *Vaccine*. 2012; 30(10): 1901–9. [PubMed Abstract](#) | [Publisher Full Text](#)
- Arguedas A, Abdelnour A, Soley C, *et al.*: **Prospective epidemiologic surveillance of invasive pneumococcal disease and pneumonia in children in San José, Costa Rica.** *Vaccine*. 2012; 30(13): 2342–8. [PubMed Abstract](#) | [Publisher Full Text](#)
- Zar HJ, Madhi SA: **Pneumococcal conjugate vaccine - A health priority.** *S Afr Med J*. 2008; 98(6): 463–7. [PubMed Abstract](#)
- Weiser JN, Ferreira DM, Paton JC: **Streptococcus pneumoniae: Transmission, colonization and invasion.** *Nat Rev Microbiol*. 2018; 16(6): 355–367. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mitchell AM, Mitchell TJ: **Streptococcus pneumoniae: Virulence factors and variation.** *Clin Microbiol Infect*. 2010; 16(5): 411–8. [PubMed Abstract](#) | [Publisher Full Text](#)
- Ramirez M: **Streptococcus pneumoniae.** *Molecular Medical Microbiology: Second Edition*. 2014.
- AlonsoDeVelasco E, Verheul AF, Verhoeft J, *et al.*: **Streptococcus pneumoniae: virulence factors, pathogenesis, and vaccines.** *Microbiol Rev*. 1995; 59(4): 591–603. [PubMed Abstract](#) | [Free Full Text](#)
- Tettelin H, Nelson KE, Paulsen IT, *et al.*: **Complete genome sequence of a virulent isolate of Streptococcus pneumoniae.** *Science*. 2001; 293(5529): 498–506. [PubMed Abstract](#) | [Publisher Full Text](#)
- Kadioglu A, Weiser JN, Paton JC, *et al.*: **The role of Streptococcus pneumoniae virulence factors in host respiratory colonization and disease.** *Nat Rev Microbiol*. 2008; 6(4): 288–301. [PubMed Abstract](#) | [Publisher Full Text](#)
- Bogaert D, De Groot R, Hermans PWM: **Streptococcus pneumoniae colonisation: The key to pneumococcal disease.** *Lancet Infect Dis*. 2004; 4(3): 144–54. [PubMed Abstract](#) | [Publisher Full Text](#)
- Ngocho JS, Magoma B, Olomi GA, *et al.*: **Effectiveness of pneumococcal conjugate vaccines against invasive pneumococcal disease among children under five years of age in Africa: A systematic review.** *PLoS One*. 2019; 14(2): e0212295. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Moher D, Liberati A, Tetzlaff J, *et al.*: **Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement.** *PLoS Med*. 2009; 6(7): e1000097. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Onwuchekwa C, Edem BE, Oga EA, *et al.*: **Detailed search.** *figshare*. Dataset. 2020. <http://www.doi.org/10.6084/m9.figshare.12656309.v2>
- Onwuchekwa C, Edem BE, Oga EA, *et al.*: **Data extraction form.** *figshare*. Dataset. 2020. <http://www.doi.org/10.6084/m9.figshare.12608198.v1>
- Institute NHL, B: **Study Quality Assessment Tools-NHLBI.** NIH. 2016. [Reference Source](#)
- Onwuchekwa C: **Quality assessment tool.** *figshare*. Dataset. 2020. <http://www.doi.org/10.6084/m9.figshare.12608264.v1>
- Cutts FT, Zaman SMA, Enwere G, *et al.*: **Efficacy of nine-valent pneumococcal conjugate vaccine against pneumonia and invasive pneumococcal disease in The Gambia: Randomised, double-blind, placebo-controlled trial.** *Lancet*. 2005; 365(9465): 1139–46. [PubMed Abstract](#) | [Publisher Full Text](#)
- StataCorp: **Stata Statistical Software: Release 13.** College Station, TX: StataCorp LP. 2013.
- Hammit LL, Etyang AO, Morpeth SC, *et al.*: **Effect of ten-valent pneumococcal conjugate vaccine on invasive pneumococcal disease and nasopharyngeal carriage in Kenya: a longitudinal surveillance study.** *Lancet*. [Internet]. 2019 [cited 2019 Jul 22]; 393(10186): 2146–54. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Silaba M, Ooko M, Bottomley C, *et al.*: **Effect of 10-valent pneumococcal conjugate vaccine on the incidence of radiologically-confirmed pneumonia and clinically-defined pneumonia in Kenyan children: an interrupted time-series analysis.** *Lancet Glob Heal*. [Internet]. 2019; 7(3): e337–46. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gatera M, Uwimana J, Manzi E, *et al.*: **Use of administrative records to assess**

- pneumococcal conjugate vaccine impact on pediatric meningitis and pneumonia hospitalizations in Rwanda.** *Vaccine*. [Internet]. 2016 [cited 2019 Jul 16]; **34**(44): 5321–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Mackenzie GA, Hill PC, Sahito SM, *et al.*: **Impact of the introduction of pneumococcal conjugate vaccination on pneumonia in The Gambia: population-based surveillance and case-control studies.** *Lancet Infect Dis*. [Internet]. 2017 [cited 2019 Jul 20]; **17**(9): 965–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 28. O'Brien KL, Baggett HC, Brooks WA, *et al.*: **Causes of severe pneumonia requiring hospital admission in children without HIV infection from Africa and Asia: the PERCH multi-country case-control study.** *Lancet*. 2019; **394**(10200): 757–779.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 29. Von Mollendorf C, Tempia S, Von Gottberg A, *et al.*: **Estimated severe pneumococcal disease cases and deaths before and after pneumococcal conjugate vaccine introduction in children younger than 5 years of age in South Africa.** *PLoS One*. 2017; **12**(7): e0179905.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 30. Madhi S: **Clinical spectrum and complications of childhood pneumonia in the era of bacterial conjugate vaccines.** *Int J Infect Dis*. 2014; **21**(SUPPLEMENT 1): 25.
[Publisher Full Text](#)
 31. Kwambana-Adams B, Hanson B, Worwui A, *et al.*: **Rapid replacement by non-vaccine pneumococcal serotypes may mitigate the impact of the pneumococcal conjugate vaccine on nasopharyngeal bacterial ecology.** *Sci Rep*. 2017; **7**(1): 8127.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 32. Weinberger D, Malley R, Lipsitch M: **Serotype replacement in disease after pneumococcal vaccination.** *Lancet*. 2011; **378**(9807): 1962–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 33. Berical AC, Harris D, Dela Cruz CS, *et al.*: **Pneumococcal vaccination strategies. An update and perspective.** *Ann Am Thorac Soc*. 2016; **13**(6): 933–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 34. Madhi SA, Groome MJ, Zar HJ, *et al.*: **Effectiveness of pneumococcal conjugate vaccine against presumed bacterial pneumonia hospitalisation in HIV-uninfected South African children: A case-control study.** *Thorax*. 2015 [cited 2019 Jul 16]; **70**(12): 1149–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
 35. Izu A, Solomon F, Nzenze SA, *et al.*: **Pneumococcal conjugate vaccines and hospitalization of children for pneumonia: a time-series analysis, South Africa, 2006–2014.** *Bull World Health Organ*. [Internet]. 2017 [cited 2019 Jul 16]; **95**(9): 618–28.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 36. Tempia S, Wolter N, Cohen C, *et al.*: **Assessing the impact of pneumococcal conjugate vaccines on invasive pneumococcal disease using polymerase chain reaction-based surveillance: An experience from South Africa.** *BMC Infect Dis*. [Internet]. 2015 [cited 2019 Jul 16]; **15**(1): 450.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 37. Alicino C, Paganino C, Orsi A, *et al.*: **The impact of 10-valent and 13-valent pneumococcal conjugate vaccines on hospitalization for pneumonia in children: A systematic review and meta-analysis.** *Vaccine*. 2017; **35**(43): 5776–5785.
[PubMed Abstract](#) | [Publisher Full Text](#)
 38. Cherian T, Mulholland EK, Carlin JB, *et al.*: **Standardized interpretation of paediatric chest radiographs for the diagnosis of pneumonia in epidemiological studies.** *Bull World Health Organ*. 2005; **83**(5): 353–9.
[PubMed Abstract](#) | [Free Full Text](#)
 39. Bernal JL, Cummins S, Gasparrini A: **Interrupted time series regression for the evaluation of public health interventions: A tutorial.** *Int J Epidemiol*. 2017; **46**(1): 348–355.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 40. Lau J, Ioannidis JP, Terrin N, *et al.*: **The case of the misleading funnel plot.** *BMJ*. 2006; **333**(7568): 597–600.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 41. Onwuchekwa C, Edem BE, Oga EA, *et al.*: **PRISMA checklist.** *figshare*. Figure. 2020.
<http://www.doi.org/10.6084/m9.figshare.12608672.v2>
 42. Chukwuemeka O, Bassey EE, Williams V, *et al.*: **Estimating the impact of pneumococcal conjugate vaccines on childhood pneumonia in sub-Saharan Africa: A systematic review.** *bioRxiv*. 865154. 2019.
[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 02 November 2020

<https://doi.org/10.5256/f1000research.27840.r73233>

© 2020 Safari D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Dodi Safari

Molecular Bacteriology Unit, Eijkman Institute for Molecular Biology, Jakarta, Indonesia

In this systematic review with the title "Estimating the impact of pneumococcal conjugate vaccines on childhood pneumonia in sub-Saharan Africa", the authors showed evidence of the impact of pneumococcal conjugate vaccines (PCV10 and PCV13) in Kenya, Gambia, Rwanda, and South Africa is inconclusive based on clinical pneumonia incidence.

1. The authors may need to elaborate how the impact of the first introduction of PCV7 vaccine on pneumonia incidence in this systematic review since most of the study location had PCV7 introduced before PCV10/PCV13, except in Kenya (page 5).
2. The authors need to add the information on the immunization doses and schedule in each location and to discuss further how these differences had an impact on the outcomes of PCV10 and PCV13 vaccination.
3. The authors need to discuss further the health and economic status in each location that can affect the outcomes of PCV10 and PCV13 vaccination.
4. 4. Please add the new update of pneumococcal polysaccharide capsule (Introduction part): <https://mbio.asm.org/content/11/3/e00937-20>¹

References

1. Ganaie F, Saad J, McGee L, van Tonder A, et al.: A New Pneumococcal Capsule Type, 10D, is the 100th Serotype and Has a Largecps Fragment from an Oral Streptococcus. *mBio*. 2020; **11** (3). [Publisher Full Text](#)

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Partly

Are sufficient details of the methods and analysis provided to allow replication by others?

Yes

Is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

Are the conclusions drawn adequately supported by the results presented in the review?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Medical microbiology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 03 Nov 2020

chukwuemeka Onwuchekwa, Institute of Tropical Medicine, Antwerp, Antwerp, Belgium

Thank you for reviewing our work and for the constructive comments and suggestions. We have carefully considered your comments and address them below:

1. As stated in our original submission:

"it impossible to separate the effect of the 7-valent from the 10- and 13-valent vaccines since the 7-valent PCV might have influenced the pre-existing disease trend".

We have gone further to stress that separating the effect of the 7-valent from subsequent 13-valent vaccination is further compounded by the short time lapse between the two vaccines (2009 and 2011). We therefore added the statement:

"However, because of the short time lapse between the 7-valent and 10- or 13-valent vaccine roll-out in these countries, it is unlikely that a significant change in pneumonia trend would be demonstrable"

2. Thank you for drawing our attention to the influence of dosing schedule, this comment was very helpful. We have included the dosing schedule used in each country as described by the authors in Table 2 (Description of included studies). In the introduction section, we have also included a sentence describing the current vaccine schedule:

"The vaccines are usually administered as three doses in early infancy (3 + 0 schedule), or two doses in early infancy plus a booster in late infancy (2 + 1 schedule)"

While there is evidence that pneumococcal vaccine schedules which include a booster (e.g. 2 + 1 or 3 + 1 schedules) are associated with a more robust immune response, it has not been shown that these schedules have any effect of carriage or indeed pneumonia incidence(1). We have therefore not explored this further as we considered this to be outside the

boundaries of our review, and we have too few studies to make any type of comparison.

3. Thank you for this comment. We agree with your statement about the influence of Health and economic situation of the populations and how this could have influence childhood pneumonia and indeed uptake of vaccination. We also indicate here that the uptake of vaccination in all the reports were above 90% in the relevant population. Furthermore, the population based impact studies compared populations before and after vaccine roll-out, with a relatively short time lapse between the baseline and post-vaccination periods. It is our belief that socioeconomic variables are unlikely to have changed significantly at population-level, and is likely to have minimal influence on the observed findings. More importantly, most of the population based studies (time-series and before-after studies) included a control group or control condition, which would allow for assessment of the influence of historical bias (including changes in socioeconomic conditions)(2)

4. Thank you for drawing our attention to the most recent information of pneumococcal serotypes. We have noted this comment and made changes to the manuscript to reflect this, also included is the reference you suggested.

References

1. Whitney CG, Goldblatt D, O'Brien KL. Dosing schedules for pneumococcal conjugate vaccine: considerations for policy makers. *Pediatr Infect Dis J*. 2014 Jan;33 Suppl 2:S172-81.
2. Bernal JL, Cummins S, Gasparrini A. The use of controls in interrupted time series studies of public health interventions. *Int J Epidemiol*. 2018;47(6):2082-93.

Competing Interests: No competing interest to declare.

Reviewer Report 06 October 2020

<https://doi.org/10.5256/f1000research.27840.r67910>

© 2020 Mmbaga B. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Blandina Theophil Mmbaga

Kilimanjaro Christian Medical Center, Kilimanjaro Clinical Research Institute, Moshi, Tanzania

Thank you for the opportunity to review this manuscript. The author works on the important topic taking into account pneumonia still remains as one of the major causes of childhood morbidity and mortality. Assessing the impact of vaccines while differentiating clinical, radiological outcome support the evidence on the vaccine impact.

The manuscript is written in good English language.

The methodology is clear on how the review was conducted and how they reached final articles for

inclusion.

No major specific comments and would recommend the manuscript acceptance.

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Yes

Are sufficient details of the methods and analysis provided to allow replication by others?

Yes

Is the statistical analysis and its interpretation appropriate?

Yes

Are the conclusions drawn adequately supported by the results presented in the review?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Paediatric infectious disease

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 06 Oct 2020

chukwuemeka Onwuchekwa, Institute of Tropical Medicine, Antwerp, Antwerp, Belgium

Thank you for the favourable review.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research