Check for updates

REVIEW

Recent advances in understanding and managing acne

[version 1; peer review: 2 approved]Ichiro Kurokawa¹, Keisuke Nakase²¹Department of Dermatology, Acne Clinical Research Center, Meiwa Hospital, Nishinomiya, Hyogo, 663-8186, Japan²Department of Microbiology, Tokyo University of Pharmacy and Life Sciences, Hachioji, Tokyo, 192-0392, Japan**v1** First published: 29 Jul 2020, 9(Faculty Rev):792
<https://doi.org/10.12688/f1000research.25588.1>Latest published: 29 Jul 2020, 9(Faculty Rev):792
<https://doi.org/10.12688/f1000research.25588.1>**Abstract**

Multidisciplinary investigations into the pathogenesis of acne have significantly progressed over the past three years. Studies of the etiology of acne from the perspectives, for example, of sebaceous gland biology, sebum, genetics, keratinization, differentiation, hair cycles, immunology, bacteriology, and wound healing have elucidated its pathogenesis. This has led to the development of new therapies and paved the way for advanced studies that will enable the further evolution of acne treatment.

Keywords

Acne, biology, sebaceous gland, hair cycle, gene, immunology, wound healing, post inflammatory hyperpigmentation, vitamin C

Open Peer Review**Approval Status**

	1	2
version 1 29 Jul 2020

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Dae Hun Suh**, Seoul National University
College of Medicine, Seoul, South Korea
2. **Mauro Picardo**, San Gallicano Dermatologic
Institute, IRCCS, Rome, Italy

Any comments on the article can be found at the end of the article.

Corresponding author: Ichiro Kurokawa (kurokawa.i@meiwa-hospital.com)

Author roles: **Kurokawa I:** Conceptualization, Project Administration, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Nakase K:** Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Kurokawa I and Nakase K. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Kurokawa I and Nakase K. **Recent advances in understanding and managing acne [version 1; peer review: 2 approved]** F1000Research 2020, 9(Faculty Rev):792 <https://doi.org/10.12688/f1000research.25588.1>

First published: 29 Jul 2020, 9(Faculty Rev):792 <https://doi.org/10.12688/f1000research.25588.1>

Introduction

Acne vulgaris (acne) is an inflammatory disease of the pilosebaceous gland¹. It initially forms invisible micro (histopathological) comedones, then often appears in adolescents on the forehead as visible, clinically recognized blackheads or whiteheads (comedones) that develop into inflammatory red papules or pustules. Such lesions can become complicated with either atrophic or hypertrophic scars. Acne can develop into refractory cysts, nodules, and subcutaneous fistulas that are resistant to therapy. Acne most commonly appears on the face, neck, chest, and upper back, where sebaceous follicles predominate. In addition to acne vulgaris, related disorders include follicular occlusive diseases such as acne conglobata, perifolliculitis capitis abscedens et suffodiens (PCAS) (dissecting cellulitis or Hoffman disease), and hidradenitis suppurativa (acne inversa). These are all refractory and therapy-resistant diseases and commonly manifest as hypertrophic scars, cysts, and nodules.

The following factors are considered to be important to the classical etiology of acne vulgaris¹: increased sebum excretion rates, endocrinological factors such as androgens, abnormal keratinization of the follicular infundibulum, bacterial proliferation, and subsequent inflammation. Recent genetic and immunological studies have now elucidated the pathogenesis of acne (Figure 1).

Recent findings in the pathogenesis of acne

Histopathology of acne

Acne is a disease of sebaceous hair follicles that comprises multilobulated large sebaceous acini, tiny vellus hairs, and dilated follicular channels². Sebaceous follicles are located in the face, chest, and upper back. The sebaceous gland is atrophic in acne lesions (Figure 1), meaning that sebum has already been discharged into follicular channels because of undifferentiated sebocytes becoming mature and differentiated. Cunliffe proposed cyclical comedo growth and explained the natural resolution of comedones³.

Recent biology of the sebaceous gland

Stem cells in hair bulges can differentiate into sebocytes and outer root sheaths⁴. The progenitors of sebaceous glands in the junctional zone between the sebaceous duct and outer root sheath are leucine-rich repeat and immunoglobulin-like domain 1 (LRIG1)⁺ cells⁵ that can potentially differentiate into isthmus (epithelial) and sebaceous glands. These cells can also differentiate into sebocytes, sebaceous duct cells, and infundibular keratinocytes depending on the influence of diverse factors. We speculate that undifferentiated sebocytes in acne differentiate into sebaceous duct cells and infundibular keratinocytes instead of mature sebocytes, resulting in the abnormal keratinization of follicular channels. Saurat refers to this phenomenon as

Figure 1. Recent advances in acne pathogenesis. Atrophy of the sebaceous gland is induced by signal transduction due to various factors. In the hair cycle, anagen in normal skin changes to telogen or catagen in the “comedo switch”. Keratin (K) 75 and K79, which are present in the companion layer of normal hair follicles, are reduced in comedo. Immunological aspects show a T helper type 1 (Th1) and Th17 shift in acne. Diversified multiple genes are implicated in acne pathogenesis. According to the above-mentioned etiology, immunity induction therapy, molecular target therapy targeted on innate immunity, and the regulation of signal transduction could be alternative candidates for acne treatment in the future. AHR, aryl hydrocarbon receptor; Cx2, cyclooxygenase-2; EGF, epidermal growth factor; FGF, fibroblast growth factor; FOXO1A, forkhead box transcription factor class O1A; GH, Growth hormone; IFN, interferon; IL, interleukin; Lef, lymphoid enhancer-binding factor; MC5R, melanocortin 5 receptor; MMP, matrix metalloprotease; PPAR, peroxisome proliferation-activated receptor; Tcf, transcription factor; TLR, Toll-like receptor; TNF, tumor necrosis factor; Wnt, Wingless. This image was produced from the author's clinic for this review.

a “comedo switch”⁵. Saurat also stated that not all follicles are involved in acne of the face, neck, and trunk and that only 0.25% of sebaceous glands are involved in visible acne lesions⁵.

With respect to the molecular network signaling pathway, β -catenin, transcription factor (Tcf) 3, and lymphoid enhancer-binding factor (Lef)-1 are important⁴. High β -catenin levels stimulate hair follicle formation, whereas low levels stimulate the epidermis and sebaceous gland. Lef-1 cooperates with Indian hedgehog to control the proliferation and differentiation of sebaceous gland progenitors⁴. Peroxisome proliferation-activated receptor (PPAR), c-Myc, and Cx-2 differentiate towards sebocytes. The repression of β -catenin and activation of c-Myc and the hedgehog signaling cascade promote the differentiation and maturation of sebocytes⁶.

In addition, Wingless (Wnt) and hedgehog signals are important because the activation of Wnt signaling in LGR5⁺ bulge cells promotes hair growth. Loss of Wnt signaling in bulge cells expressing keratin (K) 15 promotes migration and differentiation into sebocytes. Wnt signaling regulates the proliferation and specification of junctional LRIG1⁺ cells, resulting in acne pathogenesis⁷. Androgens might promote sebocyte differentiation and inhibit Wnt signaling⁷. Aryl hydrocarbon receptors (AHRs) are linked to chloracne⁷. Exposure to AHR causes comedo formation, sebaceous gland atrophy, and the upregulation of AHR expression in humans and mice⁷. AHR inhibit sebaceous differentiation by promoting the differentiation of junctional zone stem cells into infundibular keratinocytes⁷. Fibroblast growth factor (FGF) 2 stimulates proliferation through the pilosebaceous unit⁷, and epidermal growth factor receptors (EGFRs) can enlarge the hyperproliferation of sebaceous glands and increase sebum production⁷.

Sebum

The formation of acne is attributable to a decrease in linoleic acid in classical etiology¹. Squalene peroxidized by lipoperoxidase and decreased vitamin E can induce inflammation. Lipoperoxidase can induce pro-inflammatory cytokines and keratinocyte proliferation and activate PPAR⁸. Monosaturated fatty acids can induce changes in keratinocyte proliferation and differentiation. AHR is metabolized to tetrachlorodibenzo-dioxin, which induces sebocytes to undergo epithelial-type differentiation. Insulin-like growth factor (IGF) 1 stimulates the formation of unsaturated lipids and neosynthesis of lipids⁹. The IGF axis is involved in acne pathogenesis¹⁰.

Recently, 11 β -hydroxysteroid dehydrogenase type I (11 β -HSD1) has been observed to promote lipid synthesis. Tumor necrosis factor (TNF)- α promotes lipogenesis in human sebocytes¹¹. Transforming growth factor (TGF)- β maintains sebocytes in an undifferentiated state and decreases lipid accumulation¹².

Genetic findings

Recent genetic findings have clarified that genes encoding the enzymes 21-hydroxylase (*CYP21A2*), steroid 5 α -reductase type I (*SRD5A1*), and androgen receptor (*AR*), the somatotrophic axis (*GHI*, *GHR*, *IGF1*, *IGFBP3*, and *IGF1R*), and the forkhead box transcription factor class O1A (*FOXO1A*), peroxisome

proliferator-activated receptor (*PPARA*, *PPARB*, *PPARG*, *PPARD*), FGF-2 (*FGF2*), melanocortin receptor (*MC5R*, *MC1R*), matrix metalloprotease (*MMP1*, *MMP2*, *MMP3*, *MMP9*, *MMP13*), TNF- α (*TNF*), IL-1 α (*IL1A*), and Toll-like receptors (*TLR2* and *TLR4*) are implicated in the pathogenesis of acne^{13,14}.

Keratinization and hair cycles in acne

Abnormal keratinization is an important factor in acne pathogenesis¹. Expression of the hyperproliferative keratins (K6, K16, and K17) is increased in acne lesions¹⁵. Significant filaggrin expression in the infundibulum is closely associated with the abnormal keratinization involved in acne¹⁶. IL-1 α is involved in abnormal keratinization¹⁷, and inflammation precedes keratinization¹.

Hair cycles in acne have not been studied in detail. van Scott *et al.* stated that the hair cycles in acne are almost always either telogens or catagens¹⁸. However, keratin expression in the hair cycle in acne lesions has not been studied. A low micro-comedo index in acne is associated with significantly higher K75 expression¹⁹. K75 is expressed in the companion layer between the inner (Henle) layer and outer root sheaths in the lower portion of normal hair follicles²⁰. A companion layer is found in anagen but not in either telogen or catagen hair follicles. The prevalence of K75 depends on the hair cycle. Therefore, the hair cycle in acne with microcomedones is directed towards the follicular infundibulum and sebaceous duct instead of the lower portion of hair follicles that comprise companion layers. In addition, K79 is downregulated in comedonal acne lesions²¹ and is expressed in companion layers in normal anagen hair follicles¹⁹. Considering homeostasis in hair cycles in acne, progenitor cells in the junctional zone might differentiate not towards the outer root sheath of hair follicles below the sebaceous duct but towards the infundibulum and sebaceous duct cells in acne.

Immunological aspects including cytokines

The immunological aspects of acne have become noteworthy. *Malassezia* and *Demodex* are related to the pathogenicity of folliculitis and rosacea, respectively. *Cutibacterium acnes* (*C. acnes*) and normal flora are involved in acne pathogenesis due to overgrowth in closed follicles. In acne, *C. acnes* in the follicular channel stimulates Langerhans cells in the infundibulum via TLR-2, resulting in the production of IL-12 and IL-8 by activating Th1 cells. *C. acnes* also stimulates follicular keratinocytes in the infundibulum via TLR-2, resulting in the production of IL-6 and IL-8^{1,22} followed by the formation of inflammatory lesions such as red papules and pustules. Jeremy *et al.* proposed that initial inflammation caused by CD4, CD3, and macrophages induces comedones²³: inflammation precedes keratinization²³. A T helper type 1 (Th1) shift occurs in acne lesions, and Th1-positive cells are more prevalent in acne lesions than in normal skin²⁴. From the aspect of host responses to *C. acnes* in acne pathogenesis, host immunological factors against *C. acnes* produced by PBMCs can be attributed more to bacteriological factors²⁵.

In addition to Th1 cytokines, Th17 is also involved in acne pathogenesis. *C. acnes* is a potent inducer of Th17 and Th1,

and significant numbers of cells express IL-17 in acne lesions²⁶. IL-17 is reduced by vitamins A and D. IL-1 β and TNF- α are involved in acne inflammation¹¹. Corticotropin-releasing hormone (CRH) can increase IL-6 and IL-8 levels *in vitro*¹². Therefore, regulating these cytokines might offer an alternative strategy for treating acne.

Immunity induction therapy

Acne has been treated with benzoyl peroxide (BPO) formulations, adapalene, antimicrobials, anti-androgen agents, and isotretinoin, which control abnormal follicular keratinization in the infundibulum, have bactericidal and bacteriostatic effects, inhibit inflammation, and decrease sebum excretion, according to the pathomechanism of classical etiological factors. From these immunological aspects in acne, Th1, Th17, and TNF- α are upregulated in acne.

Antimicrobials are useful in immunoregulation because they show not only antibacterial but also anti-inflammatory activities. However, antimicrobial use might increase bacterial resistance to antimicrobials and cause dysbiosis as well as side effects. Novel medicines without these adverse effects have been developed. Vaccination with antibodies against the Christie-Atkins-Munch-Petersen (CAMP) factor, which is associated with *C. acnes* cytotoxicity, decreases the growth of *C. acnes* and the production of murine MIP-2²⁷. The ability of a vaccine produced by *Staphylococcus capitis* E12 to prevent *C. acnes* overgrowth has been investigated²⁸. A vaccine produced by *S. capitis* is expected for preventing *C. acnes* overgrowth and killing overgrown *C. acnes*. To suppress the inflammation completely, monoclonal antibodies blocking cytokines are needed because killed *C. acnes* induces inflammatory cytokines²⁹.

Th1 and Th17 shift inhibitors and antibodies against IL-17 and TNF- α might offer alternative approaches to treating acne. Notably, a TNF- α antibody is presently used to treat acne conglobata³⁰, hidradenitis suppurativa³¹, and PCAS³². Antibodies to IL-17, IL-23, and IL-1 α will be used to treat acne conglobata, PCAS, and hidradenitis suppurativa, which will also be treated with apremilast³³. Nitrate oxide can be an alternative treatment for acne in humans by reducing IL-1 β , IL-8, TNF- α , and IL-6 induced by monocytes and IL-8 and IL-6 induced by keratinocytes via innate immunity^{34,35}.

Wound healing

The most important complication in acne is scar formation. The rupture and breakdown of inflammatory red papules, pustules, and deep-seated subcutaneous abscesses in the deep dermis can lead to erosion and ulceration, resulting in scar formation. If ulceration is superficial, re-epithelialization proceeds without scarring. However, ulcers located deep below the reticular dermis form scars like deep dermal burns. Atrophic scars have been classified as icepick, boxcar, and rolling scars³⁶, whereas elevated hypertrophic scars like acne conglobata sometimes form. Wound healing in acne should be taken into account when considering scar formation³⁷. Scars form because of persistent inflammation and are associated with the depth of inflammation³⁸. Scar formation is also associated with MMP,

IL-6, TGF- β , macrophages, and B cells. Atrophic scars are dependent on B cells and macrophages³⁹. Sebaceous duct cells can differentiate into epidermal keratinocytes and sebocytes in wounds. Thus, sebaceous ducts are bimodal, which is consistent with keratin expression between the infundibulum and sebocytes¹⁶.

Recent non-surgical therapy for postinflammatory hyperpigmentation

Postinflammatory hyperpigmentation (PIH) is an important complication of acne vulgaris that occurs via damage to the basal cell layer. Non-surgical chemical glycolic acid (GA) peels and subsequent iontophoresis using vitamin C, vitamin A, and vitamin E are used to treat PIH, postinflammatory erythema (PIE), and atrophic scars³⁷. GA loosens cellular adhesion, promotes loss from the cornified layer, as well as the regeneration of epidermal and dermal tissues, removes follicular cast in the infundibulum, and de-roofs pustules and red papules³⁷. Vitamin C promotes re-epithelialization while inhibiting melanogenesis and reactive oxygen³⁷. Vitamin C can induce self-renewal of the mesenchymal cell cycle program and fibroblast motility, promote fibroblast migration, confer anti-inflammatory effects, and induce macrophage inflammation⁴⁰. Both PIH and PIE are treated using the vitamin C derivative, amphipathic vitamin C^{41,42}. The mechanisms through which vitamin C improves atrophic scars are thought to be via self-renewal cell cycle progression, promoted fibroblast migration, matrix deposition and neo-vascularization, anti-inflammatory effects in macrophages, and attenuation mediators in wounds via IL-1 β and TNF- α ⁴⁰. In addition, basic FGF (bFGF), an important factor in wound healing, can promote epithelialization and thus improve atrophic scars, resulting in flattening of the epidermis⁴³. bFGF supplies epidermal defects with proliferating keratinocytes.

Alternative therapy for refractory cysts and nodules in Japan

The pathogenesis of refractory nodulocystic lesions remains unclear. Significant amounts of filaggrin are expressed in cyst formation with retention hyperkeratosis⁴⁴. Nodulocystic acne and acne conglobata are treated with isotretinoin⁴⁵, but not in Japan. Kampo (traditional Japanese herbal medicine) such as Saireito can be very effective sometimes for treating nodulocystic acne⁴⁶ and PCAS⁴⁷. It works on cystic and alopecia lesions in PCAS, resulting in hair growth⁴⁷. Hair cycles in PCAS are telogens or catagens as in acne vulgaris. Saireito exerts multiple effects on endogenous corticosteroids, inflammation, reactive oxygen species, coagulation, macrophages, neutrophils, and endothelial cells⁴⁶. However, the mechanism of Saireito in acne pathogenesis awaits investigation in a basic research study.

Comprehensive multiplexed therapy in Japan

Isotretinoin and anti-androgen therapy have not been applied in the treatment of acne in Japan. Therefore, severe acne is treated with oral antimicrobials combined with Kampo, topical BPO, and local steroid injections⁴⁸. Although the mechanism of Kampo is not clear, Saireito can suppress Th1 shift and suppress B cell function, regulating Th1/Th2 balance in mice⁴⁹. Jyumihaidokuto is available for acne vulgaris in Japan⁵⁰. It

contains Bokusoku, which inhibits rat 5 α -reductase activity and testosterone-induced sebum synthesis in hamster sebocytes⁵¹. Keigairengyoto is also available for acne vulgaris in Japan⁵⁰. It has anti-bacterial effects against *C. acnes*⁵² and inhibits reactive oxygen⁵³. Comprehensive multiplexed therapy is recommended for therapy-resistant refractory acne.

Future perspectives for acne treatment

This overview describes recent advances in acne pathogenesis. Understanding the multiple diverse, interdependent, and complicated etiological factors and the biology of sebaceous glands is extremely important. Progenitor cells in the junctional zone can differentiate into infundibulum, sebocytes, and outer root sheaths depending on the influence of genes and molecular signals. The altered hair cycle in acne remains of interest from the viewpoint of the biological homeostasis of pilosebaceous units. Immunological studies of acne have generated innovative immunity induction therapies such as antibodies to

TNF- α and various cytokines. The most refractory complication of acne is scar formation, which involves the production of pro-inflammatory cytokines such as IL-1 α , IL-1 β , IL-6, TNF- α , and TGF- β ⁵⁴.

According to the etiopathogenesis of acne, potential treatments for acne in the future are topical anti-androgens, melanocortin receptor antagonists, IGF-1 inhibitors, PPAR modulators, acetylcholine inhibitors, topical retinoic and metabolism-blocking agents, monoclonal antibodies, antimicrobial peptides, antioxidants, phosphodiesterase inhibitors, IL-1 β inhibitors, vitamin D analogues, dapson, systemic antiandrogens, and immunotherapy⁵⁵.

Immunity induction and wound healing therapies will become key strategies for preventing acne scars. Acne is a disease of the pilosebaceous unit. Based on acne pathogenesis, regulation of sebocyte differentiation is a novel therapeutic procedure^{56,57}.

References

- Kurokawa I, Danby FW, Ju Q, et al.: **New developments in our understanding of acne pathogenesis and treatment.** *Exp Dermatol.* 2009; **18**(10): 821–32.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Plewig G, Kligman AM: **Acne & Rosacea.** Springer-Verlag, 2005.
- Cunliffe WJ, Holland DB, Clark SM, et al.: **Comedogenesis: Some new aetiological, clinical and therapeutic strategies.** *Br J Dermatol.* 2000; **142**(6): 1084–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zouboulis CC, Picardo M, Ju Q, et al.: **Beyond acne: Current aspects of sebaceous gland biology and function.** *Rev Endocr Metab Disord.* 2016; **17**(3): 319–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Saurat JH: **Strategic Targets in Acne: The Comedone Switch in Question.** *Dermatology.* 2015; **231**(2): 105–11.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Niemann C: **Differentiation of the sebaceous gland.** *Dermatoendocrinol.* 2009; **1**(2): 64–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Clayton RW, Göbel K, Niessen CM, et al.: **Homeostasis of the sebaceous gland and mechanisms of acne pathogenesis.** *Br J Dermatol.* 2019; **181**(4): 677–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
- Zouboulis CC, Dessinioti C: **Acne and inflammation.** Zouboulis et al. (eds) *Pathogenesis and treatment of Acne and Rosacea*, Springer-Verlag, Berlin, Heidelberg, 2014.
[Publisher Full Text](#)
- Mastrofrancesco A, Ottaviani M, Cardinali G, et al.: **Pharmacological PPAR γ modulation regulates sebogenesis and inflammation in SZ95 human sebocytes.** *Biochem Pharmacol.* 2017; **138**: 96–106.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Osher E, Macaulay VM: **Therapeutic Targeting of the IGF Axis.** *Cells.* 2019; **8**(8): 895.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Choi JJ, Park MY, Lee HJ, et al.: **TNF- α increases lipogenesis via JNK and PI3K/Akt pathways in SZ95 human sebocytes.** *J Dermatol Sci.* 2012; **65**(3): 179–88.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Suh DH, Kwon HH: **What's new in the physiopathology of acne?** *Br J Dermatol.* 2015; **172** Suppl 1: 13–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Melnik BC: **Acne and Genetics.** Zouboulis et al. (eds) *Pathogenesis and treatment of Acne and Rosacea*, Springer-Verlag, Berlin, Heidelberg, 2014.
[Publisher Full Text](#)
- Common JEA, Barker JN, van Steensel MAM: **What does acne genetics teach us about disease pathogenesis?** *Br J Dermatol.* 2019; **181**(4): 665–76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
- Hughes BR, Morris C, Cunliffe WJ, et al.: **Keratin expression in pilosebaceous epithelia in truncal skin of acne patients.** *Br J Dermatol.* 1996; **134**(2): 247–56.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kurokawa I, Mayer-da-Silva A, Gollnick H, et al.: **Monoclonal antibody labeling for cytokeratins and filaggrin in the human pilosebaceous unit of normal, seborrheic and acne skin.** *J Invest Dermatol.* 1988; **91**(6): 566–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Guy R, Green MR, Kealey T: **Modeling Acne in Vitro.** *J Invest Dermatol.* 1996; **106**(1): 176–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
- van Scott EJ, MacCardle RC: **Keratinization of the duct of the sebaceous gland and growth cycle of the hair follicle in the histogenesis of acne in human skin.** *J Invest Dermatol.* 1956; **27**(6): 405–29.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fontao F, von Engelbrechten M, Seilaz C, et al.: **Microcomedones in non-lesional acne prone skin New orientations on comedogenesis and its prevention.** *J Eur Acad Dermatol Venerol.* 2019; **34**(2): 357–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
- Kurokawa I, Takahashi K, Moll I, et al.: **Expression of keratins in cutaneous epithelial tumors and related disorders—distribution and clinical significance.** *Exp Dermatol.* 2011; **20**(3): 217–28.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Veniaminova NA, Vagnozzi AN, Kopinke D, et al.: **Keratin 79 identifies a novel population of migratory epithelial cells that initiates hair canal morphogenesis and regeneration.** *Development.* 2013; **140**(24): 4870–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Kim J: **Review of the innate immune response in acne vulgaris: Activation of Toll-like receptor 2 in acne triggers inflammatory cytokine responses.** *Dermatology.* 2005; **211**(3): 193–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jeremy AHT, Holland DB, Roberts SG, et al.: **Inflammatory events are involved in acne lesion initiation.** *J Invest Dermatol.* 2003; **121**(1): 20–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mouser PE, Seaton ED, Chu AC, et al.: **Propionibacterium acnes-Reactive T Helper-1 Cells in the Skin of Patients with Acne Vulgaris.** *J Invest Dermatol.* 2003; **121**(5): 1226–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Sugisaki H, Yamanaka K, Kakeda M, et al.: **Increased interferon-gamma, interleukin-12p40 and IL-8 production in Propionibacterium acnes-treated peripheral blood mononuclear cells from patient with acne vulgaris: Host response but not bacterial species is the determinant factor of the disease.** *J Dermatol Sci.* 2009; **55**(1): 47–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
- Agak GW, Kao S, Ouyang K, et al.: **Phenotype and Antimicrobial Activity**

- of Th17 Cells Induced by *Propionibacterium acnes* Strains Associated with Healthy and Acne Skin. *J Invest Dermatol*. 2018; 138(2): 316–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
27. Wang Y, Hata TR, Tong YL, *et al.*: The Anti-Inflammatory Activities of *Propionibacterium acnes* CAMP Factor-Targeted Acne Vaccines. *J Invest Dermatol*. 2018; 138(11): 2355–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 28. O'Neill AM, Nakatsuji T, Hayachi A, *et al.*: Identification of a Human Skin Commensal Bacterium that Selectively Kills *Cutibacterium acnes*. *J Invest Dermatol*. 2020.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 29. Lyte P, Sur R, Nigam A, *et al.*: Heat-killed *Propionibacterium acnes* is capable of inducing inflammatory responses in skin. *Exp Dermatol*. 2009; 18(12): 1070–2.
[PubMed Abstract](#) | [Publisher Full Text](#)
 30. Yiu ZZN, Madan V, Griffiths CEM: Acne conglobata and adalimumab: Use of tumour necrosis factor- α antagonists in treatment-resistant acne conglobata, and review of the literature. *Clin Exp Dermatol*. 2015; 40(4): 383–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 31. Zouboulis CC, Desai N, Emtestam L, *et al.*: European S1 guideline for the treatment of hidradenitis suppurativa/acne inversa. *J Eur Acad Dermatol Venereol*. 2015; 29(4): 619–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
 32. Takahashi T, Yamasaki K, Terui H, *et al.*: Perifolliculitis capitis abscedens et suffodiens treatment with tumor necrosis factor inhibitors: A case report and review of published cases. *J Dermatol*. 2019; 46(9): 802–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 33. Vossen ARJ, van Doorn MBA, van der Zee HH, *et al.*: Apremilast for moderate hidradenitis suppurativa: Results of a randomized controlled trial. *J Am Acad Dermatol*. 2019; 80(1): 80–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 34. Qin M, Landriscina A, Rosen JM, *et al.*: Nitric Oxide-Releasing Nanoparticles Prevent *Propionibacterium acnes*-Induced Inflammation by Both Clearing the Organism and Inhibiting Microbial Stimulation of the Innate Immune Response. *J Invest Dermatol*. 2015; 135(11): 2723–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 35. Dréno B: What is new in the pathophysiology of acne, an overview. *J Eur Acad Dermatol Venereol*. 2017; 31 Suppl 5: 8–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 36. Jacob CI, Dover JS, Kaminer MS: Acne scarring: A classification system and review of treatment options. *J Am Acad Dermatol*. 2001; 45(1): 109–17.
[PubMed Abstract](#) | [Publisher Full Text](#)
 37. Kurokawa I, Oiso N, Kawada A: Adjuvant alternative treatment with chemical peeling and subsequent iontophoresis for postinflammatory hyperpigmentation, erosion with inflamed red papules and non-inflamed atrophic scars in acne vulgaris. *J Dermatol*. 2017; 44(4): 401–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
 38. Holland DB, Jeremy AHT, Roberts SG, *et al.*: Inflammation in acne scarring: A comparison of the responses in lesions from patients prone and not prone to scar. *Br J Dermatol*. 2004; 150(1): 72–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
 39. Carlván I, Bertino B, Rivier M, *et al.*: Atrophic scar formation in patients with acne involves long-acting immune responses with plasma cells and alteration of sebaceous glands. *Br J Dermatol*. 2018; 179(4): 906–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 40. Mohammed BM, Fisher BJ, Kraskauskas D, *et al.*: Vitamin C promotes wound healing through novel pleiotropic mechanisms. *Int Wound J*. 2016; 13(4): 572–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. Kurokawa I, Yoshioka M, Ito S: Split-face comparative clinical trial using glyceryl-octyl-ascorbic acid/ascorbyl 2-phosphate 6-palmitate/DL- α -tocopherol phosphate complex treatment for postinflammatory hyperpigmentation, postinflammatory erythema and atrophic scar in acne vulgaris. *J Dermatol*. 2019; 46(10): e347–e348.
[PubMed Abstract](#) | [Publisher Full Text](#)
 42. Kurokawa I, Watanabe M, Hayashibe K: Effect of a Cosmetic Use with 2% Isostearyl-L-Ascorbic Acid Gel for Postinflammatory Hyperpigmentation and Postinflammatory Erythema in Acne Vulgaris. *JCDISA*. 2020; 10: 28–32.
[Publisher Full Text](#)
 43. Kurokawa I: Non-surgical treatment with basic fibroblast growth factor for atrophic scars in acne vulgaris. *J Dermatol*. 2018; 45(9): e238–e239.
[PubMed Abstract](#) | [Publisher Full Text](#)
 44. Kurokawa I, Umeda K, Nishimura K, *et al.*: Filaggrin expression and the pathogenesis of epidermal cysts. *Br J Dermatol*. 2007; 157(2): 415–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 45. Peck GL, Olsen TG, Yoder FW, *et al.*: Prolonged remissions of cystic and conglobate acne with 13-cis-retinoic acid. *N Engl J Med*. 1979; 300(7): 329–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
 46. Kurokawa I: Successful adjuvant alternative treatment with Saireito (Japanese herbal medicine) for nodulocystic acne. *J Nutr Disorders Ther*. 2017; 7(3): 215.
[Publisher Full Text](#)
 47. Kurokawa I: Perifolliculitis capitis abscedens et suffodiens successfully treated with Saireito. *J Dermatol*. 2019; 46(8): e299–e301.
[PubMed Abstract](#) | [Publisher Full Text](#)
 48. Kurokawa I: Comprehensive Multiplexed Therapy for Severe Nodulocystic Acne in Puberty. Comprehensive multiplex therapy for cystic acne. *Health Care: Current Reviews*. 2016; 4: 1.
[Publisher Full Text](#)
 49. Ito T, Seo N, Yagi H, *et al.*: Unique therapeutic effects of the Japanese-Chinese herbal medicine, Sairei-to, on Th1/Th2 cytokines balance of the autoimmunity of MRL/lpr mice. *J Dermatol Sci*. 2002; 28(3): 198–210.
[PubMed Abstract](#) | [Publisher Full Text](#)
 50. Hayashi N, Akamatsu H, Iwatsuki K, *et al.*: Japanese Dermatological Association Guidelines: Guidelines for the treatment of acne vulgaris 2017. *J Dermatol*. 2018; 45(8): 898–935.
[PubMed Abstract](#) | [Publisher Full Text](#)
 51. Koseki J, Matsumoto T, Matsubara Y, *et al.*: Inhibition of Rat 5 α -Reductase Activity and Testosterone-Induced Sebum Synthesis in Hamster Sebocytes by an Extract of *Quercus acutissima* Cortex. *Evid Based Complement Alternat Med*. 2015; 2015: 853846.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 52. Higaki S, Hasegawa Y, Morohashi M, *et al.*: The correlation of Kampo formulations and their ingredients on anti-bacterial activities against *Propionibacterium acnes*. *J Dermatol*. 1995; 22(1): 4–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 53. Akamatsu H, Asada Y, Horio T: Effect of keigai-rengyo-to, a Japanese kampo medicine, on neutrophil functions: A possible mechanism of action of keigai-rengyo-to in acne. *J Int Med Res*. 1997; 25(5): 255–65.
[PubMed Abstract](#) | [Publisher Full Text](#)
 54. Ogawa R: Keloid and Hypertrophic Scars Are the Result of Chronic Inflammation in the Reticular Dermis. *Int J Mol Sci*. 2017; 18(3): 606.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 55. Bhat YJ, Latief I, Hassan I: Update on etiopathogenesis and treatment of Acne. *Indian J Dermatol Venereol Leprol*. 2017; 83(3): 298–306.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 56. Ottaviani M, Flori E, Mastrofrancesco A, *et al.*: Sebocyte Differentiation as a New Target for Acne Therapy: An *in Vivo* Experience. *J Eur Acad Dermatol Venereol*. 2020.
[PubMed Abstract](#) | [Publisher Full Text](#)
 57. Harder J, Tsuruta D, Murakami M, *et al.*: What is the role of antimicrobial peptides (AMP) in acne vulgaris? *Exp Dermatol*. 2013; 22(6): 386–91.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Mauro Picardo**
Laboratory of Cutaneous Physiopathology and Integrated Center of Metabolomics Research, San Gallicano Dermatologic Institute, IRCCS, Rome, Italy
Competing Interests: No competing interests were disclosed.
2. **Dae Hun Suh**
Department of Dermatology, Seoul National University College of Medicine, Seoul, South Korea
Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research