Check for updates

REVIEW

The third model of Bax/Bak activation: a Bcl-2 family feud finally resolved? [version 1; peer review: 3 approved]

Xu Luo^{1,2}, Katelyn L. O'Neill¹, Kai Huang^{1,2}

¹Eppley Institute for Research in Cancer and Allied Diseases, Fred & Pamela Buffett Cancer Center, University of Alaska Medical Center, Omaha, NE, 68198-7696, USA

²Department of Pathology and Microbiology, University of Nebraska Medical Center, Omaha, NE, 68198-6805, USA

V1 First published: 06 Aug 2020, 9(Faculty Rev):935
<https://doi.org/10.12688/f1000research.25607.1>

Latest published: 06 Aug 2020, 9(Faculty Rev):935
<https://doi.org/10.12688/f1000research.25607.1>

Abstract

Bax and Bak, two functionally similar, pro-apoptotic proteins of the Bcl-2 family, are known as the gateway to apoptosis because of their requisite roles as effectors of mitochondrial outer membrane permeabilization (MOMP), a major step during mitochondria-dependent apoptosis. The mechanism of how cells turn Bax/Bak from inert molecules into fully active and lethal effectors had long been the focal point of a major debate centered around two competing, but not mutually exclusive, models: direct activation and indirect activation. After intensive research efforts for over two decades, it is now widely accepted that to initiate apoptosis, some of the BH3-only proteins, a subclass of the Bcl-2 family, directly engage Bax/Bak to trigger their conformational transformation and activation. However, a series of recent discoveries, using previously unavailable CRISPR-engineered cell systems, challenge the basic premise that undergirds the consensus and provide evidence for a novel and surprisingly simple model of Bax/Bak activation: the membrane (lipids)-mediated spontaneous model. This review will discuss the evidence, rationale, significance, and implications of this new model.

Keywords

Apoptosis, Bcl-2 family, Bax, Bak, BH3-only proteins, Bid, Bim, Bad, Bcl-xL, Mcl-1, Activator, Sensitizer, Direct activation, Indirect activation, Membrane-mediated Spontaneous activation, Mitochondrial outer membrane, auto-activation, de novo activation, retro-translocation, BH3 mimetics

Open Peer Review

Approval Status

	1	2	3
version 1 06 Aug 2020			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Andreas Strasser**, The Walter and Eliza Hall Institute of Medical Research, Parkville, Melbourne, Australia
2. **Andrew Gilmore**, University of Manchester, Manchester, UK
3. **Vishva Dixit**, Genentech, South San Francisco, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Xu Luo (xuluo@unmc.edu)

Author roles: Luo X: Conceptualization, Project Administration, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; O'Neill KL: Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; Huang K: Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by a pilot grant from the Nebraska Center for Cellular Signaling (P30GM106397) and National Institutes of Health (NIH) Grants R03CA205496 and R01GM118437 to Xu Luo.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2020 Luo X *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Luo X, O'Neill KL and Huang K. **The third model of Bax/Bak activation: a Bcl-2 family feud finally resolved?** [version 1; peer review: 3 approved] F1000Research 2020, 9(Faculty Rev):935 <https://doi.org/10.12688/f1000research.25607.1>

First published: 06 Aug 2020, 9(Faculty Rev):935 <https://doi.org/10.12688/f1000research.25607.1>

Introduction

Apoptosis is a highly regulated form of cell death playing important roles in embryonic development, tissue homeostasis, and disease progression¹. In response to various death stimuli, cells often initiate a mitochondria-dependent apoptotic pathway that involves mitochondrial outer membrane permeabilization (MOMP), formation of the apoptosome, and the activation of effector caspases². Generally considered the “point of no return”, MOMP irreversibly commits cells to apoptosis through the actions of Bax and Bak, two functionally similar effector proteins of the Bcl-2 family³.

Since the identification of Bcl-2 in human follicular lymphoma^{4,5} and its discovery as the first anti-apoptotic oncogene⁶ over three decades ago, 17–18 proteins that share at least one of the four Bcl-2 homology domains (BH1–4) have been identified in the mammalian system as Bcl-2 family proteins⁷, with the last member, Bmf, identified in 2001⁸. The Bcl-2 family proteins are classified into anti-apoptotic (Bcl-2, Bcl-xL, Bcl-w, Mcl-1, and A1), pro-apoptotic effector (Bax, Bak, and Bok), and pro-apoptotic BH3-only proteins (Bad, Bid, Bik, Bim, Bmf, Hrk, Noxa, and Puma)⁹ (Figure 1). Three types of interactions—1) BH3-only–anti-apoptotic, 2) anti-apoptotic–Bax/Bak, and 3) BH3-only–Bax/Bak—appear to govern the functional relationships among the Bcl-2 family members during apoptosis^{10,11}.

As the requisite effectors of MOMP, both Bax and Bak are kept in check under normal conditions. Bax is primarily in the cytosol as a globular-shaped monomeric protein consisting of nine α -helices, with helix 9 fitting into a hydrophobic groove^{12–14}. Bak has a similar structure¹⁵ but is localized to the mitochondrial outer membrane (MOM) through its helix 9. Both Bax and Bak are activated during apoptosis¹⁶. The active forms of Bax/Bak are known to homo-oligomerize in the MOM, forming proteinaceous/lipidic pores that are responsible for the release of cytochrome c and other apoptogenic factors^{17,18}. The BH3-only and the anti-apoptotic Bcl-2 proteins regulate MOMP by promoting or suppressing the activation of Bax/Bak, respectively^{9,11,19}.

Bok has recently been shown to complement Bax/Bak in developmental cell death²⁰ and to function as a non-canonical effector of MOMP in response to proteasome inhibition²¹. However, since the loss of both Bax and Bak is sufficient to block apoptosis induced by most apoptotic stimuli²², the role of Bok in apoptosis remains unclear.

As both Bax and Bak undergo a dramatic transformation from benign proteins to lethal effectors during apoptosis, the mechanism of their activation has been the subject of intense debate and is considered to be the “holy grail” of apoptosis research^{16,23}. Two competing but not mutually exclusive models, direct and indirect activation, had previously been proposed.

Direct activation

The first report of the direct interaction between a BH3-only protein and Bax/Bak was the identification of Bid as a BH3-containing death ligand that binds Bax. This finding led to the hypothesis that BH3-only proteins bind and activate Bax/Bak

to initiate apoptosis²⁴. Subsequently, it was found that some BH3 domains, i.e. BidBH3 or BimBH3, have the ability to directly activate Bax/Bak and cause the release of cytochrome c from purified mitochondria. On the other hand, BH3 peptide from Bad (BadBH3) did not directly activate Bax/Bak but was able to bind to Bcl-2, displace BidBH3, and sensitize mitochondria to BidBH3-induced cytochrome c release²⁵. These findings formed the basis for the “direct activation” model, in which the BH3-only proteins are classified into two groups: the activators, which can directly activate Bax/Bak but can also be sequestered by the anti-apoptotic Bcl-2 proteins, and the sensitizers, which cannot directly activate Bax/Bak but are able to bind to the anti-apoptotic Bcl-2 proteins and sensitize cells to apoptosis. To induce apoptosis, the sensitizers bind the anti-apoptotic Bcl-2 proteins and displace the activators, which in turn directly bind to Bax/Bak, allosterically triggering their activation^{25–28} (Figure 2A).

Although the initial classification listed Bid and Bim as direct activators because of the ability of the BH3 domain or the full-length protein to release dyes of large molecular weight from liposomes in the presence of inactive Bax^{25,28–32}, the list of direct activators has grown over the years to include Puma, Noxa, Bik, Bmf, and Hrk, although a range of potency has been reported^{26,33–39}. On the other hand, Bad has been a classic sensitizer, as the BadBH3 peptide has been used as a negative control for direct activators in liposome assays^{25,28}.

The interaction between a direct activator and Bax/Bak has been difficult to detect. This led to the hypothesis that activators only transiently engage Bax/Bak and activate them through a “hit and run” mechanism. The binding interface between an activator and Bax/Bak, while of great interest, has been controversial. NMR analysis provided evidence that the BH3 peptide from Bim transiently binds to a “trigger site” between helix 1 and helix 6^{40,41}. However, other studies using Bax/Bak mutants that lack helix 9 found that a BidBH3 peptide binds the hydrophobic groove^{42–45}. The discrepancy was largely resolved by new studies suggesting that the initial binding is at the trigger site and that the ensuing conformational changes allow a more stable interaction between the BH3 peptide and the hydrophobic groove^{46,47}. However, the biological significance of binding at the trigger site has recently been challenged^{48,49}.

Over the years, the direct activation model has evolved into several variants. The embedded together model emphasizes the importance of the membrane as a platform for all of the interactions among the Bcl-2 family proteins during Bax/Bak activation^{50,51}, the unified model emphasizes that the anti-apoptotic Bcl-2 proteins not only sequester the activator BH3-only proteins but also suppress or sequester the activated Bax/Bak³⁰, and the interconnected model is highly analogous to the unified model but also contains a feed-forward mechanism of auto-activation³⁶. One unifying feature of these models is the direct attack and engagement of Bax/Bak by the activator BH3-only proteins as the initial triggering event leading to Bax/Bak activation.

Since the beginning of the last decade, direct activation has been widely accepted as the mechanism of Bax/Bak activation

Figure 1. Schematic of Bcl-2 family proteins. Bcl-2 family proteins (human) are classified into three groups: the pro-apoptotic BH3-only proteins, the pro-apoptotic effector proteins Bax, Bak, and Bok, and the anti-apoptotic Bcl-2 proteins. Each family member contains at least one of the four Bcl-2 homology (BH) domains, BH1–4. Bax has nine α -helices, with helix 9 serving as a mitochondrial-targeting sequence¹³. Helices 2–5 constitute the homo-oligomerization domain of Bax^{18,42,52}. a.a., amino acids; TM, transmembrane.

during apoptosis^{10,53,54}. However, as much of the supporting evidence has come from *in vitro* studies^{25,28,31,32,41–43}, the physiological significance of this model has not been fully addressed.

Indirect activation

Indirect activation, or the “displacement” model, was first proposed in two important studies. First, Bak was shown to be normally sequestered by Bcl-xL and Mcl-1 but displaced by BH3-only proteins⁵⁵. Second, in direct opposition to the direct activation model, Willis *et al.* demonstrated that apoptosis can be induced by “sensitizer” BH3-only proteins or mutants of “activators” that are unable to bind Bax/Bak, even in cells that are deficient for the three activator BH3-only proteins Bid, Bim, and Puma⁵⁶. These results gave rise to the

indirect activation model, which proposed that the effectors Bax/Bak are constitutively active but are normally sequestered by the anti-apoptotic Bcl-2 proteins. During apoptosis, BH3-only proteins bind to anti-apoptotic Bcl-2 proteins and indirectly activate Bax/Bak by displacing and releasing sequestered active Bax/Bak⁵⁷ (Figure 2B, top).

Although this model is consistent with the activation of Bak, it has difficulty explaining the activation of Bax. It was therefore hypothesized that a small fraction of Bax is constitutively active but is sequestered by the anti-apoptotic Bcl-2 proteins on the mitochondria. The BH3-only proteins release the active Bax, which in turn recruits cytosolic Bax to the mitochondria and activates it through an auto-activation mechanism⁵⁸ (Figure 2B, bottom). However, this hypothesis does not address the

Figure 2. Comparison of the three models of BH3-only mediated Bax/Bak activation. **A)** The direct activation model classifies BH3-only proteins as either activators (i.e. Bid, Bim, and Puma) or sensitizers (Bad). When not sequestered by anti-apoptotic Bcl-2 proteins, the activators directly bind to and activate Bax/Bak. Sensitizers cannot directly activate Bax/Bak but can bind to anti-apoptotic Bcl-2 proteins. To initiate apoptosis, the sensitizers bind to the anti-apoptotic Bcl-2 proteins, which release the activators from sequestration and allow them to directly activate Bax/Bak. **B)** The indirect activation model hypothesizes that Bax/Bak are constitutively active but are sequestered by the anti-apoptotic Bcl-2 proteins. During apoptosis, the BH3-only proteins bind to the anti-apoptotic Bcl-2 proteins and displace active Bax/Bak, which then damage the mitochondria. It is also hypothesized that active Bax can recruit inactive cytosolic Bax to the mitochondria through auto-activation. **C)** The membrane (lipids)-mediated spontaneous activation model hypothesizes that during apoptosis, the BH3-only proteins bind to and neutralize the anti-apoptotic Bcl-2 proteins. This allows the mitochondrial outer membrane (MOM) lipids to serve as the direct activator of Bax/Bak, as Bax freely migrates to the MOM through free diffusion. Helix 9 mediates membrane association for both Bax/Bak, and such association results in spontaneous activation of Bax/Bak in the lipid bilayer of the MOM. Similar to the indirect activation model, the BH3-only proteins' only function is to inactivate the anti-apoptotic Bcl-2 proteins. However, in the membrane (lipids)-mediated spontaneous (MMS) model, both Bax and Bak are directly activated by the MOM. Figure 2c is adapted from Figure 5b of our previous paper in Cell Research⁵⁹.

origin of the fraction of active Bax and the physiological relevance of auto-activation. Furthermore, as the indirect activation model was based on the key observation of Bax/Bak activation in Bid/Bim/Puma-deficient cells, the existence of additional activator BH3-only proteins made it more difficult to fully dissociate from the direct activation model^{26,33–36}. It is worth mentioning that in an elegant genetic knockin study in mice, Merino *et al.* compared the pro-apoptotic activities of wild-type Bim and its BH3-replacement mutants and presented data to suggest that neutralization of the anti-apoptotic Bcl-2 proteins is required but not sufficient for maximal apoptosis, suggesting the presence of both indirect and direct activation *in vivo*⁶⁰.

Functional redundancy and functional/physical connections: the Achilles' heel of Bcl-2 research

A salient feature of the Bcl-2 family proteins is functional redundancy due to the presence of eight BH3-only, five anti-apoptotic, and two effector proteins. With multiple proteins sharing similar targets and functions in the same cell, it is difficult to pinpoint the real target, function, and mechanism of action of a particular protein^{23,59,62}.

As Bax and Bak are functionally redundant in many situations, our understanding of their functions was severely limited until a seminal study in which mouse embryonic fibroblasts (MEFs) doubly deficient for Bax and Bak were shown to be highly

resistant to multiple apoptotic stimuli²². This breakthrough ended years of frustration and established Bax and Bak as functionally equivalent and critical effectors of MOMP. However, the investigation into the functions and mechanisms of BH3-only proteins had been more challenging owing to the presence of eight BH3-only proteins, with seven of them potentially functioning as direct activators^{33–36}. In efforts to differentiate direct and indirect activation, loss-of-function studies have been carried out to eliminate the direct activators by generating Bid/Bim double knockout (DKO), Bid/Bim/Puma triple knockout (TKO), Bid/Bim/Puma/p53 quadruple knockout (QKO), and Bid/Bim/Puma/Noxa QKO cells^{36,55,63–65}. However, these cell models are insufficient to address this critical issue because of the presence of additional direct activators (i.e. Bik, Hrk, and Bmf) in those cells^{33–35}.

The second challenge is the complex functional and physical connections among the Bcl-2 family proteins. The anti-apoptotic Bcl-2 proteins interact with at least some of the BH3-only proteins constitutively. Therefore, in cells deficient for certain BH3-only proteins—notably, the Bid/Bim/Puma TKO and Bid/Bim/Puma/Noxa QKO cells^{36,63}—the loss of the BH3-only proteins should effectively increase the concentration of “unsequestered” or “active” Bcl-xL, Bcl-2, and Mcl-1, etc., whose total protein levels may not change. This increase would mean that these cells should have a greater capacity to sequester or suppress the active Bax/Bak and should at least in part explain the resistance of these TKOs and QKOs to apoptosis stimulation or the expression of the sensitizer protein Bad. Because of such complexity, vastly different conclusions have been drawn to either support or disprove the role of the activator proteins and the distinction between activators and sensitizers^{26,36,59,66}.

“OctaKO” and “Bcl-2 allKO” cells: essential tools to reveal that direct activation is not required for BH3-only-mediated Bax/Bak activation and to establish the membrane (lipid)-mediated spontaneous model

In the past decade, the list of direct activator BH3-only proteins has grown to include Bid, Bim, Puma, Hrk, Noxa, Bik, and Bmf, which bind both Bax/Bak and the anti-apoptotic Bcl-2 proteins. Such a functional redundancy makes it near impossible to separate the functions of the direct activators and the sensitizers unless all of the direct activators are eliminated⁶¹. The advent of CRISPR technology made it possible to remove this roadblock.

Through a sequential use of the CRISPR/Cas9 technology, HCT116 cells that lack all eight BH3-only proteins were generated and named OctaKO cells⁶⁶. This allowed for the interrogation of the function of individual BH3-only proteins without the participation of other BH3-only proteins. In this cell model, elimination of the two anti-apoptotic Bcl-2 proteins Bcl-xL and Mcl-1 by siRNA, pharmacological inhibition, or genetic elimination induced Bax/Bak activation and apoptosis efficiently. These results, for the first time, demonstrated that none of the direct activator BH3-only proteins is necessary for Bax/Bak activation once Bcl-xL/Mcl-1 are neutralized.

Similar strategies were used to generate the Bcl-2 allKO cells, which lost all eight pro-apoptotic BH3-only proteins, Bnip3 and Nix (non-apoptotic BH3-only proteins), the anti-apoptotic Bcl-2 proteins (Bcl-2, Bcl-xL, Mcl-1, A1, and Bcl-w), and Bax/Bak⁶⁶. These cells for the first time provided a clean slate to examine the functional relationship, through reconstitution, among Bcl-2 family proteins. Surprisingly, even minute amounts of Bax or Bak, re-introduced through a Dox-inducible expression system, were sufficient to kill these cells, and such killing was easily blocked by Bcl-xL. The highly efficient activation of Bax/Bak in Bcl-2 allKO cells is in stark contrast with the prediction from the direct activation model, which suggests that Bax should stay in the cytosol in the absence of an activator BH3-only protein. Importantly, mutants of Bax and Bak that lack their respective helix 9 (transmembrane [TM] domain) are incapable of homo-oligomerization and killing, and yet such activity was completely restored by the attachment of a heterologous TM domain, strongly suggesting that MOM association is sufficient for Bax/Bak activation⁶⁶.

These results formed the basis for a novel model of Bax/Bak activation, the membrane (lipid)-mediated spontaneous model (MMS), in which the BH3-only proteins activate Bax/Bak by inactivation or neutralization of the anti-apoptotic Bcl-2 proteins, instead of direct activation. This allows the Bax/Bak molecules to freely migrate to and diffuse in the MOM, homo-oligomerize, and form pores (Figure 2C, Figure 3). In this model, the driving force of Bax/Bak activation is essentially free diffusion and the well-documented MOM association through helix 9⁶⁶. Consistent with this model, cytosolic Bax was reported to be constitutively targeted to the mitochondria in non-apoptotic cells^{67,68}.

What are the targets of BH3-only proteins: anti-apoptotic Bcl-2 proteins, Bax/Bak, or both?

Our understanding of the functions of the BH3-only proteins was significantly enhanced when Chen *et al.* defined the binding specificity of various BH3-only proteins toward different anti-apoptotic Bcl-2 proteins⁶⁹. Using peptides of the BH3 domains of the BH3-only proteins, Chen *et al.* found that while Bid, Puma, and Bim are promiscuous in binding all five anti-apoptotic Bcl-2 proteins, Bad only selectively binds to Bcl-w, Bcl-xL, and Bcl-2, and Noxa binds only Mcl-1 and A1. The combination of Bad and Noxa potentially induced apoptosis⁶⁹. Using derivatives of the OctaKO cells that also lack Bcl-xL or Mcl-1, the Huang *et al.* study confirmed and refined these findings by demonstrating that while Bad and Noxa preferentially bind Bcl-xL and Mcl-1, respectively, tBid, Bim, Puma, Bmf, Bik, and Hrk are promiscuous binders toward both⁵⁹ (Figure 4A).

The critical role of Bcl-xL and Mcl-1 in cell survival was first demonstrated in HeLa cells in which siRNA knockdown of both Bcl-xL and Mcl-1 resulted in robust apoptosis^{70,71}. This assay was subsequently carried out in multiple cell lines with deficiencies in several BH3-only proteins and ultimately the OctaKO cells. Without exception, the double siRNA knockdown induced robust apoptosis, strongly suggesting that Bcl-xL

Figure 3. Membrane (lipids)-mediated spontaneous activation of Bax/Bak by the BH3-only proteins. In this model, upon apoptotic stimulation, the BH3-only proteins neutralize/inactivate the anti-apoptotic Bcl-2 proteins Bcl-xL/Mcl-1. Such inactivation allows Bax to freely migrate to the mitochondrial outer membrane (MOM) through the interaction between helix 9 and the lipid bilayer. Anchored to the MOM through helix 9, both Bax and Bak spontaneously homo-oligomerize and form apoptotic pores within the lipid milieu. This figure is adapted from Figure 7 of our previous paper in *Genes & Development*⁶⁵.

and Mcl-1 are the major targets of the BH3-only proteins during apoptosis^{36,64,65,70,71}. In support of Bcl-xL and Mcl-1's critical role as gatekeepers against Bax/Bak activation, an elegant mouse genetic study found that while *Mcl-1^{Bcl-2} +/-* and *Bcl-xL^{-/-}Bcl-2^{+/-}* mice are almost normal, *Mcl-1^{+/-}Bcl-xL^{+/-}* mice displayed craniofacial defects⁷².

Strikingly, Huang *et al.* found that Bad, the sensitizer protein, induced apoptosis as efficiently as the activator proteins tBid and Bim in the OctaKO derivative cells⁵⁹. Additionally, mutant activator BH3-only proteins with their BH3 domain replaced by that of Bad showed no difference from the wild-type activators in the induction of apoptosis in the OctaKO cells. Furthermore, tBid and Bim (both activators) as well as Bad (sensitizer) were found to induce apoptosis without distinguishing Bax from Bak, consistent with the conclusions of a recent *in vitro* study³⁵ but in contrast to an earlier study⁷³. Lastly, tBid and Bim are unable to accelerate the spontaneous activation of Bax in the Bcl-2 allKO cells, suggesting that there is no functional/physical interaction between the activator

BH3-only proteins and Bax/Bak after the neutralization of the anti-apoptotic Bcl-2 proteins⁵⁹. These results are incompatible with the direct activation model and argue strongly that there is no distinction between direct activators and sensitizers and that the only target of the BH3-only proteins is the anti-apoptotic Bcl-2 proteins. These findings unify the functions of BH3-only proteins and provide an elegantly simple model for apoptotic signaling (Figure 4B).

Of note, loss-of-function studies in mice have clearly demonstrated differential requirements for various BH3-only proteins *in vivo*^{74–84}. These observations are consistent with the notion that the BH3-only proteins' roles are determined by their expression level, location, various regulations, and affinity with different anti-apoptotic Bcl-2 proteins, etc.

A bigger role for MOM in Bax and Bak activation

MOM has traditionally been viewed as a passive victim of activated Bax/Bak. However, some *in vitro* studies have suggested that MOM can serve as a platform for multiple interactions

Figure 4. Apoptotic signaling through membrane (lipids)-mediated spontaneous activation of Bax/Bak. A The intrinsic (mitochondrial-dependent) apoptotic pathway based on the membrane (lipids)-mediated spontaneous (MMS) Bax/Bak activation model. **B** The intrinsic (mitochondrial-dependent) and extrinsic (death receptor-mediated) apoptotic pathways based on the MMS activation model. Apaf-1, apoptotic protease activating factor-1; Casp, caspase; Cyt c, cytochrome c; FADD, Fas associated death domain protein; MOM, mitochondrial outer membrane; SMAC, second mitochondria-derived activator of caspase; XIAP, X-linked inhibitor of apoptosis protein.

among the Bcl-2 family proteins, including tBid-Bax, tBid-Bcl-xL interactions, etc.^{31,85,86}. Results from the new series of studies, especially the finding that the TD (helix 9) of Bax/Bak was required and sufficient for Bax/Bak activation in the absence of all BH3-only and anti-apoptotic Bcl-2 proteins, suggested a much bigger role for the MOM in the regulation of Bax/Bak: the direct activator^{59,65,66}. In other words, the MOM itself functions as the instigator of Bax/Bak's attack by directly recruiting and activating Bax/Bak. These results also suggested a much bigger role for helix 9: the trigger site. Importantly, the interaction between helix 9 and the MOM is spontaneous, stable, and easily detectable in the cell. On the other hand, the transient activator BH3-only-Bax/Bak interaction observed *in vitro* has been shown to be non-essential in cellular assays.

How does the MOM activate Bax? First, as Bax has been shown to exist in a dynamic equilibrium between the cytosol and

mitochondria, and that Bcl-xL constantly retro-translocates Bax/Bak into the cytoplasm^{67,68,87}, it is essential that the anti-apoptotic Bcl-2 proteins are inactivated by the BH3-only proteins or other inhibitors before the MOM becomes a direct activator^{59,66}. Second, a systematic mapping of the helices of Bax for their capacity to associate with the MOM indicated that helices 4, 5, 6, and 9 contain mitochondrial targeting signals (MTS)⁸⁸. It is not difficult to imagine that the MOM has the capacity to reconfigure the whole molecule through these multiple lipid-protein interactions to promote Bax/Bak homo-oligomerization upon the initial contact with helix 9. Third, as helix 9 of Bax was recently shown to form dimers in the MOM, such dimerization through helix 9 should provide further impetus for homo-oligomerization and activation of Bax/Bak molecules^{52,89–91}.

Thus, lipids are most likely the direct activator of Bax/Bak. Consistent with this suggestion, cardiolipin was found to be

required for Bax activation^{29,92}. Interestingly, it has been shown that non-ionic detergents, which somewhat mimic the lipid bilayer environment, are excellent activators of Bax/Bak^{12,93}. It is therefore of great interest to test if certain lipid compositions can support Bax activation without the activator BH3-only proteins, even though the commonly used preparations prohibit such activity. A liposome composition that supports the constitutive activation of Bax in the absence of tBid should recapitulate what is observed in the cell.

MOM proteins may help recruit Bax to the MOM. VDAC2, despite earlier observations that it sequesters and suppresses Bak^{94,95}, has recently been shown to be required for the mitochondrial translocation of Bax^{96,97}. In addition, the mitochondrial fission regulator Drp1 has been shown to play a positive role in the regulation of Bax/Bak activation^{98–100}. As both VDAC2 and Drp1 are present in the Bcl-2 allKO cells, it is possible that these proteins play a positive role in recruiting Bax/Bak along with the lipids to allow spontaneous localization^{59,66}.

De novo activation versus auto-activation of Bax/Bak

Based on results from a liposome study with an extended BH3 peptide and Bax, it has been hypothesized that active Bax uses its BH3 domain to directly bind and activate an inactive Bax to accelerate Bax activation. This putative feed-forward mechanism is termed “auto-activation”¹⁰¹. In support of this hypothesis, during apoptosis, Bax/Bak undergo conformational changes and expose the BH3 domain^{17,40}. Auto-activation has recently been suggested to be a critical feed-forward mechanism for Bax/Bak activation in response to DNA damage and other death stimuli^{36,102}. However, the origin of the first active Bax/Bak molecule and the physiological significance of auto-activation are unclear. Furthermore, there are no experimental data to support the hypothesis that active Bax can recruit cytosolic Bax to the mitochondria.

Both the direct activation and MMS model probe the mechanism of Bax/Bak’s *de novo* activation. Although there is potential involvement of auto-activation as a feed-forward mechanism to sustain Bax/Bak activation through homo-oligomerization, it is not the triggering mechanism that initiates the activation of Bax/Bak *per se*. For example, although the “interconnected” model contains an element of auto-activation, the origin of the first active Bax/Bak molecules was attributed to direct activation³⁶.

How do the BH3 mimetics work?

Thanks to a close collaboration between basic researchers and the pharmaceutical industry, numerous small molecules that mimic the structures of the BH3 domains of the BH3-only proteins have been developed to kill cancer cells and overcome resistance¹⁰³. For example, through NMR-based fragment screen and structure-guided design, ABT-737 was developed to mimic BadBH3, and it is highly selective against Bcl-xL, Bcl-2, and Bcl-w¹⁰⁴. Using similar strategies, Kotschy *et al.* designed S63845, a highly selective inhibitor of Mcl-1¹⁰⁵. Remarkably, venetoclax, a BH3 mimetic and the

first small-molecule drug that disrupts a protein–protein interaction, was FDA approved for the treatment of a sub-type of chronic lymphocytic lymphoma^{103,106–109}. However, the obvious question is this: which interaction does it disrupt in the cell? According to the direct activation model, the Bcl-2 mimetic disrupts two interactions: activator BH3-only–Bcl-2 and Bax/Bak–Bcl-2. The liberated activator BH3-only proteins then directly activate the liberated Bax or Bak¹⁰⁶. However, this scenario is difficult to prove experimentally and is not consistent with the observation that combinations of BH3 mimetics against Bcl-xL and Mcl-1 are sufficient to kill OctaKO HCT116 cells^{66,110}. Such an observation is explained by the MMS model, in which the BH3 mimetics neutralize Bcl-xL/Mcl-1 and allow the MOM to directly activate Bax/Bak. It is of interest, however, that over-expression of Bcl-xL confers resistance to the apoptotic effects of Bcl-xL-targeting BH3 mimetics, suggesting limitations of these small-molecule inhibitors^{111,112}.

How do the anti-apoptotic Bcl-2 suppress Bax/Bak activation?

It is well established that Bax exists mostly as an inactive and monomeric protein in cytosol¹⁴. However, the spontaneous activation of Bax in the Bcl-2 allKO strongly suggests that Bax is constantly trying to land on the mitochondria through its C-terminal tail ($\alpha 9$)^{59,66}, consistent with the observation of a natural tendency of Bax to move to the mitochondria^{67,68}. It is also clear that Bcl-xL and Mcl-1 play a critical role in keeping Bax in the cytosol. How do Bcl-xL and Mcl-1 achieve this feat? Edlich *et al.* demonstrated that Bcl-xL has the ability to push the mitochondrially localized Bax molecules into the cytosol⁶⁷. Interestingly, Todt *et al.* demonstrated that Bak can also be retro-translocated by Bcl-xL, albeit at a much slower rate⁸⁷. However, the molecular mechanisms of such a retro-translocation activity remain unclear. Nevertheless, helix 9 of Bcl-xL may play a role in mediating this mysterious activity¹¹³, through which the Bax molecule may be re-packed into a globular shape so that helix 9 can fit into the hydrophobic groove. Once this re-packing is finished, Bax is expected to leave the mitochondria and become cytosolic. The molecular details of such extraordinary activity will be a focus of future studies.

The membrane (lipids)-mediated spontaneous model versus indirect and direct activation models

The MMS model combines some important elements from both the direct and the indirect activation models yet for the first time introduced a new interaction as the trigger for Bax/Bak activation (Figure 2 and Figure 3, Table 1). First, in agreement with indirect activation, the MMS model proposes that there is no hierarchy among the BH3-only proteins and the only target of the BH3-only proteins are the anti-apoptotic Bcl-2 proteins. Second, similar to the direct activation model, the MMS model proposes that a *de novo* direct activation, instead of de-repression, is the mechanism of Bax/Bak activation. However, the MOM, instead of the activator BH3-only proteins, is the direct activator. Third, the MMS model proposes that

Table 1. Comparison of the three models of BH3-only-mediated Bax/Bak activation.

Model of Bax/Bak Activation	Targets of the BH3-only proteins	BH3-only–Bax/Bak Interaction	Activation Mechanism	Direct Activator	Trigger Site on Bax/Bak	Driving Force
Indirect Activation (Displacement)	Anti-apoptotic Bcl-2 proteins	None	Displacement and De-repression	None	None	Stable BH3-only–Bcl-xL/Mcl-1 interaction
Direct Activation	Bax/Bak and anti-apoptotic Bcl-2 proteins	Yes	<i>De novo</i> and direct activation	Activator BH3-only proteins	Helices 1/6 or the hydrophobic groove	Transient BH3 (BH3-only)–groove (Bax/Bak) interaction
Membrane (lipids)-mediated Spontaneous Activation	Anti-apoptotic Bcl-2 proteins	None	<i>De novo</i> and direct activation	Mitochondrial outer membrane (lipids)	C-terminal tail (helix 9)	Stable BH3-only–Bcl-xL/Mcl-1 interaction, free diffusion, and stable Bax/Bak α 9-membrane interaction

helix 9 is the trigger site for Bax/Bak activation. Instead of a transient, “hit-and-run” interaction as proposed in the direct activation model, the interaction between Bax/Bak and the MOM is visible and stable and consists of multiple interfaces once homo-oligomerization initiates.

Some questions remain to be answered: what are the molecular details of MOM-mediated spontaneous Bax/Bak activation? What are the dynamics of the helix 9–hydrophobic groove interaction in the cytosol? Can *in vitro* liposome assays recapitulate the MMS model? What are the mechanisms of retro-translocation?

Conclusion

The balance among the Bcl-2 family members, namely the relative amounts of the BH3-only proteins and the anti-apoptotic Bcl-2 proteins, determines the fate of the cell through the activation of Bax and Bak^{59,66}. According to the MMS model, during apoptotic signaling, once all the Bcl-xL/Mcl-1 molecules are neutralized by the BH3-only proteins, Bax/Bak spontaneously (through free diffusion) associate with the MOM, which acts as a direct activator to trigger Bax/Bak's

self-aggregation and pore formation^{59,66}. This remarkably efficient and simple mechanism might provide new thinking on pharmacological intervention. For example, is it possible to manipulate the lipid composition or other properties of the MOM to either accelerate or slow down the attack by the Bax/Bak molecules? Also, since helix 9's exposure and access to the MOM is essential to this process, is it possible to design small molecules to promote or suppress the dislodging of helix 9 from the hydrophobic groove of Bax/Bak, therefore modulating the targeting of these molecules to the mitochondria^{114,115}?

Acknowledgements

We are grateful to Dr. Keith Johnson (Department of Oral Biology and Eppler Institute for Research in Cancer and Allied Diseases, University of Nebraska Medical Center, USA) for his suggestions and critical reading of the manuscript. Xu Luo dedicates this work to the loving memory of his father, Dr. Shiqi Luo, one of the finest neurosurgeons ever, whose encouragement, support, and advice made the work possible.

References

- Daniel NN, Korsmeyer SJ: **Cell death: Critical control points**. *Cell*. 2004; **116**(2): 205–19.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jiang X, Wang X: **Cytochrome C-mediated apoptosis**. *Annu Rev Biochem*. 2004; **73**: 87–106.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kalkavan H, Green DR: **MOMP, cell suicide as a BCL-2 family business**. *Cell Death Differ*. 2018; **25**(1): 46–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
- Cleary ML, Smith SD, Sklar J: **Cloning and structural analysis of cDNAs for bcl-2 and a hybrid bcl-2/immunoglobulin transcript resulting from the t(14;18) translocation**. *Cell*. 1986; **47**(1): 19–28.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tsujimoto Y, Cossman J, Jaffe E, *et al.*: **Involvement of the bcl-2 gene in human follicular lymphoma**. *Science*. 1985; **228**(4706): 1440–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Vaux DL, Cory S, Adams JM: **Bcl-2 gene promotes haemopoietic cell survival and cooperates with c-myc to immortalize pre-B cells**. *Nature*. 1988; **335**(6189): 440–2.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cory S, Huang DCS, Adams JM: **The Bcl-2 family: Roles in cell survival and oncogenesis**. *Oncogene*. 2003; **22**(53): 8590–607.
[PubMed Abstract](#) | [Publisher Full Text](#)

8. Puthalakath H, Villunger A, O'Reilly LA, *et al.*: **Bmf: A proapoptotic BH3-only protein regulated by interaction with the myosin V actin motor complex, activated by anoikis.** *Science*. 2001; 293(5536): 1829–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
9. Chipuk JE, Moldoveanu T, Liambi F, *et al.*: **The BCL-2 family reunion.** *Mol Cell*. 2010; 37(3): 299–310.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Moldoveanu T, Follis AV, Kriwacki RW, *et al.*: **Many players in BCL-2 family affairs.** *Trends Biochem Sci*. 2014; 39(3): 101–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Kale J, Osterlund EJ, Andrews DW: **BCL-2 family proteins: Changing partners in the dance towards death.** *Cell Death Differ*. 2018; 25(1): 65–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
12. Suzuki M, Youle RJ, Tjandra N: **Structure of Bax: Coregulation of dimer formation and intracellular localization.** *Cell*. 2000; 103(4): 645–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Nechushtan A, Smith CL, Hsu YT, *et al.*: **Conformation of the Bax C-terminus regulates subcellular location and cell death.** *EMBO J*. 1999; 18(9): 2330–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
14. Wolter KG, Hsu YT, Smith CL, *et al.*: **Movement of Bax from the cytosol to mitochondria during apoptosis.** *J Cell Biol*. 1997; 139(5): 1281–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Moldoveanu T, Liu Q, Tocilj A, *et al.*: **The X-ray structure of a BAK homodimer reveals an inhibitory zinc binding site.** *Mol Cell*. 2006; 24(5): 677–88.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Youle RJ, Strasser A: **The BCL-2 protein family: Opposing activities that mediate cell death.** *Nat Rev Mol Cell Biol*. 2008; 9(1): 47–59.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Dewson G, Kratina T, Sim HW, *et al.*: **To trigger apoptosis, Bak exposes its BH3 domain and homodimerizes via BH3:groove interactions.** *Mol Cell*. 2008; 30(3): 369–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. George NM, Evans JJD, Luo X: **A three-helix homo-oligomerization domain containing BH3 and BH1 is responsible for the apoptotic activity of Bax.** *Genes Dev*. 2007; 21(15): 1937–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
19. Strasser A, Cory S, Adams JM: **Deciphering the rules of programmed cell death to improve therapy of cancer and other diseases.** *EMBO J*. 2011; 30(18): 3667–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Ke FFS, Vanyai HK, Cowan AD, *et al.*: **Embryogenesis and Adult Life in the Absence of Intrinsic Apoptosis Effectors BAX, BAK, and BOK.** *Cell*. 2018; 173(5): 1217–1230.e17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
21. Liambi F, Wang YM, Victor B, *et al.*: **BOK Is a Non-canonical BCL-2 Family Effector of Apoptosis Regulated by ER-Associated Degradation.** *Cell*. 2016; 165(2): 421–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Wei MC, Zong WX, Cheng EH, *et al.*: **Proapoptotic BAX and BAK: A requisite gateway to mitochondrial dysfunction and death.** *Science*. 2001; 292(5517): 727–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Chipuk JE, Green DR: **How do BCL-2 proteins induce mitochondrial outer membrane permeabilization?** *Trends Cell Biol*. 2008; 18(4): 157–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Wang K, Yin XM, Chao DT, *et al.*: **BID: A novel BH3 domain-only death agonist.** *Genes Dev*. 1996; 10(22): 2859–69.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Letai A, Bassik MC, Walensky LD, *et al.*: **Distinct BH3 domains either sensitize or activate mitochondrial apoptosis, serving as prototype cancer therapeutics.** *Cancer Cell*. 2002; 2(3): 183–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
26. Kim H, Rafiuddin-Shah M, Tu HC, *et al.*: **Hierarchical regulation of mitochondrion-dependent apoptosis by BCL-2 subfamilies.** *Nat Cell Biol*. 2006; 8(12): 1348–58.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Kim H, Tu HC, Ren D, *et al.*: **Stepwise Activation of BAX and BAK by tBID, BIM and PUMA Initiates Mitochondrial Apoptosis.** *Mol Cell*. 2009; 36: 487–99.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
28. Kuwana T, Bouchier-Hayes L, Chipuk JE, *et al.*: **BH3 domains of BH3-only proteins differentially regulate Bax-mediated mitochondrial membrane permeabilization both directly and indirectly.** *Mol Cell*. 2005; 17(4): 525–35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
29. Kuwana T, Mackey MR, Perkins G, *et al.*: **Bid, Bax, and Lipids Cooperate to Form Supramolecular Openings in the Outer Mitochondrial Membrane.** *Cell*. 2002; 111(3): 331–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
30. Liambi F, Moldoveanu T, Tait SWG, *et al.*: **A unified model of mammalian BCL-2 protein family interactions at the mitochondria.** *Mol Cell*. 2011; 44(4): 517–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
31. Lovell JF, Billen LP, Bindner S, *et al.*: **Membrane Binding by tBid Initiates an Ordered Series of Events Culminating in Membrane Permeabilization by Bax.** *Cell*. 2008; 135(6): 1074–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
32. Bogner C, Kale J, Pogmore J, *et al.*: **Allosteric Regulation of BH3 Proteins in Bcl-xL Complexes Enables Switch-like Activation of Bax.** *Mol Cell*. 2020; 77(4): 901–912.e9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
33. Dai H, Smith A, Meng XW, *et al.*: **Transient binding of an activator BH3 domain to the Bak BH3-binding groove initiates Bak oligomerization.** *J Cell Biol*. 2011; 194(1): 39–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
34. Du H, Wolf J, Schafer B, *et al.*: **BH3 domains other than Bim and Bid can directly activate Bax/Bak.** *J Biol Chem*. 2011; 286(1): 491–501.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
35. Hockings C, Anwari K, Ninnis RL, *et al.*: **Bid chimeras indicate that most BH3-only proteins can directly activate Bak and show no preference for Bak versus Bax.** *Cell Death Dis*. 2015; 6(4): e1735.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Chen HC, Kanai M, Inoue-Yamauchi A, *et al.*: **An interconnected hierarchical model of cell death regulation by the BCL-2 family.** *Nat Cell Biol*. 2015; 17(10): 1270–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
37. Edwards AL, Gavathiotis E, LaBelle JL, *et al.*: **Multimodal Interaction with BCL-2 Family Proteins Underlies the Proapoptotic Activity of PUMA BH3.** *Chem Biol*. 2013; 20(7): 888–902.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Cartron PF, Gallenne T, Bougras G, *et al.*: **The first alpha helix of Bax plays a necessary role in its ligand-induced activation by the BH3-only proteins Bid and PUMA.** *Mol Cell*. 2004; 16(5): 807–18.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Gallenne T, Gautier F, Oliver L, *et al.*: **Bax activation by the BH3-only protein Puma promotes cell dependence on antiapoptotic Bcl-2 family members.** *J Cell Biol*. 2009; 185(2): 279–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Gavathiotis E, Reyna DE, Davis ML, *et al.*: **BH3-triggered structural reorganization drives the activation of proapoptotic BAX.** *Mol Cell*. 2010; 40(3): 481–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
41. Gavathiotis E, Suzuki M, Davis ML, *et al.*: **BAX activation is initiated at a novel interaction site.** *Nature*. 2008; 455(7216): 1076–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
42. Czabotar PE, Westphal D, Dewson G, *et al.*: **Bax crystal structures reveal how BH3 domains activate Bax and nucleate its oligomerization to induce apoptosis.** *Cell*. 2013; 152(3): 519–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
43. Moldoveanu T, Grace CR, Liambi F, *et al.*: **BID-induced structural changes in BAK promote apoptosis.** *Nat Struct Mol Biol*. 2013; 20(5): 589–97.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
44. Brouwer JM, Lan P, Cowan AD, *et al.*: **Conversion of Bim-BH3 from Activator to Inhibitor of Bak through Structure-Based Design.** *Mol Cell*. 2017; 68(4): 659–672.e9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
45. Robin AY, Krishna Kumar K, Westphal D, *et al.*: **Crystal structure of Bax bound to the BH3 peptide of Bim identifies important contacts for interaction.** *Cell Death Dis*. 2015; 6(7): e1809–e1809.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. Brouwer JM, Westphal D, Dewson G, *et al.*: **Bak core and latch domains separate during activation, and freed core domains form symmetric homodimers.** *Mol Cell*. 2014; 55(6): 938–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
47. Leshchiner ES, Braun CR, Bird GH, *et al.*: **Direct activation of full-length proapoptotic BAK.** *Proc Natl Acad Sci U S A*. 2013; 110(11): E986–95.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. Dengler MA, Robin AY, Gibson L, *et al.*: **BAX Activation: Mutations Near Its Proposed Non-canonical BH3 Binding Site Reveal Allosteric Changes Controlling Mitochondrial Association.** *Cell Rep*. 2019; 27(2): 359–373.e6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
49. Peng R, Tong JS, Li H, *et al.*: **Targeting Bax interaction sites reveals that only homo-oligomerization sites are essential for its activation.** *Cell Death*

- Differ. 2013; 20(5): 744–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
50. Leber B, Lin J, Andrews DW: **Embedded together: The life and death consequences of interaction of the Bcl-2 family with membranes.** *Apoptosis*. 2007; 12(5): 897–911.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 51. Leber B, Lin J, Andrews DW: **Still embedded together binding to membranes regulates Bcl-2 protein interactions.** *Oncogene*. 2010; 29(38): 5221–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 52. Bleicken S, Jeschke G, Stegmueller C, *et al.*: **Structural model of active Bax at the membrane.** *Mol Cell*. 2014; 56(4): 496–505.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 53. Czabotar PE, Lessene G, Strasser A, *et al.*: **Control of apoptosis by the BCL-2 protein family: Implications for physiology and therapy.** *Nat Rev Mol Cell Biol*. 2014; 15(1): 49–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
 54. Westphal D, Kluck RM, Dewson G: **Building blocks of the apoptotic pore: How Bax and Bak are activated and oligomerize during apoptosis.** *Cell Death Differ*. 2014; 21(2): 196–205.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 55. Willis SN, Chen L, Dewson G, *et al.*: **Proapoptotic Bak is sequestered by Mcl-1 and Bcl-x_L, but not Bcl-2, until displaced by BH3-only proteins.** *Genes Dev*. 2005; 19(11): 1294–305.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 56. Willis SN, Fletcher JI, Kaufmann T, *et al.*: **Apoptosis initiated when BH3 ligands engage multiple Bcl-2 homologs, not Bax or Bak.** *Science*. 2007; 315(5813): 856–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 57. Willis SN, Adams JM: **Life in the balance: How BH3-only proteins induce apoptosis.** *Curr Opin Cell Biol*. 2005; 17(6): 617–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 58. Fletcher JI, Meusburger S, Hawkins CJ, *et al.*: **Apoptosis is triggered when prosurvival Bcl-2 proteins cannot restrain Bax.** *Proc Natl Acad Sci U S A*. 2008; 105(47): 18081–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 59. Huang K, O'Neill KL, Li J, *et al.*: **BH3-only proteins target BCL-x_L/MCL-1, not BAX/BAK, to initiate apoptosis.** *Cell Res*. 2019; 29(11): 942–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 60. Mérimo D, Giam M, Hughes PD, *et al.*: **The role of BH3-only protein Bim extends beyond inhibiting Bcl-2-like prosurvival proteins.** *J Cell Biol*. 2009; 186(3): 355–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 61. Borner C, Andrews DW: **The apoptotic pore on mitochondria: Are we breaking through or still stuck?** *Cell Death Differ*. 2014; 21(2): 187–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 62. García-Sáez AJ: **The secrets of the Bcl-2 family.** *Cell Death Differ*. 2012; 19(11): 1733–40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 63. Ren D, Tu HC, Kim H, *et al.*: **BID, BIM, and PUMA are essential for activation of the BAX- and BAK-dependent cell death program.** *Science*. 2010; 330(6009): 1390–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 64. Senft D, Weber A, Saathoff F, *et al.*: **In non-transformed cells Bak activates upon loss of anti-apoptotic Bcl-XL and Mcl-1 but in the absence of active BH3-only proteins.** *Cell Death Dis*. 2015; 6(11): e1996–e1996.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 65. Zhang J, Huang K, O'Neill KL, *et al.*: **Bax/Bak activation in the absence of Bid Bim, Puma, and p53.** *Cell Death Dis*. 2016; 7(6): e2266.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 66. O'Neill KL, Huang K, Zhang J, *et al.*: **Inactivation of prosurvival Bcl-2 proteins activates Bax/Bak through the outer mitochondrial membrane.** *Genes Dev*. 2016; 30(8): 973–88.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 67. Edlich F, Banerjee S, Suzuki M, *et al.*: **Bcl-xL Retrotranslocates Bax from the Mitochondria into the Cytosol.** *Cell*. 2011; 145(1): 104–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 68. Schellenberg B, Wang P, Keeble JA, *et al.*: **Bax exists in a dynamic equilibrium between the cytosol and mitochondria to control apoptotic priming.** *Mol Cell*. 2013; 49(5): 959–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 69. Chen L, Willis SN, Wei A, *et al.*: **Differential Targeting of Prosurvival Bcl-2 Proteins by Their BH3-Only Ligands Allows Complementary Apoptotic Function.** *Mol Cell*. 2005; 17(3): 393–403.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 70. Lopez H, Zhang L, George NM, *et al.*: **Perturbation of the Bcl-2 network and an induced Noxa/Bcl-xL interaction trigger mitochondrial dysfunction after DNA damage.** *J Biol Chem*. 2010; 285(20): 15016–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 71. Zhang L, Lopez H, George NM, *et al.*: **Selective involvement of BH3-only proteins and differential targets of Noxa in diverse apoptotic pathways.** *Cell Death Differ*. 2011; 18(5): 864–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 72. Grabow S, Kueh AJ, Ke F, *et al.*: **Subtle Changes in the Levels of BCL-2 Proteins Cause Severe Craniofacial Abnormalities.** *Cell Rep*. 2018; 24(12): 3285–3295.e4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 73. Sarosiek KA, Chi X, Bachman JA, *et al.*: **BID Preferentially Activates BAK while BIM Preferentially Activates BAX, Affecting Chemotherapy Response.** *Mol Cell*. 2013; 51(6): 751–65.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 74. Coultas L, Bouillet P, Loveland KL, *et al.*: **Concomitant loss of proapoptotic BH3-only Bcl-2 antagonists Bik and Bim arrests spermatogenesis.** *EMBO J*. 2005; 24(22): 3963–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 75. Coultas L, Bouillet P, Stanley EG, *et al.*: **Proapoptotic BH3-Only Bcl-2 Family Member Bik/Nbk Is Expressed in Hemopoietic and Endothelial Cells but Is Redundant for Their Programmed Death.** *Mol Cell Biol*. 2004; 24(4): 1570–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 76. Bouillet P, Metcalf D, Huang DC, *et al.*: **Proapoptotic Bcl-2 relative Bim required for certain apoptotic responses, leukocyte homeostasis, and to preclude autoimmunity.** *Science*. 1999; 286(5445): 1735–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 77. Yin XM, Wang K, Gross A, *et al.*: **Bid-deficient mice are resistant to Fas-induced hepatocellular apoptosis.** *Nature*. 1999; 400(6747): 886–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
 78. Shibue T, Takeda K, Oda E, *et al.*: **Integral role of Noxa in p53-mediated apoptotic response.** *Genes Dev*. 2003; 17(18): 2233–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 79. Jeffers JR, Parganas E, Lee Y, *et al.*: **Puma is an essential mediator of p53-dependent and -independent apoptotic pathways.** *Cancer Cell*. 2003; 4(4): 321–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 80. Villunger A, Michalak EM, Coultas L, *et al.*: **p53- and drug-induced apoptotic responses mediated by BH3-only proteins puma and noxa.** *Science*. 2003; 302(5647): 1036–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
 81. Hübner A, Cavanagh-Kyros J, Rincon M, *et al.*: **Functional Cooperation of the Proapoptotic Bcl2 Family Proteins Bmf and Bim In Vivo.** *Mol Cell Biol*. 2010; 30(1): 98–105.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 82. Hausmann M, Leucht K, Ploner C, *et al.*: **BCL-2 modifying factor (BMF) is a central regulator of anoikis in human intestinal epithelial cells.** *J Biol Chem*. 2011; 286(30): 26533–40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 83. Coultas L, Terzano S, Thomas T, *et al.*: **Hrk/DP5 contributes to the apoptosis of select neuronal populations but is dispensable for haematopoietic cell apoptosis.** *J Cell Sci*. 2007; 120(Pt 12): 2044–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 84. Ranger AM, Zha J, Harada H, *et al.*: **Bad-deficient mice develop diffuse large B cell lymphoma.** *Proc Natl Acad Sci U S A*. 2003; 100(16): 9324–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 85. Huang K, Zhang J, O'Neill KL, *et al.*: **Cleavage by Caspase 8 and Mitochondrial Membrane Association Activate the BH3-only Protein Bid during TRAIL-induced Apoptosis.** *J Biol Chem*. 2016; 291(22): 11843–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 86. García-Sáez AJ, Ries J, Orzáez M, *et al.*: **Membrane promotes tBID interaction with BCL(XL).** *Nat Struct Mol Biol*. 2009; 16(11): 1178–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
 87. Todt F, Cakir Z, Reichenbach F, *et al.*: **Differential retrotranslocation of mitochondrial Bax and Bak.** *EMBO J*. 2014; 34(1): 67–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
 88. George NM, Targy N, Evans JJD, *et al.*: **Bax contains two functional mitochondrial targeting sequences and translocates to mitochondria in a conformational change- and homo-oligomerization-driven process.** *J Biol Chem*. 2010; 285(2): 1384–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 89. Gahl RF, He Y, Yu S, *et al.*: **Conformational Rearrangements in the Pro-Apoptotic Protein, Bax, as It Inserts Into Mitochondria.** *J Biol Chem*. 2014; 289(47): 32871–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)

90. Zhang Z, Subramaniam S, Kale J, *et al.*: **BH3-in-groove dimerization initiates and helix 9 dimerization expands Bax pore assembly in membranes.** *EMBO J.* 2016; **35**(2): 208–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
91. Iyer S, Bell F, Westphal D, *et al.*: **Bak apoptotic pores involve a flexible C-terminal region and juxtaposition of the C-terminal transmembrane domains.** *Cell Death Differ.* 2015; **22**(10): 1665–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
92. Terrones O, Antonsson B, Yamaguchi H, *et al.*: **Lipidic Pore Formation by the Concerted Action of Proapoptotic BAX and tBID.** *J Biol Chem.* 2004; **279**(29): 30081–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
93. Hsu YT, Youle RJ: **Nonionic detergents induce dimerization among members of the Bcl-2 family.** *J Biol Chem.* 1997; **272**(21): 13829–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
94. Cheng EHY, Sheiko TV, Fisher JK, *et al.*: **VDAC2 inhibits BAK activation and mitochondrial apoptosis.** *Science.* 2003; **301**(5632): 513–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
95. Ren D, Kim H, Tu HC, *et al.*: **The VDAC2-BAK rheostat controls thymocyte survival.** *Sci Signal.* 2009; **2**(85): ra48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
96. Chin HS, Li MX, Tan IKL, *et al.*: **VDAC2 enables BAX to mediate apoptosis and limit tumor development.** *Nat Commun.* 2018; **9**(1): 4976.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
97. Lauterwasser J, Todt F, Zerbes RM, *et al.*: **The porin VDAC2 is the mitochondrial platform for Bax retrotranslocation.** *Sci Rep.* 2016; **6**: 32994.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
98. Karbowski M, Lee YJ, Gaume B, *et al.*: **Spatial and temporal association of Bax with mitochondrial fission sites, Drp1, and Mfn2 during apoptosis.** *J Cell Biol.* 2002; **159**(6): 931–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
99. Milani M, Beckett AJ, Al-Zabeeby A, *et al.*: **DRP-1 functions independently of mitochondrial structural perturbations to facilitate BH3 mimetic-mediated apoptosis.** *Cell Death Discov.* 2019; **5**: 117.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
100. Montessuit S, Somasekharan SP, Terrones O, *et al.*: **Membrane Remodeling Induced by the Dynamin-Related Protein Drp1 Stimulates Bax Oligomerization.** *Cell.* 2010; **142**(6): 889–901.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
101. Tan C, Dlugosz PJ, Peng J, *et al.*: **Auto-activation of the apoptosis protein Bax increases mitochondrial membrane permeability and is inhibited by Bcl-2.** *J Biol Chem.* 2006; **281**(21): 14764–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
102. Dai H, Ding H, Meng XW, *et al.*: **Constitutive BAK activation as a determinant of drug sensitivity in malignant lymphohematopoietic cells.** *Genes Dev.* 2015; **29**(20): 2140–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
103. Ashkenazi A, Fairbrother WJ, Levenson JD, *et al.*: **From basic apoptosis discoveries to advanced selective BCL-2 family inhibitors.** *Nat Rev Drug Discov.* 2017; **16**(4): 273–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
104. Oltersdorf T, Elmore SW, Shoemaker AR, *et al.*: **An inhibitor of Bcl-2 family proteins induces regression of solid tumours.** *Nature.* 2005; **435**(7042): 677–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
105. Kotschy A, Szlavik Z, Murray J, *et al.*: **The MCL1 inhibitor S63845 is tolerable and effective in diverse cancer models.** *Nature.* 2016; **538**(7626): 477–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
106. Green DR: **A BH3 Mimetic for Killing Cancer Cells.** *Cell.* 2016; **165**(7): 1560.
[PubMed Abstract](#) | [Publisher Full Text](#)
107. Cory S, Roberts AW, Colman PM, *et al.*: **Targeting BCL-2-like Proteins to Kill Cancer Cells.** *Trends Cancer.* 2016; **2**(8): 443–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
108. Merino D, Kelly GL, Lessene G, *et al.*: **BH3-Mimetic Drugs: Blazing the Trail for New Cancer Medicines.** *Cancer Cell.* 2018; **34**(6): 879–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
109. Souers AJ, Levenson JD, Boghaert ER, *et al.*: **ABT-199, a potent and selective BCL-2 inhibitor, achieves antitumor activity while sparing platelets.** *Nat Med.* 2013; **19**(2): 202–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Faculty Opinions Recommendation](#)
110. Greaves G, Milani M, Butterworth M, *et al.*: **BH3-only proteins are dispensable for apoptosis induced by pharmacological inhibition of both MCL-1 and BCL-X_L.** *Cell Death Differ.* 2019; **26**(6): 1037–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
111. Pécot J, Mailliet L, Le Pen J, *et al.*: **Tight Sequestration of BH3 Proteins by BCL-xL at Subcellular Membranes Contributes to Apoptotic Resistance.** *Cell Rep.* 2016; **17**(12): 3347–58.
[PubMed Abstract](#) | [Publisher Full Text](#)
112. Inoue-Yamauchi A, Jeng PS, Kim K, *et al.*: **Targeting the differential addiction to anti-apoptotic BCL-2 family for cancer therapy.** *Nat Commun.* 2017; **8**: 16078.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
113. Todt F, Cakir Z, Reichenbach F, *et al.*: **The C-terminal helix of Bcl-x_L mediates Bax retrotranslocation from the mitochondria.** *Cell Death Differ.* 2013; **20**(2): 333–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
114. Reyna DE, Garner TP, Lopez A, *et al.*: **Direct Activation of BAX by BTS1A1 Overcomes Apoptosis Resistance in Acute Myeloid Leukemia.** *Cancer Cell.* 2017; **32**(4): 490–505.e10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)
115. Garner TP, Amgalan D, Reyna DE, *et al.*: **Small-molecule allosteric inhibitors of BAX.** *Nat Chem Biol.* 2019; **15**(4): 322–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Faculty Opinions Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Vishva Dixit

Department of Physiological Chemistry, Genentech, South San Francisco, CA, USA

Competing Interests: No competing interests were disclosed.

2. Andrew Gilmore

Wellcome Trust Centre for Cell/Matrix Research, Faculty of Biology, Medicine and Health, University of Manchester, Manchester, UK

Competing Interests: No competing interests were disclosed.

3. Andreas Strasser

The Walter and Eliza Hall Institute of Medical Research, Parkville, Melbourne, VIC, Australia

Competing Interests: Andreas Strasser is an employee of The Walter and Eliza Hall Institute. This Institute receives milestone payments and royalties from the sale of venetoclax. Andreas Strasser also has a collaboration with Servier for the development of MCXL01 inhibitors for cancer therapy.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research