SYSTEMATIC REVIEW

Effects of stretching exercises on human gait: a systematic review and meta-analysis [version 1; peer review: 2 approved with reservations]

Thomas Vialleron ^{1,2}, Arnaud Delafontaine ^{1,2}, Sebastien Ditcharles¹⁻³, Paul Fourcade^{1,2}, Eric Yiou^{1,2}

¹CIAMS, Univ. Paris-Sud., Université Paris-Saclay, Orsay, 91405, France

²CIAMS, Université d'Orléans, Orléans, Orléans, 45067, France

³ENKRE, Saint Maurice, Ile de France, 94410, France

V1 First published: 13 Aug 2020, 9:984
<https://doi.org/10.12688/f1000research.25570.1>
 Latest published: 30 Oct 2020, 9:984
<https://doi.org/10.12688/f1000research.25570.2>

Abstract

Background: Stretching is commonly used in physical therapy as a rehabilitation tool to improve range of motion and motor function. However, is stretching an efficient method to improve gait, and if so, for which patient category?

Methods: A systematic review of randomized and non-randomized controlled trials with meta-analysis was conducted using relevant databases. Every patient category and every type of stretching programs were included without multicomponent programs. Data were meta-analysed where possible. Estimates of effect sizes (reported as standard mean difference (SMD)) with their respective 95% confidence interval (95% CI) were reported for each outcome. The PEDro scale was used for the quality assessment.

Results: Twelve studies were included in the analysis. Stretching improved gait performance as assessed by walking speed and stride length only in a study with a frail elderly population, with small effect sizes (both SMD= 0.49; 95% CI: 0.03, 0.96; PEDro score: 3/10). The total distance and the continuous walking distance of the six-minute walking test were also improved only in a study in an elderly population who had symptomatic peripheral artery disease, with large effect sizes (SMD= 1.56; 95% CI: 0.66, 2.45 and SMD= 3.05; 95% CI: 1.86, 4.23, respectively; PEDro score: 5/10). The results were conflicting in healthy older adults or no benefit was found for most of the performance, spatiotemporal, kinetic and angular related variables. Only one study (PEDro score: 6/10) showed improvements in stance phase duration (SMD=-1.92; 95% CI: -3.04, -0.81), swing phase duration (SMD=1.92; 95% CI: 0.81, 3.04), double support phase duration (SMD= -1.69; 95% CI: -2.76, -0.62) and step length (SMD=1.37; 95% CI: 0.36, 2.38) with large effect sizes.

Conclusions: There is no strong evidence supporting the beneficial

Open Peer Review

Approval Status

	1	2	3
version 2			
(revision)
30 Oct 2020	view		view

version 1			
13 Aug 2020
	view	view	

1. **Fabrice Mégrot** , Centre de Médecine Physique et de Réadaptation pour Enfants Bois-Larris, Croix-rouge française, Lamorlaye, France
 Université de Technologie de Compiègne, Sorbonnes Université, Compiègne, France
2. **Xu Wei** , Shanghai Jiao Tong University School of Medicine, Shanghai, China
Guoning Zhang, Shanghai Jiao Tong University School of Medicine, Shanghai, China
3. **Maria Garcia Escudero**, School of Medicine and Health Sciences, Universidad Católica de Valencia San Vicente Mártir, Valencia, Spain

effect of using stretching to improve gait. Further randomized controlled trials are needed to understand the impact of stretching on human gait.

Keywords

stretching, gait, performance, balance, physical therapy

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Arnaud Delafontaine (arnaud_94150@hotmail.fr)

Author roles: Vialleron T: Conceptualization, Methodology, Writing – Original Draft Preparation; Delafontaine A: Conceptualization, Methodology, Writing – Original Draft Preparation; Ditcharles S: Writing – Review & Editing; Fourcade P: Writing – Review & Editing; Yiu E: Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2020 Vialleron T *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Vialleron T, Delafontaine A, Ditcharles S *et al.* **Effects of stretching exercises on human gait: a systematic review and meta-analysis [version 1; peer review: 2 approved with reservations]** F1000Research 2020, 9:984 <https://doi.org/10.12688/f1000research.25570.1>

First published: 13 Aug 2020, 9:984 <https://doi.org/10.12688/f1000research.25570.1>

Introduction

Gait is a highly complex motor skill that is classically considered as an integrative measure and a predictor of health in older adults (e.g. 1; cf. also 2 and 3 for recent research topics on this matter). The loss of gait or its alteration with pathological conditions are known to be related to mortality, especially in the elderly (e.g. 4,5), stressing the importance of addressing gait disorders in physiotherapy. Gait requires body propulsion and balance control for safe progression, two “subtasks” that require the coordination of multiple skeletal muscles and the integration of sensory information arising from the vestibular, visual and somatosensory systems^{6–8}. As such, gait may expose populations with sensory or motor deficits to the risk of falling with serious consequences for health and autonomy. For these reasons, improving gait is a major aim in rehabilitation for most neurological/orthopaedic disorders, such as stroke or Parkinson’s disease, and for frail older adults. Various therapeutic methods have been used to improve gait, such as resistance training⁹, endurance training¹⁰, balance training¹¹, whole body vibrations (for a complete review, see Fischer *et al.*, 2019¹²), multi-component exercise programs¹³ and stretching¹⁴.

The successful completion of numerous daily life activities is conditioned by the ability to move efficiently through a sufficient range of motion (ROM)¹⁵. Recent studies on gait initiation^{16–18} and seat-to-stand task^{19,20} showed that the experimental restriction of postural chain ROM induced by orthosis wear in young healthy adults led to instability and lower motor performance. It is well established that ROM significantly decreases with aging^{21–26} and more generally with reduced functional demand (e.g. sedentarity, immobilization, disease etc.)¹⁵. Consequently, stretching has become an important part of many sport and rehabilitation programs to maintain or improve ROM, reduce stiffness and promote physical activity. This method has been applied in older adults^{27,28}, patients with stroke²⁹, Parkinson’s disease³⁰, multiple sclerosis³¹, plantar fasciitis³² and spastic paraplegia³³, for example. In sport programs, the influence of stretching on motor performance remains an issue of debate, although recent reviews conclude that maximal muscle performance (e.g. force, power, jump height, reaction time, etc.) is impaired primarily immediately after long durations of stretch (>90 seconds)^{34,35}. To date, no review has collected results on the relationship between stretching and locomotor performance in rehabilitation programs.

Hence, the purpose of this article is to analyse the effects of a stretching program on gait in each patient category by means of a systematic literature review and meta-analysis, comparing the gait outcomes of the intervention groups with the control groups. It will contribute to provide evidence-based practice from scientific data in order to integrate stretching in rehabilitation programs in a reasoned manner.

Methods

Design and literature screening

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) methodology was employed in this systematic review³⁶. A completed PRISMA checklist was submitted to an online repository (*Reporting guidelines*).

PubMed, Science Direct, Springer and Sage databases were used for a comprehensive systematic literature search for articles published prior to 28 April 2020 with no time limit. In addition, a manual search was conducted using the reference list of selected studies. The keywords used for the search strategy in PubMed were: “stretching” AND (gait OR walk). We included only articles published in English or French.

The selection procedure was conducted by two experts in rehabilitation (TV and AD). Disagreements were discussed with a third expert in a group until a mutual consensus was reached. First, a review was performed on all available titles obtained from the literature search with the selected keywords. All relevant or potentially relevant titles were included in the subsequent phase. Then, the abstracts were reviewed with all relevant or potential articles included in the following phase. Finally, full-text articles were reviewed to ensure that only relevant studies were included. In the same way, reference lists of all included articles were reviewed to possibly include articles through cross-referencing.

Inclusion and exclusion criteria

We included randomized controlled trials (RCT) and controlled clinical trials (CCT) published in peer-reviewed journals that aimed to explore the effects of stretching on gait parameters. We included all categories of subjects, all stretching techniques and different durations of treatment since standardized protocols are lacking in the purpose of the present study. Gait could be evaluated by functional tests, electromyographic (EMG) or biomechanical analysis. The following exclusion criteria were used: lack of gait assessment, non-application of muscle stretching, multimodal exercise programs, no control group, case report and review.

Data extraction and main measurements examined

Data were extracted from the selected articles by one of the authors (TV). The extracted data were checked by another author (AD) and disagreements were resolved with a third (EY).

The following data were extracted for each selected article: (1) the names of the authors and the date of publication; (2) the number of subjects involved in the experiment with their characteristics and breakdown in each group; (3) stretching training details (in the following order: number of participants, stretching technique, muscle groups stretched, number of sets, duration of stretch, frequency, protocol duration); (4) control group details; and (5) the main outcomes related to gait with the main results. When information could not be provided, it was indicated by a “?”.

Quality and risk of bias assessment

The PEDro scale was used to assess the risk of bias, and thus the methodological quality of the selected studies³⁷. This scale was chosen for its ability to provide an overview of the external (criterion 1), internal (criteria 2–9) and statistical (criteria 9 and 10) validity of clinical trials. The scale is divided into 11 criteria, but the first is not calculated in the total score. The output of each criterion could be either “yes” (y), “no” (n) or “do not know” (?). A “y” was given a score of one point, while

an “n” or “?” was assigned zero points. Studies with a total score of 5–10/10 ($\geq 50\%$) were considered to be of high quality, and scores of 0–4/10 ($<50\%$) of low quality³⁸. Two evaluators independently assessed the quality of the included studies. In the event of disagreement, a group discussion was held with a third expert to reach a consensus.

Statistical analysis

Estimates of effect sizes (comparing the intervention group and the control group) accompanied with a measure of statistical uncertainty (95% confidence interval [95% CI]) were calculated for each outcome when sufficient data were reported. Estimates of effect sizes were reported by standard mean difference (SMD) and their respective 95% CI. In this way, the magnitude of the overall effect can be quantified as trivial (<0.2), small (0.2–0.49), moderate (0.5–0.79) or large (≥ 0.8)^{39,40}. When data were lacking to calculate estimates of effect sizes, exact p values were reported.

When at least two studies used the same outcome, meta-analysis was performed, comparing the intervention groups with the control groups. When outcomes were identified in only one study, no meta-analysis could be performed but the effect of intervention was still calculated, reporting the estimate of effect size and its 95% confidence interval. Statistical analysis and figures (i.e. forest plot to facilitate the visualization of values) were produced using a random-effect model in Review Manager software (RevMan, v 5.3, Cochrane Collaboration, Oxford UK). A random-effect model was used to take into account heterogeneity between study effects. Statistical heterogeneity was calculated using the I^2 and Cochrane Q statistic tests³⁹. Statistical significance was set at $p < 0.05$.

Level of evidence

The strength of evidence of primary outcomes was established as described by Van Tulder *et al.* 2003⁴¹ based on effect size estimates with a measure of statistical uncertainty (SMD; 95% CI), statistical heterogeneity (I^2) when applicable (multiple studies) and risk of bias (PEDro scale). The level of evidence was considered strong with consistent findings among multiple high-quality RCT (at least two RCT with a PEDro score $\geq 5/10$ that were statistically homogenous: $I^2 p \geq 0.05$). The level of evidence was considered moderate with consistent findings among multiple low-quality RCT and/or CCT (two trials with a PEDro score $< 5/10$ that were statistically homogenous) and/or one high quality RCT. The level of evidence was considered limited when only one low quality RCT and/or CCT was identified. The level of evidence was conflicting when there was inconsistency among multiple trials ($I^2 p < 0.05$).

Results

Included studies

A total of 821 titles were screened in the first search stage, one more was included through cross-referencing, and 671 were excluded because they did not concern our research question. Following exclusion, 150 studies were considered for an abstract review. A further 105 were excluded in this second stage

because they did not meet the inclusion criteria. Finally, 45 full-text articles were assessed for eligibility with 33 not accepted (Figure 1).

Thus, 12 articles were ultimately included in this systematic review. Six studies evaluated the effects of stretching in healthy older adults^{14,42–46}, one in a frail elderly population⁴⁷, one study in an elderly population with stable symptomatic peripheral artery disease⁴⁸, one in stroke patients⁴⁹, one study in adults with limited ankle ROM associated with a history of lower limb overuses injury⁵⁰, one study in healthy adults with limited ankle dorsiflexion range of motion⁵¹ and one in healthy young adults⁵². A summary of the studies selected is provided in Table 1, and their quality assessment is reported in Table 2. The results in different patient categories are reported below.

Results in different patient categories

Healthy older adults

Description of the studies and quality assessment

Six studies examined the effects of stretching on healthy elderly subjects^{14,42–46}. Regarding the characteristics of the subjects, the average sample size was 46.6 ± 33.9 subjects (ranging from 19¹⁴ to 96 subjects⁴⁴) and the mean age was 70.1 ± 3.6 years (ranging from 65.40¹³ to 75.30 years¹⁴). Regarding the characteristics of the training programs, the average training duration was 8.6 ± 2.7 weeks (ranging from 4⁴³ to 12 weeks⁴⁵), with an average frequency of 8.3 ± 6.2 sessions per week (ranging from 2⁴⁵ to 14 sessions^{42,44,46}). The average number of sets per session was 4.5 ± 2.8 sets (ranging from 2⁴⁶ to 10 sets¹⁴), with an average stretching time of 45.0 ± 18.9 seconds (ranging from 15¹⁴ to 60 seconds^{43,45,46}). Static stretching was provided in all studies. The muscle groups stretched were the hip flexors^{42–44,46}, ankle plantar flexors^{14,42,43,45}, ankle dorsiflexors⁴⁵, hip extensors⁴³, knee extensors and flexors⁴⁵. There was great heterogeneity in gait outcomes. Angular variables during gait included peak hip extension^{42,44,46}, ankle plantar flexion during gait⁴⁴, ankle range of motion during gait⁴³, anterior pelvis tilt^{43,46}, knee range of motion⁴³, pelvic rotation⁴³, lateral pelvic tilt⁴³ and hip range of motion⁴³. Spatiotemporal variables were: gait speed^{42,43,46}, stance and swing durations⁴³, double support phases⁴³, step length⁴³ and stride length^{43,46}. Kinetic variables were hip torque⁴⁴ and ankle plantar flexion power⁴⁴. Finally, two functional tests were used: the 10-meter walk test (10MWT)¹⁴ and the 6-minute walk test (6MWT)⁴⁵. Regarding the quality of the studies, the average PEDro score was 4.6 ± 1.6 and one study was identified as a non-randomized trial⁴⁵. The range of score varied from 3^{45,46} to 7¹⁴.

Meta-analyses

Four meta-analysis were conducted for the following outcomes (Figure 2): gait speed, stride length, hip extension during gait and anterior pelvic tilt.

Gait speed: For gait speed (Figure 2A), two studies were included in the meta-analysis^{42,43}. One study was excluded because intervention and control groups were not similar at baseline⁴⁶. Statistical analysis showed no significant difference between

Figure 1. PRISMA flow chart of study selection process.

groups (SMD= 0.45; 95% CI: -1.15, 2.06), with heterogeneous results ($I^2=86\%$, $p=0.007$). Thus, the level of evidence was conflicting.

Stride length: For stride length (Figure 2B), two studies were included in the meta-analysis^{42,46}. Statistical analysis showed no significant difference between groups (SMD= 0.22; 95% CI: -0.44, 0.88), with consistent results ($I^2=59\%$, $p=0.12$). Only one study was of high quality⁴², thus a moderate level of evidence supports the lack of beneficial effect of stretching to improve stride length in the elderly.

Hip extension: For hip extension during gait (kinematic data) (Figure 2C), three studies were included in the meta-analysis^{42,44,46}. Statistical analysis showed no significant difference between groups (SMD= 0.20; 95% CI: -0.06, 0.47), with consistent results ($I^2=0\%$, $p=0.99$). Two studies were of high quality^{42,44}, thus a strong level of evidence supports the lack of beneficial effect of stretching to improve hip range of motion during gait in the elderly.

Anterior pelvic tilt: For anterior pelvic tilt (Figure 2D), three studies were included in the meta-analysis^{42,44,46}. Statistical analysis showed no significant difference between groups

(SMD= -0.70; 95% CI: -1.60, 0.21), with heterogeneous results ($I^2=87\%$, $p<0.01$). Thus, the level of evidence was conflicting.

Effects of interventions in other outcomes

For the outcomes below, no meta-analysis could be performed because only one study was identified. Nevertheless, for each outcome, effect size estimates with a measure of statistical uncertainty (95% CI) were provided.

Angular variables during gait initiation: The study of Christiansen *et al.* (2008)⁴² showed no significant difference between stretching and control groups for ankle dorsiflexion during gait (SMD=0.29; 95% CI: -0.36, 0.94) with a moderate level of confidence (PEDro score: 5/10). The study of Kerrigan *et al.* (2003)⁴⁴ showed no significant difference between groups for ankle plantar flexion (SMD=-0.05; 95% CI: -0.45, 0.35), with a moderate level of confidence (PEDro score: 6/10). The study of Cristopoliski *et al.* (2009)⁴³ showed no significant difference between groups for lateral pelvic tilt (SMD= 0.93; 95% CI: -0.02, 1.88) and knee range of motion (SMD= 0.23; 95% CI: -0.67, 1.12), with a moderate level of confidence (PEDro score: 6/10).

Table 1. Summary of the included studies.

Studies	Population	Stretching group	Control group	Outcomes and main results
Kerrigan <i>et al.</i> , 2003 ⁴⁴	96 healthy older adults, ? (≥65 years)	n= 47, static stretching, hip flexors, 4 sets, 30 seconds, twice daily, 10 weeks	n=49, static shoulder deltoid-stretching, same protocol	No significant difference between groups for hip extension (SMD= 0.22; 95% CI: -0.18, 0.62), hip torque (SMD= 0.35; 95% CI: -0.06, 0.75), anterior pelvic tilt (SMD= -0.35; 95% CI: -0.76, 0.05), ankle plantar flexion ROM (SMD= -0.05, 95% CI: -0.45, 0.35), ankle plantar flexion power (SMD= 0.00; 95% CI: -0.40, 0.40), hip extension (SMD= 0.22; 95% CI: -0.19, 0.62) and hip torque (SMD= 0.35; 95% CI: -0.06, 0.75)
Gajdosik <i>et al.</i> , 2005 ¹⁴	19 community dwelling older women, ? (65–89 years)	n=10, static stretching, ankle plantar flexors, 10 sets, 15 seconds, 3 times per week, 8 weeks	n=9, no exercise.	No significant difference between groups for 10MWT (SMD= -0.76; 95% CI: -1.70, 0.18)
Christiansen, 2008 ⁴²	40 healthy older adults, 72.10±4.70 years	N= 20, static stretching, hip flexors, 3 sets, 45 seconds, twice daily, 8 weeks	n=20, maintain their current level of physical activity	No significant difference between groups for gait speed (SMD= -0.32; 95% CI: -0.97, 0.33), hip extension (SMD= 0.22; 95% CI: -0.43, 0.86), stride length (SMD= -0.14; 95% CI: -0.79, 0.50), ankle dorsiflexion (SMD= 0.29; 95% CI: -0.36, 0.94)
Cristopoliski <i>et al.</i> , 2009 ⁴³	20 healthy elderly women, 65.90±4.20 years	n=12, static stretching, hip flexors and extensors, ankle plantar flexors, 4 sets, 60 seconds, 3 sessions per week, 12 sessions	n=8, no specific activity in this period	Significant improvement in favor of stretching group for gait speed (SMD= 1.32; 95% CI: 0.32, 2.32), anterior pelvic tilt (SMD= -2.52; 95% CI: -3.77, -1.27), stand phase duration (SMD= -1.92; 95% CI: -3.04, -0.81), swing phase duration (SMD= 1.92; 95% CI: 0.81, 3.04), double support phase duration (SMD= -1.69; 95% CI: -2.76, -0.62), step length (SMD= 1.37; 95% CI: 0.36, 2.38) and pelvic rotation (SMD= 1.37; 95% CI: 0.36, 2.38) No significant difference between groups for cycle duration (SMD= -0.24; 95% CI: -1.14, 0.66), heel-contact velocity (SMD= -0.46; 95% CI: -1.37, 0.45), toe clearance (SMD= 0.91; 95% CI: -0.04, 1.86), lateral pelvic tilt (SMD= 0.93; 95% CI: -0.02, 1.88) and knee range of motion (SMD= 0.23; 95% CI: -0.67, 1.12)
Watt <i>et al.</i> , 2011 ⁴⁶	82 healthy elderly subjects, 72.6±6 years	n= 43, static stretching, hip flexors, 2 sets, 60 seconds, 2 sessions daily stretching, 10 weeks	N= 39, shoulder abductor static stretching, same protocol	Significant improvement in favor of stretching group for gait speed (SMD= 0.47; 95% CI: 0.03, 0.91) No significant difference between groups for hip extension (SMD= 0.18; 95% CI: -0.25, 0.62), anterior pelvic tilt (SMD= 0.07; 95% CI: -0.36, 0.51), stride length (SMD= 0.54; 95% CI: -0.01, 1.08)
Locks <i>et al.</i> , 2012 ⁴⁵	23 healthy older individuals, 67.5±2.12 years	n=10, static stretching, knee extensors, ankle dorsiflexor, knee flexors, ankle plantar flexors, 4 sets, 60 seconds, twice a week, 12 weeks	n=13 a one-hour seminar on healthy living every four weeks and did not perform any physical or therapeutic exercise.	No significant difference between groups for 6MWT (SMD= -0.04; 95% CI: -0.86, 0.79)
Watt <i>et al.</i> , 2011 ⁴⁷	74 frail elderly subjects, 77.00±8.00 years	n=33, static stretching, hip flexors, 2 sets, 60 seconds, 2 sessions per day, 10 weeks	n=41, shoulder abductor stretching program, same protocol	No significant difference between groups in peak hip extension, (SMD= 0.22; 95% CI: -0.24, 0.68), anterior pelvic tilt (SMD= -0.05; 95% CI: -0.51, 0.41) and cadence (SMD= 0.13; 95% CI: -0.33, 0.59) Significant improvements in favor of the stretching group in walking speed and stride length (both SMD= 0.49; 95% CI: 0.03, 0.96)

Studies	Population	Stretching group	Control group	Outcomes and main results
Hotta <i>et al.</i> , 2019 ⁴⁸	13 elderly patients with symptomatic peripheral artery disease, ?	n=13, static stretching, ankle plantar flexor stretching, 1 set, 30 minutes, 5 sessions per week, 4 weeks	n=13, no stretching intervention (cross-over intervention)	Significant improvements in favor of the stretching group for both total walking distance and continuous walking distance with large effect sizes (SMD= 1.56; 95% CI: 0.66, 2.45 and SMD= 3.05; 95% CI: 1.86, 4.23 respectively)
Kim <i>et al.</i> , 2013 ⁴⁹	24 patients with stroke, 53.30±3.16 years	n=12, static stretching, ankle plantar flexors, 1 set, 20 minutes, 4 times a week, 4 times a week, 6 weeks	n=12, conventional physical therapy as in the stretching group	No significant difference between groups in sway of the center of pressure (SMD=0.75; 95% CI: -0.09, 1.58)
Johanson <i>et al.</i> , 2006 ⁵⁰	19 adults with limited passive ankle-dorsiflexion range of motion (less than 8 degrees) and a history of lower limb overuse injury, 30.30± 9.80 years	n=11, static stretching, ankle plantar flexors, 5 sets, 30 seconds, 2 times daily, 3 weeks	n=8, continue all of their usual activities	No significant difference between groups in ankle dorsiflexion during gait in both right and left ankle (SMD= 0.50; 95% CI: -0.42, 1.43 and SMD= 0.41; 95% CI: -0.52, 1.33 respectively) and for time-to-heel-off during the stance phase of gait in both right and left ankle (SMD= -0.50; 95% CI: -1.43, 0.43 and SMD= -0.48; 95% CI: -1.41, 0.45 respectively)
Johanson <i>et al.</i> , 2009 ⁵¹	16 healthy adults with limited passive ankle-dorsiflexion range of motion (less than 5 degrees), 27.40±8.20 years	n=8, static stretching, ankle plantar flexors, 4 sets, 30 seconds, 2 times daily, 3 weeks	n=8, no physical activity or stretching programs involving the lower extremities for 3 weeks	No significant difference between groups in ankle dorsiflexion (SMD= 0.53; 95% CI: -0.48, 1.53), maximum knee extension (SMD= -0.07; 95% CI: -1.05, 0.91), medial and lateral gastrocnemius activities (SMD= 0.37; 95% CI: -0.62, 1.36 and SMD= 0.00; 95% CI: -0.98, 0.98 respectively)
Godges <i>et al.</i> , 1993 ⁵²	16 healthy, athletic, male college students, 21.00±1.00 years	n=9, static stretching, hip flexors, 3 sets, 2 minutes, 2 sessions per week, 3 weeks	n=7, continue their current activity levels	No significant difference between groups in gait economy (SMD= 0.83; 95% CI: -0.21, 1.87)

SMD: standard mean difference, CI: confidence interval, 10MWT: 10-meter walk test, 6MWT: 6-minute walk test, ROM: range of motion, ?: information not provided.

Table 2. Quality assessment of the included studies.

Study	Eligibility criteria specified	Random allocation	Concealed allocation	Groups similar at baseline	Participant blinding	Therapist blinding	Assessor blinding	<15% dropouts	Intention-to-treat analysis	Between-group difference reported	Point estimate and variability reported	Total score	Patient category
Kerrigan <i>et al.</i> , 2003	y	y	n	y	n	n	y	y	n	y	y	6	
Gajdosik <i>et al.</i> , 2005	y	y	y	y	n	n	n	y	y	y	y	7	
Christiansen, 2008	y	y	n	y	n	n	n	y	n	y	y	5	Healthy older adults
Cristopoloski <i>et al.</i> , 2009	y	y	n	y	n	n	n	y	y	y	y	6	
Watt <i>et al.</i> , 2011	y	y	n	n	n	n	y	n	n	n	y	3	
Locks <i>et al.</i> , 2012	y	n	n	y	n	n	n	n	n	y	y	3	
Watt <i>et al.</i> , 2011	y	y	n	?	n	n	y	n	n	n	y	3	Frail older adults
Hotta <i>et al.</i> , 2009	y	y	n	y	n	n	n	y	n	y	y	5	Peripheral artery disease
Kim <i>et al.</i> , 2013	n	n	n	y	n	n	n	n	n	y	y	3	Stroke
Johanson <i>et al.</i> , 2006	y	y	n	n	n	n	n	y	y	y	y	5	Lower limb overuse injury
Johanson <i>et al.</i> , 2009	y	y	n	y	n	n	n	y	y	y	y	6	Limited ankle ROM
Godges <i>et al.</i> , 1993	y	y	n	?	n	n	n	y	y	y	y	5	Healthy adults

n: criterion not fulfilled; y: criterion fulfilled; ?: criterion not mentioned; total score: each item (except the first) contributes 1 point to the total score, yielding a PEDro scale score that can range from 0 to 10.

Figure 2. Comparisons between intervention and control groups for gait speed (A), stride length (B), hip extension (C) and anterior pelvic tilt (D) in healthy older adults.

Kinetic variables: The study of Kerrigan *et al.* (2003)⁴⁴ showed no significant difference between groups for hip torque (SMD= 0.35; 95% CI: -0.06, 0.75) and ankle plantar flexion power (SMD=0.00; 95% CI: -0.40, 0.40), with a moderate level of confidence (PEDro score: 6/10).

Spatiotemporal variables: The study of Cristopoliski *et al.* (2009)⁴³ showed no significant difference between groups for cycle duration (SMD= -0.24; 95% CI: -1.14, 0.66), heel contact velocity (SMD= -0.46; 95% CI: -1.37, 0.45) and toe clearance (SMD= 0.91; 95% CI: -0.04, 1.86). However, the study showed significant decreases with large effect sizes in stance phase duration (SMD=-1.92; 95% CI: -3.04, -0.81), double support phase duration (SMD= -1.69; 95% CI: -2.76, -0.62) in favour of the stretching group as compared to the control group. Additionally, the authors found significant increases with large effect sizes of swing phase duration (SMD=1.92; 95% CI: 0.81, 3.04) and step length (SMD=1.37; 95% CI: 0.36, 2.38)

in favour of the stretching group as compared to the control group. The study obtained a PEDro score of 6/10, thus, the level of evidence for these outcomes was moderate.

Functional tests: The study of Gajdosik *et al.* (2005)¹⁴ showed no significant difference between groups for the 10MWT (SMD= -0.76; 95% CI= -1.70, 0.18), with a moderate level of confidence (PEDro score: 7/10). The study of Locks *et al.* (2012)⁴⁵ showed no significant improvement of the 6MWT in favour of the stretching group as compared to the control group (SMD= -0.04; 95% CI:-0.86, 0.79) with a limited level of confidence (low quality CCT with a PEDro score of 3/10).

Frail elderly

Description of the study and quality assessment

The study of Watt *et al.* 2011 examined the effects of stretching on frail elderly subjects⁴⁷. Regarding the characteristics of the subjects, 74 subjects were included, and the mean age

was 77.0 ± 8.0 years. Regarding the characteristics of the training programs, the stretching program lasted ten weeks, with a frequency of 14 sessions per week (two sessions per day). Participants performed two sets per session, holding the stretch for 60 seconds (static stretching), alternating the right and left limb (four minutes in total). The muscle group stretched was the hip flexors. The outcomes were cadence (steps/minute), walking speed (meters/second), stride length (meters) peak hip extension (degree) and peak anterior pelvic tilt (degree). Regarding the quality assessment, the study was identified as RCT and had an average PEDro score of 3 (low level of evidence).

Effects of intervention

The study of Watt *et al.* (2011) showed no significant difference between groups in angular variables, i.e. peak hip extension and anterior pelvic tilt (SMD= 0.22; 95% CI: -0.24, 0.68 and SMD= -0.05; 95% CI: -0.51, 0.41 respectively). There was also no significant difference for cadence (SMD= 0.13; 95% CI: -0.33, 0.59). However, the study showed significant improvements in favour of the stretching group with small effect sizes in some performance-related variables, i.e. walking speed and stride length (both SMD= 0.49; 95% CI: 0.03, 0.96).

Elderly with symptomatic peripheral artery disease

Description of the study and quality assessment

The study of Hotta *et al.* (2019) examined the effects of stretching in elderly with symptomatic peripheral artery disease⁴⁸. Regarding the characteristics of the subjects, 13 subjects were included and the mean age was not mentioned. Regarding the characteristics of the training programs, the stretching program lasted four weeks, with a frequency of five sessions per week. Participants performed one set daily, holding the stretch for 30 minutes (static stretching with splints). The muscle group stretched was ankle plantar flexors. The gait outcome was 6MWT. Regarding the quality assessment, the study was identified as RCT and had an average PEDro score of 5 (moderate level of evidence).

Effects of intervention

The study of Hotta *et al.* (2019) showed significant improvements in favour of the stretching group for both total walking distance and continuous walking distance with large effect sizes (SMD= 1.56; 95% CI: 0.66, 2.45 and SMD= 3.05; 95% CI: 1.86, 4.23 respectively).

Stroke

Description of the study and quality assessment

The study of Kim *et al.* (2013) examined the effects of stretching on stroke patients⁴⁹. Only a static muscle stretching training group and control group were included in the analysis. Regarding the characteristics of the subjects, 24 subjects were included, and the mean age was 53.3 ± 3.1 years. Regarding the characteristics of the training programs, the stretching program lasted six weeks, with a frequency of four sessions per week. Participants performed one set per session, holding the stretch for 20 minutes (static stretching). The muscle group stretched was ankle plantar flexors. The outcome was the sway of the centre of pressure during the stance phase.

Regarding the quality assessment, the study was identified as CCT and had an average PEDro score of 3 (low level of evidence).

Effects of intervention

The study of Kim *et al.* (2013) showed no significant difference between groups in the sway of the centre of pressure (SMD=0.75; 95% CI: -0.09, 1.58).

Young adults with limited ankle range of motion and a history of lower limb overuse injury

Description of the study and quality assessment

The study of Johanson *et al.* (2006) examined the effects of stretching on healthy adults with limited passive ankle-dorsiflexion range of motion (less than eight degrees) and a history of lower limb overuse injury⁵⁰. Regarding the characteristics of the subjects, 19 subjects were included and the mean age was 30.3 ± 9.8 years. Regarding the characteristics of the training programs, the stretching program lasted three weeks, with a frequency of two sessions per day. Participants performed five sets per session, holding the stretch for 30 seconds (static stretching). The muscle group stretched was ankle plantar flexors. The outcomes were ankle dorsiflexion and time-to-heel-off during the stance phase of gait. Regarding the quality assessment, the study was identified as RCT and had an average PEDro score of 5 (moderate level of evidence).

Effects of intervention

The study of Johanson *et al.* (2006) showed no significant difference between groups in ankle dorsiflexion during gait in both the right and left ankle (SMD= 0.50; 95% CI: -0.42, 1.43 and SMD= 0.41; 95% CI: -0.52, 1.33 respectively). There was also no significant difference between groups for time-to-heel-off during the stance phase of gait in both the right and left ankle (SMD= -0.50; 95% CI: -1.43, 0.43 and SMD= -0.48; 95% CI: -1.41, 0.45 respectively).

Young adults with limited ankle range of motion

Description of the study and quality assessment

The study of Johanson *et al.* (2009) examined the effects of stretching on young adults with limited passive ankle-dorsiflexion range of motion (less than five degrees)⁵¹. Regarding the characteristics of the subjects, 16 subjects were included, and the mean age was 27.4 ± 8.2 years. The characteristics of the training programs were the same as described above⁵⁰. The muscle group stretched was the ankle plantar flexors. The outcomes were maximum ankle dorsiflexion, maximum knee extension and EMG amplitude of the gastrocnemius during the stance phase of gait. Regarding the quality assessment, the study was identified as RCT and had an average PEDro score of 6 (moderate level of evidence).

Effects of intervention

The study of Johanson *et al.* (2009) showed no significant difference between groups in angular variables during gait, i.e. maximum ankle dorsiflexion and maximum knee extension (SMD= 0.53; 95% CI: -0.48, 1.53 and SMD= -0.07; 95% CI: -1.05, 0.91 respectively). There was also no significant

difference between groups for EMG variables, i.e. medial and lateral gastrocnemius activity (SMD= 0.37; 95% CI: -0.62, 1.36 and SMD= 0.00; 95% CI=: -0.98, 0.98 respectively).

Healthy young adults

Description of the study and quality assessment

The study of Godges *et al.* (1993) examined the effects of stretching on healthy young adults⁵². Only a static hip extension stretching group and control group were included in the analysis. Regarding the characteristics of the subjects, 16 subjects were included, and the mean age was 21.0±1.0 years. Regarding the characteristics of the training programs, the stretching program lasted three weeks, with a frequency of two sessions per week. Participants performed three sets per session, holding the stretch for two minutes (static stretching). The muscle group stretched was the hip flexors. The outcome was walking economy (ml/kg/min). Regarding the quality assessment, the study was identified as RCT and had an average PEDro score of 5 (moderate level of evidence).

Effects of intervention

The study of Godges *et al.* (1993) showed no significant difference between groups in gait economy in terms of oxygen consumption (SMD= 0.83; 95% CI: -0.21, 1.87).

Discussion

The aim of this systematic review was to determine the effects of a stretching program on human gait by means of a systematic literature review and meta-analysis. Twelve studies were identified in six different patient categories. Statistical analyses showed no strong level of evidence supporting the beneficial effect of a stretching program to improve any gait outcome. The major issue in conducting meta-analyses and establishing strong level of evidences was the great heterogeneity in gait variables. The results obtained in the different patient categories are discussed in detail below.

Healthy older adults

The healthy older adult population was the most studied. Two muscle groups were systematically stretched in the six identified studies: hip flexors^{42–44,46,47} and ankle plantar flexors^{14,42,43}. Hip flexor stiffness, associated with reduced hip extension during gait has been demonstrated in the elderly and may alter gait^{53,54}. In the same way, decreased calf muscle length associated with restricted dorsiflexion range of motion is well documented in older adults^{26,55,56}. A decreased ankle dorsiflexion ROM has been correlated with poorer balance test scores in the elderly⁵⁷ and may contribute to an increased risk of falls⁵⁸. All the studies included in the present analysis showed that specific stretching programs were efficient to improve passive range of motion of the targeted joints, but results are more heterogeneous regarding gait performance and dynamic ROM. This led to inconsistency in the results or the impossibility to conclude with a strong level of evidence that a stretching program improves gait in healthy older adults. Moreover, when improvement in ROM or gait performance occurred, it was not associated with a significant increase in dynamic hip extension or ankle dorsiflexion. Only trends toward increased

dynamic ROM after stretching interventions were observed^{42,44,46}. This observation was consistent in young adults.

Stroke patients

In stroke patients, ankle plantar flexor stretching has been successfully used to improve ankle stiffness^{59–62}. Decreased plantar flexors stiffness may have a beneficial effect on postural control during gait because *triceps surae* is known to play an important role during gait^{63–65} and an increase in muscle stiffness might alter synergistic muscle activities during human gait. However, only one non-randomized study⁴⁹ was identified and included in the current systematic review. Other studies that used stretching in multicomponent programs^{66–68} or in control groups^{69,70} were identified but excluded because of the addition of resistance training or the lack of a control group. Nevertheless, it should be noted that some studies showed improvements between pre- and post-stretching conditions. Forrester *et al.* (2014) showed that both robotic ankle mobilizations and manual ankle stretching improved gait velocity in stroke patients at hospital discharge compared to baseline⁶⁹. Similarly, Park *et al.* (2018) showed that both static ankle stretching and ankle mobilizations improved gait speed after four weeks of treatment compared to baseline⁷⁰. Other authors showed that one week of immobilization in dorsiflexed position (casting) followed by one week of plantar flexor stretching and gait training improved gait performances in 10MWT and 6MWT⁶⁶. Hence, these encouraging results suggest that further randomized controlled trials of good quality are needed to explore the ability of ankle stretching to improve gait parameters in stroke or in other neurological diseases exposing patients to joint stiffness, e.g. Parkinson's disease⁷¹.

Young adults

In healthy adults, the interest of practicing stretching to improve gait seems limited as they are assumed to have sufficient mobility for walking. Moreover, the included study involved athletic males⁵², a population that is known to be more flexible than inactive persons⁷². Even in young adults with limited ankle ROM, stretching did not improve dynamic dorsiflexion during gait^{50,51}. Stretching programs in apparently healthy adults should be more indicated after a prolonged period of reduced functional demand (e.g. immobilization, sedentarity), when ROM is insufficient to practice a specific activity or when high levels of flexibility are required for sport performance (e.g. gymnastics or dance) and in sports that involve stretch-shortening cycles (e.g. basketball, volleyball)¹⁵.

Conclusion

Twelve studies were identified, involving a total of 442 subjects. Despite some improvements in isolated studies, statistical analyses showed no strong level of evidence supporting the beneficial effect of using stretching alone to improve gait outcomes in rehabilitation programs. The major obstacle in conducting meta-analyses and establishing strong levels of evidence were the great heterogeneity in gait variables and the low quality of the included studies. Because the effects of stretching are not clear, further randomized controlled trials of good

quality are needed to understand the impact of stretching on human gait. Currently, stretching is more recommended to maintain and improve ROM rather than improve gait parameters and should be integrated in multicomponent programs.

Data availability

Underlying data

All data underlying the results are available as part of the article and no additional source data are required.

Reporting guidelines

Harvard Dataverse: PRISMA checklist and PRISMA flow diagram for 'Effects of stretching exercises on human gait: a systematic review and meta-analysis', <https://doi.org/10.7910/DVN/N8ZXNB>⁷³.

Data are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

References

- LeBrasseur NK: **Gait as an Integrative Measure and Predictor of Health Across Species.** *J Gerontol A Biol Sci Med Sci.* 2019; **74**(9): 1411–1412.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Blumen HM, Cavallari P, Mourey F, et al.: **Editorial: Adaptive Gait and Postural Control: from Physiological to Pathological Mechanisms, Towards Prevention and Rehabilitation.** *Front Aging Neurosci.* 2020; **12**: 45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Yiou E, Hamaoui A, Allali G: **Editorial: The Contribution of Postural Adjustments to Body Balance and Motor Performance.** *Front Hum Neurosci.* 2018; **12**: 487.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- White DK, Neogi T, Nevitt MC, et al.: **Trajectories of Gait Speed Predict Mortality in Well-Functioning Older Adults: The Health, Aging and Body Composition Study.** *J Gerontol A Biol Sci Med Sci.* 2013; **68**(4): 456–464.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rodríguez-Molinero A, Herrero-Larrea A, Miñarro A, et al.: **The spatial parameters of gait and their association with falls, functional decline and death in older adults: a prospective study.** *Sci Rep.* 2019; **9**(1): 8813.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Allum JH, Bloem BR, Carpenter MG, et al.: **Proprioceptive control of posture: a review of new concepts.** *Gait Posture.* 1998; **8**(3): 214–242.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Horak FB: **Postural orientation and equilibrium: what do we need to know about neural control of balance to prevent falls?** *Age Ageing.* 2006; **35** Suppl 2: ii7–ii11.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mirelman A, Shema S, Maidan I, et al.: **Gait.** *Handb Clin Neurol.* 2018; **159**: 119–134.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Arcolin I, Pisano F, Delconte C, et al.: **Intensive cycle ergometer training improves gait speed and endurance in patients with Parkinson's disease: A comparison with treadmill training.** *Restor Neurol Neurosci.* 2015; **34**(1): 125–138.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lustosa LP, Silva JP, Coelho FM, et al.: **Impact of resistance exercise program on functional capacity and muscular strength of knee extensor in pre-frail community-dwelling older women: a randomized crossover trial.** *Rev Bras Fisioter.* 2011; **15**(4): 318–324.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Halvarsson A, Franzén E, Ståhle A: **Balance training with multi-task exercises improves fall-related self-efficacy, gait, balance performance and physical function in older adults with osteoporosis: a randomized controlled trial.** *Clin Rehabil.* 2015; **29**(4): 365–375.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fischer M, Vialleron T, Laffaye G, et al.: **Long-Term Effects of Whole-Body Vibration on Human Gait: A Systematic Review and Meta-Analysis.** *Front Neurol.* 2019; **10**: 627.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Freiberger E, Häberle L, Spiriduso WW, et al.: **Long-term effects of three multicomponent exercise interventions on physical performance and fall-related psychological outcomes in community-dwelling older adults: a randomized controlled trial.** *J Am Geriatr Soc.* 2012; **60**(3): 437–446.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gajdosik RL, Vander Linden DW, McNair PJ, et al.: **Effects of an eight-week stretching program on the passive-elastic properties and function of the calf muscles of older women.** *Clin Biomech (Bristol, Avon).* 2005; **20**(9): 973–983.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Portero P, McNair P: **Les étirements musculo-tendineux: des données scientifiques à une pratique raisonnée.** *Kinésithérapie, la Revue.* 2015; **15**(164–165): 32–40.
[Publisher Full Text](#)
- Delafontaine A, Gagey O, Colnaghi S, et al.: **Rigid Ankle Foot Orthosis Deteriorates Mediolateral Balance Control and Vertical Braking during Gait Initiation.** *Front Hum Neurosci.* 2017; **11**: 214.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Delafontaine A, Fourcade P, Honeine JL, et al.: **Postural adaptations to unilateral knee joint hypomobility induced by orthosis wear during gait initiation.** *Sci Rep.* 2018; **8**(1): 830.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Delafontaine A, Honeine JL, Do MC, et al.: **Comparative gait initiation kinematics between simulated unilateral and bilateral ankle hypomobility: Does bilateral constraint improve speed performance?** *Neurosci Lett.* 2015; **603**: 55–59.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hamaoui A, Alami-Rodrigues C: **Influence of Cervical Spine Mobility on the Focal and Postural Components of the Sit-to-Stand Task.** *Front Hum Neurosci.* 2017; **11**: 129.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Alami-Rodrigues C, Hamaoui A: **Effect of three different lumbar splints on posturokinetic capacity when performing the sit-to-stand task.** *Ann Phys Rehabil Med.* 2017; **60**(6): 406–409.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Grimston SK, Nigg BM, Hanley DA, et al.: **Differences in ankle joint complex range of motion as a function of age.** *Foot Ankle.* 1993; **14**(4): 215–222.
[PubMed Abstract](#) | [Publisher Full Text](#)
- James B, Parker AW: **Active and passive mobility of lower limb joints in elderly men and women.** *Am J Phys Med Rehabil.* 1989; **68**(4): 162–167.
[PubMed Abstract](#) | [Publisher Full Text](#)
- McKay MJ, Baldwin JN, Ferreira P, et al.: **Normative reference values for strength and flexibility of 1,000 children and adults.** *Neurology.* 2017; **88**(1): 36–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Roach KE, Miles TP: **Normal hip and knee active range of motion: the relationship to age.** *Phys Ther.* 1991; **71**(9): 656–665.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Soucie JM, Wang C, Forsyth A, et al.: **Range of motion measurements: reference values and a database for comparison studies.** *Haemophilia.* 2011; **17**(3): 500–507.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Vandervoort AA, Chesworth BM, Cunningham DA, et al.: **Age and sex effects on mobility of the human ankle.** *J Gerontol.* 1992; **47**(1): M17–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lee PG, Jackson EA, Richardson CR: **Exercise Prescriptions in Older Adults.** *Am Fam Physician.* 2017; **95**(7): 425–432.
[PubMed Abstract](#)
- Mora JC, Valencia WM: **Exercise and Older Adults.** *Clin Geriatr Med.* 2018; **34**(1): 145–162.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Beyaert C, Vasa R, Frykberg GE: **Gait post-stroke: Pathophysiology and rehabilitation strategies.** *Neurophysiol Clin.* 2015; **45**(4–5): 335–355.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Abbruzzese G, Marchese R, Avanzino L, et al.: **Rehabilitation for Parkinson's disease: Current outlook and future challenges.** *Parkinsonism Relat Disord.* 2016; **22** Suppl 1: S60–S64.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bisht B, Darling WG, White EC, et al.: **Effects of a multimodal intervention on gait and balance of subjects with progressive multiple sclerosis: a**

- prospective longitudinal pilot study.** *Degener Neurol Neuromuscul Dis.* 2017; **7**: 79–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Thong-On S, Bovonsunthonchai S, Vachalathiti R, *et al.*: **Effects of Strengthening and Stretching Exercises on the Temporospacial Gait Parameters in Patients With Plantar Fasciitis: A Randomized Controlled Trial.** *Ann Rehabil Med.* 2019; **43**(6): 662–676.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 33. van Lith BJH, den Boer J, van de Warrenburg BPC, *et al.*: **Functional effects of botulinum toxin type A in the hip adductors and subsequent stretching in patients with hereditary spastic paraplegia.** *J Rehabil Med.* 2019; **51**(6): 434–441.
[PubMed Abstract](#) | [Publisher Full Text](#)
 34. Behm DG, Chaouachi A: **A review of the acute effects of static and dynamic stretching on performance.** *Eur J Appl Physiol.* 2011; **111**(11): 2633–2651.
[PubMed Abstract](#) | [Publisher Full Text](#)
 35. Kay AD, Blazevich AJ: **Effect of acute static stretch on maximal muscle performance: a systematic review.** *Med Sci Sports Exerc.* 2012; **44**(1): 154–164.
[PubMed Abstract](#) | [Publisher Full Text](#)
 36. Moher D, Liberati A, Tetzlaff J, *et al.*: **Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement.** *PLoS Med.* 2009; **6**(7): e1000097.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 37. de Morton NA: **The PEDro scale is a valid measure of the methodological quality of clinical trials: a demographic study.** *Aust J Physiother.* 2009; **55**(2): 129–133.
[PubMed Abstract](#) | [Publisher Full Text](#)
 38. Lindberg J, Carlsson J: **The effects of whole-body vibration training on gait and walking ability - A systematic review comparing two quality indexes.** *Physiother Theory Pract.* 2012; **28**(7): 485–98.
[PubMed Abstract](#) | [Publisher Full Text](#)
 39. Higgins J, Green S: **Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [Updated March 2011].** The Cochrane Collaboration, 2011.
[Reference Source](#)
 40. Cohen J: **Statistical Power Analysis for the Behavioral Sciences.** 2nd ed. L. Erlbaum Associates; 1988.
[Publisher Full Text](#)
 41. van Tulder M, Furlan A, Bombardier C, *et al.*: **Updated Method Guidelines for Systematic Reviews in the Cochrane Collaboration Back Review Group.** *Spine (Phila Pa 1976).* 2003; **28**(12): 1290–1299.
[PubMed Abstract](#) | [Publisher Full Text](#)
 42. Christiansen CL: **The effects of hip and ankle stretching on gait function of older people.** *Arch Phys Med Rehabil.* 2008; **89**(8): 1421–1428.
[PubMed Abstract](#) | [Publisher Full Text](#)
 43. Cristopoliski F, Barela JA, Leite N, *et al.*: **Stretching Exercise Program Improves Gait in the Elderly.** *Gerontology.* 2009; **55**(6): 614–620.
[PubMed Abstract](#) | [Publisher Full Text](#)
 44. Kerrigan DC, Xenopoulos-Oddsson A, Sullivan MJ, *et al.*: **Effect of a hip flexor-stretching program on gait in the elderly.** *Arch Phys Med Rehabil.* 2003; **84**(1): 1–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 45. Locks RR, Costa TC, Koppe S, *et al.*: **Effects of strength and flexibility training on functional performance of healthy older people.** *Rev Bras Fisioter.* 2012; **16**(3): 184–190.
[PubMed Abstract](#) | [Publisher Full Text](#)
 46. Watt JR, Jackson K, Franz JR, *et al.*: **Effect of a Supervised Hip Flexor Stretching Program on Gait in Elderly Individuals.** *PM R.* 2011; **3**(4): 324–329.
[PubMed Abstract](#) | [Publisher Full Text](#)
 47. Watt JR, Jackson K, Franz JR, *et al.*: **Effect of a Supervised Hip Flexor Stretching Program on Gait in Frail Elderly Patients.** *PM R.* 2011; **3**(4): 330–335.
[PubMed Abstract](#) | [Publisher Full Text](#)
 48. Hotta K, Batchelor WB, Graven J, *et al.*: **Daily Passive Muscle Stretching Improves Flow-Mediated Dilation of Popliteal Artery and 6-minute Walk Test in Elderly Patients with Stable Symptomatic Peripheral Artery Disease.** *Cardiovasc Revasc Med.* 2019; **20**(8): 642–648.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 49. Kim TH, Yoon JS, Lee JH: **The Effect of Ankle Joint Muscle Strengthening Training and Static Muscle Stretching Training on Stroke Patients' C.O.P Sway Amplitude.** *J Phys Ther Sci.* 2013; **25**(12): 1613–1616.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 50. Johanson MA, Wooden M, Catlin PA, *et al.*: **Effects of gastrocnemius stretching on ankle dorsiflexion and time-to heel-off during the stance phase of gait.** *Phys Ther Sport.* 2006; **7**(2): 93–100.
[Publisher Full Text](#)
 51. Johanson MA, Cuda BJ, Koontz JE, *et al.*: **Effect of stretching on ankle and knee angles and gastrocnemius activity during the stance phase of gait.** *J Sport Rehabil.* 2009; **18**(4): 521–534.
[PubMed Abstract](#) | [Publisher Full Text](#)
 52. Godges JJ, MacRae PG, Engelke KA: **Effects of Exercise on Hip Range of Motion, Trunk Muscle Performance, and Gait Economy.** *Phys Ther.* 1993; **73**(7): 468–477.
[PubMed Abstract](#) | [Publisher Full Text](#)
 53. Kerrigan DC, Lee LW, Collins JJ, *et al.*: **Reduced hip extension during walking: healthy elderly and fallers versus young adults.** *Arch Phys Med Rehabil.* 2001; **82**(1): 26–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
 54. Kerrigan DC, Todd MK, Della Croce U, *et al.*: **Biomechanical gait alterations independent of speed in the healthy elderly: evidence for specific limiting impairments.** *Arch Phys Med Rehabil.* 1998; **79**(3): 317–322.
[PubMed Abstract](#) | [Publisher Full Text](#)
 55. Gajdosik RL, Linden VWD, Williams AK: **Influence of Age on Length and Passive Elastic Stiffness Characteristics of the Calf Muscle-Tendon Unit of Women.** *Phys Ther.* 1999; **79**(9): 827–838.
[PubMed Abstract](#) | [Publisher Full Text](#)
 56. Gajdosik RL, Vander Linden DW, McNair PJ, *et al.*: **Slow passive stretch and release characteristics of the calf muscles of older women with limited dorsiflexion range of motion.** *Clin Biomech (Bristol, Avon).* 2004; **19**(4): 398–406.
[PubMed Abstract](#) | [Publisher Full Text](#)
 57. Mecagni C, Smith JP, Roberts KE, *et al.*: **Balance and ankle range of motion in community-dwelling women aged 64 to 87 years: a correlational study.** *Phys Ther.* 2000; **80**(10): 1004–1011.
[PubMed Abstract](#) | [Publisher Full Text](#)
 58. Gehlsen GM, Whaley MH: **Falls in the elderly: Part II Balance, strength, and flexibility.** *Arch Phys Med Rehabil.* 1990; **71**(10): 739–741.
[PubMed Abstract](#)
 59. Gao F, Ren Y, Roth EJ, *et al.*: **Effects of repeated ankle stretching on calf muscle-tendon and ankle biomechanical properties in stroke survivors.** *Clin Biomech (Bristol, Avon).* 2011; **26**(5): 516–522.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 60. Yeh CY, Tsai KH, Chen JJ: **Effects of prolonged muscle stretching with constant torque or constant angle on hypertonic calf muscles.** *Arch Phys Med Rehabil.* 2005; **86**(2): 235–241.
[PubMed Abstract](#) | [Publisher Full Text](#)
 61. Yeh CY, Chen JJ, Tsai KH: **Quantitative analysis of ankle hypertonía after prolonged stretch in subjects with stroke.** *J Neurosci Methods.* 2004; **137**(2): 305–314.
[PubMed Abstract](#) | [Publisher Full Text](#)
 62. Bressel E, McNair PJ: **The effect of prolonged static and cyclic stretching on ankle joint stiffness, torque relaxation, and gait in people with stroke.** *Phys Ther.* 2002; **82**(9): 880–887.
[PubMed Abstract](#) | [Publisher Full Text](#)
 63. Honeine JL, Schieppati M, Gagey O, *et al.*: **By counteracting gravity, triceps surae sets both kinematics and kinetics of gait.** *Physiol Rep.* 2014; **2**(2): e00229.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 64. Honeine JL, Schieppati M, Gagey O, *et al.*: **The Functional Role of the Triceps Surae Muscle during Human Locomotion.** *PLoS One.* 2013; **8**(1): e52943.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 65. Neptune RR, Kautz SA, Zajac FE: **Contributions of the individual ankle plantar flexors to support, forward progression and swing initiation during walking.** *J Biomech.* 2001; **34**(11): 1387–1398.
[PubMed Abstract](#) | [Publisher Full Text](#)
 66. Carda S, Invernizzi M, Baricich A, *et al.*: **Casting, taping or stretching after botulinum toxin type A for spastic equinus foot: a single-blind randomized trial on adult stroke patients.** *Clin Rehabil.* 2011; **25**(12): 1119–1127.
[PubMed Abstract](#) | [Publisher Full Text](#)
 67. Ghasemi E, Khademi-Kalantari K, Khalkhali-Zavieh M, *et al.*: **The effect of functional stretching exercises on functional outcomes in spastic stroke patients: A randomized controlled clinical trial.** *J Bodyw Mov Ther.* 2018; **22**(4): 1004–1012.
[PubMed Abstract](#) | [Publisher Full Text](#)
 68. Moore SA, Jakovljevic DG, Ford GA, *et al.*: **Exercise Induces Peripheral Muscle But Not Cardiac Adaptations After Stroke: A Randomized Controlled Pilot Trial.** *Arch Phys Med Rehabil.* 2016; **97**(4): 596–603.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 69. Forrester LW, Roy A, Krywonis A, *et al.*: **Modular Ankle Robotics Training in Early Sub-Acute Stroke: A Randomized Controlled Pilot Study.** *Neurorehabil Neural Repair.* 2014; **28**(7): 678–687.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 70. Park D, Lee JH, Kang TW, *et al.*: **Four-week training involving ankle mobilization with movement versus static muscle stretching in patients with chronic stroke: a randomized controlled trial.** *Top Stroke Rehabil.* 2019; **26**(2): 81–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 71. Raza C, Anjum R, Shakeel NA: **Parkinson's disease: Mechanisms, translational models and management strategies.** *Life Sci.* 2019; **226**: 77–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
 72. Haff G, Triplett NT, National Strength & Conditioning Association (U.S.), eds.: **Essentials of Strength Training and Conditioning.** Fourth edition. Human Kinetics; 2016.
[Reference Source](#)
 73. Delafontaine A: **Effects of stretching exercises on human gait: A systematic review and meta-analysis.** Harvard Dataverse, V1. 2020.
<http://www.doi.org/10.7910/DVN/N8ZXNB>

Open Peer Review

Current Peer Review Status: ? ?

Version 1

Reviewer Report 09 October 2020

<https://doi.org/10.5256/f1000research.28219.r69875>

© 2020 Wei X et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Xu Wei

Department of Orthopaedic Surgery, Tongren Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China

Guoning Zhang

Department of Orthopaedic Surgery, Tongren Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China

1. The purpose of this article is to analyse the effects of a stretching program on gait in each patient category by means of a systematic literature review and meta-analysis, comparing the gait outcomes of the intervention groups with the control groups. This is a very interesting research direction and it has good innovation. But different ages and diseases have different gait results and different stretching training modes and intervention methods. How to eliminate the bias caused by these differences in meta-analysis and comparison?
2. Four meta-analysis were conducted for the following outcomes (Figure 2): gait speed, stride length, hip extension during gait and anterior pelvic tilt. Why choose these four dimensions? Is there any theoretical basis?
3. Kinetic variables: The study of Kerrigan et al. (2003)⁴⁴ showed no significant difference between groups for hip torque (SMD= 0.35; 95% CI: -0.06, 0.75) and ankle plantar flexion power (SMD=0.00; 95% CI: -0.40, 0.40), with a moderate level of confidence (PEDro score: 6/10).: It seems that gait analysis cannot directly measure muscle power. sEMG can only evaluate muscle recruitment signals to evaluate muscle fiber contraction. Therefore, how is ankle flexion power measured by gait analysis?
4. Twelve studies were included in the analysis. Stretching improved gait performance as assessed by walking speed and stride length only in a study with a frail elderly population, with small effect sizes. There is no strong evidence supporting the beneficial effect of using stretching to improve gait. I think that since stretching can improve gait parameters for adults with special weakness, the effect of stretching on gait is significant, although it has

little effect on healthy adults or young adults. Should it be explained? In general, this study made a meta-analysis on the effect of stretching on gait, which has good innovation. However, whether gait analysis has guiding significance for different groups of rehabilitation training is still controversial especially for healthy adults, so the research conclusion of this paper has practical significance in clinical guidance.

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Yes

Are sufficient details of the methods and analysis provided to allow replication by others?

Yes

Is the statistical analysis and its interpretation appropriate?

Partly

Are the conclusions drawn adequately supported by the results presented in the review?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Sports medicine; osteoporosis; osteosarcoma; Spine surgery; Scoliosis

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however we have significant reservations, as outlined above.

Author Response 20 Oct 2020

ARNAUD DELAFONTAINE, Univ. Paris-Sud., Université Paris-Saclay, Orsay, France

Dear Reviewer

We greatly thank you for scrutinizing our manuscript and for his relevant comments. We feel that the manuscript has improved.

Comment 1. Different ages and diseases have different gait results and different stretching training modes and intervention methods. How to eliminate the bias caused by these differences in meta-analysis and comparison?

Reply: We add some precisions in the discussion section (healthy older adults paragraph) to specify how we have limited the risk of bias:

Please see L353-366: When data were meta-analyzed, we ensured that the groups and the training characteristics were similar to limit the risk of bias. This explains that a limited number of studies was included in the meta-analysis. It is worth noticing that the stretching technic was the same (i.e. static stretching), but that details of interventions varied across these studies. For example, both studies selected for the meta-analysis of gait speed included hip flexors and plantar flexors stretching, but, one study included hip extensor stretching⁵² whereas the other did not⁵¹. This difference may partially explain the

heterogeneous results in the meta-analysis ($I^2=86\%$, $p=0.007$). In the same way, the heterogeneous results observed in the meta-analysis of anterior pelvic tilt ($I^2=87\%$, $p<0.01$) may be explained by the stretching of additional muscle groups (hip extensors and plantar flexors) in the study of Cristopoliski et al. (2009) compared to the other studies (in which only hip flexors were stretched)^{53,55}. Nevertheless, heterogeneity in the results was not systematically observed between studies that used slightly different protocols, as showed by the consistent results in the meta-analyses of stride length ($I^2=59\%$, $p=0.12$) and hip extension ($I^2=0\%$, $p=0.99$). Thus, we assume that we have limited the risk of bias in the meta-analyses.

Comment 2. Four meta-analyses were conducted for the following outcomes (Figure 2): gait speed, stride length, hip extension during gait and anterior pelvic tilt. Why choose these four dimensions? Is there any theoretical basis?

Reply: Only these four outcomes fitted with the inclusion criteria for meta-analysis. Meta-analyses were performed only when more than one trial was identified for each outcome. Additionally, to reduce the risk of bias, we ensured that the groups were similar. For example, in the study of Watt et al. (2011), the intervention group had a significantly higher gait speed than the control group, so the trial was excluded of the meta-analysis.

Comment 3. Kinetic variables: The study of Kerrigan et al. (2003)⁴⁴ showed no significant difference between groups for hip torque (SMD= 0.35; 95% CI: -0.06, 0.75) and ankle plantar flexion power (SMD=0.00; 95% CI: -0.40, 0.40), with a moderate level of confidence (PEDro score: 6/10).: It seems that gait analysis cannot directly measure muscle power. sEMG can only evaluate muscle recruitment signals to evaluate muscle fiber contraction. Therefore, how is ankle flexion power measured by gait analysis?

Reply: In the study of Kerrigan et al. (2003), "joint torque and power calculations were based on the mass and inertial characteristics of each lower-extremity segment, the derived linear and angular velocities and accelerations of each lower-extremity segment, and the ground reaction force and joint center position estimates. Joint torques and powers were normalized for body weight and height and were reported as external in newton meters per kilogram meters (N.m/kg.m) and watts per kilogram meters (W/kg.m), respectively". Unless special recommendation of the reviewer or the Editor, we feel it is not necessary to add how the ankle flexion power was measured in the study of Kerrigan et al. (2003).

Comment 4. I think that since stretching can improve gait parameters for adults with special weakness, the effect of stretching on gait is significant, although it has little effect on healthy adults or young adults. Should it be explained?

Reply: We add some precisions in the discussion section (young adults paragraph) to specify why stretching is less interesting to improve gait parameters in young healthy adults:

Please see L387-390: In healthy adults, the interest of practicing stretching to improve gait seems limited as they are assumed to have sufficient mobility for walking. Moreover, the included study involved athletic males⁶¹, a population that is known to be more flexible

than inactive persons⁸¹. Stretching should be more indicated when range of motion is limited²⁴.

Best regards
Arnaud Delafontaine
on behalf of all the authors

Competing Interests: None

Reviewer Report 14 September 2020

<https://doi.org/10.5256/f1000research.28219.r69874>

© 2020 Mégrot F. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Fabrice Mégrot

¹ Unité Clinique d'Analyse de la Marche et du Mouvement, Centre de Médecine Physique et de Réadaptation pour Enfants Bois-Larris, Croix-rouge française, Lamorlaye, France

² BioMécanique et BioIngénierie (BMBI) - UMR 7338, Université de Technologie de Compiègne, Sorbonnes Université, Compiègne, France

Authors propose a systematic review and meta-analysis to investigate the effects of stretching programs on gait and to determine how stretching might be valuable for rehabilitation. From 150 studies identified through systematic searches and critical appraisal of the literature, 12 articles were considered by the authors.

The various demographics of the participants in the reported studies implied a mixed range of age and aetiology, even if old persons, either with an asymptomatic or a pathological health status, were the most represented individuals.

In young adults, two studies by Johanson are reported. Is it not clear if both groups of young adults with limited range of motion were identical in both studies and had the same characteristics? If yes, merging the two studies should make the deal, if not please provide specification about the second group.

Overall, the methodology is correct, and the meta-analysis well described. The results show weak effects of stretching on gait parameters.

The main issue arises from the various parameters which are considered in all the studies, both during gait initiation and straight walking (spatiotemporal parameters, kinematics, joint strength and dynamics, muscle activity, etc...). Therefore, in the results and the discussion, grouping the types of parameters according to their role in the process of walking would add value to this manuscript. The effects, demonstrated or not, have not all the same meaning depending on

whether functional parameters or kinematics angles or even muscle strength and activity are considered.

Gait should be defined in a better way, especially how it is related to the different measures in terms of processes, even if there is anyway no real benefit demonstrated by stretching used as a unique therapy. Surprisingly, no article in children with cerebral palsy has been uncovered, while the clinical care of these children is mostly based on muscle stretching.

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Yes

Are sufficient details of the methods and analysis provided to allow replication by others?

Yes

Is the statistical analysis and its interpretation appropriate?

Yes

Are the conclusions drawn adequately supported by the results presented in the review?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: motor control, biomechanics, neurosciences, nonlinear dynamics, clinical gait/movement analysis and gross motor function of children with cerebral palsy

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 20 Oct 2020

ARNAUD DELAFONTAINE, Univ. Paris-Sud., Université Paris-Saclay, Orsay, France

Reviewer 1 :

Dear Reviewer

We greatly thank you for scrutinizing our manuscript and for his relevant comments. We feel that the manuscript has improved.

Comment 1. In young adults, two studies by Johanson are reported. Is it not clear if both groups of young adults with limited range of motion were identical in both studies and had the same characteristics? If yes, merging the two studies should make the deal, if not please provide specification about the second group. The participants in both studies were different. In the study of Johanson et al. (2006), the participants were healthy adults with limited passive ankle-dorsiflexion range of motion (less than 8 degrees) **and a history of lower limb overuse injury**. In the study Johanson et al. (2009), the characteristics of the stretching program were the same as in Joahnson et al. (2006), but the participants did not

have any history of lower limb overuse injury.

Reply: To make this difference clearer, precisions were added in the “included studies” and “results” sections:

Please see L158-163: Thus, 12 articles were ultimately included in this systematic review. Six studies evaluated the effects of stretching in healthy older adults^{23,51–55}, one in a frail elderly population⁵⁶, one study in an elderly population with stable symptomatic peripheral artery disease⁵⁷, one in stroke patients⁵⁸, one study in adults with limited ankle range of motion (less than 8 degrees) associated with a history of lower limb overuse injury⁵⁹, one study in healthy adults with limited ankle dorsiflexion range of motion (less than 5 degrees)⁶⁰ and one in healthy young adults⁶¹

Please see L285-288: “The study of Johanson et al. (2006) examined the effects of stretching on healthy adults with limited passive ankle-dorsiflexion range of motion (less than 8 degrees) and a history of lower limb overuse injury⁵⁰. Regarding the characteristics of the subjects, 19 subjects were included and the mean age was 30.3±9.8 years.”

Please see L302-306: “The study of Johanson et al. (2009) examined the effects of stretching on young adults with limited passive ankle-dorsiflexion range of motion (less than 5 degrees)⁵¹. It is worth noticed that these participants were not the same than in the study of Johanson et al. (2006). In contrast, the characteristics of the training programs were the same as in Johanson et al. (2006)”.

Comment 2. In the results and the discussion, grouping the types of parameters according to their role in the process of walking would add value to this manuscript. The effects, demonstrated or not, have not all the same meaning depending on whether functional parameters or kinematics angles or even muscle strength and activity are considered.

Reply: In the results and the discussion, we chose to group the participants by patient categories because differences in ages and health parameters may result in different training effects. However, in each patient category, each variable was considered separately in an organized way.

Comment 3. Gait should be defined in a better way, especially how it is related to the different measures in terms of processes, even if there is anyway no real benefit demonstrated by stretching used as a unique therapy.

Reply: We agree. We add a whole paragraph at the beginning of the introduction section to specify how gait can be related to the different variables seen in the review:

Please see L35-44: “Gait is the medical term used to describe the human whole body movement of walking¹. Gait involves internal and external forces that act on the body to move the center of mass (COM) across a given distance². It depends on many biomechanical features that can be observed during gait analysis such as center of mass shifts, joint range of motion, forces, muscle activity, joint moments, and joint powers³. Spatiotemporal features (e.g. velocity, step length, stride length, step with, step variability) and kinematics parameters (range of motion) can be observed subjectively with functional

evaluations by clinicians(e.g. the Tinetti test⁴ or the timed up and go test⁵), but, it can be further objectified with biomechanical analysis in a laboratory². Kinetics variables (the forces that cause the body to move) must be collected in a laboratory environment with force plates (e.g.⁶⁻⁹ for recent studies that used this technic”).

Comment 4. Surprisingly, no article in children with cerebral palsy has been uncovered, while the clinical care of these children is mostly based on muscle stretching.

Reply: We agree. However, no article in children with cerebral palsy fitted our inclusion criteria during the systematic search of the literature. We add a whole paragraph in a specific section “limitations of the study”:

Please see L397-406: Some patient categories were not included in the present review, although muscle stretching is commonly indicated in their clinical care to reduce spasticity⁸². This is for example the case for children with cerebral palsy⁸³. In fact, we were able to identify studies in the literature focusing on the effects of stretching on gait in this population during the first phase of the present review, but the protocol of these studies combined stretching with another form of training (e.g.^{84,85}) or there was no control group (e.g.⁸⁶). These studies therefore did not fit with the inclusion criteria of the present systematic review and were consequently excluded. Now, it should be stressed that the effectiveness of static stretching to improve motor function in children with cerebral palsy is still controversial⁸⁷, although some authors showed that functional stretching exercises may be effective to improve gait⁸⁶. Further randomized controlled trials are needed to explore the impact of stretching on gait in this population.

Best regards
Arnaud Delafontaine
on behalf of all the authors

Competing Interests: None

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research